



## Biodistribution in Sarcoma 180-Bearing Mice of N-Succinyl-Chitosan Nanoparticles for Tumor Targeting

Chengyun YAN <sup>1\*</sup>, Jiwei GU <sup>1</sup>, Yan CHENG <sup>1</sup>, Xi WANG <sup>1</sup>, Zhan HUANG <sup>1</sup> & Dawei CHEN <sup>2</sup>

<sup>1</sup> College of Pharmacy, Jiamusi of University, No. 48 Xuefu street, Jiamusi 154007, P. R. China.

<sup>2</sup> Department of Pharmaceutics, Shenyang Pharmaceutical University, Shenyang 110016, P. R. China.

**SUMMARY.** In the present study, we sought to systemically evaluate the biodistribution and the tumor-accumulation of N-succinyl-chitosan nanoparticles (Suc-Chi-NPs, 200 nm in diameter) in model tumor-bearing mice and also the binding of Suc-Chi-NPs to k562 cells was evaluated by flow cytometry. *In vitro* studies showed that all Suc-Chi-NPs displayed high affinity to k562 cells. After intravenous injection of Suc-Chi-NPs via the tail vein, a small amount of Suc-Chi-NPs was found in liver and spleen for 4 days, whereas a negligible quantity was detected in heart and lung. The distributed amount of Suc-Chi-NPs in blood was as a high level throughout the 4 days. The distributed amount of Suc-Chi-NPs (> 15 % of dose) was accumulated at 1 day and gradually increased in tumor, as blood circulation time increased. This result suggests that Suc-Chi-NPs accumulate passively in the tumor tissue due to the enhanced permeability and retention (EPR) effect, following intravenous administration. These findings revealed the promising potential of Suc-Chi-NPs on the basis of Suc-Chi as a carrier for cancer therapy.

**KEY-WORDS:** Biodistribution, Nanoparticles, N-succinyl-chitosan, Tumor targeting.

\* Author to whom correspondence should be addressed. *E-mail:* yczyzy@yahoo.com.cn