



Short communication Received: January 3, 2010 Revised version: May 13, 2010 Accepted: May 14, 2010

In Vitro Study on Interaction of Ketotifen Fumarate with Metformin Hydrochloride

Mohammed Aktar SAYEED 1, S.M. Raquibul HASAN 2* & Md. Sohel RANA 3

- ¹ Department of Pharmacy, International Islamic University Chittagong, Bangladesh
- ² Department of Pharmacy, BRAC University, 66 Mohakhali, Dhaka-1212, Bangladesh
- ³ Department of Pharmacy, Jahangirnagar University, Savar, Dhaka-1342, Bangladesh

SUMMARY. The present study was designed to study the in vitro interaction of ketotifen fumarate and metformin hydrochloride in aqueous media at different pH values using Job's continuous-variation analysis and Ardon's spectrophotomeric measurement. Job's analysis indicated the formation of 1:1 complex between ketotifen fumarate and metformin hydrochloride at the studied pH values. On the other hand, the values of stability constants (between negative values and 1), as calculated from Ardon's plot, mean that the formation of complex due to interaction between the drugs is readily dissociable. Thus the findings of the study suggest that both drugs can be safely co-administered. However, further studies involving animal models are needed to ascertain the exact nature of interaction between them in vivo.

KEY WORDS: Ardon's mehod, Job's method, ketotifen, metformin, stability constant. Author to whom correspondence should be addressed. E-mail: smraquib_1114@yahoo.com

189 ISSN 0326-2383