



## Evaluation of the Pharmacotherapeutic Profile of HIV/AIDS Bearers

Matheus M. de O. MONTEIRO <sup>1</sup>, Jorge E.G. SANTANA <sup>1</sup>,  
Carlos H.T.B. VICENTE <sup>1</sup>, Valter BESSA FILHO <sup>1</sup>, Everton R. BEZERRA <sup>1</sup>,  
Aluísio M. FERREIRA <sup>2</sup>, Daniele I. JANEIRO <sup>\*2</sup> & Maria do S.R. QUEIROZ <sup>1</sup>

<sup>1</sup> *Departamento de Farmácia, Universidade Estadual da Paraíba,  
CEP 58429-600, Campina Grande, PB, Brasil.*

<sup>2</sup> *Faculdade Maurício de Nassau, CEP 58410-050, Campina Grande, PB, Brasil.*

---

**SUMMARY.** In this work was evaluated the pharmacotherapeutic profile of HIV/AIDS bearers registered in the Service of Specialized Attendance located in Campina Grande–PB, Brazil. The research was of the type traverse, documental, descriptive and analytical and was realized in the period of August to October 2010. Were appraised the patient records of 188 people being 66 % males and 34% females. The most part (36 %) just studied the education fundamental level and presented age group between 40-49 years old. Antiretrovirals more prescribed were nucleoside reverse transcriptase inhibitors (NRTIs) and non-nucleoside reverse-transcriptase inhibitors (NNRTIs). Schemes no recommended by the Ministry of Health also were evidenced. The Negative Results associated to the use of medicine was registered in the three supra-categories. It is necessary a larger integration of the multidisciplinary team of prescribers (physicians and dentists) and pharmacists in the evaluation of the pharmacotherapy to guarantee the reduction of the morbidity and the presence of opportunists infections, guaranteeing so a better surviving for the seropositive patients.

---

**KEY WORDS:** Acquired Immune Deficiency Syndrome (AIDS), antiviral, Human Immunodeficiency Virus (HIV).

\*Author to whom correspondence should be addressed. *E-mail:* dijanebro@yahoo.com.br