



Journal of the Science of
Food and Agriculture

**Impact assessment of mechanical harvest on fruit
physiology and consequences on oil physicochemical and
sensorial quality from 'Manzanilla de Sevilla' and
'Manzanilla Cacereña' super-high density hedgerows. A
preliminary study**

Journal:	<i>Journal of the Science of Food and Agriculture</i>
Manuscript ID:	JSFA-14-1781.R1
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	15-Oct-2014
Complete List of Authors:	Morales-Sillero, Ana; Universidad de Sevilla, Departamento de Ciencias Agroforestales Garcia, Jose; Instituto de la Grasa, Fisiología y Tecnología de Productos Vegetales
Key Words:	Olea europaea L, SHD orchards, grape straddle harvester, ethylene production, oxidative stability, phenols

SCHOLARONE™
Manuscripts

1
2
3 **Impact assessment of mechanical harvest on fruit physiology and consequences on**
4 **oil physicochemical and sensorial quality from ‘Manzanilla de Sevilla’ and**
5 **‘Manzanilla Cacereña’ super-high density hedgerows. A preliminary study**
6
7
8
9

10
11 **Running title: Impact of grape harvester on fruit physiology and oil quality from**
12 **SHD olive hedgerows**
13
14

15
16
17
18 Ana Morales-Sillero,^a José M^a García^{b*}
19

20 * Correspondence to: José M^a García ^b *Dpto. Fisiología y Tecnología de Productos*
21 *Vegetales. Instituto de la Grasa (CSIC). Avda. Padre García Tejero 4, 41012 Sevilla,*
22 *Spain. E-mail: jmgarcia@cica.es*
23
24
25
26
27

28
29 ^a *Dpto. Ciencias Agroforestales, ETSIA, Universidad de Sevilla, Carretera de Utrera,*
30 *km 1, 41013 Sevilla, Spain.*
31

32
33 ^b *Dpto. Fisiología y Tecnología de Productos Vegetales. Instituto de la Grasa (CSIC).*
34 *Avda. Padre García Tejero 4, 41012 Sevilla, Spain.*
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

BACKGROUND: Super-intensive cultivation facilitates olive mechanized harvesting, determining a substantial savings in the production cost of virgin olive oil (VOO). However, the number of varieties adapted to this type of cultivation is reduced. This paper explores the impact that harvest with grape straddle harvester of 'Manzanilla de Sevilla' and 'Manzanilla Cacereña' olives, grown in a super-intensive cultivation, has on the physiology of the fruit and the quality of the oil, subsequently extracted.

RESULTS: In both cultivars, the fruits mechanically harvested presented higher respiration and ethylene production and a lower firmness than the ones harvested by hand. Their oils exhibited lower phenol contents, oxidative stability and lower presence of positive sensory attributes. However, in these oils the values of parameters used to assess the level of quality of the VOO were kept within the limits required for the best commercial category.

CONCLUSION: Mechanical harvesting of 'Manzanilla de Sevilla' and 'Manzanilla Cacereña' super-high density hedgerows induced physiological alterations in the fruits and a reduction in the contents of natural antioxidants and flavour components in the oils, although it did not determine a loss of the "Extra" level of quality

Keywords: *Olea europaea* L., SHD orchards, grape straddle harvester, ethylene production, oxidative stability, phenols

INTRODUCTION

Super-high density (SHD) hedgerows, with ca. 1500-2000 trees ha⁻¹, are nowadays the most common designs in new olive (*Olea europaea* L.) orchards for oil production, in particular in non-traditional olive producing countries. Main advantages are early yield after planting and minimization of the costs of harvest by using of adapted grape straddle harvesters which allow a dramatic decrease in the both labour costs and time employed¹. Furthermore, if the harvest is well synchronized with the extraction process, mechanical harvesting allows producers to minimize the time between harvesting and pressing of olives, which has a very positive impact on oil quality.²

Cultivars must be of early-bearing, consistent annual yields and also of low-vigour to allow the harvesters to pass over the hedgerows and to ensure the illumination of the canopies in the long-term.³ These are the reasons by which most olive SHD orchards are cultivated with the cultivar 'Arbequina', and in less extent with 'Arbosana' and 'Koroneiki' cv. New cultivars obtained in breeding programs have also recently been introduced to the market for SHD, such as 'Sikitita', 'Askal', 'Fs-17', 'Oliana', and 'Tosca 07'.³⁻⁵ Although 'Arbequina' olive oil is otherwise highly demanded by international market due to its sensorial quality, it must be extracted at an early maturity stage of fruits and during a short period of time because of the loss of intensity in its positive attributes (fruity odour, bitter taste and green-yellow colour) and of the drastic decrease in phenolic compounds and pigment composition (chlorophylls and carotenoids).⁶ This decrease, together with that of the reduced monounsaturated/polyunsaturated fatty acids ratio, affects negatively to stability to oxidation of the 'Arbequina' oil, which in fact it is low compared with other cultivars⁷⁻⁸

1
2
3 The adapted grape straddle harvesters use rods to shake the canopy. They contact
4
5 directly with the fruit so they can cause damage which seems to increase as firmness
6
7 decreases during maturation.⁹ However, studies about the impact of these machines on
8
9 the oil quality from 'Arbequina' hedgerows orchards are really scarce. It's known that,
10
11 although the harvester may favour the physical oil extraction, it may decrease the oil
12
13 stability to oxidation in relation to the oil from fruits manually harvested. This decrease
14
15 is caused by the reduction of both the tocopherol and phenol contents. Moreover, higher
16
17 values of peroxides and also of K_{232} are also found in the oils from mechanical
18
19 harvested fruits, which indicates that mechanical harvesting by the grape straddle
20
21 harvester led to an increase of the primary steps of the oxidation process.¹⁰ These
22
23 findings are in part similar to those reported in a 'Souri' traditional orchard (with trees
24
25 at 10 x 10 m) and mechanically harvested by hand-vibrating combs fruits,¹¹ in
26
27 particular, a slight increase in peroxide level and a decrease of the total phenol content
28
29 was also found in the oils from mechanically harvested trees. An increase of the free
30
31 acidity content was also reported in these oils.
32
33
34
35

36 Therefore, regardless the type of harvester employed in an olive oil orchard, internal
37
38 fruit injures seems to cause the loss of oil quality from mechanical harvested fruits. The
39
40 selection of the harvesting method must look for the best possible conservation of the
41
42 fruit integrity, in order to obtain a VOO of high quality. If a particular method causes
43
44 bruises on the fruit surface as a result of its mechanical impact or compression, olive
45
46 respiration and the susceptibility to decay at a faster rate, in comparison to undamaged
47
48 fruit, seems to increase.¹² The oil extracted from these damaged olives can be high in
49
50 acidity, low in stability and poor in polyphenols and might develop off-flavors due to
51
52 the enzymatic activities favored by the breakdown of the cells and the contact between
53
54 enzymes and substrates that were initially compartmented differently.¹³ In the light of
55
56
57
58
59
60

1
2
3 these considerations, hand picking appears to be the best method for preventing fruit
4
5 damage,¹⁴ yet, unfortunately, olive manual harvest is quite expensive, so it is
6
7 fundamental to assess the impact of mechanical harvest on fruit physiology and
8
9 consequences on oil physicochemical and sensorial quality applied to the different olive
10
11 varieties.
12

13
14 The scarcity of suitable olive cultivars is, therefore, an important disadvantage of
15
16 SHD orchards. Olive breeding is necessary to produce new cultivars adapted to these
17
18 orchards with hedgerow management. Furthermore, it is necessary the study of the
19
20 suitability of traditional olive cultivars not only to be grown in SHD orchards but also to
21
22 produce oil with better quality than the current ones. In this sense, Morales-Sillero *et*
23
24 *al.*¹⁵ have recently showed that ‘Manzanilla de Sevilla and ‘Manzanilla Cacereña’ trees
25
26 are high-yield and may be mechanically harvested by a grape straddle harvester when
27
28 grown under SHD hedgerows. The work also highlights the efficiency of mechanical
29
30 harvesting to remove green fruits from both cultivars (98%) and the short time required
31
32 for harvesting (< 1.7 h). These cultivars are usually cultivated in intensive orchards not
33
34 only in Spain but also in USA, Israel, Argentina (‘Manzanilla de Sevilla’) and Portugal
35
36 (‘Manzanilla Cacereña’), and the productions are mainly destined for table olive. The
37
38 oil contents are not usually high, but they are quite appreciated by their stability against
39
40 oxidation, their green colour and their elevated content on the more characteristic
41
42 sensory attributes of the VOO, such as fruity, bitter or pungent.¹⁶⁻¹⁷ In fact, a substantial
43
44 part of the production (10 – 15 %), mainly corresponding to the fruit with inadequate
45
46 appearance for table market, routinely goes to olive oil extraction industries, where the
47
48 oil is habitually used for blending with other VOOs to improve its sensory quality. In a
49
50 continuation of this work, the main aim of this study is to evaluate how the grape
51
52 straddle harvester affect the physiology of fruits of both cultivars (‘Manzanilla de
53
54
55
56
57
58
59
60

1
2
3 Sevilla' and 'Manzanilla Cacereña') and the impact on the oil physicochemical
4 characteristics and sensory attributes. Results will also contribute to determine to what
5 extent the impact depends on the cultivar.
6
7
8
9

10 11 12 **EXPERIMENTAL**

13 14 **Orchard description**

15
16 The field experiment was made in 2012, in a commercial orchard of Elvas (Portugal)
17 (latitude 38,56°N; longitude 7°02'O; altitude 201 m) with five-year-old 'Manzanilla de
18 Sevilla' and 'Manzanilla Cacereña' (*Olea europaea* L.) hedgerows at a spacing of 3.75
19 m x 1.35 m (1,975 trees ha⁻¹) in a north-south orientation. The area is characterized by a
20 Mediterranean climate, with an average annual rainfall of 516.4 mm and ET_o of 1296.5
21 mm. The soil is a loam with effective depth of 0.6 m. Four random rows of trees (ca.
22 90) per cultivar ('Manzanilla de Sevilla' and 'Manzanilla Cacereña') and per harvesting
23 treatments (Hand, in which fruit were picked up by hand; and Mechanical, in which
24 trees were harvested with a grape straddle harvester), were selected for the trial.
25
26
27
28
29
30
31
32
33
34
35

36 Trees were irrigated from March to October with 280 mm ha⁻¹ with a lateral per tree
37 row with two 2.43 L h⁻¹ compensating drippers per tree, 0.60 m apart. Fertilization was
38 made according to foliar analyses to non-limiting nutrient conditions. Weeds were
39 controlled by non-residual herbicides in the tree rows and natural ground cover between
40 them. The branches growing towards the centre of the rows were removed in winter.
41
42
43
44
45
46
47 Three months before the experiment, 'Manzanilla de Sevilla' hedgerows were 2.8 m tall
48 and 2.0 m wide and those of 'Manzanilla Cacereña' were 2.4 m and 1.5 m, respectively.
49
50
51
52
53
54
55
56
57
58
59
60
Fruit yields were ca. 10,000 kg ha⁻¹ for 'Manzanilla de Sevilla' cv. and ca. 18,000 kg
ha⁻¹ for 'Manzanilla Cacereña' cv.¹⁵

Fruit harvesting and oil extraction

Harvesting was made in December 10th when fruits had a ripening index **between 3 and 4**.¹⁸ Fruit samples of 5 kg by experimental unit, each one of the four random rows, were handpicked around the canopy of 25 trees randomly selected in the manual treatment. Later, the rest of the fruits of the same replicate were mechanical harvested with a model VX 7090 (Olive, CNH Global, Belgium) and samples of 5 kg by experimental unit were also randomly taken from the total amount. **The fruits were transported the same day to the Instituto de la Grasa in Seville, stored at 5 °C during the night and analysed early in the morning.**

Fruit characteristics: Colour, firmness, respiration rate and ethylene production

Fruit skin colour was determined on the equatorial zone of 50 healthy fruits per experimental unit using a Minolta CR200 (Minolta Camera Co., Osaka, Japan) chromameter with a measuring area of 8 mm in diameter, diffuse illumination and a viewing angle of 0°. The International Commission on Illumination colour notation system (ICI L*a*b*) was applied to determine the parameters L*, a*, and b*; with L* as the lightness, a* the colour axis from green to red, and b* the colour axis from blue to yellow. The following equation (colour index, CI), previously used to monitor colour changes during olive cold storage,¹⁹ was applied:

$$CI = L^* \times (b^* - a^*) \times 10^{-2}$$

Fruit firmness was also evaluated on the equatorial zone of the same fruits, using a Zwick 3300 hand densimeter (Zwick GmbH & Co., Ulm, Germany). The consistency of each fruit was measured without rupture by the pressure in a 5 mm diameter disk.

Results were expressed **in N**.

1
2
3 Fruit samples of 500 g by replicate were placed into 2.0 L glass jar and stored at 2 °C
4
5 for 24 hours. These jars were then hermetically sealed during 3 h. Later, a G100
6
7 portable gas analyser (Geotechnical Instrument Ltd. Leamington Spa, UK) was used to
8
9 measure the CO₂ content of the head space, to estimate the respiration rate of the fruit.
10
11 The ethylene content was simultaneously evaluated, with an ICA portable ethylene
12
13 analyser (International Controlled Atmosphere Ltd., Paddock Wood, UK). Respiration
14
15 rate values were expressed as mL CO₂ L⁻¹ kg⁻¹ h⁻¹ and ethylene production mean values
16
17 were expressed as μL Ethylene L⁻¹ kg⁻¹ h⁻¹.
18
19

20 21 22 23 **Physical and chemical extraction of olive oil: VOO yield, humidity, total oil yield** 24 25 **and physical extractability** 26

27 One kilogram of olives was randomly taken from each experimental unit for oil
28
29 extraction and analysis. The oil was extracted with the Abencor analyzer (Comercial
30
31 Abengoa S.A., Seville, Spain), which simulates the industrial process of VOO yield at
32
33 laboratory scale²⁰. Each sample was milled separately. From the resulting paste in each
34
35 sample, 800 g were collected in a metallic pitcher, homogenized with a spatula and the
36
37 oil was physically extracted. After centrifugation, the oil was decanted into a graduated
38
39 tube to measure the volume, and the VOO yield so determined was expressed as
40
41 percentage of fresh weight, considering oil density at ambient temperature of 0.916 kg
42
43 L⁻¹. The extracted oil was then paper filtered and stored at -20 °C under N₂ atmosphere
44
45 until analysis. Samples of 50 g, taken from the leftover paste obtained for the physical
46
47 extraction of the oil, were dried at 105 °C up to constant weight to evaluate the humidity
48
49 and the dry weight of the olive fruits. The oil of the dried pastes were chemically
50
51 extracted with hexane, using the Soxhlet method,²¹ to evaluate the total oil yield of the
52
53 paste as the percentage referred to the fresh weight. The physical extractability was
54
55
56
57
58
59
60

1
2
3 calculated as the percentage of total oil yield that represents the physically extracted
4
5 VOO yield.
6
7

8 9 **Oil analysis**

10
11 In each oil sample, free acidity, peroxide index value, coefficients of specific extinction
12
13 at 232 and 270 nm (K_{232} and K_{270}) and the overall grading of the sensory quality of the
14
15 oils were evaluated according to the European Union Standard Methods.²² The sensory
16
17 intensities of the negative attributes (rancid, fusty, winey, musty, etc.), of the bitterness
18
19 and of the pungency were graded by a panel of eight trained tasters using a line scale,
20
21 and the results converted into a numerical score by measuring the position of the placed
22
23 mark along a 10 cm line. This allowed calculating averages among taster sensory
24
25 scores. The overall sensory quality of each oil sample was graded by the same panel
26
27 according to a structured scale of nine points, “1” being the value for the poorest quality
28
29 possible and “9” for the best, being “6.5” the lowest limit value for the best level of
30
31 sensory quality (named “Extra”) for VOOs.
32
33

34
35
36 Oxidative stability was measured by using the Rancimat method, which evaluates the
37
38 time (h) of resistance to oxidation of 3 g oil samples exposed to streams of dry air (20
39
40 L h⁻¹) heated to 100 °C.²³
41
42

43
44 The contents of chlorophyll and carotenoid pigments in the oils were evaluated by their
45
46 absorbances at 470 and 670 nm, respectively, and the results were expressed as mg kg⁻¹.
47
48

49
50 Tocopherol composition was determined by HPLC using the IUPAC method (1992).²⁵

51
52 The phenolic fraction was isolated by solid-phase extraction and analyzed by reversed-
53
54 phase high performance liquid chromatography (HPLC) using a diode-array UV
55
56 detector.²⁶ The quantification of phenolic compounds (except flavones and ferulic acid)
57
58
59
60

1
2
3 was carried out at 280 nm using *p*-hydroxyphenylacetic acid as an internal standard,
4
5 while flavones (luteoline and apigenine) and ferulic acid were quantified at 335 nm
6
7 using *o*-coumaric acid as an internal standard. The results were expressed in mg kg⁻¹.
8
9 Fatty acid composition was determined by gas chromatographic analysis of methyl
10
11 esters for each oil sample.²⁷ This was performed on an Agilent 6890 equipped with a
12
13 flame ionization detector, fitted with a fused-silica capillary column (SP-2380, 60
14
15 m/0.25 mm I.D.) coated with cyanopropylsilicone (0.20 mm film thickness). Hydrogen
16
17 was employed as carrier gas at flow of 1 mL min⁻¹. The oven temperature was
18
19 maintained at 185 °C and the injector (split 1:20) and detector at 225 °C. **The results**
20
21 **were expressed in g kg⁻¹.**
22
23
24
25
26

27 **Statistical analyses**

28
29 Analyses were performed using StatGraphics Plus 5.1 (StatGraphics Software, Inc., La
30
31 Jolla, California, USA). Data were subjected to statistical analysis using variance
32
33 analysis to determine the effect of treatments and cultivars. When necessary, to achieve
34
35 normality and homogenize the variance, data were previously transformed using the
36
37 arcsine of the square root or Box-Cox power transformations.²⁸
38
39
40
41
42

43 **RESULTS AND DISCUSSION**

44 **Fruit characteristics: Colour, firmness, respiration rate and ethylene production**

45
46 Chromatic parameters L*, a* and b* of the skin of olive fruits were not significantly
47
48 affected by the method of harvest (Table 1). Regardless the cultivar tested, no
49
50 significant effects on CI values were detected in the fruits as a consequence of the
51
52 different system of harvesting, which agrees with results of Yousfi *et al.*¹⁰ who found no
53
54
55
56
57
58
59
60

1
2
3 effect due to this factor on CI values of 'Arbequina' olives skin from a hedgerow
4
5 orchard.

6
7 Fruit firmness was clearly affected by mechanical harvesting (Table 1). Manually
8
9 harvested fruits exhibited significantly higher values of this parameter whatever the
10
11 cultivar, as it was previously found by Morales-Sillero *et al.*¹⁵ in the same experimental
12
13 orchard with green fruits. In contrast, Yousfi *et al.*¹⁰ did not find this effect on
14
15 'Arbequina' olives. It seems that the cellular tissues of both 'Manzanilla' olives are
16
17 more sensitive than those of 'Arbequina' to the harvester's mechanical action, which
18
19 would induce a deterioration of their cell walls.

20
21
22 Mechanical harvesting also induced in the fruit of both cultivars a significant
23
24 increase of the respiration rate and, specially, of the ethylene production (Table 1).
25
26 Moreover, this increase of the ethylene was greater in 'Manzanilla de Sevilla' fruits than
27
28 in those of 'Manzanilla Cacereña'. Segovia-Bravo *et al.*¹² had already found that
29
30 simulating the mechanical harvesting of 'Manzanilla de Sevilla' fruits, those which are
31
32 bruised show higher respiratory activity than the unbruised ones. This suggests that the
33
34 fruit has undergone further physical damage with mechanical harvesting than with
35
36 manual harvesting. Olive is usually classified as a non-climacteric fruit, therefore it
37
38 shows no drastic increases in their CO₂ and ethylene production rates during ripening. In
39
40 contrast, climacteric fruits, since a determined moment of the ripening process
41
42 (climacteric point) produce increased levels of ethylene, which is accompanied by a
43
44 respiration peak.²⁹ At this time, all the physiological mechanisms associated to fruit
45
46 maturation and senescence such as: colour changes, softening, chlorophyll degradation,
47
48 phenol oxidation and aroma development are accelerated in the fruits.³⁰ However,
49
50 ethylene production may also be a response of a plant to injury.³¹⁻³² Thus, the
51
52 stimulation of ethylene production found in the mechanical harvested fruits of the
53
54
55
56
57
58
59
60

1
2
3 studied cultivars seems to be a response to the internal injury caused to the fruits by the
4
5 grape straddle harvester. Moreover, this increase of ethylene production could have
6
7 favoured that of the respiration rate. A similar emission of ethylene was reported by
8
9 Yousfi *et al.*¹³ in ‘Arbequina’ fruits harvested with a similar machine. Results here
10
11 shown therefore support that mechanical harvesting causes internal damage in olive
12
13 fruits, as Dag *et al.*¹¹ and Yousfi *et al.*^{10,13} previously suggested.
14
15
16
17

18 **VOO yield, total oil yield, humidity and physical extractability**

19
20 VOO yield was significantly higher in ‘Manzanilla de Sevilla’ than in ‘Manzanilla
21
22 Cacereña’, but regardless of the cultivar, no differences in the physical oil extraction
23
24 were found in the mechanically harvested fruits when compared to the manually
25
26 harvested (Table 1). These results do not agree with those of Yousfi *et al.*¹⁰ which found
27
28 a higher virgin oil yield in ‘Arbequina’ mechanically harvested fruits by a grape
29
30 straddle harvester when compared to the handpicked ones.
31
32

33
34 Since no difference between both harvesting systems was found in the total oil yield,
35
36 which is chemically extracted, the physical extractability was also similar. In the same
37
38 way, no effect due to the different system of harvesting was detected in the fruit
39
40 humidity. Nevertheless, ‘Manzanilla Cacereña’ olives showed higher humidity values,
41
42 than ‘Manzanilla Sevillana’ fruits, although only those mechanically harvested reached
43
44 a statistical significant difference. This higher humidity could explain the significant
45
46 lower VOO yield and physical extractability values that presented this variety
47
48 (‘Manzanilla Cacereña’), because the presence of water hinders the VOO physical
49
50 extraction.³² The VOO yields did not reach high values, in agreement with previous
51
52 results related to both cultivars, although contents up to 20% have been also indicated,
53
54 in particular for the ‘Manzanilla de Sevilla’ cv.³⁴⁻³⁶. The low extractability of
55
56
57
58
59
60

1
2
3 'Manzanilla Cacereña', either by an 'Abencor' system or on an industrial scale, has also
4
5 been recently reported, and it may be even lower as fruit ripening increase.³⁵ However,
6
7 the high fruit yields of the 'Manzanilla de Sevilla' and 'Manzanilla Cacereña' trees
8
9 cultivated in the SHD hedgerows compensated at least in part the low VOO yield.
10

11 12 13 14 **Oil quality**

15
16 All the extracted oils were classified as "Extra", the best commercial category of olive
17
18 oil, given the mean values of free acidity, peroxide index, K_{232} , K_{270} and panel test
19
20 (Table 2). However, the mechanical treatment increased significantly the free acidity in
21
22 the 'Manzanilla de Sevilla' oils, from 0.3 to 0.7%, thus close to the limit (0.8%)
23
24 established by the EU-legislated standards for this category. Previously, Dag *et al.*¹¹
25
26 found that free acidity increased when 'Souri' fruits from trees under different irrigation
27
28 conditions (from non-irrigated to irrigated with 125% ET_0) were harvested by hand-
29
30 vibrating combs, compared to those manually harvested. Although this effect was
31
32 observed regardless the irrigation regime of the trees, those fruits from highly irrigated
33
34 trees reached the highest values of free acidity (0.8). In contrast, Yousfi *et al.*¹⁰ did not
35
36 find differences on the free acidity of the oils extracted from 'Arbequina' fruit manually
37
38 and mechanically harvested by a grape straddle harvester. Probably, this induction of
39
40 the free acidity increase is related with the activation of the internal olive lipase, which
41
42 would act during the process of virgin olive oil extraction. A similar increase of free
43
44 acidity values of the virgin oil is observed coinciding with the progress of fruit
45
46 ripening.³⁷ Taking in account that ethylene is the most efficient inductor of the maturity-
47
48 related enzymatic activities, it seems reasonable to think that this phenomenon, the
49
50 increase of free acidity in the virgin oil, is linked with the increase of ethylene
51
52 biosynthesis.
53
54
55
56
57
58
59
60

1
2
3 The parameters used to assess the level of oxidative deterioration in the VOO
4 (peroxide value, K_{232} and K_{270}) were not affected by the harvesting system (Table 2).
5
6
7 These results do not agree with the ones obtained by Dag *et al.*¹¹, using ‘Souri’ olives
8
9 and Yousfi *et al.*¹⁰ with ‘Arbequina’ olives, who observed that oil extracted from
10
11 mechanically harvested olives presented significantly higher peroxide values. Moreover,
12
13 Yousfi *et al.*¹³ also found significantly higher K_{232} values in these oils. Probably, these
14
15 differences were due to that the internal damage suffered by these cultivars determined
16
17 a more closed contact of the oil contained in the fruit with the atmospheric oxygen.
18
19

20
21 The oil extracted from olives mechanically harvested of both cultivars showed
22
23 systematically lower intensity of positive sensory attributes such as olive fruity or bitter,
24
25 exhibiting the oils from ‘Manzanilla Cacereña’ of this treatment, also significantly
26
27 lower intensity of pungent (Table 2). In consequence, the overall grading of the sensory
28
29 quality evaluated by panel test found that oil extracted from manually harvested olives
30
31 presented a significant higher level of sensory quality. In contrast, Yousfi *et al.*¹⁰ found
32
33 no significant differences between the panel test values of the oils extracted from
34
35 ‘Arbequina’ olives manually or mechanically harvested. This suggests that the sensory
36
37 quality of the oils contained in the cells of ‘Manzanilla de Sevilla’ and ‘Manzanilla
38
39 Cacereña’ olives is more sensitive to the internal damage caused by the mechanical
40
41 harvesting than ‘Arbequina’ oils. Enzymatic activities such as lipoxygenase,
42
43 hydroperoxide lyase, alcohol dehydrogenase or alcohol acyltransferase, related with the
44
45 release of the positive sensory attributes of the VOO, and/or other enzymes responsible
46
47 for the phenolic content of the VOO, such as glucosidases and phenoloxidas should
48
49 be implied in this fact. Probably, the higher production of ethylene in the mechanically
50
51 harvested fruits favored the inhibition of these activities. The progress of fruit
52
53 maturation coincides with the reduction of these positive attributes.³⁷
54
55
56
57
58
59
60

1
2
3 Mean values of oxidative stability were ≥ 85 h. (Table 3). They are high if compared
4 to 'Arbequina' oils, the most grown cultivar under SHD conditions worldwide, for
5 which it is usually lower than 40 h.^{5, 10, 38} The stability to oxidation, i.e the resistance to
6 rancidity, decreased significantly in the oils from mechanically harvested fruits both in
7 'Manzanilla de Sevilla' and 'Manzanilla Cacereña' oils. Yousfi *et al.*¹⁰ found that the
8 oxidative stability decreased significantly by 9 h in the 'Arbequina' oils from
9 mechanically harvested fruits as compared to the manually harvested ones. In our study
10 it decreased by 13 h in the 'Manzanilla Cacereña oils', and up to 25 h in 'Manzanilla de
11 Sevilla' ones. These decreases were probably related to the changes in the phenol
12 content in both cultivars (Table 4). Moreover, the tocopherols contents in 'Manzanilla
13 de Sevilla' (Table 3) could have probably affected the oxidative stability, given the
14 antioxidant capacity of these compounds.³⁹ In fact, the tocopherols α , β and total
15 decreased by more than 50% in the 'Manzanilla de Sevilla' oils from the mechanically
16 harvested fruits, while no differences between treatments were found for this parameter
17 in the 'Manzanilla Cacereña' oils, which otherwise had only about one third of the total
18 tocopherols as compared to 'Manzanilla de Sevilla'. No differences in tocopherols
19 contents were found by Yousfi *et al.*¹⁰ in oils from 'Arbequina' olives manually or
20 mechanically harvested, which exhibited a similar amount of total tocopherol content
21 than the oils of 'Manzanilla de Sevilla' from manually harvested olives in our
22 experiment.

23
24
25 Chlorophylls and carotenoids are the main responsible for the colour of VOO,⁴⁰ and
26 carotenoids display an antioxidant action.³⁸ The mechanical harvest did not affect the
27 chlorophylls content in any of both cultivars (Table 3). In contrast, although no
28 differences were found on the carotenoids content in the 'Manzanilla de Sevilla' oils,
29 the 'Manzanilla Cacereña' oils extracted from mechanically harvested fruits showed

1
2
3 higher values of them than the ones from manually harvested olives. Yousfi *et al.*¹⁰
4
5 previously found no effect due to the different harvesting system on these pigment
6
7 contents of 'Arbequina' oils. The pressure and the beating suffered by the fruit during
8
9 mechanized harvesting can facilitate an easier breaking of chloroplasts during the
10
11 subsequent fruit grinding and olive paste kneading of the virgin olive oil extraction,
12
13 favoring the release of a greater amount of chlorophylls and carotenoids. However, the
14
15 greater presence of ethylene, which was synthesized as response to this mechanical
16
17 damage, could stimulate enzymatic activities such as chlorophyllase or lipoxygenase
18
19 that lead to the oxidation of these pigments.⁴¹

20
21
22 The total phenol content differed clearly between cultivars (Table 4). Thus, the
23
24 'Manzanilla de Sevilla' oils from the manual treatment had more than double than those
25
26 of 'Manzanilla Cacereña' (>539 and 259 mg kg⁻¹, respectively). The high antioxidant
27
28 capacity of the 'Manzanilla de Sevilla' in relation to the phenol content oils has been
29
30 previously indicated. Several works show even more than 1000 mg kg⁻¹ of total
31
32 phenols.^{36, 42} The mechanical harvest decreased the mean values of total phenol content,
33
34 total orthodiphenols and, in particular, those of the total secoiridoids, the most abundant
35
36 phenolic compounds in VOO, which are related with the bitter and pungent attributes,
37
38 as well as the stability of VOO against oxidation.⁴³⁻⁴⁴ In both cultivars, differences were
39
40 due to the lower contents of the dialdehydic form of the decarboxymethyl oleuropeine
41
42 aglycone (3,4-DHPEA-EDA), and of the hydroxytyrosyl elenolate (3,4-DHPEA-EA) in
43
44 the oils from the mechanically harvested fruits. Both compounds simultaneously belong
45
46 to the phenolic groups of orthodiphenols and of secoiridoid derivatives. The lignans
47
48 (pinoresinol and acetoxypinoresinol) were also significantly lower in the oils of
49
50 'Manzanilla de Sevilla' olives mechanically harvested, while in the similar
51
52 'Manzanilla Cacereña' oils the pinoresinol, the *p*-HPEA-EDA, and the vanillic acid,
53
54
55
56
57
58
59
60

1
2
3 achieved a significantly lower content. Nevertheless, the contents of hydroxytyrosol and
4
5 tyrosol in ‘Manzanilla de Sevilla’, and those of hydroxytyrosolacetate in ‘Manzanilla
6
7 Cacereña’, were higher in the oils from mechanically harvested fruits. Our results
8
9 confirm those of Dag *et al.*¹¹ and Yousfy *et al.*¹⁰, therefore the idea that mechanically
10
11 harvesting affects negatively the phenol content regardless the cultivar and type of
12
13 harvester. However, when fruits are harvested by grape straddle harvesters, the response
14
15 of phenol composition differs between cultivars. Segovia-Bravo *et al.*^{12, 45} found local
16
17 tissue degradation together with an output of intracellular water and the oxidation of
18
19 phenolic compounds in ‘Manzanilla de Sevilla’ green fruits, **when simulated the**
20
21 **mechanical harvesting.**
22
23
24

25 The main phenolic compounds found in the olive fruit are oleuropein, ligstroside
26
27 and demethyloleuropein. They are glycosides that do not dissolve in the oil. The
28
29 presence of phenolic derivatives in VOO depends mainly on the action of enzyme
30
31 activities, such as glycosidases and estearases that transform these compounds in other
32
33 oil soluble molecules. Thus, the secoiridoid derivatives resulting from the enzymatic
34
35 hydrolysis of these glycosides identified as 3,4-DHPEA-EDA, the dialdehydic form of
36
37 decarboxymethyl ligstroside aglycone (*p*-HPEA-EDA), the 3,4-DHPEA-EA and the
38
39 aldehydic form ligstroside aglycone (*p*-HPEA-EA), are usually the most abundant
40
41 phenolic components found in most VOO.⁴⁶⁻⁴⁷ The differences found in this study on
42
43 phenolic content due to the different system of harvesting only can be explained by a
44
45 partial inhibition of these activities induced by the mechanical harvesting. As the
46
47 phenolic content of the VOO decreases during fruit maturation,³⁷ probably the increase
48
49 of ethylene biosynthesis induced by the grape straddle harvester in the olive cells is also
50
51 related with this fact.
52
53
54
55
56
57
58
59
60

1
2
3 Fatty acid composition was not significantly altered by the mechanization of the
4 harvest with a grape straddle harvester. In consequence, no differences in the
5 monounsaturated/polyunsaturated (MUFA/PUFA) and saturated/unsaturated
6 (SFA/UFA) ratios were found (data not shown). These results confirmed the ones
7 previously obtained by Yousfi *et al.*¹⁰ testing 'Arbequina' olives, who found no effect
8 on fatty acid composition due to the harvesting system. Greater contents of different
9 fatty acids were found in 'Manzanilla de Sevilla' than in 'Manzanilla Cacereña'. In any
10 case, it is noticeable the high contents in oleic acid in the oils from both cultivars when
11 they are cultivated under superintensive conditions. These contents are considerably
12 higher than those normally found in the oils extracted from 'Arbequina' olives (600-700
13 g kg⁻¹).⁴⁸ The elevated presence of oleic acid contributes to the high stability of the oils
14 and is also a dietary, nutritional and therapeutic improvement.⁴⁹

15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30 The current work demonstrated that the use of a grape straddle harvester in
31 'Manzanilla de Sevilla' and 'Manzanilla Cacereña' SHD hedgerow orchards induced
32 relevant changes in physiological parameters of fruits, such as decreases of firmness and
33 clear increases of the respiration rate and ethylene emission. Probably as response to
34 these physiological alterations of the fruit, the VOO extracted suffered a clear
35 deterioration of its quality characteristics. Thus, the stability against oxidation and the
36 overall rating of the sensory quality significantly decreased, likely due to a reduction of
37 the phenol contents. Furthermore, the oils extracted from mechanically harvested
38 'Manzanilla de Sevilla' olives showed a significantly higher titratable acidity and a
39 lower tocopherol content. Despite these disadvantages, the parameters of quality of the
40 oils from mechanical harvesting were kept within the established limits for the best
41 commercial category of quality. In consequence, under a commercial point of view,
42 exclusively considering the quality level of the oil extracted, the mechanical harvesting
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 of these olives grown in hedgerow would be feasible. Furthermore, taking into account
4
5 the sensorial quality and the oxidative stability of these oils, particularly interesting
6
7 from the commercial point of view would be the mixture of these ‘Manzanilla de
8
9 Sevilla’ and ‘Manzanilla Cacereña’ VOOs with those of the ‘Arbequina’, their
10
11 combination would result in a “coupage” with better balanced flavour and stability.
12

13
14 **Bearing in mind that these results are preliminary**, the complementarity of the VOOs of
15
16 these cultivars **seems to** suggest the advisability of including some hedgerows of these
17
18 cultivars in the SHD 'Arbequina' orchards, to improve the sensory quality, nutritional
19
20 value and the stability of these oils through the processing of all these olives together.
21
22

23 24 25 **ACKNOWLEDGMENTS**

26
27 This work was funded by the research project AGL2011-30371-C02-02 of the Spanish
28
29 State Secretary for Research and by the FEDER program. Special thanks to Elaia S.A.
30
31 for providing facilities in this study; Alejandro Trigo, technical assistant of Elaia S.A,
32
33 and María del Carmen Martínez, Cristina Rodríguez and Antonio Montero for technical
34
35 assistance.
36
37
38
39
40
41
42
43
44

45 46 **REFERENCES**

- 47
48 1. **Camposo S and Godini A, Preliminary observations about the performance of 13**
49
50 **varieties according to the super high density oliveculture training system in Apulia**
51
52 **(southern Italy). *Adv Horti Sci* 24:16-20 (2010).**
53
54 2. **Clodoveo ML, Hbaieb RH, Kotti F, Mugnozza GS and Gargouri M, Mechanical**
55
56 **strategies to increase nutritional and sensory quality of virgin olive oil by**
57
58
59
60

- 1
2
3 modulating the endogenous enzyme activities. *Compr Rev Food Sci S* **13**:135-154
4
5 (2014).
6
7
8 3. Connor DJ, Gómez-del-Campo M, Rousseaux MC and Searles PS, Structure,
9 management and productivity of hedgerow olive orchards: A review. *Sci hortic*
10 **169**:71-93 (2014).
11
12
13 4. Allalout A, Krichène D, Methenni K, Taamalli A, Oueslati I, Daoud D and Zarrouk
14 M, Characterization of VOO from super intensive Spanish and Greek varieties
15 grown in northern Tunisia. *Sci Hort* **120**:77-83 (2009).
16
17
18 5. Benito M, Lasa JM, Gracia P, Oria R, Abenoza M, Varona L and Sánchez-Gimeno
19 AC, Olive oil quality and ripening in super-high-density Arbequina orchard. *J Sci*
20 *Food Agric* **93**:2207-2220 (2013).
21
22
23 6. Romero A, Díaz I and Tous J, Optimal harvesting period for 'Arbequina' olive
24 cultivar in Catalonia (Spain). *Acta Hort* **586**:393-396 (2002).
25
26
27 7. Aranda F, Gomez-Alonso S, Rivera del Alamo RM, Salvador MD and Fregapane G.
28 Triglyceride, total and 2-position fatty acid composition of Cornicabra virgin olive
29 oil: Comparison with other Spanish cultivars. *Food Chem* **86**:485-492 (2004).
30
31
32 8. Rondanini DP, Castro DN, Searles PS and Rousseaux MC, Fatty acid profiles of
33 varietal virgin olive oils (*Olea europaea* L.) from mature orchards in warm arid
34 valleys of Northwestern Argentina (La Rioja). *Grasas Aceites* **62**:399-409 (2011).
35
36
37 9. Camposeo S, Alessandro GA and Gatullo CE, Ripening indices and harvesting
38 times of different olive cultivars for continuous harvest. *Sci Hort* **151**:1-10 (2013).
39
40
41 10. Yousfi K, Weiland CM and García JM, Effect of harvesting system and fruit cold
42 storage on virgin olive oil chemical composition and quality of superintensive
43 cultivated 'Arbequina' olives. *J Agric Food Chem* **60**:4743-4750 (2012).
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
11. Dag A, Ben-Gal A, Yermiyahu U, Basheer L, Nir Y, Kerem Z, The effect of irrigation level and harvest mechanization on virgin olive oil quality in a traditional rain-fed ‘Souri’ olive orchard converted to irrigation. *J Sci Food Agric* **88**:1524-1528 (2008).
 12. Segovia-Bravo KA, García-García P, López-López A and Garrido-Fernández A, Effect of bruising on respiration, superficial color, and phenolic changes in fresh Manzanilla olives (*Olea europaea pomiformis*): Development of treatments to mitigate browning. *J Agric Food Chem* **59**:5456–5464 (2011).
 13. Yousfi K, Weiland CM and García JM. Responses of fruit physiology and virgin oil quality to cold storage of mechanically harvested ‘Arbequina’ olives cultivated in hedgerow. *Grasas y Aceites* **64**:572-582 (2013).
 14. Jiménez-Jiménez F, Castro-García S, Blanco-Roldan GL, González-Sánchez EJ and Gil-Ribes J A. Isolation of table olive damage causes and bruise time evolution during fruit detachment with trunk shaker. *Spanish Journal of Agricultural Research*, **11**:65-71. (2013).
 15. Morales-Sillero A, Rallo P, Jiménez R, Casanova L and Suárez MP, Suitability of two table olive cultivars (‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’) for mechanical harvesting in super-high density hedgerows. *HortScience* **49**:1028-1033 (2014).
 16. Yousfi K, Cayuela JA and García JM, Reduction of virgin olive oil bitterness by fruit cold storage. *J Agric Food Chem* **56**:10085–10091 (2008).
 17. Franco MN, Galeano-Díaz T, Sánchez J, De Miguel C and Martín-Vertedor D, Antioxidant capacity of the phenolic fraction and its effect on the oxidative stability of olive oil varieties grown in the southwest of Spain. *Grasas Aceites* **65**: e004. doi: <http://dx.doi.org/10.3989/gya.051513> (2014).

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
18. Uceda M and Frias L, Harvest dates. Evolution of the fruit oil content, oil composition and oil quality, in *Proceeding of II Seminario Oleícola Internacional; International Olive Oil Council*, Cordoba, Spain, pp 125–130 (1975).
 19. Castellano JM, García JM, Morilla A, Gutiérrez F and Perdiguero S, Quality of ‘Picual’ olive fruits under controlled atmospheres. *J Agric Food Chem* **41**:537-539 (1993).
 20. Martinez JM, Muñoz E, Alba J and Lanzón A, Report about the use of the “Abencor” analyzer. *Grasas Aceites* **26**:379-385 (1975).
 21. Soxhlet F, Die gewichtsanalytische Bestimmung des Milchfettes. *Dingler's Polytech J* **232**: 461–465 (1879).
 22. Annexes II and IX in Official Journal European Communities n. L. 248 of 5 September, Regulation EEC/2568/91, (1991).
 23. Laübli W and Bruttel PA, Determination of the oxidative stability of fats and oils by the ‘Rancimat’ method. *J Am Oil Chem Soc* **63**:792–794 (1986).
 24. Mínguez-Mosquera MI, Rejano-Navarro L, Gándul-Rojas B, Sánchez-Gómez AH and Garrido-Fernández J, Color pigment correlation in virgin olive oils. *J Am Oil Chem Soc* **68**:332–336 (1991).
 25. IUPAC, *Standard methods for the analysis of oils, fats and derivatives*. Method 2432, 7th ed, Pergamon Press, Oxford (1992).
 26. Mateos R, Espartero JL, Trujillo M, Ríos JJ, León-Camacho M, Alcudia F and Cert A, Determination of phenols, flavones and lignans in virgin olive oil by solid-phase extraction and high-performance liquid chromatography with diodearray ultraviolet detection. *J Agric Food Chem* **49**:2185– 2192 (2001).

- 1
2
3 27. Cert A, Moreda W and Pérez-Camino MC, Methods of preparation of fatty acid
4 methyl esters (FAME). Statistical assessment of the precision characteristics from a
5 collaborative trial. *Grasas Aceites* **51**:447-456 (2000).
6
7
8
9
10 28. Box GEP and Cox DR, An analysis of transformations. *J R Stat Soc B* **26**:211–252
11 (1964).
12
13
14 29. Lelievre JM, Latche A, Jones B, Bouzayen M and Pech JC, Ethylene and fruit
15 ripening. *Physiol Plant* **101**:727–739 (1997).
16
17
18 30. Abeles EB, Morgan PW and Saltveit ME, *Ethylene in Plant Biology*, 2nd ed, San
19 Diego, Academic Press, (1992).
20
21
22 31. Yang SF and Hoffman NE, Ethylene biosynthesis and its regulation in higher plants.
23 *Ann Rev Plant Physiol* **35**:155-189 (1984).
24
25
26 32. Klotz K, Suttle JC and Campbell LG, Ethylene production and ethylene effects on
27 respiration rate of postharvest sugarbeet roots. *Postharvest Biol Tec* **56**:71-76
28 (2010).
29
30
31
32
33 33. Martinez-Moreno JM, Pereda J, Gómez-Herrera C, Janer del Valle C, Estudios
34 Físico-Químicos Sobre las Pastas de Aceitunas Molidas. XX. Viscosidades de los
35 Alpechines. *Grasas y Aceites* **15**:226–232 (1964).
36
37
38 34. Barranco D, Trujillo I and Rallo L, Libro I: Elaiografía hispánica, in *Varietades de*
39 *olivo en España*, Ed by Rallo L, Barranco D, Caballero J, Del Río C, Martín A,
40 Tous J and Trujillo I, Junta de Andalucía, MAPA, Mundi-Prensa, Madrid, pp 45-31
41 (2005).
42
43
44
45
46
47
48 35. Ardila T, Sánchez J, Osorio E, Marín J, de Miguel C and Martínez M, Performance
49 of the ‘Manzanilla Cacereña’ cultivar in the industrial olive oil elaboration, *Acta*
50 *Hortic* **949**:553-557 (2012).
51
52
53
54
55
56
57
58
59
60

- 1
2
3 36. Morales-Sillero A, García JM, Torres-Ruiz JM, Montero A, Sánchez-Ortiz A and
4
5 Fernández JE, Is the productive performance of olive trees under localized irrigation
6
7 affected by leaving some roots in drying soil? *Agr Water Manage* **123**:79– 92
8
9 (2013).
10
- 11 37. Yousfi K, Cert RM and Garcia JM, Changes in quality and phenolic compounds of
12
13 virgin olive oils during objectively described fruit maturation. *Eur Food Res*
14
15 *Technol* **223**:117-124 (2006).
16
17
- 18 38. Tovar MJ, Romero MP, Alegre S, Girona J and Motilva MJ, Composition of
19
20 organoleptic characteristics of oil from Arbequina olive (*Olea europaea* L.) trees
21
22 under deficit irrigation. *J Sci Food Agric* **82**:1755-1763 (2002).
23
24
- 25 39. Aparicio R, Roda L, Albi M and Gutierrez F, Effect of various compounds on virgin
26
27 olive oil stability measured by rancimat. *J Agr Food Chem* **47**:4150-4155 (1999).
28
29
- 30 40. Escolar D, Haro RM and Ayuso J, The color space of foods: Virgin olive oil. *J*
31
32 *Agric Food Chem* **55**:2085-2093 (2007).
33
- 34 41. Criado MN, Motilva MJ, Ramo T and Romero MP, Chlorophyll and carotenoid
35
36 profile and enzymatic activities (chlorophyllase and lipoxygenase) in olive drupes
37
38 from the fruit-setting period to harvest time. *J Am Soc Hortic Sci* **131**:593-600
39
40 (2006).
41
42
- 43 42. Morales-Sillero A, Fernández JE, Beltrán G, Jiménez MR and Troncoso A,
44
45 Influence of fertigation in ‘Manzanilla de Sevilla’ olive oil quality. *HortScience*
46
47 **42**:1157-1162 (2007).
48
- 49 43. Servili M, Esposito S, Fabiani R, Urbani S, Taticchi A, Mariucci F, Selvaggini R and
50
51 Montedoro G.F, Phenolic compounds in olive oil: antioxidant, health and
52
53 organoleptic activities according to their chemical structure. *Inflammopharmacology*
54
55 **17**:76–84 (2009).
56
57
58
59
60

- 1
2
3 44. Iconomou D, Arapoglou D and Israilides C, Improvement of phenolic antioxidants
4 and quality characteristics of virgin olive oil with the addition of enzymes and
5 nitrogen during olive paste processing. *Grasas Aceites* **61**:303–311 (2010).
6
7
8
9
10 45. Segovia-Bravo KA, Jarén-Galán M, García-García P and Garrido-Fernández A,
11 Browning reactions in olives: Mechanism and polyphenols involved. *Food Chem*
12 **114**:1380–1385 (2009).
13
14
15
16 46. Artajo LM, Romero MP, Tovar MJ and Motilva MJ, Effect of irrigation applied to
17 olive trees (*Olea europaea* L.) on phenolic compound transfer during olive oil
18 extraction. *Eur J Lipid Sci Tech* **108**:19–27 (2006).
19
20
21
22
23 47. Romero-Segura C, García-Rodríguez R, Sánchez-Ortiz A, Sanz C and Pérez AG,
24 The role of olive β -glucosidase in shaping the phenolic profile of virgin olive oil.
25 *Food Res Int* **45**:191–196 (2012).
26
27
28
29
30 48. García JM, Cuevas MV and Fernández JE, Production and oil quality in
31 ‘Arbequina’ olive (*Olea europaea*, L.) trees under two deficit irrigation strategies.
32 *Irrig Sci* **31**:359–370 (2013).
33
34
35
36 49. Parvaiz M, Hussain K, Shoaib M, William G, Tufail M, Hussain Z, Gohar D and
37 Intiaz S, A review: Therapeutic significance of olive *Olea Europaea* L. (oleaceae
38 family). *Global Journal of Pharmacology* **7**:333-336 (2013).
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Physiological and physical variables in fruits from ‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’ cultivars grown in SHD hedgerow orchards, and harvested by hand or mechanically. Each value is the mean \pm SD of four replicates.

Variables/Harvesting system	‘Manzanilla de Sevilla’		‘Manzanilla Cacereña’		Interaction ^a
	Hand	Mechanical	Hand	Mechanical	
L*	45.9 \pm 14.0	30.9 \pm 9.8	40.0 \pm 15.8	37.0 \pm 11.2	Ns
a*	3.0 \pm 14.4	8.0 \pm 8.0	13.2 \pm 12.7	11.6 \pm 8.8	Ns
b*	23.4 \pm 16.0	9.3 \pm 11.6	14.1 \pm 14.6	15.4 \pm 11.0	Ns
Color index [L*(b*-a*)/100]	13.3 \pm 17.7	1.9 \pm 9.3	3.8 \pm 14.8	3.2 \pm 9.1	Ns
Firmness (N)	32 \pm 4b	24 \pm 8a	28 \pm 4b	22 \pm 5a	Ns
Respiration rate (mL CO ₂ kg ⁻¹ h ⁻¹)	3.9 \pm 0.0aA ^b	4.9 \pm 0.0b	6.4 \pm 0.1aB	7.9 \pm 0.5b	Ns
Ethylene production (μ L kg ⁻¹ L ⁻¹ h ⁻¹)	2.1 \pm 0.2aA	12.0 \pm 1.5bB	5.4 \pm 0.1aB	13.3 \pm 0.2bB	Ns
Virgin oil yield (g kg ⁻¹)	88.2 \pm 4.1B	79.3 \pm 11.4B	45.2 \pm 8.1A	54.0 \pm 8.2A	*
Humidity (g kg ⁻¹)	645.4 \pm 21.2	660.8 \pm 12.1A	676.2 \pm 34.1	695.2 \pm 13.9B	Ns
Total oil yield (g kg ⁻¹)	119.2 \pm 17.7	97.8 \pm 6.3A	114.5 \pm 3.3	126.0 \pm 8.5B	**
Extractability (g kg ⁻¹)	754.1 \pm 98.1B	803.4 \pm 81.7B	393.8 \pm 67.0A	428.7 \pm 70.8A	Ns

^a Cultivar \times harvesting system

^b Mean values in the same row followed by different lowercase letters indicate significant differences in harvesting system for each cultivar at $P \leq 0.05$. Mean values in the same row followed by different capital letters indicate significant differences in cultivars for each harvesting system at $P \leq 0.05$. No letter means non significant effect.

Ns and *, ** indicate non-significant and significant at $P \leq 0.05$ and 0.01, respectively

Table 2. Legally established parameters to evaluate the level of quality of virgin olive oils extracted from ‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’ cultivars grown in SHD hedgerow orchards, and harvested by hand or mechanically. Each value is the mean \pm SD of four replicates.

Variables/Harvesting system	‘Manzanilla de Sevilla’		‘Manzanilla Cacereña’		Interaction ^a
	Hand	Mechanical	Hand	Mechanical	
Acidity (% Oleic acid)	0.3 \pm 0.1a ^b	0.7 \pm 0.1bB	0.3 \pm 0.0	0.3 \pm 0.0A	***
Peroxide value (meq O ₂ kg ⁻¹)	3.6 \pm 0.6	3.6 \pm 0.4	4.4 \pm 0.6	4.8 \pm 0.9	Ns
<i>K</i> ₂₃₂	1.63 \pm 0.24	1.47 \pm 0.18	1.48 \pm 0.02	1.45 \pm 0.10	Ns
<i>K</i> ₂₇₀	0.12 \pm 0.01	0.10 \pm 0.01	0.10 \pm 0.01	0.10 \pm 0.01	Ns
Sensory attributes					
Fruity	4.6 \pm 0.7bB	3.6 \pm 0.8a	3.7 \pm 0.6bA	2.6 \pm 0.8a	Ns
Bitter	3.1 \pm 0.6b	2.3 \pm 0.8a	2.6 \pm 0.9b	1.6 \pm 0.4a	Ns
Pungent	3.2 \pm 0.7	2.9 \pm 0.7B	2.7 \pm 0.2b	1.6 \pm 0.5aA	*
Overall grading (1, worst- 9, best)	7.8 \pm 0.4b	7.4 \pm 0.2a	7.4 \pm 0.4b	6.9 \pm 0.3a	Ns

^a Cultivar \times harvesting system

^bMean values in the same row followed by different lowercase letters indicate significant differences in treatments for each cultivar at $P \leq 0.05$. Mean values in the same row followed by different capital letters indicate significant differences in cultivars for each treatment at $P \leq 0.05$. No letter means non significant effect.

Ns and * and *** indicate non-significant and significant at $P \leq 0.05$ and 0.001, respectively

Table 3. Oxidative stability, tocopherol composition and photosynthetic pigments content of virgin olive oils extracted from ‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’ cultivars grown in SHD hedgerow orchards, and harvested by hand or mechanically. Each value is the mean \pm SD of four replicates.

	‘Manzanilla de Sevilla’		‘Manzanilla Cacereña’		Interaction ^a
	Hand	Mechanical	Hand	Mechanical	
Oxidative stability (h)	113.5 \pm 7.2bB ^b	89.0 \pm 10.5a	98.6 \pm 4.5bA	85.2 \pm 4.7a	Ns
α -Tocopherol (mg kg ⁻¹)	405.1 \pm 98.5bB	170.0 \pm 59.3a	144.2 \pm 47.3A	171.4 \pm 22.4	**
β -Tocopherol (mg kg ⁻¹)	41.8 \pm 9.6bB	8.8 \pm 9.9a	ND	ND	**
γ -Tocopherol (mg kg ⁻¹)	ND	ND	ND	ND	Ns
Total tocoferols (mg kg ⁻¹)	446.9 \pm 107.0bB	178.8 \pm 68.9a	144.2 \pm 47.3A	171.4 \pm 22.4	**
Chlorophylls (mg kg ⁻¹)	7.8 \pm 1.7	7.6 \pm 1.5	6.9 \pm 1.8	7.5 \pm 1.9	Ns
Carotenoids (mg kg ⁻¹)	6.0 \pm 1.1	5.4 \pm 0.6A	5.7 \pm 0.4a	7.3 \pm 1.2bB	*

^a Cultivar \times harvesting system

^b Mean values in the same row followed by different lowercase letters indicate significant differences in treatments for each cultivar at $P \leq 0.05$. Mean values in the same row followed by different capital letters indicate significant differences in cultivars for each treatment at $P \leq 0.05$. No letter means non significant effect.

ND, Ns and *, ** indicate non detected, non-significant and significant at $P \leq 0.05$ and 0.01, respectively

Table 4. Phenolic composition (mg kg⁻¹) of virgin olive oils extracted from ‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’ cultivars grown in SHD hedgerow orchards, and harvested by hand or mechanically. Each value is the mean ± SD of four replicates.

	‘Manzanilla de Sevilla’		‘Manzanilla Cacereña’		Interaction ^a
	Hand	Mechanical	Hand	Mechanical	
Hydroxytyrosol	5.7 ± 2.2aB ^x	11.2 ± 3.7b	1.5 ± 0.2A	4.7 ± 5.3	Ns
Tyrosol	9.0 ± 1.6a	13.7 ± 2.9bB	8.1 ± 0.4	8.3 ± 0.9A	*
Vanillic acid	0.6 ± 0.0B	0.6 ± 0.1B	0.5 ± 0.0bA	0.3 ± 0.0aA	*
Vanilline	ND	ND	0.1 ± 0.0	0.1 ± 0.1	Ns
p-Coumaric acid	1.7 ± 0.3B	1.8 ± 0.5B	0.3 ± 0.0A	0.3 ± 0.0A	Ns
hydroxytyrosol acetate	5.0 ± 2.1	7.3 ± 0.6B	2.8 ± 0.4a	3.5 ± 0.4bA	Ns
3, 4-DHPEA-EDA ^c	107.7 ± 25.2bB	48.5 ± 27.1a	45.1 ± 3.7bA	31.6 ± 5.9a	Ns
Tirosol acetate	ND	ND	ND	ND	Ns
p-HPEA-EDA ^d	65.9 ± 28.6	47.6 ± 8.1B	41.5 ± 4.1b	27.4 ± 3.3aA	Ns
Pinosresinol	2.7 ± 0.6bA	0.8 ± 0.9aA	3.7 ± 0.1bB	3.0 ± 0.5aB	Ns
Cinamic acid	ND	ND	ND	ND	Ns
Acetoxypinosresinol	25.5 ± 2.9b	12.5 ± 7.4a	9.6 ± 1.7	5.6 ± 3.0	Ns
3, 4-DHPEA-EA ^e	219.0 ± 57.3bB	130.9 ± 42.2a	107.1 ± 9.8bA	75.7 ± 13.7a	Ns
p-HPEA-EA ^f	21.2 ± 8.6	17.2 ± 2.4	18.4 ± 0.9	19.4 ± 0.5	Ns
Ferulic acid	70.1 ± 15.0B	86.7 ± 9.2B	10.6 ± 1.4A	10.9 ± 2.6A	Ns
Luteoline	3.2 ± 0.5A	3.5 ± 0.4A	6.3 ± 0.4B	6.3 ± 1.0B	Ns
Apigenine	1.8 ± 0.2A	1.6 ± 0.3A	3.0 ± 0.1B	3.3 ± 0.4B	Ns
Σ Phenols	539.1 ± 86.7bB	383.9 ± 79.5aB	258.6 ± 16.2bA	200.4 ± 36.0aA	Ns
ΣO-diphenols	340.6 ± 66.1bB	201.4 ± 66.7aB	162.8 ± 13.1bA	121.8 ± 13.8aA	*
ΣSecoiridoid derivatives	413.8 ± 93.5bB	244.2 ± 75.6aB	212.1 ± 14.5bA	154.1 ± 30.1aA	Ns

^a Cultivar × harvesting system

^b Mean values in the same row followed by different lowercase letters indicate significant differences in treatments for each cultivar at $P \leq 0.05$. Mean values in the same row followed by different capital letters indicate significant differences in cultivars for each treatment at $P \leq 0.05$. No letter means non significant effect.

ND, Ns and * indicate non detected, non-significant and significant at $P \leq 0.05$, respectively

^c Dialdehydic form of the decarboxymethyl oleuropeinaglycone.

^d Dialdehydic form of the decarboxymethyl ligstrosideaglycone.

^e Hydroxytyrosylelenolate.

^f Tyrosylelenolate.

Table 5. Fatty acid composition (g kg^{-1}) of virgin olive oils extracted from ‘Manzanilla de Sevilla’ and ‘Manzanilla Cacereña’ cultivars grown in SHD hedgerow orchards, and harvested by hand or mechanically. Each value is the mean \pm SD of four replicates.

Fatty acid ^a	‘Manzanilla de Sevilla’		‘Manzanilla Cacereña’		Interaction ^a
	Hand	Mechanical	Hand	Mechanical	
Palmític (16:0) ^c	140.1 \pm 2.2B ^b	139.0 \pm 1.1B	129.3 \pm 2.4A	127.3 \pm 0.1A	Ns
Palmitoleic (16:1)	13.2 \pm 0.1	14.4 \pm 0.2B	12.0 \pm 0.0	11.2 \pm 0.0A	Ns
Estearic (18:0)	28.1 \pm 0.2B	30.2 \pm 1.1B	15.4 \pm 0.3A	15.5 \pm 1.1A	Ns
Oleic (18:1)	749.3 \pm 6.3A	746.2 \pm 3.3A	784.2 \pm 1.1B	786.2 \pm 3.3B	Ns
Linoleic (18:2)	55.2 \pm 4.2B	56.0 \pm 3.3B	44.0 \pm 1.1A	43.4 \pm 2.2A	Ns
Araquídic (20:0)	4.0 \pm 0.1	4.1 \pm 0.2	3.8 \pm 0.4	4.2 \pm 0.4	Ns
Linolenic (18:3)	2.0 \pm 0.2A	2.0 \pm 0.5A	4.4 \pm 0.4B	4.3 \pm 0.4B	Ns
Gondoic (20:1)	7.0 \pm 0.3	7.2 \pm 0.4	8.0 \pm 0.3	8.2 \pm 0.3	Ns
Behenic (22:0)	1.0 \pm 0.1	1.0 \pm 0.1	1.1 \pm 0.1	1.1 \pm 0.1	Ns

^a Cultivar \times harvesting system

^b Mean values in the same row followed by different lowercase letters indicate significant differences in treatments for each cultivar at $P \leq 0.05$. Mean values in the same row followed by different capital letters indicate significant differences in cultivars for each treatment at $P \leq 0.05$. No letter means non significant effect.

Ns indicate non-significant

^c(Carbon number : unsaturation number)