



**Cite this article:** Thiel D, Bauknecht P, Jékely G, Hejnol A. 2017 An ancient FMRFamide-related peptide–receptor pair induces defence behaviour in a brachiopod larva. *Open Biol.* **7**: 170136.

<http://dx.doi.org/10.1098/rsob.170136>

Received: 5 June 2017

Accepted: 25 July 2017

#### Subject Area:

cognition/bioinformatics/cellular biology

#### Keywords:

Trochozoa, FMRFamide, neuropeptide receptor, defence behaviour, planktonic larva, brachiopod

#### Author for correspondence:

Andreas Hejnol

e-mail: [andreas.hejnol@uib.no](mailto:andreas.hejnol@uib.no)

Electronic supplementary material is available online at <https://dx.doi.org/10.6084/m9.figshare.c.3850909>.

# An ancient FMRFamide-related peptide–receptor pair induces defence behaviour in a brachiopod larva

Daniel Thiel<sup>1</sup>, Philipp Bauknecht<sup>2</sup>, Gáspár Jékely<sup>2</sup> and Andreas Hejnol<sup>1</sup>

<sup>1</sup>Sars International Centre for Marine Molecular Biology, University of Bergen, Thormøhlensgate 55, 5006 Bergen, Norway

<sup>2</sup>Max Planck Institute for Developmental Biology, Spemannstraße 35, 72076 Tübingen, Germany

 DT, 0000-0003-1398-3512; GJ, 0000-0001-8496-9836; AH, 0000-0003-2196-8507

Animal behaviour often comprises spatially separated sub-reactions and even ciliated larvae are able to coordinate sub-reactions of complex behaviours (metamorphosis, feeding). How these sub-reactions are coordinated is currently not well understood. Neuropeptides are potential candidates for triggering larval behaviour. However, although their immunoreactivity has been widely analysed, their function in trochozoan larvae has only been studied for a few cases. Here, we investigate the role of neuropeptides in the defence behaviour of brachiopod larvae. When mechanically disturbed, the planktonic larvae of *Terebratalia transversa* protrude their stiff chaetae and sink down slowly. We identified endogenous FLRFamide-type neuropeptides (AFLRFamide and DFLRFamide) in *T. transversa* larvae and show that the protrusion of the chaetae as well as the sinking reaction can both be induced by each of these peptides. This also correlates with the presence of FLRFamidergic neurons in the apical lobe and adjacent to the trunk musculature. We deorphanized the AFLRFamide/DFLRFamide receptor and detected its expression in the same tissues. Furthermore, the ability of native and modified FLRFamide-type peptides to activate this receptor was found to correspond with their ability to trigger behavioural responses. Our results show how FLRFamide-type neuropeptides can induce two coherent sub-reactions in a larva with a simple nervous system.

## 1. Background

Planktonic organisms have evolved different strategies to defend themselves from predation [1–4]. Morphological characters such as shells, spines or chaetae [5–7] and behaviours such as vertical migration, contraction, active fleeing or passive sinking [8–11] can help to cope with certain predators. This is especially true for ciliated larvae that do not possess an elaborated nervous system and face the challenge of remaining in the water column for dispersal while avoiding predation. The startle behaviour of several planktonic annelid and brachiopod larvae has often been described as a defence strategy, where they stop swimming and protrude long and pointed chaetae [12–16]. The co-occurring sub-reactions of spreading the chaetae and stopping swimming take place in spatially separated tissues: the internal trunk musculature and the ciliated apical edge, respectively. Both sub-reactions have to be coordinated within the framework of a larval nervous system. One mechanism to achieve coordination of different reactions could be the use of neuropeptides as signalling molecules. Neuropeptides are known to influence many behaviours and can be crucial in the regulation and coordination of spatially or temporally separated coherent sub-reactions. During insect ecdysis, for example, the eclosion hormone and ecdysis-triggering hormone both act as a form of master-regulator on different peripheral as well as central targets, and each coordinates several sub-reactions [17–20]. Another example is neuropeptide

Y, which stimulates appetitive as well as consummatory ingestive behaviour in the Siberian hamster [21].

The influence of neuropeptides on the behaviour of trochozoan larvae has only been demonstrated in a few studies, which show that neuropeptides can trigger settlement and influence their ciliary-based locomotion [22–25]. One of the neuropeptides that has been shown to influence ciliary beating of different trochozoan larvae is FMRFamide [22,23,25]. FMRFamide-immunoreactivity is widely used as a marker for neural substructures in morphological studies [26,27]. Furthermore, while FMRFamide-related peptides (FaRPs) have been identified in many metazoans, their phylogenetic relationships are difficult to infer [28–30]. For the comparison of larval nervous systems, it is therefore of crucial interest to understand the functional role of a neuropeptide and its versatility to trigger larval behaviours.

While experimental studies in trochozoan larvae are limited, the physiological effect of FMRFamide-like peptides has been intensively investigated in adult trochozoans, where it has been found to have various effects on muscular activity [31–37]. Depending on the species, it can increase or decrease the heartbeat [31], cause contractions or relaxation of somatic muscles [32–34] or modulate the effects of classical neurotransmitters on somatic muscles [36,37]. Many immunohistochemical analyses on trochozoan larvae of different clades show FMRFamide-like immunoreactivity associated with muscles or ciliary bands [38–46], but experimental data are most often missing and functional studies are restricted to mollusc and annelid larvae [22,23,25]. Despite the recurring association of FMRFamide-like peptides with musculature in trochozoans, only one study has shown an effect of FMRFamide on the musculature of a trochozoan larva, which describes the induction of frequent contractions of the ciliated velum of *Tritia obsoleta* veliger larvae [22].

Since neuropeptides can act over longer distances [47], the localization of the neuropeptide receptor provides more information about the tissues actually affected than the peptide secreting cells that are labelled with the peptide antibodies. The majority of neuropeptide receptors are G-protein-coupled receptors (GPCRs), with a few exceptions like insulin receptors or peptide-gated ion channels [29,48,49]. Three different receptors for FMRFamide have been deorphanized in invertebrates so far. One is an FMRFamide-gated amiloride-sensitive Na<sup>+</sup> channel (FaNaCh) that has been identified in molluscs [50–52]. The two other receptors belong to two different groups of neuropeptide GPCRs. One of these FMRFamide-GPCRs was identified in the fruit fly *Drosophila melanogaster* [53,54] and the other one in the annelid *Platynereis dumerilii* [49]. This stands in contrast to many cases in which homologous ligands also activate homologous receptors [49,55,56].

To expand the taxon sampling of functional neuropeptide studies in lophotrochozoans and to better understand the role of the widely used neuropeptide marker FMRFamide, we investigated the effects of an FaRP on the behaviour of a brachiopod larva. Recent research on brachiopods has revealed important insights in evolutionary developmental biology [57–59] and descriptions of their nervous system include the use of FMRFamide antibodies [60,61] and classical neuronal markers [62,63], as well as other molecular techniques [64]. Here, we show that the endogenous FLRFamide-like peptides induce the characteristic defence behaviour in the larvae of the brachiopod *Terebratalia transversa*, which consists of a downward sinking and the protrusion of their chaetae. Behavioural

experiments and receptor deorphanization, in combination with immunohistochemistry, and *in situ* hybridization show that both sub-reactions can be specifically triggered by a single peptide acting via an ancient FaRP receptor. Together our results show how a single neuropeptide can trigger two coherent reactions and integrate evolutionary novelties such as trochozoan chaetae [57] into the *T. transversa* larval defence behaviour.

## 2. Material and methods

### 2.1. Collection and rearing of *Terebratalia transversa* larvae


Adult *T. transversa* (Sowerby, 1846) were collected in January 2015 and 2016 by dredging at approximately 50–100 m depth close to the University of Washington's Friday Harbor Laboratories, San Juan Islands, WA, USA. Larvae were obtained according to Stricker & Reed [65] by artificial fertilization and kept at 8–10°C. Two different types of larvae were used for the experiments: early larvae (2 days post-fertilization) before chaeta formation and late larvae (4–5 days post-fertilization) with clearly developed mantle lobes and long chaetae. For immunohistochemistry and *in situ* hybridization, larvae were relaxed in 7.8% MgCl<sub>2</sub>·6H<sub>2</sub>O in distilled water for 10–15 min, fixed in 4% methanol-free formaldehyde in seawater for 1 h, subsequently washed in PBS + 0.1% Tween and transferred into 100% methanol for storage at –20°C.

### 2.2. Bioinformatics

The previously published transcriptome of *T. transversa* (SRX1307070) was searched for peptide precursor and receptor candidates using BLAST. Publicly available FMRFamide-like precursor sequences from NCBI were used as reference sequences, and the resulting candidates were checked for signal peptides, cleavage sites and amidation sites. Neuropeptide precursor genes were also searched in the transcriptome of *Novocrania anomala* (SRX1343816). As reference sequences for the peptide receptor, previously published datasets [29,49] were used as well as transcriptomes of *Xenoturbella bocki* (SRX1343818), *Nemertoderma westbladi* (SRX1343819), *Meara stichopi* (SRX1343814) and *Halicryptus spinulosus* (SRX1343820) to obtain additional sequences. The candidates were compared using the software CLANS [66], with a *p*-value cutoff of  $1 \times 10^{-70}$ . Sequences that were strongly connected in the cluster map were aligned with CLUSTAL X v. 2.1 [67], non-conserved stretches were deleted manually, and the best fitting amino acid substitution matrix was determined with MODELGENERATOR v. 0.85 [68]. The final phylogenetic analysis was calculated with PHYML v. 3.0 [69] with 500 bootstrap replicates and visualized with FIGTREE v. 1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree>).

### 2.3. Behavioural assays

The reaction of larvae to synthetic peptides (GenScript) that were predicted from the prepropeptide sequence were tested and compared with the reactions to peptides with non-native modifications. Freely swimming larvae were exposed to different concentrations of peptides in 4-well and 6-well plates (1 ml and 5 ml total volume per well, respectively), and their



**Figure 1.** Defence reaction and FLRFamide prepropeptide of the *T. transversa* larva. (a–c) sketch, (d,e) SEM photographs, anterior up, (f) sketch. (a) Larva in relaxed stance during normal swimming; (b) non-contracted larva that begins to spread its chaetae; (c) larva in defence stance with outspread chaetae; (d) non-contracted competent larva; (e) contracted competent larva with outspread chaetae; (f) schematic of *T. transversa* FLRFamide prepropeptide. al, anterior lobe; ml, mantle lobe; pl, pedicle lobe; vm, ciliated ventral midline. Scale bar, 50  $\mu\text{m}$ .

reactions were observed under a stereomicroscope. To determine the efficiency of the native and modified peptides, the larvae were tested for the necessary minimum concentration at which they fully contracted and spread their chaetae, using 30–100 larvae in each test. To get an estimation of the peptide concentration that was necessary to induce a complete contraction, the larvae were initially exposed to 50  $\text{nmol l}^{-1}$  of the respective peptide and the concentration was then increased stepwise until the larvae fully contracted or a concentration of 50  $\mu\text{mol l}^{-1}$  was reached as an upper cutoff. Larvae were considered fully contracted when their chaetae were spread in all directions (figure 1c) and when they did not further increase in contraction after an increase in peptide concentration, or after the use of the most sensitive peptide. To test the influence of the peptides on the vertical distribution of larvae in a water column, freely swimming larvae (50–100 per treatment) were exposed to peptides in transparent 4.5 ml cuvettes. About 60 s after addition of the peptides, the larvae were recorded with a digital camera (DMK 31AU03 camera, The Imaging Source) in a darkened box with artificial top-illumination. All experiments were repeated at least once with a different batch of larvae from another fertilization and the outcome was averaged. Control animals were exposed to DMSO only.

#### 2.4. Receptor deorphanization

For the receptor deorphanization, the procedure was used as described by Bauknecht & Jékely [49]: full-length open reading frames of the receptor candidate sequences were amplified by PCR from cDNA of mixed larval stages. The forward primers included a 5' attachment consisting of a spacer, an EcoRI or BamHI restriction site and a Kozak sequence and the reverse primers included a 5' attachment consisting of a spacer and a NotI restriction site (see the electronic supplementary material for primer sequences). The amplified products were cut with the corresponding restriction enzymes, cloned into pcDNA3.1(+) mammalian expression vector (Sigma-Aldrich), sequenced from both ends with a T7 forward and a bGH

reverse primer, and transfected into CHO-K1 cells together with a calcium-sensitive luminescent apoaequorin-GFP fusion protein encoding plasmid (G5A) and a promiscuous  $\text{G}\alpha\text{-16}$  protein encoding plasmid. After 2 days, Coelenterazine h (Promega, Madison, WI, USA) was added and incubated with the cells for 2 h. The luminescence response of the transfected cells was measured in a plate reader (BioTek Synergy Mx or Synergy H4, BioTek, Winooski, USA) over 45 s after addition of the neuropeptides. The response of the cells to 1 mM histamine was used as a general control in each plate. All measurements for the dose–response curves were made twice with different cell passages. Dose–response curves were calculated using PRISM 6 (GraphPad, La Jolla, USA) and normalized against the upper plateau values (100% activation).

#### 2.5. *In situ* hybridization

FLRFamide precursor and receptor sequences were amplified by PCR and cloned into pGEM T-easy vector (Promega) for *in vitro* transcription of DIG-UTP or DNP-UTP labelled RNA probes. For tropomyosin, a previously published clone [70] was used. The *in situ* hybridization protocol with an alternative hybridization buffer is published elsewhere [71]. The protocol from Hejnal [72] was adjusted with a proteinase K treatment (10  $\mu\text{g ml}^{-1}$ ) of 8 min and with a postfixation in 3.7% formaldehyde + 0.2% glutaraldehyde in PBS + 0.1% Tween 20. The hybridization buffer contained 4  $\text{mol l}^{-1}$  urea, 5 $\times$  SSC, 1% dextran, 1% SDS, 50  $\mu\text{g ml}^{-1}$  heparin, 50  $\mu\text{g ml}^{-1}$  single-stranded DNA (no formamide). The signal was developed with the TSA Plus Cy3 or Cy5 kit (Perkin Elmer) or NBT/BCIP as a substrate and detected via fluorescence or NBT/BCIP reflection [73] with a Leica SP5 confocal laser-scanning microscope.

#### 2.6. Immunohistochemistry

Customized polyclonal antibodies were raised in rabbits against CFLRFamide, coupled via a disulfide bridge to Keyhole limpet hemocyanin (GenScript<sup>®</sup>). Co-staining was either

done with mouse anti-acetylated  $\alpha$ -tubulin (Sigma, T6793) or mouse anti-actin (Seven Hills Bioreagents, LMAB-C4) antibodies. For the staining procedure, the protocol of Conzelmann & Jékely [74] was used with the following adjustments: proteinase K treatment ( $10 \mu\text{g ml}^{-1}$ ) was done for 3–5 min, and after the proteinase inactivation step with glycine ( $2 \text{ mg ml}^{-1}$ ) the samples were incubated for 2–4 h in PBS + 0.5% TritonX. Primary antibodies were incubated over three nights at  $4^\circ\text{C}$  and washed for 4–6 h with at least 10 changes of washing medium. Secondary antibodies (Alexa 555 goat anti-rabbit and Alexa 647 goat anti-mouse) were incubated overnight and an additional secondary antibody (Alexa 488 goat anti-rat) without corresponding primary antibody was included to test and subtract unspecific staining. After washing the secondary antibodies for 4–6 h with at least 10 changes of buffer, specimens were transferred into methanol and mounted in Murray's clear (2 : 1 parts benzyl benzoate : benzyl alcohol).

### 3. Results

#### 3.1. The endogenous neuropeptides DFLRFamide and AFLRFamide trigger the defence behaviour of *Terebratalia transversa* larvae

During normal swimming, the chaetae of competent *T. transversa* larvae rest against their pedicle lobe with their tips forming a bundle (figure 1a). When the larvae get disturbed (e.g. mechanical irritation with a pipette tip), they stop swimming, sink down slowly and exhibit a defensive stance by lengthwise contraction of their body to spread the four bundles of chaetae outwards (figure 1b,c,e; electronic supplementary material, Video S1). At maximal contraction, the larvae spread their chaetae in all directions to surround their soft body (figure 1c,e).

We identified an FLRFamide prepropeptide sequence in the transcriptome of *T. transversa*. The FLRFamide precursor contains a signal peptide, three copies of DFLRFamide and five copies of AFLRFamide, partially separated by intermediate sequences (figure 1f; see also the electronic supplementary material for colour-coded amino acid sequence). When we exposed larvae to synthetic FLRFamide, they contracted lengthwise, spread out their chaetae and sank down slowly. Both predicted peptides, DFLRFamide and AFLRFamide, caused the same behaviour. When exposed to  $50\text{--}100 \text{ nmol l}^{-1}$  DFLRFamide, all larvae showed initial signs of contraction, indicated by their chaetae bundles being slightly fanned out while still pointing in a posterior direction (figure 1b,d). About half of the larvae continued swimming while the other half started to sink slowly towards the bottom. An increase in peptide concentrations led to an overall increase in contraction, resulting in a stronger spreading of the chaetae and more larvae sinking. A maximum contraction of all larvae, with their body being completely surrounded by chaetae, was observed at concentrations of  $500\text{--}750 \text{ nmol l}^{-1}$  DFLRFamide. When we removed the peptides by exchanging the medium with fresh seawater, all larvae returned to normal swimming behaviour. Continuous exposure for about 2 h led to a desensitization and the larvae resumed normal swimming without removing the peptides. When the larvae were desensitized by continuous exposure to DFLRFamide, they also became insensitive to AFLRFamide and vice versa. Desensitized larvae still showed

an initial contraction when they were disturbed, but it seemed like they had to be disturbed more harshly than non-desensitized larvae. Larvae that were exposed to DFLRFamide concentrations below the threshold that is able to induce a defensive behaviour ( $50 \text{ nmol l}^{-1}$ ) seemed to be more sensitive than untreated ones and already contracted when slightly disturbed. However, we were not able to quantify the necessary strengths of disturbance. We also tested orthologues of AFLRFamide/DFLRFamide on late chaetous larvae of *P. dumerilii* and *Novocrania anomala* (FMRFamide and YMRFamide, respectively; see the electronic supplementary material for *N. anomala* precursor sequence), but even concentrations above  $50 \mu\text{mol l}^{-1}$  did not induce any defence reaction.


Taken together, we found that *T. transversa* larvae show a sustained behaviour which is similar to their startle response when we exposed them to one of the neuropeptides encoded on the endogenously expressed FLRFamide precursor.

#### 3.2. FLRFamide causes sinking of larvae independent from the protrusion of their chaetae

One part of the defence behaviour of *T. transversa* larvae is a slow downward sinking. A similar reaction can already be observed in early larvae before they develop long chaetae. Owing to the shape of the larvae and the lack of a clear restriction of the prototroch, it was not possible to directly record the ciliary beating. However, because *T. transversa* larval locomotion is purely driven by ciliary beating, we hypothesize that FLRFamide influences the ciliary movement. To measure the swimming behaviour in an unbiased manner, we recorded the position of freely swimming larvae in vertical columns and compared it to the position of larvae after exposure to DFLRFamide. To test the possibility that the sinking is caused by an increase in the water drag due to the protruded chaetae, we also recorded early larvae that already express FLRFamide in the apical lobe (electronic supplementary material, figure S1d) but do not possess long chaetae yet (electronic supplementary material, figure S1e). Both stages showed a sinking behaviour that shifted the distribution of the larvae in the water column downwards, compared to the controls (figure 2). Early larvae kept swimming more freely close to the bottom, whereas late larvae usually stayed at the bottom and moved only very slowly forward.

#### 3.3. Modified peptides trigger the defence behaviour at different concentration thresholds

We tested at which concentration modified peptides induce the contraction that leads to the erection of the chaetae, with  $50 \mu\text{mol l}^{-1}$  as a cutoff for the maximum concentration (table 1; electronic supplementary material, table S1). The larvae were most sensitive to DFLRFamide and showed full contraction (figure 1c,e), sinking and very slow movements on the bottom of the dish at concentrations between 500 and  $750 \text{ nmol l}^{-1}$  (batch dependent). Further increasing of the concentration did not lead to an obvious increase in the reaction. AFLRFamide was slightly less effective by triggering full contraction of all larvae between 1 and  $1.5 \mu\text{mol l}^{-1}$ . The reduced peptide sequence FLRFamide was effective at  $3 \mu\text{mol l}^{-1}$ . Changing the amidated C-terminal phenylalanine to an amidated tryptophan reduced the effectiveness by



**Figure 2.** Influence of FLRFamide on the vertical distribution of early and late larvae in a water column. Horizontal red bar shows average level of swimming height,  $p$ -values are calculated for difference in distribution of larvae in upper versus lower half of the column (two-tailed, unpaired  $t$ -test), blue line is the estimated trend line (not statistically supported). Distribution was measured over a period of 5 s, about 1 min after exposure to  $5 \mu\text{mol l}^{-1}$  DFLRFamide.

**Table 1.** Necessary peptide concentrations to evoke larval defence stance compared to  $\text{EC}_{50}$  values of receptor activation.

peptide	necessary concentration to induce full contraction ( $\mu\text{mol l}^{-1}$ )	$\text{EC}_{50}$ receptor assay
DFLRFamide	0.625	$26.5 \text{ nmol l}^{-1}$
AFLRFamide	1.5	$12.4 \text{ nmol l}^{-1}$
FLRFamide	3	$33.2 \text{ nmol l}^{-1}$
DFLRFamide	8.75	$0.9 \mu\text{mol l}^{-1}$


about 10-fold, with a minimum necessary concentration of  $7.5 \mu\text{mol l}^{-1}$  for DFLRFamide or  $20 \mu\text{mol l}^{-1}$  for AFLRFamide. Changing the C-terminal phenylalanine to the non-aromatic leucine only led to a very weak contraction (similar to figure 1b) in some of the larvae at  $50 \mu\text{mol l}^{-1}$  DFLRFamide, whereas  $50 \mu\text{mol l}^{-1}$  AFLRFamide gave no reaction at all. Reducing the sequence to the three C-terminal amino acids, LRFamide, also did not lead to any contraction, and nor did any of the other endogenous peptides that we tested. (A list of tested peptides is given in the electronic supplementary material, table S1.) The overall most effective versions were the ones that are encoded on the pro-peptide sequence, DFLRFamide and AFLRFamide. The reduced peptide FLRFamide was slightly less effective and a modification of the amidated C-terminus reduced the effectiveness even more.

### 3.4. Identification of the *Terebratalia transversa* FaRP receptor


Based on BLAST e-value similarities and cluster analyses, we tested four receptor candidates for their activation by FLRFamide (figure 3). One candidate belongs to a cluster of

receptors that includes the deorphanized *P. dumerilii* FMRFamide receptor [49] with related sequences in all major bilaterian groups including Xenacoelomorpha (figure 3 'I'; electronic supplementary material, figure S2). The second candidate belongs to the luqin receptors (figure 3 'II'; electronic supplementary material, figure S2). The third candidate belongs to a group of related receptors with unknown ligand (figure 3 'III'; electronic supplementary material, figure S2), and the fourth one belongs to a group that shows similarities with the *Drosophila melanogaster* FMRFamide and the *P. dumerilii* NPY-4 receptors and is not related to the other three receptors (figure 3 'IV'). Transcriptome searches for an FMRFamide-gated ion channel (FaNaCh), which has been identified in molluscs, did not reveal any orthologues in *T. transversa*, even when using FaNaCh orthologues that were identified in the brachiopods *Lingula anatina* and *Novocrania anomala*.

To test whether FLRFamide is the ligand of one of these receptors, we tested their activation by DFLRFamide in transfected CHO-K1 cells. Only the candidate that is related to the *P. dumerilii* FMRFamide receptor was activated by  $1 \mu\text{mol l}^{-1}$  DFLRFamide, but none of the other tested receptors. We therefore called this receptor the *T. transversa* FLRFamide receptor. As DFLRFamide triggered the defence stance at concentrations below  $1 \mu\text{mol l}^{-1}$  in the behavioural assay, we did not test the negative GPCRs at higher peptide doses. We further compared the luminescence response of FLRFamide receptor expressing CHO-K1 cells to  $1 \mu\text{mol l}^{-1}$  DFLRFamide, AFLRFamide, FLRFamide, DFLRFamide and DFLRFamide (figure 4a). The two native forms DFLRFamide and AFLRFamide led to the highest luminescence, followed by FLRFamide in a similar range. DFLRFamide gave a strongly decreased luminescence and the values of DFLRFamide were barely higher than the negative control. Dose-response curves were recorded for DFLRFamide, AFLRFamide, FLRFamide and DFLRFamide (figure 4b; electronic supplementary material, figure S3) and  $\text{EC}_{50}$  values (half maximal effective concentration) were determined for each peptide. AFLRFamide showed the



**Figure 3.** Clustermap of metazoan neuropeptide GPCRs related to FMRFamide receptors. Connections correspond to blastp connection with  $p$ -values of less than  $1 \times 10^{-70}$ . Groups that include the receptors I–IV that were tested for activation by FLRFamide are encircled. NPY, neuropeptide Y; NPF, neuropeptide F; PRL, prolactin releasing peptide.



**Figure 4.** Luminescence response of *T. transversa* FLRFamide receptor expressing CHO-K1 cells to different peptides. (a) Relative luminescence of *T. transversa* FLRFamide receptor expressing cells after exposure to different peptides with a fixed concentration of  $1 \mu\text{mol l}^{-1}$ . (b) Dose–response curves of *T. transversa* FLRFamide receptor expressing cells to different concentrations of DFLRFamide, AFLRFamide, FLRFamide and DFLRWamide. Luminescence values are given relative to maximum luminescence (max = 1). RLU relative luminescence.

lowest EC<sub>50</sub> value ( $1.24 \times 10^{-8} \text{ mol l}^{-1}$ ), the EC<sub>50</sub> for DFLRFamide was about two times as high ( $2.65 \times 10^{-8} \text{ mol l}^{-1}$ ), the one for FLRFamide was about three times higher ( $3.32 \times 10^{-8} \text{ mol l}^{-1}$ ) and the one for DFLRWamide was the highest of all tested peptides ( $9.06 \times 10^{-7} \text{ mol l}^{-1}$ ). The EC<sub>50</sub> values are listed in table 1, together with the concentrations necessary to trigger the defence stance in the behavioural assay.


After we deorphanized the FLRFamide receptor, we tested its phylogenetic relationship to the receptors that showed connections in the cluster analysis (figure 3 ‘I–III’). We did not include the unrelated *T. transversa* orphan receptor that is related to the insect FMRFamide and *P. dumerilii* NPY-4 receptors (figure 3 ‘IV’). As seen in the cluster analysis, the *T. transversa* FLRFamide receptor is directly related to the *P. dumerilii* FMRFamide receptor and several orphan receptors of other trochozoans (electronic supplementary material, figure S2). Orthologues to these trochozoan FMRFamide/FLRFamide receptors were found in the insect *Nilaparvata lugens*, the hemichordate *Saccoglossus kowalevskii*, and the xenacoelomorph *Meara stichopi*. Further related

GPCRs include orphan receptors from the cephalochordate *Branchiostoma floridae*, the ghost-shark *Callorhynchus milii*, and the xenacoelomorphs *M. stichopi* and *Nemertoderma westbladi*. All of these receptors form a fully supported group of neuropeptide GPCRs with homologues in all major bilaterian clades that are well separated from other neuropeptide GPCR groups (electronic supplementary material, figure S2).

In summary, we discovered that the *T. transversa* FLRFamide receptor belongs to an ancient neuropeptide receptor group and is efficiently activated by the two peptides AFLRFamide and DFLRFamide that are encoded on the *T. transversa* prepropeptide sequence.

### 3.5. *In situ* hybridization and immunohistochemistry show localization of peptide receptor in trunk musculature and apical prototroch region

The FLRFamide precursor has several expression domains within the apical lobe around the neuropile, and two domains



**Figure 5.** *In situ* hybridization and immunostaining of *Terebratalia* FLRFamide, *Terebratalia* FLRFamide receptor, musculature and tubulin. (a–e) *in situ* hybridization; (f–i) immunohistochemistry; (a,c,d,e,g,j) front view; (b,f,h,i) side view, ventral side left. (a,b) *FLRFa* and *tropomyosin* expression, arrows show *FLRFa* expression in mantle lobe. (c) *FLRFa* and *FLRFa* receptor expression, arrows show *FLRFa* expression in mantle lobe. (d) *FLRFa* receptor expression, stars show expression underneath prototroch. (e) *tropomyosin* and *FLRFa* receptor co-expression around chaetae sacs. (f) *FLRFa* and tubulin staining, star shows branching of *FLRFa*-positive nerves inside ventral trunk area, arrows show branching of dorsal *FLRFa*-positive trunk-nerve towards chaetae sacs. (g) *FLRFa* staining, stars show branching of *FLRFa*-positive nerves inside dorsal trunk area, arrows show projections into secretory cells underneath the prototroch. (h) *FLRFa* and tubulin staining, arrows show nerve projecting from neuropile into ventral part of the trunk. (i) *FLRFa* and actin staining, arrows show lining of the musculature by *FLRFa*-midgergic nerves, projecting into mantle and posterior part of the trunk. (j) Schematic drawing of *FLRFa*-midgergic cells and nerves, *FLRFa* receptor and musculature. cs, chaetae sacs; g, gut; lc, locomotory cilia of the prototroch; m, musculature; np, neuropile; r, receptor; sc, secretory cells; sv, secretory vesicles. Colour code is indicated at the bottom of the figure plate: magenta, *FLRFamide* ((a,b,c) precursor expression, (f–i) peptide antibody staining); green, *FLRFamide* receptor; blue, musculature ((a,b,e) *tropomyosin* expression, (i) actin antibody staining); yellow, nerve tracks and cilia; grey, DAPI. Scale bar, 50  $\mu$ m.

on the ventral side at the anterior border of the mantle lobe (figure 5a–c; electronic supplementary material, figure S1b,c). The number of domains in the apical lobe varies between three and five (figure 5a,c) and each domain consists of approximately three to seven cells. The combined *in situ* hybridization with *tropomyosin* as a marker for the musculature shows that the *FLRFamide* precursor expression in the mantle lobe is adjacent to the ventral side of the trunk musculature (figure 5b). The *FLRFamide* receptor is expressed in a left and a right stripe in the trunk musculature (figure 5c,d; electronic supplementary material, figure S1a) as well as in the musculature that projects towards and surrounds the chaetae sacs (figure 5e; electronic supplementary material, figure S1a). Apart from the expression in the musculature, the receptor is also expressed in the apical lobe in a broad stripe underneath the ciliated prototroch (figure 5c,d; electronic supplementary material, figure S1a).

Since *in situ* hybridization only reveals where the peptide precursor is expressed, we used antibody stainings to visualize the nerves that secrete the active peptides. The customized *FLRFamide* antibody revealed immunoreactive longitudinal nerves that project from the apical neuropile (figure 5f,g,h,i) pairwise along the ventral (figure 5h) and dorsal (figure 5f) side into the trunk after branching off into the mantle towards the chaetae sacs at the border of the apical lobe and the mantle lobe (figure 5f,h,i). The nerves in the trunk are

branching off strongly on the ventral side (figure 5f,g) and are, at least partially, directly adjacent to the musculature (figure 5i). The neuropile shows generally strong *FLRFamide* immunoreactivity with some nerves projecting towards the apical ciliary band (figure 5f) and into the secretory cells that continue into secretory vesicles outside the apical lobe underneath the prototrochal region (figure 5g,i). The secretory cells and vesicles themselves are prone to antibody trapping so no statement can be made as to whether they in fact contain *FLRFamide* (compare figure 5g without background subtraction and figure 5f,h,i with background subtraction in secretory cells and secretory vesicles).

## 4. Discussion

### 4.1. *FLRFamide* triggers two coherent reactions via an ancient FaRP receptor

The receptor deorphanization and phylogenetic analysis shows that the *Terebratalia* *FLRFamide* receptor belongs to the ancient FaRP-GPCR group with closely related trochozoan GPCRs that include the deorphanized *P. dumerilii* FMRFamide receptor [49] and related orphan GPCRs in all major bilaterian groups. The comparable sensitivity to different peptide modifications of the larvae in the behavioural assay and the EC<sub>50</sub>

values of the receptor cell assay suggests that the larval response is triggered via the FLRFamide receptor. The expression of the FLRFamide receptor in the longitudinal trunk musculature and the musculature adjacent to the chaetae sacs in the mantle supports a direct mode of signalling, whereby FLRFamide is able to trigger the protrusion of the chaetae by inducing a muscle contraction. The expression of the receptor in a broad stripe underneath the ciliary band and the sinking of early and late larvae, independent of the presence or the absence of chaetae, also support a direct effect of FLRFamide on the ciliated cells to induce the sinking behaviour. While a direct influence of FMRFamide on the ciliary movement of trochozoan larvae has already been suggested before [23,25], the combination of this reaction with the muscular contraction observed in the *T. transversa* larval startle response consists of two different behavioural actions. In the context of the natural *T. transversa* defence behaviour, the two FLRFamide-like peptides are probably not the main neurotransmitter of their peptidergic neurons, as larvae that are desensitized to AFLRFamide/DFLRFamide are still able to show a defensive stance, although it seems like the stress level has to be increased. A possible explanation for the role of FLRFamide might be that it acts as a co-transmitter, to modify or support the signals in the different tissues that are necessary for this defence behaviour.

#### 4.2. The advantage of coherent sub-reactions during *Terebratalia transversa* defence behaviour and their control by a single peptide

Neuropeptides are considered to be ancient signalling molecules that are used in complex as well as simple nervous systems and are even present in *Trichoplax* which lacks neurons entirely [29,75]. There are a few examples of complex behaviours that involve coherent sub-reactions like insect ecdysis or feeding, which are known to deploy single neuropeptides to act on several targets as a form of master-regulator [17–21]. When a single neuropeptide is able to trigger or support the erection of chaetae and sinking, it might be involved in coordinating the startle reaction independently from a direct neuronal wiring between these two structures.

While many zoo-planktonic organisms escape potential predators by a sudden increase in velocity, some species have been observed to use passive sinking as an efficient escape strategy instead [3,76]. Passive sinking seems efficient for slow animals to escape quicker predators such as copepods that do not detect their prey by vision but by sensing water disturbance [3,11,77]. It has been described that other brachiopod and annelid larvae seem to have a similar startle behaviour as *T. transversa* [12–16]. Direct observations showed that small fish spit out *Sabellaria* larvae with their spines erected [15] and experimental data showed that *Sabellaria* larvae with long chaetae have a higher survival rate when exposed to different predators compared with younger larvae without chaetae [14]. The combination of a passive sinking behaviour while actively erecting chaetae might increase the chance to escape different predators when compared with either behaviour alone. The increased water drag due to the erected chaetae would probably also make an active fleeing inefficient.

Our results demonstrate a case in which a single receptor–ligand pair can trigger two coherent reactions that integrate

evolutionary novelties such as trochozoan chaetae [57] and ancient traits such as ciliary-based locomotion [78] into the *T. transversa* larval startle behaviour.

#### 4.3. The involvement of a specific neuropeptide in certain behavioural traits is not necessarily conserved during evolution

While the FaRP receptor–ligand pair in *P. dumerilii* and *T. transversa* is conserved, the involvement of FaRPs in trochozoan larval behaviour seems to vary. Several studies on trochozoan larvae have shown that FMRFamide-like immunoreactive nerves can be associated with different structures in a single animal and often include a combination of the apical organ, ciliary bands and the musculature [38–41,43,79]. In this context, it is also important to mention that the antibodies against FMRFamide that are commonly used in morphological studies can cross-react with other peptides ending in RFamide, even within the same species [80,81]. Inter-species comparisons and homologizations of such labelled neurons, especially across larger evolutionary distances, are therefore problematic. Only a few experimental studies exist on the effect of FMRFamide on trochozoan larvae and those focus on the regulation of the ciliary-based locomotion, which ultimately influences the vertical swimming direction [22,23,25]. These experiments on trochozoan larvae showed a taxon-specific up- or downregulation of the ciliary beating or the ciliary arrests. The ciliary beating alone, however, can be influenced by more than one peptide in a single species [23], as different neuronal circuits seem to trigger similar or opposing effects of the same effector organ and might thereby fine-tune the reaction, based on different neuronal inputs. Studies on adult trochozoans show diverse effects of FMRFamide on various muscles [32–35] and further taxon-specific functions such as osmoregulation [82], chromatophore expansion [83] or suppression of salivary gland activity [84]. Even experiments on different adult bivalve species showed species specific up- or downregulation of the heartbeat by FMRFamide [31]. The seemingly ubiquitous presence of FaRPs in trochozoan species with various taxon-specific effects and association with different tissues suggest that the FMRFamide-like peptides proved to be generally useful as a regulatory signalling system and were probably redeployed several times during trochozoan evolution, rather than having a strictly conserved role that is always associated with similar behavioural traits. The observation that the presumed orthologues in the annelid *P. dumerilii* and the brachiopod *N. novocrania* (FMRFamide and YMRFamide, respectively) do not trigger their respective defence behaviours supports the hypothesis that the involvement of a specific neuropeptide in similar behavioural traits is not necessarily conserved.

**Data accessibility.** Sequences of the here described genes and the accession numbers of publically available sequences that were used in this study are listed in the electronic supplementary material. Nucleotide sequences of the FLRFamide precursor and FLRFamide receptor are available on GenBank (MF543007, MF543008). Additional material with confocal stacks of stained larvae and videos of the influence of DFLRFamide on swimming larvae in the vertical assay is available at <https://doi.org/10.6084/m9.figshare.c.3830605.v1>. The original datasets generated and analysed during this study are available from the corresponding author on reasonable request.

**Authors' contributions.** D.T., A.H. and G.J. designed the study. D.T. and A.H. collected the animals and drafted the paper. D.T. performed



behavioural experiments, gene cloning, *in situ* hybridization and immunohistochemistry. D.T. and P.B. performed receptor deorphanization experiments. All authors commented on the drafted manuscript and contributed to the final manuscript version.

**Competing interests.** We declare we have no competing interests.

**Funding.** This research was supported by the FP7-PEOPLE-2012-ITN grant no. 317172 'NEPTUNE' and received further support by the DFG—Deutsche Forschungsgemeinschaft to G.J. (Reference no. JE 777/3-1).

## References

- Herzog Q, Laforsch C. 2013 Modality matters for the expression of inducible defenses: introducing a concept of predator modality. *BMC Biol.* **11**, 113. (doi:10.1186/1741-7007-11-113)
- Higginson AD, Ruxton GD. 2010 Adaptive changes in size and age at metamorphosis can qualitatively vary with predator type and available defenses. *Ecology* **91**, 2756–2768. (doi:10.1890/08-2269.1)
- Ohman MD. 1988 Behavioral responses of zooplankton to predation. *Bull. Mar. Sci.* **43**, 530–550.
- Vaughn D, Allen JD. 2010 The peril of the plankton. *Integr. Comp. Biol.* **50**, 552–570. (doi:10.1093/icb/icq037)
- Morgan SG. 1989 Adaptive significance of spination in estuarine crab zoeae. *Ecology* **70**, 464–482. (doi:10.2307/1937551)
- Petrusek A, Tollrian R, Schwenk K, Haas A, Laforsch C. 2009 A 'crown of thorns' is an inducible defense that protects *Daphnia* against an ancient predator. *Proc. Natl Acad. Sci. USA* **106**, 2248–2252. (doi:10.1073/pnas.0808075106)
- Vaughn D. 2007 Predator-induced morphological defenses in marine zooplankton: a larval case study. *Ecology* **88**, 1030–1039. (doi:10.1890/06-0689)
- Brandl Z. 1998 Feeding strategies of planktonic cyclopoids in lacustrine ecosystems. *J. Mar. Syst.* **15**, 87–95. (doi:10.1016/S0924-7963(97)00042-0)
- Echevarria ML, Wolfe GV, Taylor AR. 2016 Feast or flee: bioelectrical regulation of feeding and predator evasion behaviors in the planktonic alveolate *Favella* sp. (Spirotrichia). *J. Exp. Biol.* **219**, 445–456. (doi:10.1242/jeb.121871)
- Gliwicz MZ. 1986 Predation and the evolution of vertical migration in zooplankton. *Nature* **320**, 746–748. (doi:10.1038/320746a0)
- Kerfoot WC. 1978 Combat between predatory copepods and their prey: *Cyclops*, *Epischura*, and *Bosmina*. *Limnol. Oceanogr.* **23**, 1089–1102. (doi:10.4319/lo.1978.23.6.1089)
- Chuang SH. 1977 Larval development in *Discinisca* (inarticulate brachiopod). *Am. Zool.* **17**, 39–53. (doi:10.1093/icb/17.1.39)
- Nielsen C. 1991 The development of the brachiopod *Crania (Neocrania) anomala* (O. F. Müller) and its phylogenetic significance. *Acta Zool-Stockholm.* **72**, 7–28. (doi:10.1111/j.1463-6395.1991.tb00312.x)
- Pennington JT, Chia FS. 1984 Morphological and behavioral defenses of trochophore larvae of *Sabellaria cementarium* (Polychaeta) against four planktonic predators. *Biol. Bull.* **167**, 168–175. (doi:10.2307/1541345)
- Wilson DP. 1929 The larvæ of the British sabellarians. *J. Mar. Biol. Assoc. UK* **16**, 221–268. (doi:10.1017/S0025315400029787)
- Wilson DP. 1928 The Larvae of *Polydora ciliata* Johnston and *Polydora hoplura* Claparède. *J. Mar. Biol. Assoc. UK* **15**, 567–603. (doi:10.1017/S0025315400009553)
- Hewes RS, Truman JW. 1991 The roles of central and peripheral eclosion hormone release in the control of ecdysis behavior in *Manduca sexta*. *J. Comp. Physiol. A* **168**, 697–707. (doi:10.1007/BF00224359)
- Kim YJ, Žitňan D, Galizia CG, Cho KH, Adams ME. 2006 A command chemical triggers an innate behavior by sequential activation of multiple peptidergic ensembles. *Curr. Biol.* **16**, 1395–1407. (doi:10.1016/j.cub.2006.06.027)
- Taghert PH, Nitabach MN. 2012 Peptide neuromodulation in invertebrate model systems. *Neuron* **76**, 82–97. (doi:10.1016/j.neuron.2012.08.035)
- Truman JW. 2005 Hormonal control of insect ecdysis: endocrine cascades for coordinating behavior with physiology. *Vitam. Horm.* **73**, 1–30. (doi:10.1016/S0083-6729(05)73001-6)
- Dailey MJ, Bartness TJ. 2009 Appetitive and consummatory ingestive behaviors stimulated by PVH and perifornical area NPY injections. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **296**, R877–R892. (doi:10.1152/ajpregu.90568.2008)
- Braubach OR, Dickinson AJG, Evans CCE, Croll RP. 2006 Neural control of the velum in larvae of the gastropod, *Ilyanassa obsoleta*. *J. Exp. Biol.* **209**, 4676–4689. (doi:10.1242/jeb.02556)
- Conzelmann M, Offenburger SL, Asadulina A, Keller T, Münch TA, Jékely G. 2011 Neuropeptides regulate swimming depth of *Platynereis* larvae. *Proc. Natl Acad. Sci. USA* **108**, E1174–E1183. (doi:10.1073/pnas.1109085108)
- Conzelmann M, Williams EA, Tunaru S, Randel N, Shahidi R, Asadulina A, Berger J, Offermanns S, Jékely G. 2013 Conserved MIP receptor–ligand pair regulates *Platynereis* larval settlement. *Proc. Natl Acad. Sci. USA* **110**, 8224–8229. (doi:10.1073/pnas.1220285110)
- Penniman JR, Doll MK, Pires A. 2013 Neural correlates of settlement in veliger larvae of the gastropod, *Crepidula fornicata*. *Invertebr. Biol.* **132**, 14–26. (doi:10.1111/ivb.12014)
- Schmidt-Rhaesa A, Harzsch S, Puschke G. 2016 *Structure & evolution of invertebrate nervous systems*, 1st edn. New York, NY: Oxford University Press.
- D'Aniello B, Polese G, Luongo L, Scandurra A, Magliozzi L, Aria M, Pinelli C. 2016 Neuroanatomical relationships between FMRFamide-immunoreactive components of the nervus terminalis and the topology of olfactory bulbs in teleost fish. *Cell Tissue Res.* **364**, 43–57. (doi:10.1007/s00441-015-2295-4)
- Walker RJ, Papaioannou S, Holden-Dye L. 2009 A review of FMRFamide- and RFamide-like peptides in metazoa. *Invert. Neurosci.* **9**, 111–153. (doi:10.1007/s10158-010-0097-7)
- Jékely G. 2013 Global view of the evolution and diversity of metazoan neuropeptide signaling. *Proc. Natl Acad. Sci. USA* **110**, 8702–8707. (doi:10.1073/pnas.1221833110)
- Espinoza E, Carrigan M, Thomas SG, Shaw G, Edison AS. 2000 A statistical view of FMRFamide neuropeptide diversity. *Mol. Neurobiol.* **21**, 35–56. (doi:10.1385/MN:21:1-2:035)
- Painter SD, Greenberg MJ. 1982 A survey of the responses of bivalve hearts to the molluscan neuropeptide FMRFamide and to 5-hydroxytryptamine. *Biol. Bull.* **162**, 311–332. (doi:10.2307/1540986)
- Lehman HK, Greenberg MJ. 1987 The actions of FMRFamide-like peptides on visceral and somatic muscles of the snail *Helix aspersa*. *J. Exp. Biol.* **131**, 55–68.
- Krajniak KG, Greenberg MJ. 1992 The localization of FMRFamide in the nervous and somatic tissues of *Nereis virens* and its effects upon the isolated esophagus. *Comp. Biochem. Phys. C* **101**, 93–100. (doi:10.1016/0742-8413(92)90205-L)
- Muneoka Y, Saitoh H. 1986 Pharmacology of FMRFamide in *Mytilus* catch muscle. *Comp. Biochem. Phys. C* **85**, 207–214. (doi:10.1016/0742-8413(86)90075-7)
- Moulis A. 2006 The action of RFamide neuropeptides on molluscs, with special reference to the gastropods *Buccinum undatum* and *Busycon canaliculatum*. *Peptides* **27**, 1153–1165. (doi:10.1016/j.peptides.2005.07.031)
- Norris BJ, Calabrese RL. 1987 Identification of motor neurons that contain a FMRFamidelike peptide and the effects of FMRFamide on longitudinal muscle in the medicinal leech, *Hirudo medicinalis*. *J. Comp. Neurol.* **266**, 95–111. (doi:10.1002/cne.902660108)

37. Raffa RB, Bianchi CP. 1986 Further evidence for a neuromodulatory role of FMRFamide involving intracellular  $Ca^{2+}$  pools in smooth muscle of *Mytilus edulis*. *Comp. Biochem. Phys. C* **84**, 23–28. (doi:10.1016/0742-8413(86)90159-3)
38. Dyachuk V, Odintsova N. 2009 Development of the larval muscle system in the mussel *Mytilus trossulus* (Mollusca, Bivalvia). *Dev. Growth Differ.* **51**, 69–79. (doi:10.1111/j.1440-169X.2008.01081.x)
39. Helm C, Vöcking O, Kourtesis I, Hausen H. 2016 *Owenia fusiformis*—a basally branching annelid suitable for studying ancestral features of annelid neural development. *BMC Evol. Biol.* **16**, 129. (doi:10.1186/s12862-016-0690-4)
40. Hay-Schmidt A. 1995 The larval nervous system of *Polygordius lacteus* Scheinder, 1868 (Polygordiidae, Polychaeta): immunocytochemical data. *Acta Zool-Stockholm.* **76**, 121–140. (doi:10.1111/j.1463-6395.1995.tb00987.x)
41. Dyachuk V, Wanninger A, Voronezhskaya EE. 2012 Innervation of bivalve larval catch muscles by serotonergic and FMRFamidergic neurons. *Acta Biol. Hung.* **63**, 221–229. (doi:10.1556/ABiol.63.2012.Suppl.2.30)
42. Gruhl A. 2009 Serotonergic and FMRFamidergic nervous systems in gymnolaemate bryozoan larvae. *Zoomorphology* **128**, 135–156. (doi:10.1007/s00435-009-0084-x)
43. Hindinger S, Schwaha T, Wanninger A. 2013 Immunocytochemical studies reveal novel neural structures in nemertean pilidium larvae and provide evidence for incorporation of larval components into the juvenile nervous system. *Front. Zool.* **10**, 31. (doi:10.1186/1742-9994-10-31)
44. Shun'kina KV, Starunov VV, Zaitseva OV, Ostrovskii AN. 2013 Serotonin and FMRFamide immunoreactive elements in the nervous system of freshwater bryozoans (Bryozoa: Phylactolaemata). *Dokl. Biol. Sci.* **451**, 244–247. (doi:10.1134/S0012496613040108)
45. Hay-Schmidt A. 1990 Distribution of catecholamine-containing, serotonin-like and neuropeptide FMRFamide-like immunoreactive neurons and processes in the nervous system of the actinotroch larva of *Phoronis muelleri* (Phoronida). *Cell Tissue Res.* **259**, 105–118. (doi:10.1007/Bf00571435)
46. Hay-Schmidt A. 1990 Catecholamine-containing, serotonin-like, and FMRFamide-like immunoreactive neurons and processes in the nervous-system of the early actinotroch larva of *Phoronis vancouverensis* (Phoronida): distribution and development. *Can. J. Zool.* **68**, 1525–1536. (doi:10.1139/z90-226)
47. van den Pol AN. 2012 Neuropeptide transmission in brain circuits. *Neuron* **76**, 98–115. (doi:10.1016/j.neuron.2012.09.014)
48. Mirabeau O, Joly JS. 2013 Molecular evolution of peptidergic signaling systems in bilaterians. *Proc. Natl Acad. Sci. USA* **110**, E2028–E2037. (doi:10.1073/pnas.1219956110)
49. Bauknecht P, Jékely G. 2015 Large-scale combinatorial deorphanization of *Platynereis* neuropeptide GPCRs. *Cell Rep.* **12**, 684–693. (doi:10.1016/j.celrep.2015.06.052)
50. Lingueglia E, Champigny G, Lazdunski M, Barbry P. 1995 Cloning of the amiloride-sensitive FMRFamide peptide-gated sodium channel. *Nature* **378**, 730–733. (doi:10.1038/378730a0)
51. Niu YY *et al.* 2016 Exploration of the peptide recognition of an amiloride-sensitive FMRFamide peptide-gated sodium channel. *J. Biol. Chem.* **291**, 7571–7582. (doi:10.1074/jbc.M115.710251)
52. Perry SJ, Straub VA, Schofield MG, Burke JF, Benjamin PR. 2001 Neuronal expression of an FMRFamide-gated  $Na^+$  channel and its modulation by acid pH. *J. Neurosci.* **21**, 5559–5567.
53. Cazzamali G, Grimmelikhuijzen CJ. 2002 Molecular cloning and functional expression of the first insect FMRFamide receptor. *Proc. Natl Acad. Sci. USA* **99**, 12 073–12 078. (doi:10.1073/pnas.192442799)
54. Meeusen T, Mertens I, Clynen E, Baggerman G, Nichols R, Nachman RJ, Huybrechts R, De Loof A, Schoofs L. 2002 Identification in *Drosophila melanogaster* of the invertebrate G protein-coupled FMRFamide receptor. *Proc. Natl Acad. Sci. USA* **99**, 15 363–15 368. (doi:10.1073/pnas.252339599)
55. Hauser F, Grimmelikhuijzen CJ. 2014 Evolution of the AKH/corazonin/ACP/GnRH receptor superfamily and their ligands in the Protostomia. *Gen. Comp. Endocrinol.* **209**, 35–49. (doi:10.1016/j.ygcen.2014.07.009)
56. Froomincx L, Van Rompay L, Temmerman L, Van Sinay E, Beets I, Janssen T, Husson SJ, Schoofs L. 2012 Neuropeptide GPCRs in *C. elegans*. *Front. Endocrinol. (Lausanne)* **3**, 167. (doi:10.3389/fendo.2012.00167)
57. Schiemann SM, Martín-Durán JM, Børve A, Vellutini BC, Passamaneck YJ, Hejnal A. 2017 Clustered brachiopod Hox genes are not expressed collinearly and are associated with lophotrochozoan novelties. *Proc. Natl Acad. Sci. USA* **114**, E1913–E1922. (doi:10.1073/pnas.1614501114)
58. Martín-Durán JM, Passamaneck YJ, Martindale MQ, Hejnal A. 2016 The developmental basis for the recurrent evolution of deuterostomy and protostomy. *Nat. Ecol. Evol.* **1**, 0005. (doi:10.1038/s41559-016-0005)
59. Vellutini BC, Hejnal A. 2016 Expression of segment polarity genes in brachiopods supports a non-segmental ancestral role of *engrailed* for bilaterians. *Sci. Rep.* **6**, 32387. (doi:10.1038/srep32387)
60. Temereva EN, Tsitrin EB. 2015 Modern data on the innervation of the lophophore in *Lingula anatina* (Brachiopoda) support the monophyly of the lophophorates. *PLoS ONE* **10**, e0123040. (doi:10.1371/journal.pone.0123040)
61. Hay-Schmidt A. 1992 Ultrastructure and immunocytochemistry of the nervous system of the larvae of *Lingula anatina* and *Glottidia* sp. (Brachiopoda). *Zoomorphology* **112**, 189–205. (doi:10.1007/Bf01632817)
62. Santagata S. 2011 Evaluating neurophylogenetic patterns in the larval nervous systems of brachiopods and their evolutionary significance to other bilaterian phyla. *J. Morphol.* **272**, 1153–1169. (doi:10.1002/jmor.10975)
63. Altenburger A, Wanninger A. 2010 Neuromuscular development in *Novocrania anomala*: evidence for the presence of serotonin and a spiralian-like apical organ in lecithotrophic brachiopod larvae. *Evol. Dev.* **12**, 16–24. (doi:10.1111/j.1525-142X.2009.00387.x)
64. Santagata S, Resh C, Hejnal A, Martindale MQ, Passamaneck YJ. 2012 Development of the larval anterior neurogenic domains of *Terebratalia transversa* (Brachiopoda) provides insights into the diversification of larval apical organs and the spiralian nervous system. *Evodevo* **3**, 3. (10.1186/2041-9139-3-3)
65. Stricker SA, Reed CG. 1985 The ontogeny of shell secretion in *Terebratalia transversa* (Brachiopoda, Articulata) 1. Development of the mantle. *J. Morphol.* **183**, 233–250. (doi:10.1002/jmor.1051830302)
66. Frickey T, Lupas A. 2004 CLANS: a Java application for visualizing protein families based on pairwise similarity. *Bioinformatics* **20**, 3702–3704. (doi:10.1093/bioinformatics/bth444)
67. Larkin MA *et al.* 2007 Clustal W and Clustal X version 2.0. *Bioinformatics* **23**, 2947–2948. (doi:10.1093/bioinformatics/btm404)
68. Keane TM, Creevey CJ, Pentony MM, Naughton TJ, McInerney JO. 2006 Assessment of methods for amino acid matrix selection and their use on empirical data shows that ad hoc assumptions for choice of matrix are not justified. *BMC Evol. Biol.* **6**, 29. (doi:10.1186/1471-2148-6-29)
69. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010 New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321. (doi:10.1093/sysbio/syq010)
70. Passamaneck YJ, Hejnal A, Martindale MQ. 2015 Mesodermal gene expression during the embryonic and larval development of the articulate brachiopod *Terebratalia transversa*. *Evodevo* **6**, 10. (doi:10.1186/s13227-015-0004-8)
71. Sinigaglia C, Thiel D, Hejnal A, Houliston E, Leclère L. In press. A safer, urea-based in situ hybridization method improves detection of gene expression in diverse animal species. *bioRxiv* (doi:10.1101/133470)
72. Hejnal A. 2008 *In situ* protocol for embryos and juveniles of *Convolutriloba longifissura*. *Protoc. Exch.*, 201. (doi:10.1038/nprot.2008.201)
73. Jékely G, Arendt D. 2007 Cellular resolution expression profiling using confocal detection of NBT/BCIP precipitate by reflection microscopy. *Biotechniques* **42**, 751–755. (doi:10.2144/000112462)
74. Conzelmann M, Jékely G. 2012 Antibodies against conserved amidated neuropeptide epitopes enrich the comparative neurobiology toolbox. *Evodevo* **3**, 23. (doi:10.1186/2041-9139-3-23)
75. Nikitin M. 2015 Bioinformatic prediction of *Trichoplax adhaerens* regulatory peptides. *Gen. Comp. Endocr.* **212**, 145–155. (doi:10.1016/j.ygcen.2014.03.049)
76. Burdick DS, Hartline DK, Lenz PH. 2007 Escape strategies in co-occurring calanoid copepods. *Limnol. Oceanogr.* **52**, 2373–2385. (doi:10.4319/lo.2007.52.6.2373)

77. Williamson CE. 1983 Behavioral interactions between a cyclopoid copepod predator and its prey. *J. Plankton Res.* **5**, 701–711. (doi:10.1093/plankt/5.5.701)
78. Jékely G. 2011 Origin and early evolution of neural circuits for the control of ciliary locomotion. *Proc. R. Soc. B* **278**, 914–922. (doi:10.1098/rspb.2010.2027)
79. Nezlin LP. 2010 The golden age of comparative morphology: laser scanning microscopy and neurogenesis in trochophore animals. *J. Dev. Biol.* **41**, 381–390. (doi:10.1134/S1062360410060056)
80. Vilim FS *et al.* 2010 Distinct mechanisms produce functionally complementary actions of neuropeptides that are structurally related but derived from different precursors. *J. Neurosci.* **30**, 131–147. (doi:10.1523/Jneurosci.3282-09.2010)
81. Peymen K, Watteyne J, Frooninckx L, Schoofs L, Beets I. 2014 The FMRFamide-like peptide family in nematodes. *Front. Endocrinol.* **5**, 90. (doi:10.3389/fendo.2014.00090)
82. Salzet M, Bulet P, Watzet C, Malecha J. 1994 FMRFamide-related peptides in the sex segmental ganglia of the Pharyngobdellid leech *Erpobdella octoculata*: identification and involvement in the control of hydric balance. *Eur. J. Biochem.* **221**, 269–275. (doi:10.1111/j.1432-1033.1994.tb18738.x)
83. Loi PK, Tublitz N. 1997 Molecular analysis of FMRFamide- and FMRFamide-related peptides (FaRPS) in the cuttlefish *Sepia officinalis*. *J. Exp. Biol.* **200**, 1483–1489.
84. Bulloch AGM, Price DA, Murphy AD, Lee TD, Bowes HN. 1988 FMRFamide peptides in *Helisoma*: identification and physiological actions at a peripheral synapse. *J. Neurosci.* **8**, 3459–3469.