

Evaluation of Antioxidant Properties in Thirteen Fijian Medicinal Plants Used in Alzheimer's Disease and Related Illness

Ranish Nitesh Chand^{1,2}, Romila Devi Gopalan^{1*} and Ketan Christi¹**Ranish Nitesh Chand^{1,2},
Romila Devi Gopalan^{1*} and
Ketan Christi¹**¹School of Biological and Chemical Sciences, Faculty of Science, Technology and Environment, The University of the South Pacific, Suva, FIJI.²Senior Education Officer Research, Ministry of Education Heritage and Arts, FIJI.**Correspondence****Romila D Gopalan**, School of Biological and Chemical Sciences, Faculty of Science, Technology and Environment, The University of the South Pacific, Suva, FIJI.

Phone no: +679 32 32446

E-mail: devi_rd@usp.ac.fj**History**

- Submission Date: 18-04-2017;
- Review completed: 27-05-2017;
- Accepted Date: 15-06-2017.

DOI : 10.5530/fra.2018.1.3**Article Available online**<http://www.antiox.org/v8/i1>**Copyright**

© 2018 Phcog.Net. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

**ABSTRACT**

Objective: The present study aims to evaluate antioxidant properties of decoction and ethanol extracts of Fijian medicinal plants using 2,2-diphenyl-1-picryl-hydrazyl (DPPH) scavenging assay. **Method:** Thirteen plant species belonging to Melastomataceae, Asteraceae, Apiaceae, Rutaceae, Goodeniaceae, Loganiaceae, Araliaceae, Solanaceae, Polygonaceae, Zingiberaceae and Anacardiaceae families were tested at 0.2 mg/mL, 0.4 mg/mL, 0.6 mg/mL, 0.8 mg/mL, 1.0 mg/mL, 1.5 mg/mL and 2.0 mg/mL concentrations for antioxidant properties. The antioxidant capabilities were compared with ascorbic acid standard. **Results and Discussions:** Among the decoction and ethanol extracts tested, all plants showed DPPH scavenging activity. The most potent antioxidant activity was seen in *C. hirta* with an IC₅₀ value of 0.64 mg/mL. The activity of *C. hirta* was twofold more potent than the standard ascorbic acid (IC₅₀ = 1.33 mg/mL) indicating that polar extracts of *C. hirta* contains compounds with relatively better antioxidant properties than ascorbic acid. **Conclusion:** The plant extracts used in this study were crude extracts, as it is envisaged that if the phytochemicals were isolated and purified from these plants, more prominent results could be expected. These plants could prove leads to safer and better candidates for the future selection of antioxidant.

Key words: DPPH, Reactive Oxygen Species, Radical scavenging activity, Antioxidants, *C. hirta*, Ethanolic extracts, Decoction.

Key Message: Polar extracts from medicinal plants have antioxidant properties apart from its traditional use. These plants can be investigated to understand the full potential of these plants.

INTRODUCTION

Oxygen is essential for respiration; however, under certain conditions it can have a negative effect on our health due to the formation of reactive oxygen species (ROS) or free radicals. ROS are formed under normal physiological conditions but become deleterious when not being eliminated by the endogenous antioxidant systems. The existence of ROS leads to oxidative stress¹ and is often cytotoxic and gives rise to tissue injuries.² ROS can attack lipids and proteins, destroy membranes, damage DNA, and lead to chromosomal damage.¹ Thus, ROS can contribute to development of numerous diseases, such as cardiovascular diseases,³ neural disorders,⁴ mild cognitive impairment,⁵ Alzheimer's disease,⁶ alcohol induced liver disease,⁷ Parkinson's disease,⁸ ulcerative colitis,⁹ ageing,¹⁰ atherosclerosis,¹¹ and cancer.¹² For these reasons, antioxidants are of interest for the treatment of many kinds of cellular degeneration including Alzheimer's disease.¹³⁻¹⁴

Numerous recent studies have shown that medicinal plants are an important source of antioxidants.¹⁵⁻¹⁹ Plants contain a wide variety of free radical scavenging molecules, such as flavonoids, anthocyanins, carotenoids, dietary glutathione, alkaloids, tannins,

saponins, steroids, terpenoids and rotenoids.¹⁹⁻²¹ Therefore, medicinal plants becomes the target to explore cost effective new antioxidants with fewer side effects. In this study, medicinal plants used by traditional healers in Fiji to treat Alzheimer's disease and related illnesses were screened for their antioxidant potential.

METHODOLOGY**Selection of plants**

Thirteen plant species listed in Table 1 were selected for antioxidant activity based on the traditional knowledge of using polar extracts (aqueous and ethanolic extracts) were used in this study. The fresh plant specimens were collected from their natural habitats from Vanua Levu and Viti Levu in Fiji.

Extraction

The plant samples were thoroughly washed with tap water to remove dust and were dried according to the method described by Mohd *et al.*²² in a convection oven at 40 °C for 48 h until there was no change in weight. The dried plant parts were ground separately

Cite this article: Chand RN, Gopalan RD and Christi K. Evaluation of Antioxidant Properties in Thirteen Fijian Medicinal Plants Used in Alzheimer's Disease and Related Illness. Free Radicals and Antioxidants. 2017;8(1):11-7.

Table 1: Fijian Medicinal plants used in this study

No	Scientific name	Family	Fijian Name
1.	<i>Clidemia hirta</i>	Melastomataceae	Bona bulmakau
2.	<i>Bidens pilosa</i>	Asteraceae	Batimadramadra
3.	<i>Centella asiatica</i>	Apiaceae	Totondro
4.	<i>Ageratum conyzoides</i>	Asteraceae	Botebotokeoro
5.	<i>Micromelum minutum</i>	Rutaceae	Qiqilia kau
6.	<i>Scaevola koenigii</i>	Goodeniaceae	Vevedu
7.	<i>Neuburgia collina</i>	Loganiaceae	Qiqilia wawa
8.	<i>Polycias fruticosa</i>	Araliaceae	Danidani
9.	<i>Persicaria perfoliata</i>	Polygonaceae	Wabosucu
10.	<i>Zingiber officinalis</i>	Zingiberaceae	Layalaya
11.	<i>Eudia hortensis</i>	Rutaceae	Udi
12.	<i>Physalis angulata</i>	Solanaceae	Cevucevu
13.	<i>Spondias pinnata</i>	Anacardiaceae	Stone breaker

into powder using an electric blender and stored in an air tight container. The samples were kept in dark until ready for extraction.

Ethanolic Extraction

12 g of powdered plant sample was extracted with 100 mL of 96% ethanol using soxhlet extraction unit at 60 °C for 18-20 hours. Filtered extract from the soxhlet extractor was concentrated using roto evaporator to yield a dark semi-solid residue. The extract was stored in a refrigerator at 4 °C in air-tight bottles until further use.

Aqueous Extract (Decoction)

10 g of powdered plant material was added to 100 mL of deionised water in a beaker and boiled for 10 minutes. The decoction was cooled and filtered through Whatman No. 1 filter paper. The filtrate was further concentrated by freeze-drying and stored at 4 °C in a sealed bottle prior to analysis.

Anti-oxidative Analysis using DPPH radical scavenging activity

Free radical scavenging ability of the samples, based on the scavenging activity of DPPH free radical, was evaluated using the procedure described previously by Blois²³ with slight modification. Different dilutions (0.2, 0.4, 0.6, 0.8, 1.0, 1.5 and 2.0 mg/mL) of 50 µL plant extract (ethanolic/decoction) were added to 5 mL of 0.002% ethanolic solution of DPPH in a vial covered with aluminium foil. The samples containing DPPH was incubated in the dark for 30 minutes. The absorbance was then measured at 517 nm using a double beam UV spectrophotometer (UV Winlab Version 2.85.04, Perkin Elmer).²⁴ The absorbance of plant extracts dissolved in ethanol (without DPPH) was measured and subtracted from the absorbance of the DPPH for colour correction. Negative control contained all components apart from the plant extracts, whereas positive control ascorbic acid was used in the assay as standard. Ethanol was used as a blank. The assay was repeated in triplicates and percentage inhibition was calculated using the formula:

$$\% \text{ scavenging effect} = \frac{\text{control absorbance} - \text{sample absorbance}}{\text{control absorbance}} \times 100$$

Control absorbance = absorbance of the control reaction containing all reagents except the plant extracts.

Sample absorbance = absorbance of the plant extracts.

The interpretations of the results were done using the IC₅₀ values (concentration of substrate that causes 50% loss of the DPPH activity) as well as the % radical scavenging activity. A lower IC₅₀ value corresponds to a larger scavenging power and a DPPH radical scavenging activity of the crude extract of less than 20 mg/mL was considered significant.

Statistical Analysis

The assays were conducted in triplicates. The data were expressed as means ± standard error (SE). IC₅₀ values were calculated in MS Excel linear regression analysis among the percent inhibition against the extract concentrations.

RESULTS AND DISCUSSION

Free radical scavenging activity in medicinal plants has attracted a lot of interest recently. It is assumed that high antioxidant properties of the plants is linked to effective treatment of illnesses. In this study, thirteen medicinal plants were evaluated for antioxidant properties using DPPH assay. The DPPH• is stable organic nitrogen centered free radical with a dark purple color which becomes colorless when it reacts with antioxidants to form non-radicals.²⁵ Thus, in the analysis, lower absorbance of the reaction mixture indicated the higher free radical scavenging activity.

DPPH radical scavenging activity using water (decoction) extraction of plant samples

The water extracts of all plant species showed free radical scavenging activity, shown in Table 2. Scavenging activity increases with increasing concentration of crude extracts. The most potent activity was shown by *C. hirta*, which displayed an activity of 94.98% to 98.23% between concentrations of 0.6 to 2.0 mg/mL, followed by *B. pilosa* and *C. asiatica* with 22.84% and 20.48% respectively at 2.0 mg/mL. The remaining plants displayed the scavenging activity lower than 20% at 2.0 mg/mL.

DPPH radical scavenging activity using ethanolic extracts of the plant samples

Ethanolic extracts of all plants demonstrated free radical scavenging activity as shown in Table 3. The scavenging activity increases with the increase in the concentration. Highest scavenging activity was displayed at 2.0 mg/mL for majority of plants. All plant species demonstrated free radical scavenging activity higher than that of the water extract except for *C. hirta*. The scavenging activity of *C. hirta* (72.19%) was the highest amongst the plant tested for ethanol extract but lower than that of the water extract (97.93%) at 2 mg/mL. Therefore, *C. hirta* had the most promising result for antioxidant properties in water and ethanol extracts.

Inhibition of DPPH at 50% concentration (IC₅₀)

Half maximal inhibitory concentration (IC₅₀) refers to the concentration of the plant extract where the response is reduced by half and thus determines the effectiveness of the antioxidant. Those plant extracts with higher free radical scavenging activity had lower IC₅₀ value; in this case it was considered lower than 20 mg/mL to select the effective sample. IC₅₀ values of the thirteen Fijian plants used in this study are given in Tables 2 and 3. For the water extract, the most potent antioxidant exhibiting plants were *C. hirta*, *B. pilosa* and *C. asiatica* with IC₅₀ values of 0.64, 18.24 and 18.24 mg/mL respectively. For the ethanol extracts, five most potent plant species are *C. hirta* (1.45 mg/mL), *A. conyzoides* (13.22), *B. pilosa* (13.79 mg/mL), *C. asiatica* (15.45 mg/mL) and *S. koenigii* (17.55).

The DPPH radical scavenging abilities of the extracts were significantly comparable to those of ascorbic acid (standard), showing that some extracts have the hydrogen donating ability and could serve as free radical inhibitors or scavengers, acting possibly as primary antioxidants.

Table 2: Percentage free radical scavenging activity of water extracts (decoction) of plants.

Conc/mg/mL	<i>C. hirta</i>	<i>B. pilosa</i>	<i>C. asiatica</i>	<i>A. conyzoides</i>	<i>M. minutum</i>	<i>S. koenigii</i>	<i>N. collina</i>
0.2	43.11 ± 4.81	5.55 ± 0.12	2.05 ± 1.12	2.44 ± 6.11	3.13 ± 0.76	0.39 ± 7.06	1.95 ± 1.28
0.4	73.72 ± 1.04	7.47 ± 0.07	2.46 ± 1.10	4.53 ± 4.07	5.02 ± 0.51	1.74 ± 6.07	4.03 ± 0.91
0.6	94.98 ± 1.23	10.36 ± 1.10	5.90 ± 1.39	7.17 ± 4.50	5.42 ± 0.44	3.71 ± 6.39	4.24 ± 0.06
0.8	94.98 ± 0.53	11.67 ± 0.41	7.66 ± 1.46	8.30 ± 4.21	6.79 ± 0.34	5.77 ± 6.18	6.09 ± 0.99
1.0	97.05 ± 0.42	11.91 ± 0.11	8.89 ± 0.97	9.30 ± 3.79	8.62 ± 0.79	6.89 ± 7.75	5.58 ± 0.44
1.5	98.23 ± 0.13	17.63 ± 0.57	14.91 ± 1.25	12.18 ± 4.26	10.52 ± 0.78	10.79 ± 3.98	8.47 ± 0.83
2.0	97.93 ± 0.36	22.84 ± 0.44	20.48 ± 1.36	16.68 ± 4.35	12.21 ± 0.65	10.07 ± 6.76	9.89 ± 1.03
IC ₅₀	0.64	18.24	18.24	23.25	32.79	26.95	40.40

Conc/mg/mL	<i>P. fruticosa</i>	<i>P. angulata</i>	<i>S. pinnata</i>	<i>P. perfoliata</i>	<i>Z. officinalis</i>	<i>E. hortensis</i>	Ascorbic Acid
0.2	4.34 ± 2.43	1.25 ± 1.03	0.68 ± 1.21	4.69 ± 1.92	2.48 ± 2.39	3.68 ± 2.54	34.33 ± 0.54
0.4	4.46 ± 0.76	2.74 ± 1.54	0.89 ± 2.59	8.24 ± 3.93	2.77 ± 0.73	5.77 ± 1.87	52.70 ± 0.98
0.6	5.03 ± 1.97	3.40 ± 1.22	2.75 ± 1.59	3.51 ± 1.45	0.90 ± 2.52	7.43 ± 2.64	77.80 ± 0.83
0.8	6.24 ± 0.71	5.04 ± 2.43	3.43 ± 2.12	2.41 ± 1.14	1.02 ± 1.22	8.33 ± 3.21	83.78 ± 0.77
1.0	6.93 ± 1.85	6.36 ± 2.66	3.84 ± 2.97	3.02 ± 1.22	0.49 ± 1.95	10.27 ± 2.43	87.22 ± 0.22
1.5	7.09 ± 0.64	8.01 ± 2.71	5.20 ± 0.43	1.81 ± 1.25	-6.27 ± 1.36	13.44 ± 4.68	88.25 ± 0.31
2.0	7.22 ± 0.77	11.41 ± 1.23	7.48 ± 2.87	4.83 ± 1.35	-7.71 ± 2.03	15.25 ± 3.25	88.78 ± 0.63
IC ₅₀	82.15	32.36	95.45	98.27	25.62	45.72	1.33

Each value represents the mean ± SE

n = 3

Table 3: Percentage free radical scavenging activity of ethanolic extracts of plants.

Conc.mg/mL	<i>C. hirta</i>	<i>B. pilosa</i>	<i>C. asiatica</i>	<i>A. conyzoides</i>	<i>M. minutum</i>	<i>S. koenigii</i>	<i>N. collina</i>
0.2	46.51 ± 14.70	11.38 ± 4.53	13.79 ± 0.82	7.84 ± 0.76	6.31 ± 1.25	11.45 ± 2.39	17.82 ± 2.47
0.4	54.69 ± 0.91	19.02 ± 8.46	17.72 ± 0.05	11.30 ± 4.07	9.12 ± 1.90	14.33 ± 0.93	20.93 ± 2.35
0.6	57.32 ± 0.93	8.50 ± 13.51	18.45 ± 0.23	16.89 ± 0.68	10.51 ± 1.68	15.90 ± 3.95	22.20 ± 2.92
0.8	58.63 ± 3.51	17.03 ± 5.98	22.43 ± 0.12	15.78 ± 2.49	14.77 ± 6.66	18.15 ± 2.33	25.25 ± 4.94
1.0	60.73 ± 1.35	20.59 ± 3.23	23.22 ± 0.36	18.72 ± 6.83	12.17 ± 2.11	18.26 ± 1.61	20.58 ± 3.52
1.5	67.23 ± 1.00	16.34 ± 6.08	25.30 ± 2.36	24.15 ± 5.25	20.15 ± 5.98	21.76 ± 0.93	20.58 ± 1.34
2.0	72.19 ± 0.34	38.87 ± 3.96	30.30 ± 1.91	31.09 ± 1.14	15.92 ± 6.07	27.61 ± 0.40	21.58 ± 2.68
IC ₅₀	1.45	13.79	15.45	13.22	23.87	17.55	93.76

Conc. mg/mL	<i>P. fruticosa</i>	<i>P. angulata</i>	<i>S. pinnata</i>	<i>P. perfoliata</i>	<i>Z. officinalis</i>	<i>E. hortensis</i>	Ascorbic acid
0.2	28.53 ± 0.66	8.78 ± 2.41	0.99 ± 2.14	6.53 ± 6.06	11.64 ± 0.81	12.45 ± 2.56	34.33 ± 0.54
0.4	28.11 ± 1.23	9.49 ± 3.03	3.35 ± 1.33	14.78 ± 2.32	12.92 ± 1.07	15.23 ± 2.33	52.70 ± 0.98
0.6	21.50 ± 4.69	10.40 ± 1.62	5.44 ± 3.09	10.09 ± 3.15	14.08 ± 3.22	16.55 ± 1.35	77.80 ± 0.83
0.8	18.87 ± 0.37	11.53 ± 0.57	6.06 ± 2.64	8.58 ± 8.08	13.56 ± 16.89	18.03 ± 3.34	83.78 ± 0.77
1.0	17.70 ± 3.08	12.11 ± 0.81	7.30 ± 4.90	6.54 ± 2.04	11.79 ± 0.52	18.57 ± 0.20	87.22 ± 0.22
1.5	9.37 ± 1.51	14.56 ± 0.18	10.11 ± 3.76	7.36 ± 2.43	14.38 ± 0.45	17.79 ± 4.13	88.25 ± 0.31
2.0	14.20 ± 12.08	17.75 ± 2.04	10.22 ± 2.54	6.50 ± 6.73	13.10 ± 3.76	18.70 ± 2.60	88.78 ± 0.63
IC ₅₀	-6.05	31.63	-58.69	2110.45	39.95	32.47	1.33

Each value represents the mean ± SE

n = 3




Figure 1: Graph of IC₅₀ values of free radical scavenging activity of decoction and ethanolic extracts.




Figure 2: Graph of % free radical scavenging activity of ethanolic extract and decoction of *N. Collina*.




Figure 3: Graph showing plants with reversible free radical scavenging activity.

Among different fractions of lyophilized samples of decoction and ethanolic extract, *C. hirta* belonging to melastomataceae family; showed highest activity which can be attributed to its high antioxidant potential. *C. hirta* had the IC₅₀ value of 0.64 mg/mL for decoction while 1.45 mg/mL for extraction with ethanol. On the other hand, the IC₅₀ value of ascorbic acid was 1.33 mg/mL. This shows that there were more polar constituents in *C. hirta* which are responsible for the antioxidant activity.

Comparison of IC₅₀ for decoction and ethanolic extract

Higher antioxidant activity was noted for ethanolic extract than decoction as illustrated in Figure 1. The DPPH free radical scavenging potential of the tested water extract fractions were in a descending order of *C. hirta* > *B. pilosa* > *C. asiatica* > *A. conyzoides* > *S. koenigii* > *M. minutum* > *E. hortensis* > *P. angulata* > *N. collina* > *S. pinnata* > *P. fruticosa* > *P. perfoliata* > *Z. officinalis*. *C. hirta* had similar IC₅₀ to positive control (ascorbic acid) indicating a high concentration of antioxidant components. The antioxidant potential for ethanolic extracts are almost 2 fold higher than that obtained for decoction.

N. collina belonging to Loganiaceae family displayed higher free radical activity with ethanol extract compared to the decoction as shown in Figure 2 in the current study. At lower concentrations, ethanolic extract showed 5-8 fold higher free radical activity, which decreases yet higher activity at higher concentrations. This is consistent with the study carried out by Sudhira *et al.*²⁶ with *Strychnos colubrina* also from Loganiaceae family, which showed more potent activity with the methanol extract than the aqueous extract. The study also demonstrated the presence of alkaloids, phenols, glycosides, steroids, and flavonoids in *S. colubrina*, which is thought to be responsible for antioxidant and antimicrobial properties against Gram-positive and Gram-negative bacteria.²⁶ Similarly, *B. pilosa* and *A. conyzoides* both belong to Asteraceae family had shown similar antioxidant activity in this study. According to Shukla *et al.*,²⁷ a group of plants belonging to the family Asteraceae contained phenols, sterols and flavonoids as active constituents. These phytochemicals may also be present in *B. pilosa* and *A. conyzoides* that are responsible for anti-oxidative property.

Additionally, the methanolic extract of *M. edule* and *M. umbellatum* belonging to Melastomataceae family also showed remarkable scavenging capacity on free radicals in a concentration dependent manner.²⁸⁻²⁹ Rutin, a compound isolated from leaf extract of *M. edule* had strong reductive capacity on DPPH radicals with an IC₅₀ value of 17.06 µg/mL and ferric ion chelation (IC₅₀ value 17.29 µg/mL) potential. The methanolic extract of *M. umbellatum* was found with high content of phenol. Rutin and phenol involved in the antioxidant activity might also be present in *C. hirta* (Melastomataceae) as it exhibits similar antioxidant properties and belongs to the same family.

Other studies have shown the presence of glycosides, carbohydrates, tannins and coumarins present in *Dissotis longipetala* (Melastomataceae) in the bark extract.³⁰ The authors postulated high antioxidant and reducing power to high total phenol and low total flavonoids contents. The results obtained for *C. hirta* in our study could possibly have high levels of anti-oxidant compounds, which might be the reason behind its traditional uses in the treatment of a range of illnesses (headache, thrush, haemostasis and skin infections) in Fiji.

Reversible Inhibition

Most plants showed activity in a concentration dependent inhibition; however, there were a few plant extracts that showed decreased percentage activity with the increase in the concentration. This could be due to (1) the decreased solubility of the plant extract and (2) the turbidity or intense colour of the plant extract with higher concentration that affected absor-

bance. It was noted that *P. perfoliata* and *Z. officinalis* inhibited reversibly upon dilution with water and ethanolic extracts as shown in Figure 3. Water extracts of *Z. officinalis* showed a 5 fold decrease in free radical scavenging activity at 1.0 mg/mL when compared to the activity at 0.2 mg/mL. Similarly, DPPH free radical scavenging activity for ethanolic extract of *P. fruticosa* showed a 2 fold decrease in inhibition at 2 mg/mL. *P. fruticosa*, which belongs to the family of Araliaceae showed an IC₅₀ value of 82.15 mg/mL DPPH free radical activity in the water extract. Similar result was reported for *Cussonia arborea*, a plant that belongs to the same family as *P. fruticosa* (with IC₅₀ value 91.93 mg/mL in methanol extract)³⁰ indicating poor antioxidant properties.

Activity with the most diluted solution

In decoction extracts, *C. hirta* showed the maximum free radical scavenging potential (43.11%) at a concentration of 0.2 mg/mL, indicating the potency greater than ascorbic acid (34.33%). Decoction extracts of *B. pilosa*, *P. perfoliata*, *P. fruticosa*, *M. minutum*, *Z. officinalis*, *A. conyzoides*, *C. asiatica* and *N. collina* displayed antioxidant properties between 1% to 10% while *S. pinnata* and *S. koenigii* showed low potency (activity below 1%) indicating *C. hirta* may contain compounds with better scavenging ROS species than ascorbic acid.

Similarly, ethanolic extracts produced more potent radical scavenging activity at 0.2 mg/mL compared to that of decoction extract with several plant species displaying activity above 10%. The most active plant species at the concentration of 0.2 mg/mL were from *C. hirta* (46.51%), *P. fruticosa* (28.53%), *N. collina* (17.82%), *C. asiatica* (13.79%), *E. hortensis* (12.45%), *Z. officinalis* (11.64%), *S. koenigii* (11.45%) and *B. pilosa* (11.38%). All other plant extracts had produced an activity lower than 10% but higher than 1% unlike with decoction where two of the plant extracts displayed activity of less than 1%.

CONCLUSION

This is the first reported study for antioxidant activity of the thirteen plants used except for *C. asiatica*. The results of the present work showed that both water and ethanolic extracts of all plants demonstrated DPPH activity in concentration dependent manner except for *P. perfoliata*, *Z. officinalis* and *P. fruticosa*. The majority of plants showed higher DPPH radical scavenging activity of their ethanolic extracts compared to their water extracts. The higher antioxidant potential of the ethanol extract may suggest that ethanol is able to extract more active compounds with possible DPPH radical scavenging activity than water. The most potent antioxidant extract was from *C. hirta* which had 97.93% free radical activity with water extract while 72.19% with ethanol extract at 2.0 mg/mL suggesting that *C. hirta* contains efficient antioxidant compounds. All the plants tested in this study may offer potential for the treatment of different diseases with their anti-oxidant properties, thus can be promising candidates for more detailed *in-vitro* and *in-vivo* studies.

ACKNOWLEDGEMENT

Authors acknowledge the research grant from Faculty of Science, Technology and Environment, The University of the South Pacific.

CONFLICT OF INTEREST

Authors hereby confirm no conflict of interest associated with publication of this research work.


ABBREVIATION USED

DPPH: 2,2-diphenyl-1-picryl-hydrazyl; IC₅₀: Half maximal inhibitory concentration; ROS: Reactive Oxygen Species.

REFERENCES

- Chandra S., Dave R. *In-vitro* models for antioxidant activity evaluation and some medicinal plants possessing antioxidant properties: An overview. *Afr J Microbiol Res.* 2009;3(13):981-96.
- Jainu M, Devi CS. *In vitro* and *in vivo* evaluation of free radical scavenging potential of *Cissus quadrangularis*. *Afr J Biomed Res.* 2005;8:95-9.
- Singh U, Jialal I. Oxidative stress and atherosclerosis. *Pathophysiology.* 2006;13(3):129-42.
- Sas K, Robotka H, Toldi J, Vécsei L. Mitochondria, metabolic disturbances, oxidative stress and the kynurenine system, with focus on neurodegenerative disorders. *Journal of the Neurological Sciences.* 2007;257(1):221-39.
- Guidi I, Galimberti D, Lonati S, Novembrino C, Bamonti F, *et al.* Oxidative imbalance in patients with mild cognitive impairment and Alzheimer's disease. *Neurobiology of Aging.* 2006;27(2):262-9.
- Smith MA, Rottkamp CA, Nunomura A, Raina AK, Perry G. Oxidative stress in Alzheimer's disease. *Biochim Biophys Acta.* 2000;1502:139-44.
- Arteel GE. Oxidants and antioxidants in alcohol-induced liver disease. *Gastroenterology.* 2003;124(3):778-90.
- Bolton JL, Trush MA, Penning TM, Dryhurst G, Monks TJ. Role of quinones in toxicology. *Chem Res Toxicol.* 2000;13:135-60.
- Ramakrishna BS, Varghese R, Jayakumar S, Mathan M, Balasubramanian KA. Circulating antioxidants in ulcerative colitis and their relationship to disease severity and activity. *J Gastroenterol Hepatol.* 1997;12:490-4.
- Hyun DH, Hernandez JO, Mattson MP, de Cabo R. The plasma membrane redox system in aging. *Ageing Research Reviews.* 2006;5(2):209-20.
- Upston JM, Kritharides L, Stocker R. The role of vitamin E in atherosclerosis. *Prog Lipid Res.* 2003;42:405-22.
- Kinnula VL, Crapo JD. Superoxide dismutases in malignant cells and human tumors. *Free Radic Biol Med.* 2004;36:718-44.
- Geber M, Boutron-Ruault MC, Hercberg S, Riboli E, Scalbert A, *et al.* Food and cancer: state of the art about the protective effect of fruits and vegetables. *Bull Cancer.* 2002;89:293-312.
- Frank B, Gupta S. A review of antioxidants and Alzheimer's disease. *Annals of Clinical Psychiatry.* 2005;17(4):269-86.
- Rice-Evans, C. Flavonoids and isoflavones: absorption, metabolism and bioactivity. *Free Radic Biol Med.* 2004;36:827-8.
- Sikarwar MS, Hui BJ, Subramaniam K, Valeisamy BD, Yean LK, Balaj, K. Antioxidant activity of *Artocarpus altilis* (Parkinson) Fosberg leaves. *Free Rad Antiox.* 2014;4:33-9.
- Mandade R, Sreenivas SA, Choudhury A. Radical scavenging and antioxidant activity of *Carthamus tinctorius* extracts. *Free Radicals and Antioxidants.* 2011;1(3):87-93.
- Chothani DL, Mishra SH. *In vitro* anti-oxidant activity of *Ruellia tuberosa* root extracts. *Free Radicals and Antioxidants.* 2012;2(4):38-44.
- Kaliyappan M, Nallasamy N, Thirumurthy SPK. *In vitro* anti-oxidant activity in methanolic extracts of five *Pogostemon* species. *Rad Antiox.* 2017;7:190-4.
- Assob J, Nde C, Nsagha PF, Njimoh DS, Nfor DL, *et al.* The incidence of feco-oral parasites in street-food vendors in Buea, south-west region Cameroon. *Afr Health Sci.* 2012;12(3):376-80.
- Zou YP, Lu YH, Wei DZ. Protective effects of flavonoids rich extract of *Hypericum perforatum L.* against hydrogen peroxide-induced apoptosis in PC12 cells. *Phytother Res.* 2010;24(1):6-10.
- Zin ZM, Abdul-Hamid A, Osman A. Antioxidative activity of extracts from Mengkudu (*Morinda citrifolia L.*) root, fruit and leaf. *Food Chemistry.* 2002;78(2):227-31.
- Blois MS. Antioxidant determinations by the use of a stable free radical. *Nature.* 1958;181:1199-200.
- Soare J, Dinis T, Cunha A, Almeida L. Antioxidant activities of some extracts of *Thymus zygis*. *Free Radic Res.* 1997;26(5):469-78.
- Hseu YC, Chang WH, Chen CS, Liao JW, Huang CJ, *et al.* Antioxidant activities of *Toona Sinensis* leaves extracts using different antioxidant models. *Food and Chemical Toxicology.* 2008;46(1):105-14.
- Sudhira L, Rao SV, Kamakshamma J. Phytochemical Screening, Antioxidant and Antibacterial Activity of *Strychnos colubrina L.* as an important Endangered Medicinal Species in Eastern Ghats. *Journal of Pharmaceutical Sciences and Research.* 2015;7(5):242-7.
- Shukla S, Mehta A, Vivek KB, Shukla S. *In vitro* antioxidant activity and total phenolic content of ethanolic leaf extract of *Stevia rebaudiana Bert.* *Food Chem Toxicol.* 2009;47:2338-43
- Sridevi H, Jayaraman P, Pachaiyappan P. Anti-inflammatory and antioxidant activities of *Memecylon umbellatum. Burm. f* leaf extract under *in vitro*. *J Pharm Biol Sci.* 2014;9(5):61-8.
- Srinivasan R, Natarajan D, Shivakumar MS. Antioxidant compound quercetin-3-O- α -L-rhamnoside (1 \rightarrow 6)- β -D-glucose (Rutin) isolated from ethyl acetate leaf extracts of *Memecylon edule* Roxb (Melastamataceae). *Free Rad Antiox.* 2015;5(1):35-42.
- Assob JCN, Njouendou AJ, Nkeng PEA, Chouna JR, Badami SM, Verapur VP, *et al.* *In-vitro* screening of antioxidant properties of ten Cameroonian medicinal plants. *J Adv Biotechnol.* 2014;3:2-15.

GRAPHICAL ABSTRACT



SUMMARY

- The present study explored the anti-oxidant activity for decoction and ethanol extracts of Fijian medicinal plants.
- The plants from thirteen species were tested at 0.2 mg/mL, 0.4 mg/mL, 0.6 mg/mL, 0.8 mg/mL, 1.0 mg/mL, 1.5 mg/mL and 2.0 mg/mL concentrations.
- All plants demonstrated free radical scavenging activity of the water extracts as well as ethanol extracts.
- The scavenging activity increases with the increase in the concentration from 0.2 to 2.0 mg/mL.
- The findings encourage studying these plants further in isolating and identifying the active ingredients for their antioxidant properties.

ABOUT AUTHORS



Mr. Ranish N Chand: Senior Education Officer Research, Ministry of Education, Heritage and Arts, Suva, Fiji.



Dr. Romila Devi Gopalan: Senior Lecturer, School of Biological and Chemical Sciences, University of the South Pacific, Suva, Fiji



Dr. Ketan Christi: Senior Lecturer, School of Biological and Chemical Sciences, University of the South Pacific, Suva, Fiji

Cite this article: Chand RN, Gopalan RD and Christi K. Evaluation of Antioxidant Properties in Thirteen Fijian Medicinal Plants Used in Alzheimer's Disease and Related Illness. *Free Radicals and Antioxidants*. 2017;8(1):11-7.