



## Propofol Exhibits Inhibitory Effect Towards Human Liver Microsomes (HLMs)- Catalyzed Glucuronidation of Thienorphine

Shu-Yao ZHANG <sup>1#</sup>, Lei CHEN <sup>2#</sup>, Chao-Xian LIN <sup>3,\*</sup>, Zhi-Wei ZHU <sup>1</sup>,  
Ling FANG <sup>1</sup>, Dai-Shan XIN <sup>1</sup>, & Dai-Nian GUO <sup>3</sup>

<sup>1</sup> Intravenous drug use deployment center, Cancer Hospital of Medical College,  
Shantou University, Shantou City, Guangdong Province, 515031, China

<sup>2</sup> Shantou University Medical College Cancer Hospital, No. 7, Raoping Road, Shantou, 515041, China

<sup>3</sup> Office of the new drug clinical trials, Cancer Hospital of Medical College,  
Shantou University, Shantou City, Guangdong Province, 515031, China

**SUMMARY.** Drug-drug interaction (DDI) is a challenging problem in the process of drug utilization. Inhibition of glucuronidation reaction of drugs is a major reason for DDI. The aim of the present study is to predict propofol-thienorphine interaction from the perspective of propofol's inhibition towards thienorphine glucuronidation. The human liver microsomes (HLMs) incubation system supplemented with uridine 5'-diphosphoglucuronic acid (UDPGA) was used. The results showed that propofol inhibited HLMs-catalyzed thienorphine glucuronidation in a concentration-dependent manner. Both Dixon plot and Lineweaver-Burk plot showed that the inhibition of thienorphine glucuronidation by propofol was best fit to competitive inhibition, and the second plot using slopes from Lineweaver-Burk plot versus thienorphine concentration was used to determine the inhibition kinetic parameter ( $K_i$ ) value to be 365.9  $\mu$ M. Whether the *in vitro* inhibition of propofol towards thienorphine glucuronidation can induce the *in vivo* propofol-thienorphine interaction might be influenced by many factors, including various pharmacokinetic factors influencing the *in vivo* concentration of propofol. These data should be carefully explained due to complicated factors influencing the *in vitro-in vivo* extrapolation (IVIVE) results.

**KEY WORDS:** Drug-drug interaction (DDI), Propofol, Thienorphine.

\* Author to whom correspondence should be addressed. E-mail: linchaoxian123456@163.com; # These two authors equally contributed to this work.