



This is a repository copy of *Correction: Telomerase is required for zebrafish lifespan.*

White Rose Research Online URL for this paper:
<https://eprints.whiterose.ac.uk/130922/>

Version: Published Version

Article:

Henriques, C.M. orcid.org/0000-0003-1882-756X, Carneiro, M.C., Tenente, I.M. et al. (2 more authors) (2017) Correction: Telomerase is required for zebrafish lifespan. PLoS Genetics, 13 (3). e1006652. ISSN 1553-7390

<https://doi.org/10.1371/journal.pgen.1006652>

© 2017 Henriques et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. (<http://creativecommons.org/licenses/by/4.0/>)

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

CORRECTION

Correction: Telomerase Is Required for Zebrafish Lifespan

Catarina M. Henriques, Madalena C. Carneiro, Inês M. Tenente, António Jacinto, Miguel Godinho Ferreira

In panel A of [Fig 1](#), a duplication of the *tert*^{-/-} Skin lane appears where the *tert*^{-/-} Fin lane should be. Please view the correct [Fig 1](#) here with the correct *tert*^{-/-} Fin lane shown. Further clarification and copies of the original gels can be found in [S1 File](#).



OPEN ACCESS

Citation: Henriques CM, Carneiro MC, Tenente IM, Jacinto A, Ferreira MG (2017) Correction: Telomerase Is Required for Zebrafish Lifespan. PLoS Genet 13(3): e1006652. doi:10.1371/journal.pgen.1006652

Published: March 13, 2017

Copyright: © 2017 Henriques et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 1. Telomerase mutant zebrafish have shorter telomeres than WT siblings. A) Representative image of TRAP assay showing that telomerase is not active in the *tert*^{-/-} zebrafish, as compared to *tert*^{+/+} siblings. Here shown are caudal fin and skin protein extracts. HeLa cell extract is shown as positive control. N = 4. B) Representative image of restriction fragment analysis of caudal fin genomic DNA of 3 different individuals at different ages, by southern blot (random primer-labelled telomeric probe (CCCTAA)₁₂-³²P-dCTP). *tert*^{+/+} Zebrafish have heterogeneous telomeres, with two distinct peaks of different lengths. In *tert*^{+/+} the highest peak (~16 Kb, top red arrow) becomes more distinct after 1 months of age and decreases in length over-time (B and D). The lowest peak of telomere intensity also decreases in length (bottom red arrow, B and D). *tert*^{-/-} zebrafish have shorter telomeres than *tert*^{+/+} siblings in different tissues (see also Figure S1A and S1B), observed by the decrease in length of the higher TRF peak. The shortest TRF peaks accompany those of *tert*^{+/+} siblings, and decrease over-time at similar rates. C) Testes fractionation in *tert*^{+/+} reveals the two-telomere length populations in whole testes, whereas mature sperm only shows the shorter TRF smear of about 6 Kb, suggesting different telomere lengths in different cells within a tissue. D) TRF mean sizes were calculated as described in [50]. E) Telomere PNA-FISH in 6-month-old gut tissue shows cells with different telomere intensities in the wild type, mainly localizing to the proliferative niche. In contrast *tert*^{-/-} mutants display cells with less bright and more homogeneous telomere intensity.

doi:10.1371/journal.pgen.1006652.g001

Supporting information

S1 File. Correction and copies of the original gel scans from where lanes were selected. (DOCX)

Reference

1. Henriques CM, Carneiro MC, Tenente IM, Jacinto A, Ferreira MG (2013) Telomerase Is Required for Zebrafish Lifespan. *PLoS Genet* 9(1): e1003214. doi:[10.1371/journal.pgen.1003214](https://doi.org/10.1371/journal.pgen.1003214) PMID: [23349637](https://pubmed.ncbi.nlm.nih.gov/23349637/)