

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



LSHTM Research Online

Shaheed, A; Rathore, S; Bastable, A; Bruce, J; Cairncross, S; Brown, J; (2018) Adherence to Point-of-Use Water Treatment over Short-Term Implementation: Parallel Crossover Trials of Flocculation-Disinfection Sachets in Pakistan and Zambia. *Environmental science & technology*. ISSN 0013-936X
DOI: <https://doi.org/10.1021/acs.est.8b00167>

Downloaded from: <http://researchonline.lshtm.ac.uk/4647771/>

DOI: <https://doi.org/10.1021/acs.est.8b00167>

Usage Guidelines:

Please refer to usage guidelines at <https://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

<https://researchonline.lshtm.ac.uk>

1 **Adherence to point-of-use water treatment over short-term**
2 **implementation: parallel crossover trials of flocculation-disinfection**
3 **sachets in Pakistan and Zambia**
4

5 Shaheed, A.¹, Rathore, S.², Bastable, A.³, Bruce, J.¹, Cairncross, S.¹, Brown, J.^{4*}

6 ¹Department of Disease Control, Faculty of Infectious and Tropical Diseases, London
7 School of Hygiene & Tropical Medicine, Keppel Street, London WC1E 7HT, United
8 Kingdom

9 ²Mehran University of Engineering and Technology, Jamshoro, Sindh 76062, Pakistan

10 ³Oxfam GB, Oxfam House, John Smith Drive, Oxford, OX4 2JY, United Kingdom

11 ⁴School of Civil and Environmental Engineering, Georgia Institute of Technology

12 *corresponding author.

13 Contact details: School of Civil and Environmental Engineering, Georgia Institute of
14 Technology, 311 Ferst Drive, Atlanta, GA 30332. Tel: +1 (404) 385-4579. Email:
15 joe.brown@ce.gatech.edu



16
17 [TOC art]

18 **ABSTRACT**

19 The health benefits of point-of-use (POU) water treatment can only be realized through
20 high adherence: correct, consistent, and sustained use. We conducted parallel randomized,
21 longitudinal crossover trials measuring short-term adherence to two single-use flocculant-
22 disinfectant sachets in Pakistan and Zambia. In both trials, adherence declined sharply for
23 both products over the eight-week surveillance periods, with overall lower adherence to
24 both products in Zambia. There was no significant difference in adherence between the two
25 products. Estimated median daily production of treated water dropped over the crossover
26 period from 2.5 to 1.4 l person⁻¹ day⁻¹ (46% decline) in Pakistan, and from 1.4 to 1.1 l
27 person⁻¹ day⁻¹ (21% decline) in Zambia. The percentage of surveillance points with
28 detectable total chlorine in household drinking water declined from 70% to 49% in Pakistan
29 and rose marginally from 28 to 30% in Zambia. The relatively low and decreasing
30 adherence observed in this study suggests that these products would have provided little
31 protection from waterborne disease risk in these settings. Our findings underscore the
32 challenge of achieving high adherence to POU water treatment, even under conditions of
33 short-term adoption with intensive follow-up.

34

35 **INTRODUCTION**

36 Water quality improvements, including point-of-use (POU) water treatment, are intended
37 to deliver health benefits by reducing exposure to waterborne pathogens¹⁻³. POU water
38 treatment is often recommended for short-term deployment, such as in emergency response,
39 where interim strategies are required to reduce potentially elevated waterborne disease
40 risks when safe water supplies are unavailable⁴⁻⁶. The degree to which POU methods
41 provide protection against disease depends on several factors, including (i) whether
42 drinking water is an important source of pathogen exposure and (ii) effectiveness of the
43 technology in reducing the presence or viability of waterborne pathogens under real-world
44 use conditions.

45

46 Protective effects are also a function of the consistency of treatment over time, since even
47 brief periods of exposure to high risk water can control overall risk⁷⁻⁹. POU *compliance* or
48 *adherence* has been defined as the correct and consistent adoption of a given method^{10 10},

49 or the percentage of total water consumed that is treated^{7, 8}. Previous studies have explored
50 the relationship between POU adherence and health outcomes using Quantitative Microbial
51 Risk Assessment (QMRA), modeling probabilities of infection and estimating the resulting
52 burden of disease⁷⁻⁹. Under most modeling scenarios where waterborne disease risk is high,
53 POU interventions require exclusive or nearly exclusive use to deliver substantial health
54 benefits.

55
56 Despite the critical role of adherence in achieving health gains via water quality
57 interventions, adherence has not been consistently measured in field trials^{2, 3, 11, 12}. Where
58 measured, adherence ranges from very low (<30%) to nearly exclusive use¹³⁻¹⁵. Reviews
59 have found relatively greater disease reductions in studies reporting higher adherence^{3, 11},
60 lower health impact in longer-term studies,¹⁶ and declining adherence overall in
61 longitudinal trials¹⁷⁻²⁰. The effects of adherence on the health impact of water quality
62 interventions are unclear from the epidemiological evidence base, however. Various
63 methods for measuring adherence have been used across a relatively small number of
64 studies.

65
66 Achieving high adherence to POU interventions can be challenging, often requiring
67 substantial changes to individual or collective behaviors and strategies² that can exert a
68 burden on users; changes may be difficult to implement over short-term periods, such as in
69 humanitarian response. These settings may represent the most compelling contexts for
70 POU treatment, however²¹.

71
72 In this study, we examined short-term adherence to POU flocculant-disinfectant sachets,
73 as commonly recommended options for improving drinking-water quality in short-term
74 implementation. Products were a previously characterized flocculant-disinfectant sachet¹⁵
75 and a new product intended to be more acceptable to users by reducing treatment time,
76 streamlining treatment steps, and producing water expected to have a less pronounced
77 chlorine taste, developed because taste and treatment effort may be key barriers to
78 adherence for this type of intervention^{22, 23}. We conducted randomized, longitudinal
79 crossover trials at two sites: flood-prone, rural Sindh, Pakistan, and a cholera-impacted

80 urban area of Lusaka, Zambia. Each trial – using an identical design – was intended to
81 replicate typical short-term deployment in terms of setting and support provided to users.
82 We hypothesized that both treatment options would attain high adherence during short-
83 term, intensive implementation in both settings. We further hypothesized that the taste-
84 improved flocculant-disinfectant sachet would result in increased adherence as potentially
85 more acceptable to users.

86

87 **METHODS**

88 **Study setting and population.** We conducted trials of the two products at two sites: urban
89 Lusaka, Zambia (2012) and rural Sindh, Pakistan (2013). Study site criteria were (1)
90 primary use of water sources lacking adequate disinfection, (2) high prevalence of
91 household-level water storage, (3) recent (but not current) water-related emergencies, and
92 (4) community-level support for the project. We worked with Oxfam country offices to
93 identify potential study sites as typical of those where short-term implementation of POU
94 treatment would be considered. The Zambian trial site was a low-income settlement in
95 Lusaka of over 100,000 inhabitants with a history of inadequate sanitation, water, solid
96 waste management, and seasonal cholera outbreaks in the rainy season^{24, 25}. No cholera
97 cases were reported during the trial period, which included the end of the dry season and
98 the onset of the rainy season. The Pakistan trial was located in a community situated on the
99 edge of a small rural town adjacent to two industrial sites in Sindh province. More than
100 98% of households reported experiencing one or both of the two major floods that affected
101 the area in 2010²⁶ and Sindh in 2011²⁷.

102

103 At each site, we randomly selected households to determine eligibility for participation in
104 the study. Eligible households were any living in the study area who stated an expectation
105 that the household would be present in the community for the duration of the eight week
106 study. We enrolled all eligible, consenting households until the *a priori* sample size
107 criterion was met, intended to detect a difference of 20% in outcomes of adherence,
108 accounting for clustering, loss to follow-up, and missing data. We used standard formulae
109 for sample size calculations,²⁸⁻³⁰ further details on sample size calculations are provided
110 in Supporting Information. The primary respondent for households was an adult (usually

111 female) with responsibility for household water management, including collection, storage,
112 and treatment.

113

114 **Interventions.** We tested two single-use flocculant-disinfectant sachets intended for batch
115 POU treatment of 10 liter volumes: the Purifier of Water (PoW), which has been previously
116 studied under field use conditions^{13, 31-33} and a new flocculant-disinfection sachet, Pureit,
117 developed with the intention of reducing treatment effort and a less pronounced “chlorine
118 taste” in treated water³⁴. Both products are used similarly. Users add sachet contents to 10
119 liter volumes of untreated water. Flocs form and settle as chlorine is released; treated water
120 is decanted through a cotton cloth filter into a storage container. Differences in use are
121 described in Supporting Information. Total time needed for batch treatment per
122 manufacturer recommendations was 27 minutes for PoW and 22 minutes for Pureit. The
123 proprietary Pureit formulation was intended to result in less noticeable chlorine taste in
124 post-treatment water, an innovation designed to promote increased uptake and adherence
125 (Supporting Information). Pureit contains the same coagulant (ferric sulfate) and chlorine-
126 based disinfectant (calcium hypochlorite) as PoW. Its performance under controlled
127 laboratory conditions has been previously characterized³⁴. We supplied all households with
128 sufficient sachets to treat all household drinking water for the duration of the study period,
129 along with the other required materials: a 10 liter bucket, a safe water storage container
130 fitted with a tap and lid, a stirring utensil, and a cotton cloth of the type recommended for
131 use with the products. We informed all participating households that additional sachets
132 were available for any reason throughout the trial, according to households’ needs and
133 preferences, at no cost. We asked that households retain all used and unused sachets
134 throughout the study, and provided each household with containers for this purpose. We
135 recorded the number of sachets provided and the number of used and unused sachets at
136 each household visit.

137

138 The study implementation team aimed to provide guidance to users consistent with Oxfam
139 practice recommendations for POU deployment in emergency response (Supporting
140 Information). The study team trained groups of households in use of the methods before
141 distribution, holding structured training sessions for this purpose. Trainings included step-

142 by-step instructions, demonstrations, and dialogue with participants about the project and
143 the POU methods. Enumeration team members were also available to answer questions
144 and provide further instructions to users at weekly follow-up visits throughout the study.
145 No specific, intensive behavior change component was included in the intervention; we
146 conveyed simple messaging about water risks, available water treatment options, and
147 explicitly described the intent of the study to measure adherence to these interventions over
148 time under actual household use conditions.

149

150 **Study design.** We conducted randomized, longitudinal crossover trials of the two products
151 over eight week periods using identical methods at both study sites (Figure 1). Crossover
152 trials, where all households receive each technology in randomized order, minimize the
153 possibility that an observed effect would be attributable to between-arm differences³⁵⁻³⁷.
154 This study design has been used previously in comparing technology use *in situ*³⁸, allowing
155 for households to serve as their own control³⁹, and enabling within and between-group
156 comparisons on study outcomes. Briefly, we randomly allocated products to half of
157 participating households for four weeks (“Period 1”), after which they were switched to
158 the alternate product for another four weeks (“Period 2”). Pre-defined primary measures of
159 adherence were: (i) self-reported daily use of the product, measured via weekly surveys;
160 (ii) per-capita daily sachet use, measured by counting households’ used sachets at each
161 follow-up visit (also used to calculate the volume of water treated per person per day by
162 the household); and (iii) detection of total chlorine in household drinking water samples.


163

164 At weekly, unannounced visits, enumerators administered surveys collecting information
165 on the household and its water management practices, adherence outcomes, and
166 observations on household hygiene. The survey team collected samples of any water
167 respondents indicated as having been treated. We tested household drinking water at the
168 point of sampling in duplicate for free and total chlorine using a colorimetric *N,N*-diethyl-
169 *p*-phenylenediamine (DPD) method with a detection limit of 0.2 mg l⁻¹ (Palintest Standard
170 Comparator Kit ® PT 220).

171

172

173 **Figure 1.** Crossover trial design.



174

175

176 The enumerator team double entered all data in Epidata 3.1 (Epidata Association,
177 Denmark). After cleaning and checking for internal consistency of data, we conducted all
178 statistical analyses in Stata 12 (StataCorp, TX, USA). We obtained ethical clearance from
179 the London School of Hygiene & Tropical Medicine, and No Objection certifications from
180 the Lusaka City Council and the Office of the Deputy Commissioner in our study district
181 in Sindh.

182

183 **Statistical analysis.** All outcome data were characterized by non-normal distributions and
184 required non-parametric statistical methods for hypothesis testing appropriate to crossover
185 trials.⁴⁰⁻⁴² All analyses accounted for clustering of repeat visits within households on
186 adherence outcomes. We used Somer's D non-parametric analysis of variance for
187 estimating two-way and stratified differences in usage measures across categories such as
188 product, crossover period, and weekly visits²⁴. We used ordered and generalized ordered
189 logistic regression to assess trends in *per capita* consumption over time and between
190 products, with consumption calculated from counts of used sachets and household size; we
191 considered 2.5 l person⁻¹ day⁻¹ the minimum target for meeting household safe water
192 needs.⁴³ We also used logistic regression to test binary outcomes including the presence of

193 detectable total chlorine in water samples and further regression methods for additional
194 analyses (Supporting Information). *A priori* covariates included in models were: crossover
195 period, order of product allocation, reported use of untreated water, household size, and
196 days between visits.

197

198 **RESULTS**

199 Cohort characteristics varied between sites (Table 1), reflecting differences around water
200 access and sources, sanitation, reported education and literacy, and other variables. In
201 Zambia, approximately 8% of recruited households were lost to follow-up due to
202 households leaving the study site. In Pakistan, approximately 10% of recruited households
203 were lost to follow-up, due to having either left the study site or stated lack of interest in
204 the products for water treatment. Of the latter, half returned to the study site after seeing
205 the continued use in the rest of the community; we included data from these households in
206 the analysis. Other descriptive data from both trial sites are provided in Table 1.

207

208 In Zambia, the primary water sources for over 90% of households were public standpipes
209 serving the community. Water delivered to standpipes was reportedly treated by municipal
210 authorities; we periodically tested sources before and during this study and found no
211 evidence of chlorine residual (detection limit: 0.2 mg l⁻¹). Shallow dug wells accounted for
212 the main secondary water source, and were used regularly by households, mostly for
213 washing, cleaning, and cooking, though also for supplementary drinking water.

214 When asked about previous use of household water treatment, 1% of respondents reported
215 ever using filtration, 14% reported occasional boiling, and 58% occasionally used liquid
216 chlorine solution, which had been previously distributed in the community during cholera
217 outbreaks.

218

219 In Pakistan, the primary water source for all households – the Indus River – was accessed
220 via a rudimentary piped supply delivering water to either on-plot taps (68% of households)
221 or community standpipes (32% of households). The only treatment step was mechanical
222 filtration of large particles via screening at the river intake and further settling in the storage
223 tank. As in Zambia, we tested sources before and during this study and found no evidence

224 of chlorine residual. In reporting previous use of household water treatment, 36% of
 225 participating households reported boiling their water at least some of the time, 27%
 226 reporting using alum when turbidity was high, and 82% reported using simple cloth
 227 filtration to strain particulates from water before use.

228

229 **Table 1.** Selected key descriptive characteristics of households enrolled at both trial sites.

Variable	Zambia	Pakistan
Households enrolled	214	247
Households lost to follow-up (%)	17 (8%)	25 (10%)
Median household size (range)	6 (2 – 17)	5 (1 – 13)
Individuals enrolled	1211	1218
Female (%)	51%	51%
Children under 5 at trial start (%)	17%	20%
Median age (range)	17 (<1 – 88)	20 (<1 – 90)
Adults fully literate (%)	60%	5%
Self-reported household daily expenditure (\$USD)		
≤2	18%	19%
>2 – 5	41%	31%
>5 – 8	25%	31%
> 8	17%	19%
Household primary drinking water source		
Public standpipe	92%	32%
On-plot piped water	7%	68%
Shallow well	1%	-
Household sanitation		
None/open defecation	10%	3%
Own pit latrine	14%	48%
Shared pit latrine	76%	49%

230

231 We present intervention use across three measures of adherence: (i) used sachet counts and
 232 calculated volume of treated water per person per day, (ii) detection of chlorine residual in
 233 household drinking water, and (iii) self-reported daily use. Results are summarized in Table
 234 2 for both products and both trial sites collapsed by crossover period. Figures 2 and 3
 235 present adherence measures at each surveillance point for both trial sites. As general trends,
 236 we noted decreases in adherence in the second month of exposure, after households
 237 switched products at the crossover point, overall statistically comparable usage between

238 the two products, and some variability in adherence across the four visits in each crossover
239 period. Results specific to each adherence measure are described further below.

240
241 **Sachet usage and calculated volume of treated water.** Sachet usage, indicated by counts
242 of used and unused sachets retained by households, has been used previously to measure
243 use of batch flocculant-disinfectant POU methods.^{13, 44, 45} By this measure, adherence was
244 higher in Pakistan than it was in Zambia across all time points (Table 2). Weekly household
245 sachet usage in Pakistan dropped from a median of 9 in the first crossover period to near 5
246 sachets household⁻¹ week⁻¹ in the second. In Zambia, median usage per visit dropped from
247 6 sachets household⁻¹ week⁻¹ in the first crossover period to 4 sachets in the second period.

248
249 To translate sachet count data into a readily interpretable measure indicating the potential
250 for treated water to meet households' basic drinking water needs, we used retained sachet
251 counts to calculate the daily per capita volume of treated water available to household
252 residents. We calculated this by counting the number of used sachets since the previous
253 surveillance point, multiplying by 10 liters of treated water per sachet, and dividing by the
254 number of days and number of individuals in the household. Examining this measure,
255 differences in calculated per capita production of treated water over time were greatest
256 between crossover periods, though there was also a slight but statistically significant
257 difference across the four visits in the second crossover period in Pakistan ($p=0.001$, Table
258 S1, Supporting Information), and across the first four visits of the first crossover period in
259 Zambia ($p=0.029$, Table S1, Supporting Information). Volume treated did not differ based
260 on which product was used, in either Pakistan ($p=0.36$, Table S1, Supporting Information)
261 and Zambia ($p=0.91$, Table S1, Supporting Information).

262 We estimated production of treated water to be approximately 2.5 l person⁻¹ day⁻¹ in the
263 first crossover period in Pakistan (Table 2), dropping significantly ($p<0.001$, Table S1,
264 Supporting Information) by approximately 44% to 1.4 l person⁻¹ day⁻¹ in the second
265 crossover period. Overall, estimated use was lower in Zambia: 1.4 l person⁻¹ day⁻¹ in the
266 first crossover period, dropping by 21% to 1.1 l person⁻¹ day⁻¹ (Table 2) in the second
267 crossover period ($p<0.001$). Figures 2 and 3 illustrate the decrease in treated water

268 consumption over the crossover period and the changes by surveillance point during each
269 crossover period.

270 We compared our calculated quantity of treated water per capita to the Sphere-
271 recommended minimum guideline value for daily water consumption in emergencies : 2.5
272 l person⁻¹ day⁻¹.⁴⁶ In Pakistan, 52% of households consumed at least 2.5 l person⁻¹ day⁻¹ in
273 the first crossover period, dropping to 31% in the second period (Table 2). In Zambia, 30%
274 of households consumed at least 2.5 l person⁻¹ day⁻¹ in the first crossover period, dropping
275 to 20% in the second crossover period. We used generalized ordered logistic regression to
276 assess whether crossover period, product, or consumption of untreated water was
277 associated with achieving $\geq 50\%$ of the Sphere-recommended minimum volume for
278 drinking water across sites; results are presented in Table 3. Accounting for clustering of
279 adherence outcomes by repeated household measures, household size, and order of product
280 allocation, we estimated reduced odds of meeting this threshold in the second crossover
281 period for each product and trial site, compared with the first crossover period: aOR = 0.56
282 (95% CI 0.49 – 0.69) in Zambia and aOR = 0.31 (95% CI 0.25 – 0.40) in Pakistan. Although
283 one product (PoW) was associated with borderline-significant increased odds of meeting
284 this threshold in the Pakistan trial, we observed no clear differences in products for this
285 measure. Self-reported untreated water consumption was associated with decreased odds
286 of treatment sufficient to reach the Sphere minimum: aOR = 0.79 (95% CI 0.64 – 0.97) in
287 Zambia and aOR = 0.71 (95% CI 0.57 – 0.89) in Pakistan.

288

289
290

Table 2. Adherence measures by product, crossover period, and trial site.

	<u>Zambia</u>			<u>Pakistan</u>		
Crossover period 1	Pureit	Purifier of Water	Both products	Pureit	Purifier of Water	Both products
Median number sachets used daily per capita	0.80	0.86	0.83	1.3	1.3	1.3
Calculated per capita daily volume water treated: median, l person ⁻¹ day ⁻¹ (range)	1.4 (0-28)	1.6 (0-27)	1.4 (0-28)	2.5 (0-21)	2.5 (0-24)	2.5 (0-24)
Drinking water total chlorine ≥ 0.2 mg l ⁻¹ , % total household visits	30%	27%	29%	72%	67%	70%
Reported untreated water consumption, % total household visits	49%	49%	49%	23%	28%	25%
Calculated daily per capita water treated ≥ 2.5 l person ⁻¹ day ⁻¹ , % total household visits	28%	31%	30%	52%	52%	52%
Crossover period 2	Pureit	Purifier of Water	Both products	Pureit	Purifier of Water	Both products
Median number sachets used daily per capita	0.63	0.60	0.60	0.67	0.80	0.73
Calculated per capita daily volume water treated: median, l person ⁻¹ day ⁻¹ (range)	1.1 (0-15)	1.0 (0-10)	1.1 (0-15)	1.4 (0-25)	1.6 (0-33)	1.4 (0-33)
Drinking water total chlorine ≥ 0.2 mg l ⁻¹ , % total household visits	31%	25%	28%	47%	50%	49%
Reported untreated water consumption, % total household visits	60%	63%	61%	40%	31%	36%
Calculated daily per capita water treated ≥ 2.5 l person ⁻¹ day ⁻¹ , % total household visits	21%	20%	20%	31%	32%	31%

291
292


293
294
295
296
297

Table 3. Adjusted odds ratios (aORs) describing associations between selected variables and adherence, using the threshold of $\geq 50\%$ of the Sphere-recommended minimum volume for drinking water as calculated from used sachet counts.

Variable	Zambia		Pakistan	
	aOR* (95% CI)	P-value	aOR (95% CI)	P-value
Crossover period				
Period 1	1		1	
Period 2	0.56 (0.49 – 0.69)	<0.001	0.31 (0.25 – 0.40)	<0.001
Product				
Pureit	1		1	
Purifier of Water	0.97 (0.79 – 1.2)	0.73	1.3 (1.0 – 1.6)	0.064
Reported untreated water consumption				
No	1		1	
Yes	0.79 (0.64 – 0.97)	0.026	0.71 (0.57 – 0.89)	0.0003
*Logistic regression models adjusted for time between surveillance points, household size, order of product allocation, clustering of repeat measures, as well as crossover period, product, and reported untreated water consumption as appropriate.				


298
299
300
301
302

303 **Figure 2.** Median calculated per capita production of treated drinking water, from used
 304 sachet counts, collapsed across both products. The Sphere-recommended minimum is
 305 2.5 l person⁻¹ day⁻¹ for meeting drinking water needs only.⁴⁶
 306



307

308 **Figure 3.** Median self-reported daily use and presence of chlorine residual (total chlorine
 309 $\geq 0.2 \text{ mg l}^{-1}$) in household drinking water, collapsed across both products.
 310



311

312 **Chlorine residual as indicator of use.** Direct detection of chlorine residual in
313 household drinking water is an unambiguous, objectively measurable indicator of past
314 treatment¹¹ using either product we assessed. In Zambia, detectable chlorine (total
315 chlorine ≥ 0.2 mg l⁻¹) was observed in between 18% and 44% of households across
316 surveillance points, without apparent large difference between the two crossover
317 periods (Figure 3). Approximately 4% of households reported having treated drinking
318 water on hand at the time of unannounced visits across all eight surveillance points
319 (Table S3, Supporting Information). Less than 60% of samples indicated to have been
320 treated by respondents in the 24 hours preceding the household visit were observed to
321 have detectable chlorine. In Pakistan, detectable chlorine was observed in between 64%
322 and 76% of households' drinking water during the first crossover period, dropping to
323 between 43% and 58% in the second crossover period (Figure 3); 19% of households
324 had samples of reportedly treated water across all eight surveillance points (Table S3).
325 When water was indicated by the survey respondent to have been treated in the 24 hours
326 preceding the household visit, detectable chlorine was found in 90% of samples.

327

328 Self-reported intervention use was not associated with presence of detectable chlorine
329 at either trial site: aOR = 0.86 (95% CI 0.59 – 1.2) in Zambia and aOR = 1.1 (95% CI
330 0.27 – 4.2) in Pakistan (Table 4). Counts of ≥ 1 used sachet per day were associated
331 with increased odds of detection of chlorine in Zambia but not Pakistan. Self-report of
332 untreated water consumption was not associated with lower probability of chlorine
333 detection in household drinking water at either trial site.

334

335 **Self-reported use and consumption of untreated water.** In contrast to more objective
336 measures of adherence, self-reported use of both products was relatively high at both
337 trial sites (Table 2, Figure 3). We asked household respondents to estimate their
338 adherence over the week preceding each follow-up point, to compare with observed
339 adherence measures. In Zambia, the median percentage of respondents indicating daily
340 use of the intervention varied between 66% and 86%; in Pakistan, median values were
341 between 87% and 98% throughout the trial (Figure 3). Self-report of drinking untreated
342 water was also common, however. Households in Zambia reported consuming
343 untreated water alongside treated water throughout the study, increasing from
344 approximately 49% in the first crossover period to 61% in the second period (Table 2).
345 Self-report of untreated water consumption was associated with lower adherence (Table

346 3). Approximately 25% of households reported consuming untreated water in the first
 347 crossover period in Pakistan, increasing to 36% in the second period (Table 2).

348

349

350 **Table 4.** Adjusted odds ratios describing associations between selected variables and
 351 presence of detectable total chlorine ($\geq 0.2 \text{ mg l}^{-1}$) in household drinking water, both
 352 products.

353

Variable	Zambia		Pakistan	
	aOR* (95% CI)	P-value	aOR (95% CI)	P-value
Self-reported daily usage				
No	1		1	
Yes	0.86 (0.59 – 1.2)	0.43	1.1 (0.27 – 4.2)	0.93
Sachet count				
<1 per household per day	1		1	
≥ 1 per household per day	1.6 (1.2 – 2.2)	0.004	1.0 (0.66 – 1.5)	0.97
Reported untreated water consumption				
No	1		1	
Yes	0.76 (0.58 – 1.0)	0.053	1.3 (0.78 – 2.1)	0.32
*Logistic regression models adjusted for time between surveillance points, household size, order of product allocation, clustering of repeat measures, as well as crossover period, product, and reported untreated water consumption as appropriate.				

354

355

356

357

DISCUSSION

Our objective was to assess adherence to two similar POU treatment options during short-term implementation via crossover trials using the same design in two different contexts. We found no evidence of a significant difference in adherence between products, suggesting that differences between products (e.g., taste, smell, user burden) were not meaningful in determining adherence. We found variable adherence at both sites, with use decreasing over the surveillance period for both products via all measures, with the exception of self-reported daily use in Zambia, which increased overall during the trial. Calculated volume of water treated per capita per day (from used sachet counts) decreased markedly following crossover at both trial sites, with a greater reduction in Pakistan. We hypothesize that this reduction could be a period effect⁴⁷, resulting from habituation to the product during the first crossover period and subsequent resistance to uptake of the new product following crossover. Exploring this and other explanatory hypotheses will require further statistical analysis of potential quantitative and qualitative determinants to adherence in the context of these trials.

Besides decreasing over the trial duration, overall adherence was relatively low. The highest average per capita treatment estimates (Pakistan in the first crossover period) met the minimum Sphere-recommended guidelines of 2.5 l person⁻¹ day⁻¹, suggesting that these water treatment methods may provide sufficient treated water to meet basic daily drinking water requirements under some conditions. Overall estimated production of treated water by this measure decreased by more than 40% in the second crossover period in Pakistan, however. Per capita consumption was well below 2 l person⁻¹ day⁻¹ in Zambia during both crossover periods, suggesting that the level of observed use would be insufficient for meeting minimum needs. The Sphere guideline value is a conservative estimate for drinking water only (not including other consumptive uses such as cooking), below the World Health Organization-recommended 7.5 l person⁻¹ day⁻¹ to provide for hydration and food preparation in non-emergency contexts.^{48, 49} Moreover, respondents at both trial sites reported consuming untreated water throughout the trial. There is an emerging consensus that to deliver health impact, safe drinking-water must represent a high proportion of total water consumption, given that overall waterborne disease risks can be dominated by brief

389 periods of exposure when treatment is inconsistent and untreated water is of moderate to
390 high risk.⁷⁻⁹ Given that consistent treatment is central to realizing the health benefits of
391 POU interventions,⁷ our findings indicate that the protective effect of these interventions
392 would have been limited if waterborne pathogen risks had been present in these contexts.

393

394 Our findings of variable and generally low adherence are consistent with several studies
395 reporting on POU adoption and use,^{4, 19, 44, 50} including reductions in adherence over time^{18,}
396 ¹⁹ and the concomitant consumption of untreated water.^{51, 52} Our findings also support the
397 hypothesis that decreases in health impact of longer duration health impact trials may be
398 due to decreased adherence over time.^{11, 16, 53} Our study questions the assertion that short
399 term, high-follow-up contexts are likely to be especially amenable to POU interventions²¹:
400 we did not observe this in either trial. Further, our findings are consistent with the few
401 available studies of POU uptake in humanitarian response^{4, 5, 54, 55} that suggest considerable
402 barriers remain to realizing benefits of POU over short-term deployment, though we stress
403 that our trial settings should not be interpreted as closely resembling the humanitarian
404 context. Our trials examined adherence to products that were distributed at no cost to the
405 user. Cost recovery might well have resulted in different levels and patterns of adherence
406 in this non-emergency intervention context.

407

408 Our study allowed us to examine the advantages and disadvantages of several measures of
409 adherence. Self-reported adherence exceeded more objective measures at both trial sites,
410 adding to a growing evidence base suggesting possible bias in self-reported measures of
411 use for POU interventions.^{4,39,42-44} The assumption that households with access to a water
412 treatment intervention actually use it consistently and correctly over time – as is assumed
413 in intention-to-treat analysis, common in POU health impact trials – may not generally
414 hold. It is advisable to build in multiple measures of adherence so that adherence can be
415 estimated empirically, consistent with WHO guidance on monitoring and evaluation in
416 POU trials.⁴⁵ Measurement of adherence is critical to evaluating interventions whose
417 impacts are closely linked with user behaviors that influence exposure risks.

418

419 These trials had a number of important limitations. First, while we intentionally focused
420 on communities with recent histories of waterborne disease risks, there were no outbreaks
421 concurrent with trials. Perception of risk can motivate water treatment and may have other
422 effects on behavior.^{56, 57} Therefore, we cannot conclude that the results from this study
423 indicate adherence in emergency response situations: when there is an obvious, immediate
424 threat to health, such as during an outbreak, increased uptake and use could realistically be
425 expected. Second, though we aimed to assess real-world short-term usage, courtesy bias
426 may have been introduced as the study was overtly a research trial without masking trial
427 intent to participants: the “implementers” in this case were also the enumerators conducting
428 interviews on use. Users may have felt compelled to respond to perceived investigator
429 biases, including reporting increased adherence. Although used sachet counts might be a
430 more objective measure of use than self-report, the measure can be manipulated and is
431 therefore not immune to bias: respondents could empty sachets intentionally, though we
432 did not observe this. Because the timing of unannounced follow-up visits followed a pattern
433 (approximately weekly) and were not always random in order on a given day, households
434 could have treated water selectively on days when visits were expected, without our
435 knowledge. Third, households were provided with all the necessary supplemental material
436 to treat their water, which could have acted as further incentive to join or continue
437 participation in the study insofar as additional sachets had value to users, or may have
438 contributed further to courtesy bias. We observed no on-selling of sachets at either site, but
439 it is possible that this occurred without our knowledge. Fourth, adherence measures – even
440 the several we have included – are imperfect measures of “true” adherence, defined as the
441 percentage of water consumed that has been effectively treated; in typical field settings,
442 this is probably impossible to measure exactly. Fifth, this study was based on two specific
443 flocculant-disinfectant sachets that may not be representative of other POU products, each
444 with characteristics that may differ meaningfully from other POU methods or technologies.
445 POU methods are subject to different perceived benefits and costs to users, with potential
446 implications for short- and long-term adherence. For example, flocculant-disinfectants
447 have been noted for their considerable time and effort requirements¹⁰ while filters may
448 require relatively less effort for regular usage in most settings.^{16, 58} Finally, our ability to

449 detect chlorine residuals was limited by the detection limit of the colorimetric test at 0.2
450 mg l⁻¹, resulting in potential underestimation of adherence by this measure.⁵⁹

451

452 Despite weekly contact with households by the study team, we did not include intensive
453 behavior change programming in these trials beyond basic training and ongoing support at
454 surveillance points. Achieving high adherence to household water treatment may require
455 significant investment of time and resources for successful implementation at scale, given
456 the complexity of human behavior and the reality of water management practices in
457 underserved settings.^{60, 61} For some interventions in some settings, however, adherence
458 may be low or may decline rapidly over time, suggesting low potential for reducing
459 waterborne disease risk. Further work is required to appropriately match water quality
460 interventions to specific settings where they have the greatest chance of impacting global
461 public health.

462

463

464 **ACKNOWLEDGEMENTS**

465 We would like to thank Oxfam GB offices in Oxford, Hyderabad, and Lusaka; the Sindh
466 government and the Lusaka City Council; the Research and Development Foundation
467 (RDF); the field staff that led data implementation and data collection; and the community
468 members whose participation, patience, and insights have been central to our work. We
469 gratefully acknowledge funding support for this study from Oxfam GB.

470

471 **SUPPORTING INFORMATION**

472 Supporting information includes instructions on use for both treatment methods, sample
473 size calculations, further product description, and additional details on intervention
474 methods. Supplementary tables provide details on additional statistical analyses.

475

REFERENCES

- 476
477
478 1. Cairncross, S.; Valdmanis, V., Water Supply, Sanitation, and Hygiene Promotion.
479 In *Disease Control Priorities in Developing Countries*, Jamison, D. T.; Breman, J. G.;
480 Measham, A. R.; Alleyne, G.; Claeson, M.; Evans, D. B.; Jha, P.; Mills, A.; Musgrove, P.,
481 Eds. World Bank. The International Bank for Reconstruction and Development/The World
482 Bank Group: Washington (DC), 2006.
- 483 2. Clasen, T., Household Water Treatment and Safe Storage to Prevent Diarrheal
484 Disease in Developing Countries. *Current environmental health reports* **2015**, *2*, (1), 69-
485 74.
- 486 3. Clasen, T. F.; Alexander, K. T.; Sinclair, D.; Boisson, S.; Peletz, R.; Chang, H. H.;
487 Majorin, F.; Cairncross, S., Interventions to improve water quality for preventing diarrhoea.
488 *The Cochrane database of systematic reviews* **2015**, (10), Cd004794.
- 489 4. Colindres, R. E.; Jain, S.; Bowen, A.; Mintz, E.; Domond, P., After the flood: an
490 evaluation of in-home drinking water treatment with combined flocculent-disinfectant
491 following Tropical Storm Jeanne -- Gonaives, Haiti, 2004. *Journal of water and health*
492 **2007**, *5*, (3), 367-74.
- 493 5. Lantagne, D. S.; Clasen, T. F., Use of household water treatment and safe storage
494 methods in acute emergency response: case study results from Nepal, Indonesia, Kenya,
495 and Haiti. *Environmental science & technology* **2012**, *46*, (20), 11352-60.
- 496 6. Mintz, E. D.; Reiff, F. M.; Tauxe, R. V., Safe water treatment and storage in the
497 home. A practical new strategy to prevent waterborne disease. *Jama* **1995**, *273*, (12), 948-
498 53.
- 499 7. Brown, J.; Clasen, T., High adherence is necessary to realize health gains from
500 water quality interventions. *PloS one* **2012**, *7*, (5), e36735.
- 501 8. Enger, K. S.; Nelson, K. L.; Rose, J. B.; Eisenberg, J. N., The joint effects of
502 efficacy and compliance: a study of household water treatment effectiveness against
503 childhood diarrhea. *Water research* **2013**, *47*, (3), 1181-90.
- 504 9. Hunter, P. R.; Zmirou-Navier, D.; Hartemann, P., Estimating the impact on health
505 of poor reliability of drinking water interventions in developing countries. *The Science of*
506 *the total environment* **2009**, *407*, (8), 2621-4.
- 507 10. Clasen, T., Scaling Up Household Water Treatment Among Low-Income
508 Populations. In WHO Press: Geneva, 2009.
- 509 11. Arnold, B. F.; Colford, J. M., Jr., Treating water with chlorine at point-of-use to
510 improve water quality and reduce child diarrhea in developing countries: a systematic
511 review and meta-analysis. *The American journal of tropical medicine and hygiene* **2007**,
512 *76*, (2), 354-64.
- 513 12. Taylor, D. L.; Kahawita, T. M.; Cairncross, S.; Ensink, J. H., The Impact of Water,
514 Sanitation and Hygiene Interventions to Control Cholera: A Systematic Review. *PloS one*
515 **2015**, *10*, (8), e0135676.
- 516 13. Chiller, T. M.; Mendoza, C. E.; Lopez, M. B.; Alvarez, M.; Hoekstra, R. M.;
517 Keswick, B. H.; Luby, S. P., Reducing diarrhoea in Guatemalan children: randomized
518 controlled trial of flocculant-disinfectant for drinking-water. *Bulletin of the World Health*
519 *Organization* **2006**, *84*, (1), 28-35.
- 520 14. Fiebelkorn, A. P.; Person, B.; Quick, R. E.; Vindigni, S. M.; Jhung, M.; Bowen, A.;
521 Riley, P. L., Systematic review of behavior change research on point-of-use water

522 treatment interventions in countries categorized as low- to medium-development on the
523 human development index. *Social science & medicine (1982)* **2012**, 75, (4), 622-33.

524 15. Thevos, A. K.; Kaona, F. A.; Siajunza, M. T.; Quick, R. E., Adoption of safe water
525 behaviors in Zambia: comparing educational and motivational approaches. *Education for*
526 *health (Abingdon, England)* **2000**, 13, (3), 366-76.

527 16. Hunter, P. R., Household water treatment in developing countries: comparing
528 different intervention types using meta-regression. *Environmental science & technology*
529 **2009**, 43, (23), 8991-7.

530 17. Albert, J.; Luoto, J.; Levine, D., End-user preferences for and performance of
531 competing POU water treatment technologies among the rural poor of Kenya.
532 *Environmental science & technology* **2010**, 44, (12), 4426-32.

533 18. Luoto, J.; Mahmud, M.; Albert, J.; Luby, S.; Najnin, N.; Unicomb, L.; Levine, D.
534 I., Learning to dislike safe water products: results from a randomized controlled trial of the
535 effects of direct and peer experience on willingness to pay. *Environmental science &*
536 *technology* **2012**, 46, (11), 6244-51.

537 19. Brown, J.; Proum, S.; Sobsey, M. D., Sustained use of a household-scale water
538 filtration device in rural Cambodia. *Journal of water and health* **2009**, 7, (3), 404-12.

539 20. Rosa, G.; Kelly, P.; Clasen, T., Consistency of Use and Effectiveness of Household
540 Water Treatment Practices Among Urban and Rural Populations Claiming to Treat Their
541 Drinking Water at Home: A Case Study in Zambia. *The American journal of tropical*
542 *medicine and hygiene* **2016**, 94, (2), 445-55.

543 21. Schmidt, W. P.; Cairncross, S., Household water treatment in poor populations: is
544 there enough evidence for scaling up now? *Environmental science & technology* **2009**, 43,
545 (4), 986-92.

546 22. Crider, Y.; Sultana, S.; Unicomb, L.; Davis, J.; Luby, S. P.; Pickering, A. J., Can
547 you taste it? Taste detection and acceptability thresholds for chlorine residual in drinking
548 water in Dhaka, Bangladesh. *The Science of the total environment* **2017**, 613-614, 840-846.

549 23. Pickering, A. J.; Crider, Y.; Amin, N.; Bauza, V.; Unicomb, L.; Davis, J.; Luby, S.
550 P., Differences in field effectiveness and adoption between a novel automated chlorination
551 system and household manual chlorination of drinking water in Dhaka, Bangladesh: a
552 randomized controlled trial. *PloS one* **2015**, 10, (3), e0118397.

553 24. Chilufya, E. Factors related to recurrence of cholera outbreaks in George compound,
554 Lusaka district. University of Zambia, Lusaka, 2013.

555 25. Grönwall, J., Groundwater dependence among poor urban people: out of sight is
556 out of mind? *International Journal of Urban Sustainable Development* **2011**, 3, (1), 26-39.

557 26. Gaurav, K.; Sinha, R.; Panda, P. K., The Indus flood of 2010 in Pakistan: a
558 perspective analysis using remote sensing data. *Natural Hazards* **2011**, 59, (3), 1815.

559 27. Haq, M.; Akhtar, M.; Muhammad, S.; Paras, S.; Rahmatullah, J., Techniques of
560 Remote Sensing and GIS for flood monitoring and damage assessment: A case study of
561 Sindh province, Pakistan. *The Egyptian Journal of Remote Sensing and Space Science*
562 **2012**, 15, (2), 135-141.

563 28. Diggle, P., Heagerty, P., Liang, K.-Y. & Zeger, S., *Analysis of longitudinal data*.
564 Oxford University Press: Oxford, 2002.

565 29. Hayes, R. J.; Bennett, S., Simple sample size calculation for cluster-randomized
566 trials. *International journal of epidemiology* **1999**, 28, (2), 319-26.

- 567 30. Leon, A. C., Sample-Size Requirements for Comparisons of Two Groups on
568 Repeated Observations of a Binary Outcome. *Evaluation & the Health Professions* **2004**,
569 27, (1), 34-44.
- 570 31. Crump, J. A.; Okoth, G. O.; Slutsker, L.; Ogaja, D. O.; Keswick, B. H.; Luby, S.
571 P., Effect of point-of-use disinfection, flocculation and combined flocculation-disinfection
572 on drinking water quality in western Kenya. *Journal of applied microbiology* **2004**, 97, (1),
573 225-31.
- 574 32. Rangel, J. M.; Lopez, B.; Mejia, M. A.; Mendoza, C.; Luby, S., A novel technology
575 to improve drinking water quality: a microbiological evaluation of in-home flocculation
576 and chlorination in rural Guatemala. *Journal of water and health* **2003**, 1, (1), 15-22.
- 577 33. Souter, P. F.; Cruickshank, G. D.; Tankerville, M. Z.; Keswick, B. H.; Ellis, B. D.;
578 Langworthy, D. E.; Metz, K. A.; Appleby, M. R.; Hamilton, N.; Jones, A. L.; Perry, J. D.,
579 Evaluation of a new water treatment for point-of-use household applications to remove
580 microorganisms and arsenic from drinking water. *Journal of water and health* **2003**, 1, (2),
581 73-84.
- 582 34. Marois-Fiset, J. T.; Shaheed, A.; Brown, J.; Dorea, C. C., Laboratory evaluation of
583 a new coagulant/disinfectant point-of-use water treatment product for emergencies.
584 *Journal of applied microbiology* **2016**, 121, (3), 892-902.
- 585 35. Mills, E. J.; Kelly, S.; Wu, P.; Guyatt, G. H., Epidemiology and reporting of
586 randomized trials employing re-randomization of patient groups: a systematic survey.
587 *Contemporary clinical trials* **2007**, 28, (3), 268-75.
- 588 36. Senn, J., *Cross-over trials in clinical research*. John Wiley: Chichester, 2002.
- 589 37. Louis, T. A.; Lavori, P. W.; Bailar, J. C., 3rd; Polansky, M., Crossover and self-
590 controlled designs in clinical research. *The New England journal of medicine* **1984**, 310,
591 (1), 24-31.
- 592 38. Lozier, M. J.; Sircar, K.; Christensen, B.; Pillarisetti, A.; Pennise, D.; Bruce, N.;
593 Stanistreet, D.; Naeher, L.; Pilishvili, T.; Farrar, J. L.; Sage, M.; Nyagol, R.; Muoki, J.;
594 Wofchuck, T.; Yip, F., Use of Temperature Sensors to Determine Exclusivity of Improved
595 Stove Use and Associated Household Air Pollution Reductions in Kenya. *Environmental*
596 *science & technology* **2016**, 50, (8), 4564-71.
- 597 39. Brown, B. W., Jr., The crossover experiment for clinical trials. *Biometrics* **1980**, 36,
598 (1), 69-79.
- 599 40. Koch, G. G., The use of non-parametric methods in the statistical analysis of the
600 two-period change-over design. *Biometrics* **1972**, 28, (2), 577-84.
- 601 41. Putt, M. E.; Chinchilli, V. M., Nonparametric Approaches to the Analysis of
602 Crossover Studies. *Statistical Science* **2004**, 19, (4), 712-719.
- 603 42. Öhrvik, J., Nonparametric Methods in Crossover Trials. *Biometrical Journal* **1998**,
604 40, (7), 771-789.
- 605 43. Project, T. S., *The Sphere Handbook*. Practical Action Publishing, Schumacher
606 Centre for Technology and Development: Rugby, UK, 2011.
- 607 44. Luby, S. P.; Mendoza, C.; Keswick, B. H.; Chiller, T. M.; Hoekstra, R. M.,
608 Difficulties in bringing point-of-use water treatment to scale in rural Guatemala. *The*
609 *American journal of tropical medicine and hygiene* **2008**, 78, (3), 382-7.
- 610 45. Reller, M. E.; Mendoza, C. E.; Lopez, M. B.; Alvarez, M.; Hoekstra, R. M.; Olson,
611 C. A.; Baier, K. G.; Keswick, B. H.; Luby, S. P., A randomized controlled trial of
612 household-based flocculant-disinfectant drinking water treatment for diarrhea prevention

613 in rural Guatemala. *The American journal of tropical medicine and hygiene* **2003**, *69*, (4),
614 411-9.

615 46. Project, S., *The Sphere Project: Humanitarian Charter and Minimum Standards in*
616 *Humanitarian Response*. The Sphere Project: 2011.

617 47. Senn, S. S., *Cross-over Trials in Clinical Research, 2nd Edition*. John Wiley &
618 Sons: New York, 2002.

619 48. Organization, W. H., *Guidelines for Drinking-Water Quality, 4th Edition*. WHO
620 Press: Geneva, 2011.

621 49. Reed, B., Shaw, R., and Chatterton, K. , *Technical notes on drinking-water,*
622 *sanitation and hygiene in emergencies* World Health Organization (WHO), Water,
623 Engineering and Development Centre (WEDC), Loughborough, UK: Geneva, 2013.

624 50. Brown, J.; Sobsey, M. D., Boiling as household water treatment in Cambodia: a
625 longitudinal study of boiling practice and microbiological effectiveness. *The American*
626 *journal of tropical medicine and hygiene* **2012**, *87*, (3), 394-8.

627 51. Boisson, S.; Kiyombo, M.; Sthresley, L.; Tumba, S.; Makambo, J.; Clasen, T.,
628 Field assessment of a novel household-based water filtration device: a randomised,
629 placebo-controlled trial in the Democratic Republic of Congo. *PloS one* **2010**, *5*, (9),
630 e12613.

631 52. Rosa, G.; Huaylinos, M. L.; Gil, A.; Lanata, C.; Clasen, T., Assessing the
632 consistency and microbiological effectiveness of household water treatment practices by
633 urban and rural populations claiming to treat their water at home: a case study in Peru. *PloS*
634 *one* **2014**, *9*, (12), e114997.

635 53. Clasen, T.; Schmidt, W. P.; Rabie, T.; Roberts, I.; Cairncross, S., Interventions to
636 improve water quality for preventing diarrhoea: systematic review and meta-analysis. *BMJ*
637 *(Clinical research ed.)* **2007**, *334*, (7597), 782.

638 54. Doocy, S.; Burnham, G., Point-of-use water treatment and diarrhoea reduction in
639 the emergency context: an effectiveness trial in Liberia. *Tropical medicine & international*
640 *health : TM & IH* **2006**, *11*, (10), 1542-52.

641 55. Lantagne, D.; Clasen, T., Effective use of household water treatment and safe
642 storage in response to the 2010 Haiti earthquake. *The American journal of tropical*
643 *medicine and hygiene* **2013**, *89*, (3), 426-33.

644 56. Hamoudi, A.; Jeuland, M.; Lombardo, S.; Patil, S.; Pattanayak, S. K.; Rai, S., The
645 effect of water quality testing on household behavior: evidence from an experiment in rural
646 India. *The American journal of tropical medicine and hygiene* **2012**, *87*, (1), 18-22.

647 57. Jalan, J.; Somanathan, E., The importance of being informed: Experimental
648 evidence on demand for environmental quality. *Journal of Development Economics* **2008**,
649 *87*, (1), 14-28.

650 58. Sobsey, M. D.; Stauber, C. E.; Casanova, L. M.; Brown, J. M.; Elliott, M. A., Point
651 of use household drinking water filtration: A practical, effective solution for providing
652 sustained access to safe drinking water in the developing world. *Environmental science &*
653 *technology* **2008**, *42*, (12), 4261-7.

654 59. Meade, R. D.; Murray, A. L.; Mittelman, A. M.; Rayner, J.; Lantagne, D. S.,
655 Accuracy, precision, usability, and cost of portable silver test methods for ceramic filter
656 factories. *Journal of water and health* **2017**, *15*, (1), 72-82.

- 657 60. Lilje, J.; Mosler, H. J., Effects of a behavior change campaign on household
658 drinking water disinfection in the Lake Chad basin using the RANAS approach. *The*
659 *Science of the total environment* **2018**, (619-620), 1599-1607.
- 660 61. Ojomo, E.; Elliott, M.; Goodyear, L.; Forson, M.; Bartram, J., Sustainability and
661 scale-up of household water treatment and safe storage practices: Enablers and barriers to
662 effective implementation. *International journal of hygiene and environmental health* **2015**,
663 *218*, (8), 704-13.
- 664