



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Early Blood Pressure Lowering Does Not Reduce Growth of Intraventricular Hemorrhage following Acute Intracerebral Hemorrhage

Citation for published version:

Chan, E, Anderson, CS, Wang, X, Arima, H, Saxena, A, Moullaali, TJ, Delcourt, C, Wu, G, Wang, J, Chen, G, Lavados, PM, Stapf, C, Robinson, T, Chalmers, J & INTERACT Investigators 2016, 'Early Blood Pressure Lowering Does Not Reduce Growth of Intraventricular Hemorrhage following Acute Intracerebral Hemorrhage: Results of the INTERACT Studies' *Cerebrovascular diseases extra*, vol 6, no. 3, pp. 71-75. DOI: 10.1159/000448897

Digital Object Identifier (DOI):

[10.1159/000448897](https://doi.org/10.1159/000448897)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Cerebrovascular diseases extra

Publisher Rights Statement:

This article is licensed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND).

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Original Paper

Early Blood Pressure Lowering Does Not Reduce Growth of Intraventricular Hemorrhage following Acute Intracerebral Hemorrhage: Results of the INTERACT Studies

Edward Chan^{a, b} Craig S. Anderson^{a, b} Xia Wang^{a, b} Hisatomi Arima^{a, b}
Anubhav Saxena^{a, b} Tom J. Moullaali^a Candice Delcourt^{a, b} Guojun Wu^c
Jinchao Wang^c Guofang Chen^d Pablo M. Lavados^{e, f} Christian Stapf^g
Thompson Robinson^h John Chalmers^{a, b} The INTERACT Investigators

^aThe George Institute for Global Health, Royal Prince Alfred Hospital, and ^bCentral Clinical School, University of Sydney, Sydney, N.S.W., Australia; ^cDepartment of Neurology, Hebei Yutian Hospital, Tangshan, and ^dDepartment of Neurology, Xuzhou Central Hospital, Xuzhou, China; ^eServicio de Neurología, Departamento de Medicina, Clínica Alemana, Universidad del Desarrollo, and ^fDepartamento de Ciencias Neurológicas, Facultad de Medicina, Universidad de Chile, Santiago, Chile; ^gCRCHUM, Université de Montréal et Neurologie Vasculaire, Hôpital Notre-Dame – CHUM, Montréal, Que., Canada;

^hDepartment of Cardiovascular Sciences and NIHR Biomedical Research Unit in Cardiovascular Disease, University of Leicester, Leicester, UK

Key Words

Stroke · Intracerebral hemorrhage · Intraventricular hemorrhage · Hypertension · Intensive blood pressure lowering

Abstract

Background: Intraventricular hemorrhage (IVH) extension is common following acute intracerebral hemorrhage (ICH) and is associated with poor prognosis. **Aim:** To determine whether intensive blood pressure (BP)-lowering therapy reduces IVH growth. **Methods:** Pooled analyses of the Intensive Blood Pressure Reduction in Acute Cerebral Hemorrhage Trials (INTERACT1 and INTERACT2) computed tomography (CT) substudies; multicenter, open, controlled, randomized trials of patients with acute spontaneous ICH and elevated systolic BP, randomly assigned to intensive (<140 mm Hg) or guideline-based (<180 mm Hg) BP management. Participants had blinded central analyses of baseline and 24-hour CT. Association of BP lowering to IVH growth was assessed in analysis of covariance. **Results:** There was no signif-

Clinical trial registration: <http://www.clinicaltrials.gov> (unique identifier: NCT00716079).

Prof. Craig S. Anderson
The George Institute for Global Health
PO Box M201, Missenden Road
Sydney, NSW 2050 (Australia)
E-Mail canderson@george.org.au

icant difference in adjusted mean IVH growth following intensive (n = 228) compared to guideline-recommended (n = 228) BP treatment (1.6 versus 2.2 ml, respectively; p = 0.56). Adjusted mean IVH growth was nonsignificantly greater in patients with a mean achieved systolic BP ≥ 160 mm Hg over 24 h (3.94 ml; p trend = 0.26). **Conclusions:** Early intensive BP-lowering treatment had no clear effect on IVH in acute ICH.

© 2016 The Author(s)
Published by S. Karger AG, Basel

Introduction

Approximately one third of patients with acute spontaneous intracerebral hemorrhage (ICH) have intraventricular hemorrhage (IVH) extension, which confers a worse prognosis [1]. High blood pressure (BP) is associated with both hematoma growth and poor outcome [2], but limited data exist on the relationship of BP and IVH [3]. We aimed to determine whether BP-lowering therapy is associated with IVH growth among participants of the pilot and main phases of the Intensive Blood Pressure Reduction in Acute Cerebral Hemorrhage Trials (INTERACT1 and INTERACT2) [2, 4].

Methods

The INTERACT studies were international, multicenter, prospective, open, blinded-endpoint, randomized controlled trials, described in detail elsewhere [2, 4]. They involved 3,243 ICH patients (<6 h of onset; 404 INTERACT1, 2,839 INTERACT2) and elevated systolic BP (150–220 mm Hg) randomly assigned to intensive (target systolic BP <140 mm Hg) or guideline-recommended (<180 mm Hg) BP management. In predefined computed tomography (CT) substudies, 1,310 consecutive patients (346 INTERACT1, 964 INTERACT2) had a second CT at 24 ± 3 h. ICH and IVH volume analyses were performed centrally by expert assessors blind to other data using MISTar version 3.2 (Apollo Medical Imaging Technology). The ethics committees approved the study protocols and written informed consent was obtained from the patients or appropriate surrogates.

Absolute increase in IVH volume was log-transformed to remove skewness and to achieve near-normal distribution of data for analyses. The association of randomized BP-lowering treatment on the absolute increase in IVH volume over 24 h was assessed in analysis of covariance (ANCOVA), adjusted for age, sex, baseline IVH volume, ICH location and volume, and China region. The mean achieved systolic BP over 24 h was calculated from the average of 1, 6, 12, 18, and 24 h postrandomization readings; their association with absolute increase in IVH volume was assessed by ANCOVA adjusting for the same variables, and also for use of statin therapy, baseline systolic BP, and randomized treatment. A p value <0.05 was considered statistically significant and SAS version 9.3 (SAS Institute, Cary, N.C., USA) was used in all analyses.

Results

There were 456 CT substudy participants with IVH and available data (online suppl. fig. 1; for all online suppl. material, see www.karger.com/doi/10.1159/000448897). Guideline-treated patients were more likely to be female, but otherwise there were no significant differences in baseline characteristics, including time from onset to randomization, BP, as well as ICH and IVH volumes (table 1).

Table 1. Characteristics of patients by randomized treatment

Characteristic	Intensive group (n = 228)	Guideline group (n = 228)	p value
Demographics			
Age, years	67 (13)	68 (12)	0.28
Female	75 (33%)	97 (43%)	0.03
Chinese region	113 (50%)	112 (49%)	0.93
Medical history			
Previous IVH	15 (7%)	17 (8%)	0.72
Previous ischemic stroke	23 (10%)	28 (12%)	0.47
Heart disease	10 (4%)	9 (4%)	0.81
Hypertension	161 (71%)	168 (74%)	0.51
Diabetes mellitus	32 (14%)	27 (12%)	0.47
Medication history			
Antihypertensive agent	119 (52%)	118 (52%)	0.89
Oral anticoagulant	18 (8%)	14 (6%)	0.46
Antiplatelet therapy	38 (17%)	40 (18%)	0.80
Lipid-lowering therapy	27 (12%)	26 (11%)	0.87
Clinical features			
Time from onset to randomization	3.7 (2.7–4.7)	3.5 (2.8–4.7)	0.89
Systolic BP	181 (17)	181 (17)	0.81
Diastolic BP	99 (15)	100 (15)	0.37
Glasgow Coma Scale score ¹	14 (12–15)	14 (12–15)	0.35
NIHSS score ²	14 (9–18)	14 (9–19)	0.51
NIHSS score ≥15	102 (45%)	113 (50%)	0.28
CT findings			
Time from onset to first CT, h	1.8 (1.3–2.6)	1.8 (1.2–2.5)	0.61
Baseline hematoma volume, ml	13.3 (7.5–27.5)	12.3 (7.3–25.7)	0.59
24-hour hematoma volume, ml	13.9 (7.8–34.0)	13.7 (8.0–40.2)	0.36
Baseline IVH volume	2.5 (0.2–9.6)	3.2 (0.0–12.6)	0.32
24-hour IVH volume	4.3 (1.5–12.5)	6.4 (1.6–15.3)	0.15
Baseline deep location of hematoma	186 (82%)	199 (87%)	0.09
Baseline intraventricular hematoma extension	178 (78%)	171 (75%)	0.44

Data are n (%), mean ± SD, or median (IQR).

¹ Scores on the Glasgow Coma Scale range from 15 (fully conscious) to 3 (deep coma). ² Scores on the NIHSS range from 0 (normal neurologic status) to 42 (coma with quadriplegia).

No significant differences were observed in absolute or adjusted mean IVH growth between intensive and guideline BP-lowering treatment [3.4 (95% confidence interval 1.2–5.5) vs. 4.8 ml (2.7–6.9), $p = 0.35$; 1.6 (0.2–3.1) vs. 2.2 (0.8–3.7), $p = 0.56$, respectively]. Patients with an achieved systolic BP over 24 h ≥ 160 mm Hg were younger, more likely to be from China, and had a higher baseline BP (online suppl. table 1). There was a nonsignificant trend towards a greater absolute and adjusted mean IVH growth of 3.58 ml (1.56–5.72) and 3.94 ml (1.46–6.61), respectively, with a mean achieved systolic BP over 24 h ≥ 160 mm Hg (fig. 1). Patients with IVH growth were older, had higher National Institutes of Health Stroke Scale (NIHSS) scores, larger 24-hour ICH volumes, and smaller baseline IVH volumes (online suppl. table 2). The odds ratio for death or major disability was significantly greater with than without IVH growth (adjusted odds ratio 3.51, 95% confidence interval 2.05–6.02; online suppl. table 3).




Fig. 1. Intraventricular volume growth by average achieved BP between 1 and 24 h after randomization. † Assessed by ANCOVA adjusting for age, sex, China region, use of statin, baseline systolic BP, randomized treatment, baseline IVH volume, and ICH location. Because of skewed raw data, adjusted means are reported with 95% confidence intervals obtained by back-transformation.

Discussion

This post hoc analysis of 456 IVH patients from the INTERACT CT substudies showed no clear relationship between IVH growth in acute ICH and a strategy of early intensive BP management, supporting an earlier analysis of the INTERACT2 study of no association of BP and IVH volume [1]. Thus, the clinical benefits of intensive BP lowering on functional outcome seen in the INTERACT studies are likely attributable to mechanisms other than IVH expansion.

These results concur with a previous small prospective cohort study, where again no significant difference in admission systolic BP or mean BP across categories of IVH volumes within the first 24 h of ICH onset was reported [5]. However, the mean 24-hour IVH volume was still higher in our standard BP treatment group, though not significantly different from intensive BP lowering. BP should not be excluded as a contributing factor to IVH growth. Whilst baseline BP per se does not appear to be a major factor in the pathogenesis of IVH, there was a trend towards a greater IVH volume growth at higher average achieved systolic BP over the first 24 h. Indeed, further analyses of the INTERACT2 study showed that rapid and sustained systolic BP reduction was associated with less ICH hematoma growth over 24 h [6]. Another consideration relates to whether intensive BP reduction is associated with a lower risk of developing IVH. We previously reported that delayed IVH is associated with a worse prognosis, though neither intensive BP lowering nor deep location of ICH were significantly associated with delayed IVH [7].

Whilst the strengths to this study include having a robust, well-characterized cohort from large international trials with standardized care, there are several limitations that include the exclusion of patients with large ICH volumes and impaired Glasgow Coma Scale, which makes the results less generalizable to more severe ICH. Furthermore, despite the large sample size, it is still too small to provide precise estimates of a modest effect of early intensive BP lowering on IVH.

In summary, early intensive BP lowering according to the INTERACT protocol had no clear effect on IVH growth, although the lowest IVH growth was achieved in patients with sustained BP control over 24 h.

Acknowledgment

The INTERACT studies were supported by research grants from the National Health and Medical Research Council (NHMRC) of Australia, which played no role in study design, conduct, analyses, or reporting.

Disclosure Statement

T. Robinson reports consultancy payments from Bayer, Boehringer Ingelheim and Daiichi Sankyo, and his institution has received grant funding from the National Institute of Health Research, the British Heart Foundation, the Stroke Association, and the Engineering and Physical Sciences Research Council. J. Chalmers reports grants from the NHMRC during the conduct of the study, grants from Servier International outside the submitted work, and being a chief or co-chief investigator for other large stroke trials. C.S. Anderson reports grant support from the NHMRC, advisory board fees from Astra Zeneca and Medtronic, and speaker fees from Takeda China. The remaining authors have nothing to disclose.

References

- 1 Chan E, Anderson CS, Wang X, Arima H, Saxena A, Moullaali TJ, Heeley E, Delcourt C, Wu G, Wang J, Chen G, Lavados PM, Stapf C, Robinson T, Chalmers J, Huang Y; INTERACT2 Investigators: Significance of intraventricular hemorrhage in acute intracerebral hemorrhage: Intensive Blood Pressure Reduction in Acute Cerebral Hemorrhage Trial results. *Stroke* 2015;46:653–658.
- 2 Anderson CS, Heeley E, Huang Y, Wang J, Stapf C, Delcourt C, Lindley R, Robinson T, Lavados P, Neal B, Hata J, Arima H, Parsons M, Li Y, Wang J, Heritier S, Li Q, Woodward M, Simes RJ, Davis SM, Chalmers J; INTERACT2 Investigators: Rapid blood-pressure lowering in patients with acute intracerebral hemorrhage. *N Engl J Med* 2013;368:2355–2365.
- 3 Ohwaki K, Yano E, Nagashima H, Hirata M, Nakagomi T, Tamura A: Blood pressure management in acute intracerebral hemorrhage: relationship between elevated blood pressure and hematoma enlargement. *Stroke* 2004;35:1364–1367.
- 4 Anderson CS, Huang Y, Wang JG, Arima H, Neal B, Peng B, Heeley E, Skulina C, Parsons MW, Kim JS, Tao QL, Li YC, Jiang JD, Tai LW, Zhang JL, Xu E, Cheng Y, Heritier S, Morgenstern LB, Chalmers J; INTERACT Investigators: Intensive Blood Pressure Reduction in Acute Cerebral Haemorrhage Trial (INTERACT): a randomised pilot trial. *Lancet Neurol* 2008;7:391–399.
- 5 Fortes Lima TT, Prandini MN, Gallo P, Cavalheiro S: Prognostic value of intraventricular bleeding in spontaneous intraparenchymal cerebral hemorrhage of small volume: a prospective cohort study. *Neurosurgery* 2012;70:929–934.
- 6 Carcel C, Wang X, Sato S, Stapf C, Sandset EC, Delcourt C, Arima H, Robinson T, Lavados P, Chalmers J, Anderson CS; INTERACT2 Investigators: Degree and timing of intensive blood pressure lowering on hematoma growth in intracerebral hemorrhage: Intensive Blood Pressure Reduction in Acute Cerebral Hemorrhage Trial-2 results. *Stroke* 2016;47:1651–1653.
- 7 Moullaali TJ, Sato S, Wang X, Rabinstein AA, Arima H, Carcel C, Chen G, Robinson T, Heeley E, Chan E, Delcourt C, Stapf C, Cordonnier C, Lindley RI, Chalmers J, Anderson CS; INTERACT Investigators: Prognostic significance of delayed intraventricular haemorrhage in the INTERACT studies. *J Neurol Neurosurg Psychiatry* 2016, Epub ahead of print.