

Guideline for the Management of Mantle Cell Lymphoma

A British Society for Haematology Guideline

Authors: P. McKay¹, M. Leach¹, R. Jackson², S Robinson³ and S. Rule⁴

Authors' affiliations:

¹Department of Haematology, Beatson West of Scotland Cancer Centre, Glasgow

²Department of Pathology, Queen Elizabeth University Hospital, Glasgow

³Department of Haematology, University Hospitals Bristol

⁴Department of Haematology, Plymouth University Peninsula Schools of Medicine and Dentistry

Correspondence:

BSH Administrator, British Society for Haematology, 100 White Lion Street, London, N1 9PF, UK.

E-mail: bshguidelines@b-s-s.org.uk

Methodology: This guideline was compiled according to the BSH process at www.b-s-h.org.uk. The Grading of Recommendations Assessment, Development and Evaluation (GRADE) nomenclature was used to evaluate levels of evidence and to assess the strength of recommendations. The GRADE criteria can be found at <http://www.gradeworkinggroup.org>.

MEDLINE, EMBASE, DYNAMED, TRIP, NHS EVIDENCE were searched systematically for publications in English from 1980 to 2017 using the key words

lymphoma and mantle cell. References from relevant publications were also searched. Editorials, studies with < 8 cases and letters were excluded. Conference abstracts have been included if deemed to be of particular relevance.

Review of the manuscript was performed by the British Society for Haematology (BSH) Guidelines Committee Haemato-oncology Task Force, the BSH Guidelines Committee and the Haemato-oncology sounding board of BSH. It was also on the members section of the BSH website for comment. It has also been reviewed by patient representatives identified by the Lymphoma Association; these organisations do not necessarily approve or endorse the contents.

The aim is to provide healthcare professionals with guidance on the management of patients with mantle cell lymphoma (MCL). Individual patient circumstances may dictate an alternative approach. This is an update of the guidance published in 2012 (McKay, *et al* 2012), incorporating new therapeutic options including transplant data. Developments in pathology, in particular molecular pathology and use of PET/CT scanning in staging and response assessment are now covered in a separate Good Practice Paper (McKay, *et al* 2018).

Introduction

Mantle cell lymphoma (MCL) is a B-cell malignancy with unique biological, pathological and clinical features comprising 3-10% of all non-Hodgkin lymphomas (NHLs) (Swerdlow, *et al* 1983). It was recognised as a specific entity in the REAL classification (Harris, *et al* 1994) and is characterized by the chromosomal translocation t(11;14)(q13.3;q32.33) resulting in over-expression of the cell cycle protein cyclin D1 (Akiyama, *et al* 1994, Campo, *et al* 1999).

MCL arises **mainly** in older adults (median age 60-65 years) with a male predominance (Argatoff, *et al* 1997, Bosch, *et al* 1998). It often has the worst features of both high and low grade NHL; an aggressive clinical course, but with a pattern of resistant and relapsing disease, rendering it incurable with standard therapy. **Historically, median survival was 4-5 years (Herrmann, *et al* 2009) however this is now in the order of 8-12 years in younger fitter patients who are able to tolerate modern intensive therapies (Eskelund, *et al* 2016, Hermine, *et al* 2016).**

Clinical presentation of MCL is well described (Argatoff, *et al* 1997, Bosch, *et al* 1998, Tiemann, *et al* 2005, Zucca, *et al* 1995). The majority (>90%) of patients present with advanced stage disease. Lymphadenopathy is generally widespread at diagnosis. Splenomegaly, bone marrow infiltration and peripheral blood involvement are common. Bulk disease at diagnosis and B symptoms are less common.

Extranodal involvement is frequent, particularly affecting the gastro-intestinal tract (Romaguera, *et al* 2003) and liver, but infiltration of breast, lung, skin, soft tissue, salivary gland and orbit are also seen. Involvement of more than two extranodal sites occurs in 30-50% of patients (Jares and Campo 2008). Spread to the central nervous system (CNS) can occur, but is rare at diagnosis.

The clinical course is heterogeneous. Presentation can correlate with pathological sub-type; patients with blastoid histology often have an aggressive clinical course, show refractoriness to treatment and have short survival. In contrast, some cases present with more indolent disease (Eve, *et al* 2009a, Martin, *et al* 2009) characterized by splenomegaly, peripheral blood lymphocytosis, little or no nodal disease (Nodit, *et al* 2003, Orchard, *et al* 2003) and survival of the order of 5-12

years. The entity of MCL in situ is now recognized, indicating early foci of disease in otherwise reactive nodes in asymptomatic patients (Carvajal-Cuenca, *et al* 2012).

CNS disease

CNS relapse rates of 4.1 to 7.8% are reported, with high Ki67 and blastoid histology identified as risk factors in multivariate analysis (Cheah, *et al* 2016, Chihara, *et al* 2015, Conconi, *et al* 2013). CNS relapse tends to occur early (median 15 to 20 months) and survival is poor (3 to 8 months). A high incidence of leptomeningeal disease (91% by flow cytometry) has been reported (Cheah, *et al* 2013) with parenchymal disease being uncommon (12% of cases).

Diagnosis, Initial Investigations, Staging and Prognostic Scores – please refer to the Good Practice Paper (McKay, *et al* 2018).

Management

Patients should be offered a clinical trial if available.

Assessment of performance status (PS) using the Eastern Co operative Oncology Group performance status tool (ECOG) is recommended.

Geriatric tools such as the Cumulative Illness Rating Scale (CIRS) and activities of daily living (ADL) assessment may be useful in elderly patients with lymphoma (Nabhan, *et al* 2012).

First-line management

Early stage MCL

A minority of patients present with localised (stage 1A and 2A) MCL. Evidence for management of this group is scarce. Involved field radiotherapy may result in good responses with 64–80% complete remission (CR) and long-term remission, with possible cure in some patients (Rosenbluth and Yahalom 2006, Vandenberghe, *et al* 1997). If early-stage disease is suspected, staging should be confirmed with bone marrow examination and gastro-intestinal investigations. **The role of FDG-PET in early stage MCL remains unproven but may be considered if radical radiotherapy is being proposed** - please refer to Good Practice Paper (McKay, *et al* 2018).

Advanced stage MCL

The majority of patients require treatment at diagnosis but a small subgroup of patients, often with leukaemic presentation +/- splenomegaly, have more indolent disease. Some patients presenting with widespread low volume lymphadenopathy may also run an indolent course. A watch and wait approach is reasonable in well patients with leukemic presentation +/- splenomegaly and in elderly, asymptomatic patients with advanced stage disease. It is not clear whether early treatment of young, asymptomatic patients with advanced stage disease alters long term outcome. A retrospective study of 97 patients demonstrated that for 30% of cases managed initially by observation, treatment was deferred for a median of 12 months (range 4 to 128) with acceptable outcomes (Martin, *et al* 2009). We advise careful assessment of the individual patient. Indications for treatment include:

- Bulky symptomatic lymphadenopathy
- B symptoms
- Symptomatic organomegaly
- GI symptoms including bleeding
- Bone marrow failure

Once treatment is deemed necessary, choice of regimen will depend on age, co-morbidities, performance status and the aim of therapy. Distinction should be made between young fit patients suitable for autologous peripheral blood stem cell transplantation (ASCT), and those for whom this is not appropriate. For those where the aim is to proceed to high dose therapy (HDT) and ASCT in first remission, intensive chemotherapy should be given with the aim of obtaining a complete response.

Rituximab has less activity in MCL than in follicular or diffuse large B-cell lymphoma. Rituximab monotherapy, although well tolerated, is not recommended, having been shown in several trials to have only modest activity (Foran, *et al* 2000, Ghielmini, *et al* 2005, Weigert, *et al* 2007). Addition of rituximab to chemotherapy, however, improves responses including CR rate (Howard, *et al* 2002, Lenz, *et al* 2005, Schulz, *et al* 2007). A recently published phase III randomised trial (Rule, *et al* 2016) confirms survival benefit when rituximab is combined with chemotherapy in first line treatment.

Patients fit for autologous stem cell transplant

CHOP (or R-CHOP) is not as efficacious as more intensive regimens for achieving maximal response prior to transplantation (Andersen, *et al* 2003, Dreyling, *et al* 2005, Lefrere, *et al* 2002). There is increasing awareness of the efficacy of cytarabine in MCL, with excellent responses, overall response rate (ORR) and CR rate, when incorporated into first line regimens. Most studies have shown an ORR of >90% and CR rate of >50%.

High-dose cytarabine was first utilized in MCL as part of the dose-intense hyperCVAD regimen, with good results in single-arm, single-centre settings (Khoury, *et al* 1998). Addition of rituximab gave CR rates of 87-92% (Ritchie, *et al* 2007, Romaguera, *et al* 2010); remissions appeared durable, even without consolidation with ASCT. The main concern with R-hyperCVAD is acute marrow toxicity and infectious complications, and a long-term 5% risk of myelodysplastic syndrome (MDS) or acute myeloid leukaemia (AML). Two multi-centre studies (Bernstein, *et al* 2013, Merli, *et al* 2012) concluded that while R-hyperCVAD is an active regimen for first-line therapy in MCL, it is difficult to deliver, with almost 40% of patients unable to complete the planned treatment schedule. As a result, hyperCVAD-like regimens are not recommended as a first line approach.

High-dose cytarabine was subsequently incorporated into other combination chemotherapies. In the Nordic MCL2 study (Geisler, *et al* 2008) excellent results were achieved with a cytarabine-containing intensive induction immunochemotherapy regimen followed by ASCT: Patients received three cycles of augmented CHOP alternating with 3 cycles of high-dose cytarabine. Rituximab was given from cycle 4 onwards with an additional dose given as in vivo purging prior to

stem-cell mobilization. Event-free survival (EFS) at 5 years was >60% suggesting possible cure, but two updates of this study (Eskelund, *et al* 2016, Geisler, *et al* 2012) have shown a continuous pattern of relapse and no plateau in the survival curve despite very durable responses. Notably, in the low and intermediate risk groups 40% of patients were still in first remission at 12 years.

In a phase 2 GELA trial, (Delarue, *et al* 2013) cytarabine and rituximab were incorporated into induction before ASCT. Treatment comprised 3 cycles CHOP(21) with rituximab included in cycle 3, and 3 cycles of R-DHAP.

Responding patients were eligible for ASCT. ORR was 93% after R-CHOP and 95% after R-DHAP with more CRs on completion of R-DHAP (57% versus 12%).

Median EFS was 83 months and median OS not reached at median follow-up 67 months. Five-year OS was 75%.

The 'MCL Younger' study evaluated use of a high-dose cytarabine- containing ASCT conditioning regimen after induction treatment with alternating courses of R-CHOP and R-DHAP (cytarabine group), compared to 6 x R-CHOP and conventional, cyclophosphamide/total body irradiation (TBI), myeloablative conditioning prior to ASCT (control group). After median follow up of 6.1 years, time to treatment failure was significantly longer in the cytarabine group (9.1 years vs 3.9 years) (Hermine, *et al* 2016).

In the phase 3 LyMa trial, 299 patients received 4 cycles of R-DHAP, with 4 cycles R-CHOP if not in CR/partial remission (PR) as induction therapy prior to ASCT with R-BEAM conditioning, and randomization to rituximab maintenance or not.

Following 4 courses of R-DHAP, ORR was 89% (CR 77%). At a median follow up of 54.4 months, median PFS and OS were not reached, with a 4 year PFS and OS of 67.8% and 78% respectively. .

A phase 2 study (Armand, *et al* 2016) tested induction with 3 cycles of R-bendamustine followed by 3 cycles of R-high dose cytarabine in 23 transplant-eligible patients. CR /Cru rate was 96%. Of 15 minimal residual disease (MRD)-evaluable patients 93% achieved MRD negativity. PFS was 96% at median follow up 13 months. Toxicity was mainly haematological and was more common after R-high dose cytarabine, but overall the regimen was well tolerated.

Despite increasing response rate and response quality, duration of response with chemoimmunotherapy alone remains disappointing. There is, therefore, an increasing tendency to consolidate with high dose therapy and peripheral blood stem cell transplantation in younger patients. This has been shown to increase EFS and possibly OS and is discussed further in the transplant section below.

The optimal approach to therapy of blastoid variant MCL has not been determined, with no studies specifically designed for this subtype. Blastoid variant MCL patients have, however, been included in most of the studies reported above: many have identified blastoid morphology as a poor prognostic factor with early relapses and short survival (Shrestha, *et al* 2015). More durable responses in patients treated with intensive up front therapy including R-hyperCVAD (Romaguera, *et al* 2010) have been reported, and those consolidating primary therapy using ASCT (Damon, *et al* 2009, Geisler, *et al* 2012, Gressin, *et al* 2010).

Conventional chemotherapy for patients unfit for high-dose regimens

The majority of MCL patients are elderly and high-dose therapy is not feasible. For this group, there is no gold-standard therapy, and it is difficult to recommend a specific regimen. Historically, MCL was grouped with low grade lymphomas in clinical studies and treated with regimens including CVP (Meusers, *et al* 1989, Teodorovic, *et al* 1995, Unterhalt, *et al* 1996), fludarabine combinations (Cohen, *et al* 2001) and CHOP (Lenz, *et al* 2005, Meusers, *et al* 1989, Nickenig, *et al* 2006). Most studies included only small numbers of patients. The general pattern was of reasonable ORR (60-88%), but short progression free interval (PFI) (7-21 months) and poor OS (40-85% at 2 years). No particular combination appeared superior in terms of OS. In particular, the addition of anthracycline in the only randomized controlled trial (RCT) (Meusers, *et al* 1989) did not confer survival benefit. Despite this being confirmed in other studies (Bosch, *et al* 1998, Weisenburger, *et al* 2000) CHOP (+R) remains a widely used chemotherapy regimen and forms the control arm in many randomized studies.

Efficacy of purine analogues in MCL has been studied in the front-line setting, with mixed results. When used as a single agent, ORRs of around 30% are seen (Ghielmini, *et al* 2005, Grillo-Lopez 2005). However, when used in combination with cytotoxic agents such as idarubicin or cyclophosphamide response rates increase to approximately 60% (Cohen, *et al* 2001, Zinzani, *et al* 1999, Zinzani, *et al* 2000). In addition to significant haematological and immunological toxicity of purine analogues their use may preclude stem-cell mobilization (Dreyling and Hiddemann 2008, Eve, *et al* 2009b). The European MCL Consortium compared 6 cycles of R-FC to 8 cycles

of R-CHOP (Kluin-Nelemans, *et al* 2012). Responding patients underwent a second randomization to maintenance treatment with either rituximab or interferon alfa, continued until disease progression. Complete response rates were similar (R-FC 40%, R-CHOP 34%), but R-FC was associated with more frequent disease progression (14% versus 5%) and shortened OS at 4 years (47% versus 62%, $p=0.005$). The R-FC regimen was associated with significantly more toxicity (also demonstrated in the study by Rule *et al* (Rule, *et al* 2016)) and is therefore no longer recommended as first line therapy for elderly patients with MCL.

Bendamustine has efficacy in MCL (Herold, *et al* 2006, Rummel, *et al* 2005) and having a favourable toxicity profile compared to CHOP may have an increasing role in treatment. The combination bendamustine and rituximab (BR) demonstrated an ORR of 75% (CR 50%) in the study by Rummel *et al* (Rummel, *et al* 2005). In the BRIGHT study, a randomized global non-inferiority phase 3 study in indolent lymphomas including MCL (Flinn, *et al* 2014), BR was non-inferior to R-CHOP or R-CVP with ORR 97% (CR 31%) for BR and 91% (CR 25%) for R-CHOP/R-CVP. Another phase 3 non-inferiority multi-centre study by the StiL group (Rummel, *et al* 2013) compared BR with R-CHOP as first line treatment for patients with indolent and mantle cell lymphomas, reporting improved PFS in the BR group (69.5 months versus 31.2 months) at a median follow up of 45 months. BR was associated with less alopecia, haematological toxicity, infections and peripheral neuropathy, although skin reactions were more common.

In a phase 3 trial (Robak, *et al* 2015) 487 patients with untreated MCL ineligible for ASCT, were randomized to receive 6-8 cycles of R-CHOP or VR-CAP where

bortezomib was substituted for vincristine in the R-CHOP regimen and given at a dose of 1.3 mg/m² on days 1, 4, 8 and 11. VR-CAP improved PFS (24.7 vs 14.4 months), CR rate (53% vs 42%) and 4 year OS (64% vs 54%) but haematological toxicity was increased. This study showed surprisingly poor PFS in the R-CHOP control arm compared to historical data and considering the tolerability of BR in the BRIGHT and StiL studies, BR appears to be an attractive option for elderly patients.

The GOELAMS group evaluated the RiPAD+C (rituximab, bortezomib, doxorubicin, dexamethasone and chlorambucil) regimen in a phase 2 study of 39 patients, median age 72 years (Houot, *et al* 2012). After 6 cycles, ORR was 74% with CR 59%. Median PFS was 26 months, median OS not reached. The intravenous delivery of bortezomib may have contributed to the high incidence of grade 3 neurotoxicity.

Use of cytarabine in the treatment of older MCL patients is less well established. Visco and colleagues (Visco, *et al* 2013) used a combination of rituximab, bendamustine and cytarabine (R-BAC) in a phase 2 study of 20 treatment-naive patients > 65 years. At a cytarabine dose of 800 mg/m² (the maximum tolerated dose established in phase 1 of the study), the main toxicity was transient grade 3-4 thrombocytopenia (87%) with febrile neutropenia occurring in 12%. ORR was 100% (95% complete response) with 2 year PFS 95%. A dose adjusted R-BAC regimen with cytarabine dose 500 mg/m² was better tolerated, with less haematological toxicity. Final results of the phase 2 study (Visco 2017) **showed CR of 91% with 2 year PFS and OS of 81% and 86% respectively.** The R-BAC combination therefore has considerable potential but results need to be substantiated in larger studies.

Radio-immunotherapy has been added to R-CHOP in an attempt to improve outcome. In a phase 2 study (Smith, *et al* 2012), 56 patients age >18 years, median age 60 years, received 4 cycles of R-CHOP followed by consolidation with yttrium-90-ibritumomab tiuxetan. ORR was 82% (55% CR/Cru) and median time to treatment failure 34.2 months (median follow up 72 months). Estimated 5-year OS was 73%. The regimen was well tolerated with results comparing favourably with historical R-CHOP alone.

Other reported regimens for elderly patients include rituximab combined with chlorambucil (Bauwens, *et al* 2005, Sachanas, *et al* 2011) or cladribine (Spurgeon, *et al* 2011) with overall response rates of 60 to 90%.

CNS prophylaxis

The ability of CNS-penetrating therapies to reduce incidence of CNS relapse is unclear, with conflicting results reported (Cheah, *et al* 2013, Chihara, *et al* 2015, Conconi, *et al* 2013). It is not possible to give clear recommendations on CNS prophylaxis. Empirical use of CNS-penetrating chemotherapy may be beneficial in patients with high Ki67 and/or blastoid morphology.

Maintenance rituximab

R-chemotherapy followed by rituximab maintenance has in some studies prolonged duration of response but not OS (Forstpointner, *et al* 2006, Kahl, *et al* 2006). In others, such as the European MCL Network Study, MCL Elderly Study (Kluin-Nelemans, *et al* 2012) a survival benefit was demonstrated: Here, in addition to

reducing the risk of progression or death by 45%, 58% of patients receiving maintenance rituximab were in remission after 4 years compared with 29% of patients receiving interferon alfa. Patients who responded well to R-CHOP induction had significantly improved OS after maintenance rituximab (87% in CR versus 63% in PR patients). **In a subgroup analysis of the StiLNHL7-2008 MAINTAIN trial, maintenance rituximab every 2 months for 2 years conferred no additional benefit (PFS and OS) compared to observation alone in patients treated with BR (Rummel, MJ, ASCO, 2017, abstract 7503.** A meta-analysis of 3 large randomized clinical trials demonstrated improved PFS but not OS with rituximab maintenance (Vidal 2016). **Maintenance rituximab is included as standard in many current clinical trials and despite the lack of definitive evidence, akin to follicular lymphoma, we recommend maintenance rituximab in patients responding (CR or PR) to R-chemotherapy and who are not proceeding to ASCT as well as in the post-ASCT setting (discussed later). It is difficult to recommend an optimum duration of maintenance therapy. In the European MCL Network Study, MCL Elderly Study (Kluin-Nelemans, *et al* 2012), treatment was continued until disease progression however many institutions limit duration to 2 years. It is therefore advised that duration should be in accordance with local regulatory approvals. In the post transplant setting, duration should be for 3 years (Le Gouill, *et al* 2017).**

Assessment of Treatment Response

Treatment response in MCL is assessed using conventional CT scanning (Cheson, *et al* 2007). As mentioned in the Good Practice Paper (McKay, *et al* 2018) the role of FDG-PET imaging in MCL is not established. **There is increasing evidence that**

end of treatment (EOT) PET/CT scan has prognostic significance (Mato, et al/ 2012 Kolstad, et al/ 2014) however as this does not affect management decisions, we do not recommend a routine EOT PET/CT scan at present unless in the context of a clinical trial.

Although minimal residual disease (MRD) monitoring (using quantitative PCR for the t(11;14) translocation) has been shown to have predictive value particularly following ASCT (Hoster and Pott 2016) the methodology is not widely available at the present time and the clinical benefit unclear.

Recommendations

- **When considering how to treat MCL, clinical presentation (with recognition of the “indolent” form), proliferation index, clinical risk scores (sMIPI and MIPI-c) and performance status, should be taken into account. (1B)**
- **For patients with confirmed early-stage MCL (stage 1A or 2A) involved field radiotherapy is recommended. (1B)**
- **For patients presenting with indolent disease, a period of observation may be appropriate, prior to initiation of definitive therapy. (2B)**
- **Addition of rituximab to combination chemotherapy regimens improves outcomes. Rituximab should therefore be included in the first line chemotherapy regime in the treatment of MCL. (1B)**
- **Patients who are unsuitable for ASCT should receive R-chemotherapy. The best outcome data is for R-CHOP followed by maintenance**

rituximab however R-bendamustine is efficacious and has a more favourable side effect profile (1B)

- **Younger, fit patients should receive rituximab with a regimen containing high dose cytarabine with a view to proceeding to ASCT in 1st remission. (1B)**
- **Single agent rituximab is not recommended. (1A)**
- **Maintenance rituximab therapy is recommended in patients achieving a response (CR or PR) following R-chemotherapy who are not proceeding to autograft. (1B)**
- **Treatment response should be assessed using conventional CT scanning. FDG-PET imaging and assessment of MRD status are not currently recommended outside a clinical trial. (1B)**

Management, second line and beyond

Excluding transplant-eligible patients, median survival after first relapse of MCL is 1-2 years. There is no standard second line chemotherapeutic regimen. Treatments generally produce fewer CRs and duration of response is shorter than following initial therapy, being of the order of 9 months for second and subsequent treatments. Many publications have examined therapeutic regimens in relapsed lymphoma, including patients with MCL. Most are single arm studies, containing fewer than 15 cases, thereby limiting the conclusions that can be drawn. Additionally, inclusion / exclusion criteria of the various relapse studies are different, making it impossible to comment on comparative efficacy.

Transplant

Where the patient has not received ASCT as part of first-line therapy but is considered fit for such therapy at relapse, consideration of consolidation of second response with ASCT is a clinical option. In young patients who have previously undergone ASCT, assessment of fitness for an allogeneic procedure (alloSCT) should be made. If the patient is sufficiently fit, the aim should be to obtain a response with second-line therapy and consolidate with transplantation. This usually involves use of standard salvage regimens as applied in the context of aggressive lymphomas. With the advent of the oral Bruton's kinase (BTK) inhibitor, ibrutinib, however, an alternative salvage option exists; ibrutinib is highly active in the majority of relapsed MCL patients, and the duration of response provides a window to plan and perform the allograft procedure.

The majority of relapsed MCL patients will not be eligible for ASCT or alloSCT. Choice of therapy will be influenced by age, performance status, co-morbidities and initial therapy. It is logical that where a patient has received one prior line of treatment, a different agent be chosen at relapse.

Role of rituximab

Based on the only randomised trial demonstrating survival benefit (Forstpointner, *et al* 2004), combination chemotherapy is usually given with rituximab in relapsed MCL. The role of maintenance rituximab following second line therapy is not clear, with only one published study (Forstpointner, *et al* 2006): 56 MCL patients treated with R-FCM were randomised between rituximab maintenance and no further therapy. Median response duration was identical between arms, but a higher proportion of patients in the maintenance arm experienced remissions beyond 2 years (45% v 9%).

Bendamustine-based regimens

The largest relapse study of bendamustine in combination with rituximab included 45 patients (Czuczman, *et al* 2015). ORR was 82%, CR 40%. Median duration of response was 1.6 years, and 3 year OS 55%. Addition of cytarabine to bendamustine and rituximab (R-BAC) seems an active regimen in relapse as well as in the front line setting (Visco, *et al* 2013); 20 relapsed or refractory patients were treated achieving 80% response rate (70% CR) and estimated 2 year PFS of 70%.

Purine analogues

Activity of fludarabine monotherapy (ORR 33%) (Decaudin, *et al* 1998) is enhanced by addition of cyclophosphamide (Cohen, *et al* 2001) and rituximab (Thomas, *et al* 2005) where ORR was 75%, with 57% CR and response duration 11 months. Single agent cladribine in the relapse setting leads to response rates of 46%, with 21% CR and median PFS 5.4 months (Inwards, *et al* 2008). Responses are probably improved by the addition of cyclophosphamide and rituximab (Robak, *et al* 2006).

Gemcitabine

In a small study of 30 patients with relapsed/refractory disease, gemcitabine was given in combination with dexamethasone (12 patients) or dexamethasone together with cisplatin (18 patients) (Morschhauser, *et al* 2007). ORR was similar, at 36% and 44% respectively, but PFS was very short at 3 months.

Thalidomide

Thalidomide as a single agent is probably less active than lenalidomide. However in a small study of 16 patients, when thalidomide was given in combination with rituximab,

ORR was 81% with 31% CR (Kaufmann, *et al* 2004). Responses were durable with PFS 20.4 months. Rituximab and thalidomide have been added to the PEP-C regimen, a low dose continuous chemotherapy schedule incorporating prednisolone, cyclophosphamide, etoposide and procarbazine: among 22 patients ORR was 82% with 46% CR. Median duration of therapy was 17 months (Coleman, *et al* 2008). Addition of thalidomide and rituximab to this regimen did not appear to improve results (Ruan, *et al* 2010).

Licensed agents including ibrutinib

Worldwide there are 4 drugs licensed for use in relapsed MCL, bortezomib, lenalidomide, temsirolimus and ibrutinib. The comparative efficacy of these drugs as single agents in trials involving relapsed and refractory patients is shown in the table. These were registration trials, with broadly comparable patient populations. Ibrutinib appears the most active of these agents with best ORR, CR rate and duration of response. There has been one randomised trial involving these drugs (Dreyling, *et al* 2016) where ibrutinib was compared with temsirolimus in 280 patients after a median of two prior lines of therapy. There was a significantly better primary endpoint, PFS, with ibrutinib (14.6 months v 6.2 months: $p < 0.0001$), and less treatment-related toxicity, but after median follow up of 20 months, no survival benefit was observed.

As with chemotherapy in MCL, novel agents have been used in combination with rituximab and results of these studies are included in the table. Whilst there is no randomised data, there is a consistent improvement in both response rate and depth following addition of rituximab for all agents, and a suggestion of improved PFS.

A small randomised trial of 46 patients incorporated a novel agent into standard chemotherapy in the relapsed setting: Bortezomib was added to standard CHOP in first relapse MCL (Furtado, *et al* 2015). There was improved ORR (83% v 48%), CR rare (35% v 22%) and OS (36% v 12%) in favour of the bortezomib combination. Of note bortezomib was given on a weekly schedule of 1.6 mg/m², which did not significantly increase neurotoxicity, even when added to vincristine.

Studies of bortezomib combined with dexamethasone and rituximab (Lamm, *et al* 2011), bendamustine and rituximab (Friedberg, *et al* 2011), high dose cytarabine (William, *et al* 2014) (Weigert, *et al* 2009), gemcitabine (Kouroukis, *et al* 2011) and radioimmunotherapy (Beaven, *et al* 2012) show response rates over 50%, but numbers are very small. Morrison *et al* studied addition of bortezomib to lenalidomide in 54 patients (Morrison, *et al* 2015) with ORR 39.5% (CR 15%) but median PFS only 7 months.

Lenalidomide is an oral immunomodulatory agent exhibiting activity in a range of haematological malignancies, and in MCL has single agent activity of around 30% (Goy, *et al* 2013). Responses may, however, be durable, on a par with those with ibrutinib. Response rates can be increased by combining with rituximab, 57% ORR (36% CR) (Wang, *et al* 2012), dexamethasone, ORR 52% (24% CR) (Zaja, *et al* 2012) or bortezomib, ORR 39.5% (15% CR) (Morrison, *et al* 2015). PFS is improved with all of these combinations but is most durable with rituximab or dexamethasone, at around 12 months.

Temsirolimus is an mTOR inhibitor with ORR in the registration study of 22% (2% CR) (Hess, *et al* 2009) .In the randomised study against ibrutinib (Dreyling, *et al* 2016), ORR was higher at 40% (1% CR) reflecting the fact that these patients were far less heavily pre-treated. Addition of rituximab increases response, especially CR rate (59% ORR, 19% CR), but there are limited published data on further combination therapies. One study of temsirolimus with bortezomib (Fenske, *et al* 2015) was active, but only included 7 patients.

Ibrutinib appears the most active single agent in treatment of relapsed refractory MCL, with response rates of 68% (21% CR) (Wang, *et al* 2013). In contrast to temsirolimus, responses were virtually identical in the licensing and randomised trials (Dreyling, *et al* 2016). Addition of rituximab increased ORR to 87% (38% CR) and, in the only other published combination study, addition of bendamustine and rituximab in 17 patients produced an ORR of 94% with 66% CR (Maddocks, *et al* 2015). Ibrutinib can cross the blood brain barrier and is effective in CNS relapse (Tucker, *et al* 2016). **The earlier ibrutinib is used the more effective it is (Rule *et al* 2017). Trials are on-going, assessing a wide range of ibrutinib combination therapies both at relapse and front line. Alternative BTK inhibitors appear to have comparable efficacy (Walter, *et al* 2016, Wang , *et al* 2017).**

Other Drugs

The newer anti-CD20 monoclonal antibodies have lower activity than single agent rituximab in relapsed MCL, (Furtado, *et al* 2014, Morschhauser, *et al* 2013) but since the majority of the patients studied were exposed to prior rituximab, comparisons are difficult to interpret.

The PI3k inhibitor, idelalisib, shows activity as a single agent in relapsed MCL, with ORR 40% (CR 5%). Responses are very short, with median duration 2.7 months and PFS 3.7 months (Kahl, *et al* 2014).

The BCL-2 inhibitor venetoclax looks extremely active. A phase I trial (Davids, *et al* 2017) included 28 patients with MCL and demonstrated an ORR 75% (21% CR) and median PFS of 14 months, almost identical to results seen with single agent ibrutinib.

Single agent flavopiridol (Kouroukis, *et al* 2003), enzastaurin (Morschhauser, *et al* 2008), everolimus (Wang, *et al* 2014) and vorinostat (Kirschbaum, *et al* 2011) have been evaluated in MCL but observed responses do not warrant further exploration.

Recommendations

- **There is no standard therapeutic approach at relapse. An individualised approach should be adopted based on age, co-morbidities, performance status, and response and toxicity with prior therapy (1B)**
- **Ibrutinib is the most active single agent in the relapse setting and should be considered as an option (1A).**
- **An alternative chemotherapeutic regimen should be given at relapse to that given front line (1A)**
- **Rituximab should be given in combination with chemotherapy at relapse (1A)**
- **The activity of novel agents is increased with co-administration of rituximab (1B)**

- **There is little evidence to support the role of maintenance rituximab following relapse therapy**

Stem Cell Transplantation in Mantle Cell Lymphoma

Autologous Stem Cell Transplantation (ASCT)

Only one prospective study compares outcome of ASCT with non-SCT strategies for first-line consolidation in MCL (Dreyling, *et al* 2005). This trial compared outcome of HDT and ASCT versus interferon after CHOP-based induction. Whilst a PFS advantage for HDT consolidation was shown (median PFS 39 months vs 17 months) no OS advantage was demonstrated.

There have been a number of phase 2, single arm studies evaluating the role of HDT consolidation after more intensive induction regimens (Damon, *et al* 2009, Delarue, *et al* 2013, Geisler, *et al* 2008, Gianni, *et al* 2003, Khouri, *et al* 1998). Whilst it is difficult to separate the beneficial effects of intensive induction from those of consolidation, HDT results are impressive, with 4-5 year PFS and OS rates of 56-73% and 64-81% respectively. Data from the recently reported MCL Younger trial which tested induction with either R-CHOP or R-CHOP/R-DHAP prior to HDT and ASCT showed that irrespective of induction used, consolidation HDT converted a significant percentage of patients to an MRD-negative state (Hermine, *et al* 2016). Further, achievement of MRD negativity was an important predictor of favourable outcome. Some investigators have questioned whether HDT consolidation is required after modern, intensive induction regimens, which alone may achieve similar results (Romaguera, *et al* 2010), but, given the excellent results obtained

following intensive induction and HDT consolidation, this approach should be considered a standard of care for suitable patients.. Whether HDT can be safely omitted from intensive first-line therapy that incorporates a BTK inhibitor will be tested in the EMCL Triangle Trial NCT02858258: ASCT After a Rituximab/Ibrutinib/Ara-c Containing iNduction in Generalized Mantle Cell Lymphoma. n.d. [cited 2017 July 7]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02858258>.

Despite such intensive first line approaches most patients will relapse (Eskelund, *et al* 2016, Geisler, *et al* 2012). Prognostic factors predicting favourable outcome of first line consolidation HDT include a low or intermediate MIPI score (Geisler, *et al* 2012), achievement of an MRD-negative state (Hermine, *et al* 2016) and disease status at transplantation (Fenske, *et al* 2014, Till, *et al* 2008). Of note, blastic or pleomorphic morphology was not an adverse predictor of outcome in the Nordic MCL2 trial (Geisler, *et al* 2010) .

In an attempt to improve outcome after first-line HDT a number of strategies have been tested: Addition of radioimmunotherapy to BEAM conditioning demonstrated acceptable toxicity but no clear improvement in disease control (Kolstad, *et al* 2014). The role of TBI in conditioning therapy has been explored only in the context of retrospective analyses, with the hypothesis that patients in PR prior to ASCT may benefit from TBI-based conditioning (Hoster 2016, Rubio 2010). There is however no clearly established optimal conditioning regimen for ASCT in MCL.

Timing of high-dose therapy and ASCT in the disease course has been studied in both The European Society for Blood and Marrow Transplantation (EBMT) and the Center for International Blood and Marrow Transplant Research (CIBMTR) in retrospective analyses. Both studies showed that results of ASCT are superior when used in the first line setting (Fenske, *et al* 2014, Vandenberghe, *et al* 2003), concurring with single centre experience (Tam, *et al* 2009).

A major challenge in the application of HDT and ASCT in MCL is the relative age of the patient population. Registry data describes transplant-related mortality of 10%, in patients aged more than 60 years treated with ASCT compared to 3% in patients under 60 (Jantunen, *et al* 2012). Thus, whilst many clinicians routinely consider HDT and

ASCT for patients over the age of 60, careful evaluation in this patient group is advised. The transplant-specific co-morbidity index has been demonstrated to be predictive of non-relapse mortality in patients undergoing allogeneic SCT (Sorró, *et al* 2005, Sorro, *et al* 2014) but has not been specifically studied in the setting of ASCT for MCL.

Maintenance therapy post-ASCT

Given the pattern of continued relapse maintenance strategies post-ASCT have been investigated. Use of bortezomib maintenance has produced conflicting results and has no current place in therapy (Doorduijn 2015, Kaplan 2015). Rituximab maintenance therapy for a period of 3 years post-ASCT has recently been shown to improve both PFS and OS following first line ASCT (**Le Gouill, *et al* 2017**). **In this large, prospective, randomized trial from the LYSA group, 4 year PFS and OS**

were **83% and 89% respectively for those receiving rituximab versus 64% and 80% in the control arm (observation only)**. We therefore recommend maintenance rituximab in the post-ASCT setting.

Allogeneic SCT (alloSCT)

AlloSCT offers the theoretical benefits of a stem cell graft neither contaminated by lymphoma cells nor damaged by exposure to mutagenic agents, and makes possible the development of a graft-versus-lymphoma effect. Data describing alloSCT for MCL are, however, relatively limited, mostly restricted to retrospective single centre or registry studies, with few prospective single-arm studies. Interpretation of data is further hampered by inclusion of heterogeneous populations of patients, including those transplanted in first-line and relapse settings and patients who have variably received prior ASCT. These studies report significant toxicity, with non-relapse mortality rates of 9-25% at 1 year (Cook, *et al* 2010, Fenske, *et al* 2014, Le Gouill, *et al* 2012, Maris, *et al* 2004, Tam, *et al* 2009). Chronic graft-versus-host disease develops in 40-50% of patients and morbidity associated with this should be considered. Relapse following alloSCT has been reported at between 15 and 38%, with a suggestion of a plateau in relapse after 4 years, even when reduced-intensity conditioning is used (Fenske, *et al* 2014, Le Gouill, *et al* 2012, Tam, *et al* 2009) suggesting that alloSCT may be able to eradicate MCL in a significant minority of patients (Fenske, *et al* 2014, Tam, *et al* 2009, Robinson, *et al* 2017). Longer term follow up is required to determine the curative potential of alloSCT in MCL and establish the efficacy of donor lymphocyte infusions (Cook, *et al* 2010).

There is a paucity of data describing the role of alloSCT in first-line consolidation. The British Society of Blood and Marrow Transplantation (BSBMT) prospective trial of first line consolidation with a BEAM-CAMPATH conditioned alloSCT reported 2 year non-relapse mortality (NRM) and PFS rates of 13.5% and 68% respectively (Peggs 2016). The CIBMTR have reported that for patients undergoing an alloSCT after only 1 or 2 lines of prior therapy (and no prior ASCT), relapse risk following a reduced intensity alloSCT was only 15% compared with 32% for those undergoing an ASCT. The benefit of a lower relapse rate after alloSCT was, however, offset by higher NRM, resulting in no difference in survival (Fenske, *et al* 2014). Whether alloSCT can produce better outcomes than ASCT in patients with high risk disease remains to be determined. Therefore, first line consolidation alloSCT should only be considered within the context of a clinical study.

Data on alloSCT used in either the relapse setting or in patients who have undergone a prior ASCT is more comprehensive (Cook, *et al* 2010, Dietrich, *et al* 2014, Fenske, *et al* 2014, Hamadani, *et al* 2013, Le Gouill, *et al* 2012, Tam, *et al* 2009). As to be expected, the results of alloSCT when used later in the disease course are not as good as when used earlier (Fenske, *et al* 2014). However, patients relapsing after prior ASCT (Dietrich, *et al* 2014) and even those with refractory disease at the time of alloSCT (Hamadani, *et al* 2013) may achieve a prolonged PFS, suggesting that a minority of such high-risk patients may be cured.

Both myeloablative and reduced-intensity conditioning regimens may be considered but there is no clear evidence of the superiority of either approach (Hamadani 2013). Rather, the conditioning regimen should be selected according to age, comorbidities,

Haematopoietic Cell Transplantation-Comorbidity Index (HCT-CI) and prior therapy, on an individual patient basis. Data from retrospective studies suggests the use of T-cell depletion is associated with a higher relapse rate (Cook, *et al* 2010, Robinson, *et al* 2017). However, the overall impact of T-cell depletion on survival post-alloSCT remains controversial (Cook, *et al* 2010, Hamadani, *et al* 2013, Robinson, *et al* 2017). Haploidentical alloSCT enables provision of multiple donors for the majority of patients, but this form of transplantation requires further evaluation in MCL (Dietrich, *et al* 2016). The sequencing of alloSCT relative to the emerging novel therapies also requires further assessment.

Recommendations

- **ASCT should be considered as consolidation of first line therapy for all patients deemed fit for intensive therapy. Patients over 60 years of age should be thoroughly assessed for the suitability of this approach (1A)**
- **ASCT to consolidate first response is most likely to benefit those who achieve a complete remission. (1A)**
- **ASCT can significantly prolong duration of disease response though at present there is insufficient data to demonstrate whether there is a significant overall survival benefit. (1A)**
- **Maintenance rituximab is recommended post-ASCT in fit patients treated with intensive induction. (1A)**
- **AlloSCT may be considered in second remission for fit patients with an appropriate donor. The intensity of the conditioning regimen should be selected on an individual patient basis. (1C)**
- **AlloSCT can be effective at rescuing patients who relapse post-ASCT. (1B)**

- **AlloSCT as part of first-line therapy should be considered only for patients with high-risk disease and preferably within the context of a clinical trial. (1C)**

Comparison of the 4 drugs licensed for use in MCL: data as single agents and in combination with rituximab.

Treatment	Number of Patients	ORR	CR	Median DOR (months)	Median PFS (months)	Median OS (months)
Ibrutinib (Wang, <i>et al</i> 2013)	111	68%	21%	17.5	13.9	Not reached
Ibrutinib + rituximab (Wang, <i>et al</i> 2016)	50	87%	38%	NR	15 month PFS 69%	15 month OS 83%
Bortezomib (Fisher, <i>et al</i> 2006)	155	33%	8%	9.2	6.5	23.5
Bortezomib + rituximab (Agathocleous, <i>et al</i> 2010)	19	58%	16%	NR	NR	NR
Lenalidomide (Goy, <i>et al</i> 2013)	134	28%	8%	16.6	4	19
Lenalidomide + rituximab (Wang, <i>et al</i> 2012)	44	57%	36%	19	11.1	24.3
Temsirolimus (Hess, <i>et al</i> 2009)	54 ^a	22%	2%	7.1	4.8	12.8
Temsirolimus + rituximab (Ansell, <i>et al</i> 2011)	69	59%	19%	10.6	9.7	29.5
CR: Complete Response. DOR: Duration of Response. ORR: Overall Response Rate. OS: Overall Survival. PFS: Progression-free Survival NR: not reported ^a : Results are presented for temsirolimus 175/75 mg dose group.						

Acknowledgements

The BSH Haemato-oncology task force members at the time of writing this guideline were Dr Gail Jones (Chair), Dr Guy Pratt (Secretary), Dr Simon Stern, Dr Jonathan Lambert, Dr

Nilima Parry-Jones, Dr Pam McKay and Dr Alastair Whiteway. The authors would like to thank them, the BSH sounding board, and the BSH guidelines committee for their support in preparing this guideline.

Declaration of Interests

The BSH paid the expenses incurred during the writing of this guideline.

All authors have made a declaration of interests to the BSH and Task Force Chairs which may be viewed on request.

Review Process

Members of the writing group will inform the writing group Chair if any new pertinent evidence becomes available that would alter the strength of the recommendations made in this document or render it obsolete. The document will be archived and removed from the BSH current guidelines website if it becomes obsolete. If new recommendations are made an addendum will be published on the BSH guidelines website www.b-s-h.org.uk.

Disclaimer

While the advice and information in this guidance is believed to be true and accurate at the time of going to press, neither the authors, the BSH nor the publishers accept any legal responsibility for the content of this guidance.

References

- Agathocleous, A., Rohatiner, A., Rule, S., Hunter, H., Kerr, J.P., Neeson, S.M., Matthews, J., Strauss, S., Montoto, S., Johnson, P., Radford, J. & Lister, A. (2010) Weekly versus twice weekly bortezomib given in conjunction with rituximab, in patients with recurrent follicular lymphoma, mantle cell lymphoma and Waldenstrom macroglobulinaemia. *Br J Haematol*, **151**, 346-353.
- Akiyama, N., Tsuruta, H., Sasaki, H., Sakamoto, H., Hamaguchi, M., Ohmura, Y., Seto, M., Ueda, R., Hirai, H., Yazaki, Y. & et al. (1994) Messenger RNA levels of five genes located at chromosome 11q13 in B-cell tumors with chromosome translocation t(11;14)(q13;q32). *Cancer Res*, **54**, 377-379.
- Andersen, N.S., Pedersen, L., Elonen, E., Johnson, A., Kolstad, A., Franssila, K., Langholm, R., Ralfkiaer, E., Akerman, M., Eriksson, M., Kuittinen, O., Geisler, C.H. & Nordic Lymphoma, G. (2003) Primary treatment with autologous stem cell transplantation in mantle cell lymphoma: outcome related to remission pretransplant. *Eur J Haematol*, **71**, 73-80.
- Ansell, S.M., Tang, H., Kurtin, P.J., Koenig, P.A., Inwards, D.J., Shah, K., Ziesmer, S.C., Feldman, A.L., Rao, R., Gupta, M., Erlichman, C. & Witzig, T.E. (2011) Temsirolimus and rituximab in patients with relapsed or refractory mantle cell lymphoma: a phase 2 study. *Lancet Oncol*, **12**, 361-368.
- Argatoff, L.H., Connors, J.M., Klasa, R.J., Horsman, D.E. & Gascoyne, R.D. (1997) Mantle cell lymphoma: a clinicopathologic study of 80 cases. *Blood*, **89**, 2067-2078.
- Armand, P., Redd, R., Bsat, J., Mayuram, S., Giardino, A., Fisher, D.C., LaCasce, A.S., Jacobson, C., Davids, M.S., Brown, J.R., Weng, L., Wilkins, J., Faham, M., Freedman, A.S., Joyce, R. & Jacobsen, E.D. (2016) A phase 2 study of Rituximab-Bendamustine and Rituximab-Cytarabine for transplant-eligible patients with mantle cell lymphoma. *Br J Haematol*, **173**, 89-95.
- Bauwens, D., Maerevoet, M., Michaux, L., Theate, I., Hagemeyer, A., Stul, M., Danse, E., Costantini, S., Vannuffel, P., Straetmans, N., Vekemans, M.C., Deneys, V., Ferrant, A. & Van Den Neste, E. (2005) Activity and safety of combined rituximab with chlorambucil in patients with mantle cell lymphoma. *Br J Haematol*, **131**, 338-340.
- Beaven, A.W., Shea, T.C., Moore, D.T., Feldman, T., Ivanova, A., Ferraro, M., Ford, P., Smith, J. & Goy, A. (2012) A phase I study evaluating ibritumomab tiuxetan (Zevalin(R)) in combination with bortezomib (Velcade(R)) in relapsed/refractory mantle cell and low grade B-cell non-Hodgkin lymphoma. *Leuk Lymphoma*, **53**, 254-258.
- Bernstein, S.H., Epner, E., Unger, J.M., Leblanc, M., Cebula, E., Burack, R., Rimsza, L., Miller, T.P. & Fisher, R.I. (2013) A phase II multicenter trial of hyperCVAD MTX/Ara-C and rituximab in patients with previously untreated mantle cell lymphoma; SWOG 0213. *Ann Oncol*, **24**, 1587-1593.
- Bosch, F., Lopez-Guillermo, A., Campo, E., Ribera, J.M., Conde, E., Piris, M.A., Vallespi, T., Woessner, S. & Montserrat, E. (1998) Mantle cell lymphoma: presenting features, response to therapy, and prognostic factors. *Cancer*, **82**, 567-575.
- Campo, E., Raffeld, M. & Jaffe, E.S. (1999) Mantle-cell lymphoma. *Semin Hematol*, **36**, 115-127.
- Carvajal-Cuenca, A., Sua, L.F., Silva, N.M., Pittaluga, S., Royo, C., Song, J.Y., Sargent, R.L., Espinet, B., Climent, F., Jacobs, S.A., Delabie, J., Naresh, K.N., Bagg, A., Brousset, P., Warnke, R.A., Serrano, S., Harris, N.L., Swerdlow, S.H., Jaffe, E.S. & Campo, E. (2012) In situ mantle cell lymphoma: clinical implications of an incidental finding with indolent clinical behavior. *Haematologica*, **97**, 270-278.
- Cheah, C.Y., George, A., Gine, E., Chiappella, A., Kluin-Nelemans, H.C., Jurczak, W., Krawczyk, K., Mocikova, H., Klener, P., Salek, D., Walewski, J., Szymczyk, M.,

- Smolej, L., Auer, R.L., Ritchie, D.S., Arcaini, L., Williams, M.E., Dreyling, M., Seymour, J.F. & European Mantle Cell Lymphoma, N. (2013) Central nervous system involvement in mantle cell lymphoma: clinical features, prognostic factors and outcomes from the European Mantle Cell Lymphoma Network. *Ann Oncol*, **24**, 2119-2123.
- Cheah, C.Y., Seymour, J.F. & Wang, M.L. (2016) Mantle Cell Lymphoma. *J Clin Oncol*, **34**, 1256-1269.
- Cheson, B.D., Pfistner, B., Juweid, M.E., Gascoyne, R.D., Specht, L., Horning, S.J., Coiffier, B., Fisher, R.I., Hagenbeek, A., Zucca, E., Rosen, S.T., Stroobants, S., Lister, T.A., Hoppe, R.T., Dreyling, M., Tobinai, K., Vose, J.M., Connors, J.M., Federico, M., Diehl, V. & International Harmonization Project on, L. (2007) Revised response criteria for malignant lymphoma. *J Clin Oncol*, **25**, 579-586.
- Chihara, D., Asano, N., Ohmachi, K., Nishikori, M., Okamoto, M., Sawa, M., Sakai, R., Okoshi, Y., Tsukamoto, N., Yakushijin, Y., Nakamura, S., Kinoshita, T., Ogura, M. & Suzuki, R. (2015) Ki-67 is a strong predictor of central nervous system relapse in patients with mantle cell lymphoma (MCL). *Ann Oncol*, **26**, 966-973.
- Cohen, B.J., Moskowitz, C., Straus, D., Noy, A., Hedrick, E. & Zelenetz, A. (2001) Cyclophosphamide/fludarabine (CF) is active in the treatment of mantle cell lymphoma. *Leuk Lymphoma*, **42**, 1015-1022.
- Coleman, M., Martin, P., Ruan, J., Furman, R., Niesvizky, R., Elstrom, R., George, P., Leonard, J. & Kaufmann, T. (2008) Low-dose metronomic, multidrug therapy with the PEP-C oral combination chemotherapy regimen for mantle cell lymphoma. *Leuk Lymphoma*, **49**, 447-450.
- Conconi, A., Franceschetti, S., Lobetti-Bodoni, C., Stathis, A., Margiotta-Casaluci, G., Ramponi, A., Mazzucchelli, L., Bertoni, F., Ghielmini, M., Gaidano, G., Cavalli, F. & Zucca, E. (2013) Risk factors of central nervous system relapse in mantle cell lymphoma. *Leuk Lymphoma*, **54**, 1908-1914.
- Cook, G., Smith, G.M., Kirkland, K., Lee, J., Pearce, R., Thomson, K., Morris, E., Orchard, K., Rule, S., Russell, N., Craddock, C., Marks, D.I., Clinical Trials Committee of the British Society for, B. & Marrow, T. (2010) Outcome following Reduced-Intensity Allogeneic Stem Cell Transplantation (RIC AlloSCT) for relapsed and refractory mantle cell lymphoma (MCL): a study of the British Society for Blood and Marrow Transplantation. *Biol Blood Marrow Transplant*, **16**, 1419-1427.
- Czuczman, M.S., Goy, A., Lamonica, D., Graf, D.A., Munteanu, M.C. & van der Jagt, R.H. (2015) Phase II study of bendamustine combined with rituximab in relapsed/refractory mantle cell lymphoma: efficacy, tolerability, and safety findings. *Ann Hematol*, **94**, 2025-2032.
- Damon, L.E., Johnson, J.L., Niedzwiecki, D., Cheson, B.D., Hurd, D.D., Bartlett, N.L., Lacasce, A.S., Blum, K.A., Byrd, J.C., Kelly, M., Stock, W., Linker, C.A. & Canellos, G.P. (2009) Immunochemotherapy and autologous stem-cell transplantation for untreated patients with mantle-cell lymphoma: CALGB 59909. *J Clin Oncol*, **27**, 6101-6108.
- Davids, M.S., Roberts, A.W., Seymour, J.F., Pagel, J.M., Kahl, B.S., Wierda, W.G., Puvvada, S., Kipps, T.J., Anderson, M.A., Salem, A.H., Dunbar, M., Zhu, M., Peale, F., Ross, J.A., Gressick, L., Desai, M., Kim, S.Y., Verdugo, M., Humerickhouse, R.A., Gordon, G.B. & Gerecitano, J.F. (2017) Phase I First-in-Human Study of Venetoclax in Patients With Relapsed or Refractory Non-Hodgkin Lymphoma. *J Clin Oncol*, **35**, 826-833.
- Decaudin, D., Bosq, J., Tertian, G., Nedellec, G., Bennaceur, A., Venuat, A.M., Bayle, C., Carde, P., Bendahmane, B., Hayat, M. & Munck, J.N. (1998) Phase II trial of

- fludarabine monophosphate in patients with mantle-cell lymphomas. *J Clin Oncol*, **16**, 579-583.
- Delarue, R., Haioun, C., Ribrag, V., Brice, P., Delmer, A., Tilly, H., Salles, G., Van Hoof, A., Casasnovas, O., Brousse, N., Lefrere, F., Hermine, O. & Groupe d'Etude des Lymphomes de l'A. (2013) CHOP and DHAP plus rituximab followed by autologous stem cell transplantation in mantle cell lymphoma: a phase 2 study from the Groupe d'Etude des Lymphomes de l'Adulte. *Blood*, **121**, 48-53.
- Dietrich, S., Boumendil, A., Finel, H., Avivi, I., Volin, L., Cornelissen, J., Jarosinska, R.J., Schmid, C., Finke, J., Stevens, W.B., Schouten, H.C., Kaufmann, M., Sebban, C., Trneny, M., Kobbe, G., Fornecker, L.M., Schetelig, J., Kanfer, E., Heinicke, T., Pfreundschuh, M., Diez-Martin, J.L., Bordessoule, D., Robinson, S. & Dreger, P. (2014) Outcome and prognostic factors in patients with mantle-cell lymphoma relapsing after autologous stem-cell transplantation: a retrospective study of the European Group for Blood and Marrow Transplantation (EBMT). *Ann Oncol*, **25**, 1053-1058.
- Dietrich, S., Finel, H., Martinez, C., Tischer, J., Blaise, D., Chevallier, P., Castagna, L., Milpied, N., Bacigalupo, A., Corradini, P., Mohty, M., Sanz, M., Hausmann, A., Montoto, S., Robinson, S., Boumendil, A., Sureda, A. & Dreger, P. (2016) Post-transplant cyclophosphamide-based haplo-identical transplantation as alternative to matched sibling or unrelated donor transplantation for non-Hodgkin lymphoma: a registry study by the European society for blood and marrow transplantation. *Leukemia*.
- Doorduyn, J. (2015) Bortezomib maintenance therapy after induction with R-CHOP, ARA-C and autologous stem cell transplantation in newly diagnosed MCL patients, results of a multicenter phase II HOVON study. In: *57th ASH annual meeting*, Vol. 126, p. 339. *Blood*.
- Dreyling, M. & Hiddemann, W. (2008) Dose-intense treatment of mantle cell lymphoma: can durable remission be achieved? *Curr Opin Oncol*, **20**, 487-494.
- Dreyling, M., Jurczak, W., Jerkeman, M., Silva, R.S., Rusconi, C., Trneny, M., Offner, F., Caballero, D., Joao, C., Witzens-Harig, M., Hess, G., Bence-Bruckler, I., Cho, S.G., Bothos, J., Goldberg, J.D., Enny, C., Traina, S., Balasubramanian, S., Bandyopadhyay, N., Sun, S., Vermeulen, J., Rizo, A. & Rule, S. (2016) Ibrutinib versus temsirolimus in patients with relapsed or refractory mantle-cell lymphoma: an international, randomised, open-label, phase 3 study. *Lancet*, **387**, 770-778.
- Dreyling, M., Lenz, G., Hoster, E., Van Hoof, A., Gisselbrecht, C., Schmits, R., Metzner, B., Truemper, L., Reiser, M., Steinhauer, H., Boiron, J.M., Boogaerts, M.A., Aldaoud, A., Silingardi, V., Kluin-Nelemans, H.C., Hasford, J., Parwaresch, R., Unterhalt, M. & Hiddemann, W. (2005) Early consolidation by myeloablative radiochemotherapy followed by autologous stem cell transplantation in first remission significantly prolongs progression-free survival in mantle-cell lymphoma: results of a prospective randomized trial of the European MCL Network. *Blood*, **105**, 2677-2684.
- Eskelund, C.W., Kolstad, A., Jerkeman, M., Raty, R., Laurell, A., Eloranta, S., Smedby, K.E., Husby, S., Pedersen, L.B., Andersen, N.S., Eriksson, M., Kimby, E., Bentzen, H., Kuittinen, O., Lauritzsen, G.F., Nilsson-Ehle, H., Ralfkiaer, E., Ehinger, M., Sundstrom, C., Delabie, J., Karjalainen-Lindsberg, M.L., Workman, C.T., Garde, C., Elonen, E., Brown, P., Gronbaek, K. & Geisler, C.H. (2016) 15-year follow-up of the Second Nordic Mantle Cell Lymphoma trial (MCL2): prolonged remissions without survival plateau. *Br J Haematol*, **175**(3), 410-418.

- Eve, H.E., Furtado, M.V., Hamon, M.D. & Rule, S.A. (2009a) Time to treatment does not influence overall survival in newly diagnosed mantle-cell lymphoma. *J Clin Oncol*, **27**, e189-190; author reply e191.
- Eve, H.E., Seymour, J.F. & Rule, S.A. (2009b) Impairment of peripheral blood stem-cell mobilisation in patients with mantle-cell lymphoma following primary treatment with fludarabine and cyclophosphamide +/- rituximab. *Leuk Lymphoma*, **50**, 463-465.
- Fenske, T.S., Shah, N.M., Kim, K.M., Saha, S., Zhang, C., Baim, A.E., Farnen, J.P., Onitilo, A.A., Blank, J.H., Ahuja, H., Wassenaar, T., Qamar, R., Mansky, P., Traynor, A.M., Mattison, R.J. & Kahl, B.S. (2015) A phase 2 study of weekly temsirolimus and bortezomib for relapsed or refractory B-cell non-Hodgkin lymphoma: A Wisconsin Oncology Network study. *Cancer*, **121**, 3465-3471.
- Fenske, T.S., Zhang, M.J., Carreras, J., Ayala, E., Burns, L.J., Cashen, A., Costa, L.J., Freytes, C.O., Gale, R.P., Hamadani, M., Holmberg, L.A., Inwards, D.J., Lazarus, H.M., Maziarz, R.T., Munker, R., Perales, M.A., Rizzieri, D.A., Schouten, H.C., Smith, S.M., Waller, E.K., Wirk, B.M., Laport, G.G., Maloney, D.G., Montoto, S. & Hari, P.N. (2014) Autologous or reduced-intensity conditioning allogeneic hematopoietic cell transplantation for chemotherapy-sensitive mantle-cell lymphoma: analysis of transplantation timing and modality. *J Clin Oncol*, **32**, 273-281.
- Fisher, R.I., Bernstein, S.H., Kahl, B.S., Djulbegovic, B., Robertson, M.J., de Vos, S., Epner, E., Krishnan, A., Leonard, J.P., Lonial, S., Stadtmauer, E.A., O'Connor, O.A., Shi, H., Boral, A.L. & Goy, A. (2006) Multicenter phase II study of bortezomib in patients with relapsed or refractory mantle cell lymphoma. *J Clin Oncol*, **24**, 4867-4874.
- Flinn, I.W., van der Jagt, R., Kahl, B.S., Wood, P., Hawkins, T.E., Macdonald, D., Hertzberg, M., Kwan, Y.L., Simpson, D., Craig, M., Kolibaba, K., Issa, S., Clementi, R., Hallman, D.M., Munteanu, M., Chen, L. & Burke, J.M. (2014) Open-label, randomized, noninferiority study of bendamustine-rituximab or R-CHOP/R-CVP in first-line treatment of advanced indolent NHL or MCL: the BRIGHT study. *Blood*, **123**, 2944-2952.
- Foran, J.M., Rohatiner, A.Z., Cunningham, D., Popescu, R.A., Solal-Celigny, P., Ghielmini, M., Coiffier, B., Johnson, P.W., Gisselbrecht, C., Reyes, F., Radford, J.A., Bessell, E.M., Souleau, B., Benzohra, A. & Lister, T.A. (2000) European phase II study of rituximab (chimeric anti-CD20 monoclonal antibody) for patients with newly diagnosed mantle-cell lymphoma and previously treated mantle-cell lymphoma, immunocytoma, and small B-cell lymphocytic lymphoma. *J Clin Oncol*, **18**, 317-324.
- Forstpointner, R., Dreyling, M., Repp, R., Hermann, S., Hanel, A., Metzner, B., Pott, C., Hartmann, F., Rothmann, F., Rohrberg, R., Bock, H.P., Wandt, H., Unterhalt, M., Hiddemann, W. & German Low-Grade Lymphoma Study, G. (2004) The addition of rituximab to a combination of fludarabine, cyclophosphamide, mitoxantrone (FCM) significantly increases the response rate and prolongs survival as compared with FCM alone in patients with relapsed and refractory follicular and mantle cell lymphomas: results of a prospective randomized study of the German Low-Grade Lymphoma Study Group. *Blood*, **104**, 3064-3071.
- Forstpointner, R., Unterhalt, M., Dreyling, M., Bock, H.P., Repp, R., Wandt, H., Pott, C., Seymour, J.F., Metzner, B., Hanel, A., Lehmann, T., Hartmann, F., Einsele, H., Hiddemann, W. & German Low Grade Lymphoma Study, G. (2006) Maintenance therapy with rituximab leads to a significant prolongation of response duration after salvage therapy with a combination of rituximab, fludarabine, cyclophosphamide, and mitoxantrone (R-FCM) in patients with recurring and refractory follicular and mantle cell lymphomas: Results of a prospective randomized study of the German Low Grade Lymphoma Study Group (GLSG). *Blood*, **108**, 4003-4008.

- Friedberg, J.W., Vose, J.M., Kelly, J.L., Young, F., Bernstein, S.H., Peterson, D., Rich, L., Blumel, S., Proia, N.K., Liesveld, J., Fisher, R.I., Armitage, J.O., Grant, S. & Leonard, J.P. (2011) The combination of bendamustine, bortezomib, and rituximab for patients with relapsed/refractory indolent and mantle cell non-Hodgkin lymphoma. *Blood*, **117**, 2807-2812.
- Furtado, M., Dyer, M.J., Johnson, R., Berrow, M. & Rule, S. (2014) Ofatumumab monotherapy in relapsed/refractory mantle cell lymphoma--a phase II trial. *Br J Haematol*, **165**, 575-578.
- Furtado, M., Johnson, R., Kruger, A., Turner, D. & Rule, S. (2015) Addition of bortezomib to standard dose chop chemotherapy improves response and survival in relapsed mantle cell lymphoma. *Br J Haematol*, **168**, 55-62.
- Geisler, C.H., Kolstad, A., Laurell, A., Andersen, N.S., Pedersen, L.B., Jerkeman, M., Eriksson, M., Nordstrom, M., Kimby, E., Boesen, A.M., Kuittinen, O., Lauritzsen, G.F., Nilsson-Ehle, H., Ralfkiaer, E., Akerman, M., Ehinger, M., Sundstrom, C., Langholm, R., Delabie, J., Karjalainen-Lindsberg, M.L., Brown, P., Elonen, E. & Nordic Lymphoma, G. (2008) Long-term progression-free survival of mantle cell lymphoma after intensive front-line immunochemotherapy with in vivo-purged stem cell rescue: a nonrandomized phase 2 multicenter study by the Nordic Lymphoma Group. *Blood*, **112**, 2687-2693.
- Geisler, C.H., Kolstad, A., Laurell, A., Jerkeman, M., Raty, R., Andersen, N.S., Pedersen, L.B., Eriksson, M., Nordstrom, M., Kimby, E., Bentzen, H., Kuittinen, O., Lauritzsen, G.F., Nilsson-Ehle, H., Ralfkiaer, E., Ehinger, M., Sundstrom, C., Delabie, J., Karjalainen-Lindsberg, M.L., Brown, P., Elonen, E. & Nordic Lymphoma, G. (2012) Nordic MCL2 trial update: six-year follow-up after intensive immunochemotherapy for untreated mantle cell lymphoma followed by BEAM or BEAC + autologous stem-cell support: still very long survival but late relapses do occur. *Br J Haematol*, **158**, 355-362.
- Geisler, C.H., Kolstad, A., Laurell, A., Raty, R., Jerkeman, M., Eriksson, M., Nordstrom, M., Kimby, E., Boesen, A.M., Nilsson-Ehle, H., Kuittinen, O., Lauritzsen, G.F., Ralfkiaer, E., Ehinger, M., Sundstrom, C., Delabie, J., Karjalainen-Lindsberg, M.L., Brown, P., Elonen, E. & Nordic Lymphoma, G. (2010) The Mantle Cell Lymphoma International Prognostic Index (MIPI) is superior to the International Prognostic Index (IPI) in predicting survival following intensive first-line immunochemotherapy and autologous stem cell transplantation (ASCT). *Blood*, **115**, 1530-1533.
- Ghielmini, M., Schmitz, S.F., Cogliatti, S., Bertoni, F., Waltzer, U., Fey, M.F., Betticher, D.C., Schefer, H., Pichert, G., Stahel, R., Ketterer, N., Bargetzi, M., Cerny, T. & Swiss Group for Clinical Cancer, R. (2005) Effect of single-agent rituximab given at the standard schedule or as prolonged treatment in patients with mantle cell lymphoma: a study of the Swiss Group for Clinical Cancer Research (SAKK). *J Clin Oncol*, **23**, 705-711.
- Gianni, A.M., Magni, M., Martelli, M., Di Nicola, M., Carlo-Stella, C., Pilotti, S., Rambaldi, A., Cortelazzo, S., Patti, C., Parvis, G., Benedetti, F., Capria, S., Corradini, P., Tarella, C. & Barbui, T. (2003) Long-term remission in mantle cell lymphoma following high-dose sequential chemotherapy and in vivo rituximab-purged stem cell autografting (R-HDS regimen). *Blood*, **102**, 749-755.
- Goy, A., Sinha, R., Williams, M.E., Kalayoglu Basisik, S., Drach, J., Ramchandren, R., Zhang, L., Cicero, S., Fu, T. & Witzig, T.E. (2013) Single-agent lenalidomide in patients with mantle-cell lymphoma who relapsed or progressed after or were refractory to bortezomib: phase II MCL-001 (EMERGE) study. *J Clin Oncol*, **31**, 3688-3695.

- Gressin, R., Caulet-Maugendre, S., Deconinck, E., Tournilhac, O., Gyan, E., Moles, M.P., El Yamani, A., Cornillon, J., Rossi, J.F., Le Gouill, S., Lepeu, G., Damaj, G., Celigny, P.S., Maisonneuve, H., Corront, B., Vilque, J.P., Casassus, P., Lamy, T., Colonna, M., Colombat, P. & French, G.G. (2010) Evaluation of the (R)VAD+C regimen for the treatment of newly diagnosed mantle cell lymphoma. Combined results of two prospective phase II trials from the French GOELAMS group. *Haematologica*, **95**, 1350-1357.
- Grillo-Lopez, A.J. (2005) Is rituximab effective as a single agent in mantle-cell lymphoma? *Nat Clin Pract Oncol*, **2**, 500-501.
- Hamadani, M., Saber, W., Ahn, K.W., Carreras, J., Cairo, M.S., Fenske, T.S., Gale, R.P., Gibson, J., Hale, G.A., Hari, P.N., Hsu, J.W., Inwards, D.J., Kamble, R.T., Klein, A., Maharaj, D., Marks, D.I., Rizzieri, D.A., Savani, B.N., Schouten, H.C., Waller, E.K., Wirk, B. & Lazarus, H.M. (2013) Allogeneic hematopoietic cell transplantation for chemotherapy-unresponsive mantle cell lymphoma: a cohort analysis from the center for international blood and marrow transplant research. *Biol Blood Marrow Transplant*, **19**, 625-631.
- Harris, N.L., Jaffe, E.S., Stein, H., Banks, P.M., Chan, J.K., Cleary, M.L., Delsol, G., De Wolf-Peeters, C., Falini, B., Gatter, K.C. & et al. (1994) A revised European-American classification of lymphoid neoplasms: a proposal from the International Lymphoma Study Group. *Blood*, **84**, 1361-1392.
- Hermine, O., Hoster, E., Walewski, J., Bosly, A., Stilgenbauer, S., Thieblemont, C., Szymczyk, M., Bouabdallah, R., Kneba, M., Hallek, M., Salles, G., Feugier, P., Ribrag, V., Birkmann, J., Forstpointner, R., Haioun, C., Hanel, M., Casasnovas, R.O., Finke, J., Peter, N., Bouabdallah, K., Sebban, C., Fischer, T., Duhrsen, U., Metzner, B., Maschmeyer, G., Kanz, L., Schmidt, C., Delarue, R., Brousse, N., Klapper, W., Macintyre, E., Delfau-Larue, M.H., Pott, C., Hiddemann, W., Unterhalt, M., Dreyling, M. & European Mantle Cell Lymphoma, N. (2016) Addition of high-dose cytarabine to immunochemotherapy before autologous stem-cell transplantation in patients aged 65 years or younger with mantle cell lymphoma (MCL Younger): a randomised, open-label, phase 3 trial of the European Mantle Cell Lymphoma Network. *Lancet*, **388**, 565-575.
- Herold, M., Schulze, A., Niederwieser, D., Franke, A., Fricke, H.J., Richter, P., Freund, M., Ismer, B., Dachselt, K., Boewer, C., Schirmer, V., Weniger, J., Pasold, R., Winkelmann, C., Klinkenstein, C., Schulze, M., Arzberger, H., Bremer, K., Hahnfeld, S., Schwarzer, A., Muller, C., Muller, C., East German Study Group, H. & Oncology (2006) Bendamustine, vincristine and prednisone (BOP) versus cyclophosphamide, vincristine and prednisone (COP) in advanced indolent non-Hodgkin's lymphoma and mantle cell lymphoma: results of a randomised phase III trial (OSHO# 19). *J Cancer Res Clin Oncol*, **132**, 105-112.
- Herrmann, A., Hoster, E., Zwingers, T., Brittinger, G., Engelhard, M., Meusers, P., Reiser, M., Forstpointner, R., Metzner, B., Peter, N., Wormann, B., Trumper, L., Pfreundschuh, M., Einsele, H., Hiddemann, W., Unterhalt, M. & Dreyling, M. (2009) Improvement of overall survival in advanced stage mantle cell lymphoma. *J Clin Oncol*, **27**, 511-518.
- Hess, G., Herbrecht, R., Romaguera, J., Verhoef, G., Crump, M., Gisselbrecht, C., Laurell, A., Offner, F., Strahs, A., Berkenblit, A., Hanushevsky, O., Clancy, J., Hewes, B., Moore, L. & Coiffier, B. (2009) Phase III study to evaluate temsirolimus compared with investigator's choice therapy for the treatment of relapsed or refractory mantle cell lymphoma. *J Clin Oncol*, **27**, 3822-3829.

- Hosein, P.J., Pastorini, V.H., Paes, F.M., Eber, D., Chapman, J.R., Serafini, A.N., Alizadeh, A.A. & Lossos, I.S. (2011) Utility of positron emission tomography scans in mantle cell lymphoma. *Am J Hematol*, **86**, 841-845.
- Hoster, E. (2016) Role Of High-Dose Cytarabine and Total Body Irradiation Conditioning before Autologous Stem Cell Transplantation In Mantle Cell Lymphoma - A Comparison Of Nordic MCL2, HOVON 45, and European MCL Younger Trials. *Leukaemia*, **30**, 1428-1430
- Hoster, E. & Pott, C. (2016) Minimal residual disease in mantle cell lymphoma: insights into biology and impact on treatment. *Hematology Am Soc Hematol Educ Program*, **2016**, 437-445.
- Houot, R., Le Gouill, S., Ojeda Uribe, M., Mounier, C., Courby, S., Dartigeas, C., Bouabdallah, K., Alexis Vigier, M., Moles, M.P., Tournilhac, O., Arakelyan, N., Rodon, P., El Yamani, A., Sutton, L., Fornecker, L., Assouline, D., Harousseau, J.L., Maisonneuve, H., Caulet-Maugendre, S., Gressin, R. & French, G.g. (2012) Combination of rituximab, bortezomib, doxorubicin, dexamethasone and chlorambucil (RiPAD+C) as first-line therapy for elderly mantle cell lymphoma patients: results of a phase II trial from the GOELAMS. *Ann Oncol*, **23**, 1555-1561.
- Howard, O.M., Gribben, J.G., Neuberger, D.S., Grossbard, M., Poor, C., Janicek, M.J. & Shipp, M.A. (2002) Rituximab and CHOP induction therapy for newly diagnosed mantle-cell lymphoma: molecular complete responses are not predictive of progression-free survival. *J Clin Oncol*, **20**, 1288-1294.
- Inwards, D.J., Fishkin, P.A., Hillman, D.W., Brown, D.W., Ansell, S.M., Kurtin, P.J., Fonseca, R., Morton, R.F., Veeder, M.H. & Witzig, T.E. (2008) Long-term results of the treatment of patients with mantle cell lymphoma with cladribine (2-CDA) alone (95-80-53) or 2-CDA and rituximab (N0189) in the North Central Cancer Treatment Group. *Cancer*, **113**, 108-116.
- Jantunen, E., Canals, C., Attal, M., Thomson, K., Milpied, N., Buzyn, A., Ferrant, A., Biron, P., Crawley, C., Schattenberg, A., Luan, J.J., Tilly, H., Rio, B., Wijermans, P.W., Dreger, P., Sureda, A., Lymphoma Working Party of the European Group for, B. & Marrow, T. (2012) Autologous stem-cell transplantation in patients with mantle cell lymphoma beyond 65 years of age: a study from the European Group for Blood and Marrow Transplantation (EBMT). *Ann Oncol*, **23**, 166-171.
- Jares, P. & Campo, E. (2008) Advances in the understanding of mantle cell lymphoma. *Br J Haematol*, **142**, 149-165.
- Kahl, B.S., Longo, W.L., Eickhoff, J.C., Zehnder, J., Jones, C., Blank, J., McFarland, T., Bottner, W., Rezazadeh, H., Werndli, J., Bailey, H.H. & Wisconsin Oncology, N. (2006) Maintenance rituximab following induction chemoimmunotherapy may prolong progression-free survival in mantle cell lymphoma: a pilot study from the Wisconsin Oncology Network. *Ann Oncol*, **17**, 1418-1423.
- Kahl, B.S., Spurgeon, S.E., Furman, R.R., Flinn, I.W., Coutre, S.E., Brown, J.R., Benson, D.M., Byrd, J.C., Peterman, S., Cho, Y., Yu, A., Godfrey, W.R. & Wagner-Johnston, N.D. (2014) A phase 1 study of the PI3Kdelta inhibitor idelalisib in patients with relapsed/refractory mantle cell lymphoma (MCL). *Blood*, **123**, 3398-3405.
- Kaplan, L. (2015) Bortezomib maintenance(BM) versus consolidation (BC) following aggressive immunochemotherapy and autologous stem cell transplant (ASCT) for untreated mantle cell lymphoma (MCL):CALGB (Alliance) 50403. In: *57th ASH annual meeting*, Vol. 126, p. 337. Blood.
- Kaufmann, H., Raderer, M., Wohner, S., Puspok, A., Bankier, A., Zielinski, C., Chott, A. & Drach, J. (2004) Antitumor activity of rituximab plus thalidomide in patients with relapsed/refractory mantle cell lymphoma. *Blood*, **104**, 2269-2271.

- Khouri, I.F., Romaguera, J., Kantarjian, H., Palmer, J.L., Pugh, W.C., Korbling, M., Hagemester, F., Samuels, B., Rodriguez, A., Giralt, S., Younes, A., Przepiorka, D., Claxton, D., Cabanillas, F. & Champlin, R. (1998) Hyper-CVAD and high-dose methotrexate/cytarabine followed by stem-cell transplantation: an active regimen for aggressive mantle-cell lymphoma. *J Clin Oncol*, **16**, 3803-3809.
- Kirschbaum, M., Frankel, P., Popplewell, L., Zain, J., Delioukina, M., Pullarkat, V., Matsuoka, D., Pulone, B., Rotter, A.J., Espinoza-Delgado, I., Nademanee, A., Forman, S.J., Gandara, D. & Newman, E. (2011) Phase II study of vorinostat for treatment of relapsed or refractory indolent non-Hodgkin's lymphoma and mantle cell lymphoma. *J Clin Oncol*, **29**, 1198-1203.
- Kluin-Nelemans, H.C., Hoster, E., Hermine, O., Walewski, J., Trnety, M., Geisler, C.H., Stilgenbauer, S., Thieblemont, C., Vehling-Kaiser, U., Doorduijn, J.K., Coiffier, B., Forstpointner, R., Tilly, H., Kanz, L., Feugier, P., Szymczyk, M., Hallek, M., Kremers, S., Lepeu, G., Sanhes, L., Zijlstra, J.M., Bouabdallah, R., Lugtenburg, P.J., Macro, M., Pfreundschuh, M., Prochazka, V., Di Raimondo, F., Ribrag, V., Uppenkamp, M., Andre, M., Klapper, W., Hiddemann, W., Unterhalt, M. & Dreyling, M.H. (2012) Treatment of older patients with mantle-cell lymphoma. *N Engl J Med*, **367**, 520-531.
- Kolstad, A., Laurell, A., Jerkeman, M., Gronbaek, K., Elonen, E., Raty, R., Pedersen, L.B., Loft, A., Bogsrud, T.V., Kimby, E., Hansen, P.B., Fagerli, U.M., Nilsson-Ehle, H., Lauritzsen, G.F., Lehmann, A.K., Sundstrom, C., Karjalainen-Lindsberg, M.L., Ralfkiaer, E., Ehinger, M., Delabie, J., Bentzen, H., Schildt, J., Kostova-Aherdan, K., Frederiksen, H., Brown Pde, N., Geisler, C.H. & Nordic Lymphoma, G. (2014) Nordic MCL3 study: 90Y-ibritumomab-tiuxetan added to BEAM/C in non-CR patients before transplant in mantle cell lymphoma. *Blood*, **123**, 2953-2959.
- Kouroukis, C.T., Belch, A., Crump, M., Eisenhauer, E., Gascoyne, R.D., Meyer, R., Lohmann, R., Lopez, P., Powers, J., Turner, R., Connors, J.M. & National Cancer Institute of Canada Clinical Trials, G. (2003) Flavopiridol in untreated or relapsed mantle-cell lymphoma: results of a phase II study of the National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol*, **21**, 1740-1745.
- Kouroukis, C.T., Fernandez, L.A., Crump, M., Gascoyne, R.D., Chua, N.S., Buckstein, R., Turner, R., Assouline, S., Klasa, R.J., Walsh, W., Powers, J. & Eisenhauer, E. (2011) A phase II study of bortezomib and gemcitabine in relapsed mantle cell lymphoma from the National Cancer Institute of Canada Clinical Trials Group (IND 172). *Leuk Lymphoma*, **52**, 394-399.
- Lamm, W., Kaufmann, H., Raderer, M., Hoffmann, M., Chott, A., Zielinski, C. & Drach, J. (2011) Bortezomib combined with rituximab and dexamethasone is an active regimen for patients with relapsed and chemotherapy-refractory mantle cell lymphoma. *Haematologica*, **96**, 1008-1014.
- Le Gouill, S., Thieblemont, C., Oberic, L., Moreau, A., Bouabdallah, K., Dartigeas, C., Damaj, G., Gastinne, T., Ribrag, V., Feugier, P., Casasnovas, O., Zerazhi, H., Haioun, C., Maisonneuve, H., Houot, R., Jardin F., Van Den Neste, E., Tournilhac, O., Le Du, K., Morschhauser, F., Cartron, G., Fornecker, L. M., Canioni, D., Callanan, M., Bene, M. C., salles, G., Tilly, H., Lamy, T., Gressin, R., Hermine, O; LYSA Group (2017) Rituximab after Autologous Stem-Cell Transplantation in Mantle-Cell Lymphoma. *N Engl J Med*, **377**, 1250-1260.

- Le Gouill, S., Kroger, N., Dhedin, N., Nagler, A., Bouabdallah, K., Yakoub-Agha, I., Kanouni, T., Bulabois, C.E., Tournilhac, O., Buzyn, A., Rio, B., Moles, M.P., Shimoni, A., Bacher, U., Ocheni, S., Milpied, N., Harousseau, J.L., Moreau, P., Leux, C. & Mohty, M. (2012) Reduced-intensity conditioning allogeneic stem cell transplantation for relapsed/refractory mantle cell lymphoma: a multicenter experience. *Ann Oncol*, **23**, 2695-2703.
- Lefrere, F., Delmer, A., Suzan, F., Levy, V., Belanger, C., Djabbari, M., Arnulf, B., Damaj, G., Maillard, N., Ribrag, V., Janvier, M., Sebban, C., Casasnovas, R.O., Bouabdallah, R., Dreyfus, F., Verkarre, V., Delabesse, E., Valensi, F., McIntyre, E., Brousse, N., Varet, B. & Hermine, O. (2002) Sequential chemotherapy by CHOP and DHAP regimens followed by high-dose therapy with stem cell transplantation induces a high rate of complete response and improves event-free survival in mantle cell lymphoma: a prospective study. *Leukemia*, **16**, 587-593.
- Lenz, G., Dreyling, M., Hoster, E., Wormann, B., Duhrsen, U., Metzner, B., Eimermacher, H., Neubauer, A., Wandt, H., Steinhauer, H., Martin, S., Heidemann, E., Aldaoud, A., Parwaresch, R., Hasford, J., Unterhalt, M. & Hiddemann, W. (2005) Immunochemotherapy with rituximab and cyclophosphamide, doxorubicin, vincristine, and prednisone significantly improves response and time to treatment failure, but not long-term outcome in patients with previously untreated mantle cell lymphoma: results of a prospective randomized trial of the German Low Grade Lymphoma Study Group (GLSG). *J Clin Oncol*, **23**, 1984-1992.
- Maddocks, K., Christian, B., Jaglowski, S., Flynn, J., Jones, J.A., Porcu, P., Wei, L., Jenkins, C., Lozanski, G., Byrd, J.C. & Blum, K.A. (2015) A phase 1/1b study of rituximab, bendamustine, and ibrutinib in patients with untreated and relapsed/refractory non-Hodgkin lymphoma. *Blood*, **125**, 242-248.
- Maris, M.B., Sandmaier, B.M., Storer, B.E., Chauncey, T., Stuart, M.J., Maziarz, R.T., Agura, E., Langston, A.A., Pulsipher, M., Storb, R. & Maloney, D.G. (2004) Allogeneic hematopoietic cell transplantation after fludarabine and 2 Gy total body irradiation for relapsed and refractory mantle cell lymphoma. *Blood*, **104**, 3535-3542.
- Martin, P., Chadburn, A., Christos, P., Weil, K., Furman, R.R., Ruan, J., Elstrom, R., Niesvizky, R., Ely, S., Diliberto, M., Melnick, A., Knowles, D.M., Chen-Kiang, S., Coleman, M. & Leonard, J.P. (2009) Outcome of deferred initial therapy in mantle-cell lymphoma. *J Clin Oncol*, **27**, 1209-1213.
- Mato, A.R., Svoboda, J., Feldman, T., Zielonka, T., Agress, H., Panush, D., Miller, M., Toth, P., Lizotte, P.M., Nasta, S., Goldberg, S., Chong, E., Schuster, S., Pecora, A.L. & Goy, A. (2012) Post-treatment (not interim) positron emission tomography-computed tomography scan status is highly predictive of outcome in mantle cell lymphoma patients treated with R-HyperCVAD. *Cancer*, **118**, 3565-3570.
- McKay, P., Leach, M., Jackson, R., Cook, G., Rule, S. & British Committee for Standards in, H. (2012) Guidelines for the investigation and management of mantle cell lymphoma. *Br J Haematol*, **159**, 405-426.
- McKay, P., Leach, M., Jackson, R., Robinson, S. & Rule, S. (2018) A British Society for Haematology Good Practice Paper on the Diagnosis and Investigation of Patients with Mantle Cell Lymphoma. *Br J Haematol*, in press.
- Merli, F., Luminari, S., Ilariucci, F., Petrini, M., Visco, C., Ambrosetti, A., Stelitano, C., Caracciolo, F., Di Renzo, N., Angrilli, F., Carella, A.M., Capodanno, I., Barbolini, E., Galimberti, S. & Federico, M. (2012) Rituximab plus HyperCVAD alternating with

- high dose cytarabine and methotrexate for the initial treatment of patients with mantle cell lymphoma, a multicentre trial from Gruppo Italiano Studio Linfomi. *Br J Haematol*, **156**, 346-353.
- Meusers, P., Engelhard, M., Bartels, H., Binder, T., Fulle, H.H., Gorg, K., Gunzer, U., Havemann, K., Kayser, W., Konig, E. & et al. (1989) Multicentre randomized therapeutic trial for advanced centrocytic lymphoma: anthracycline does not improve the prognosis. *Hematol Oncol*, **7**, 365-380.
- Morrison, V.A., Jung, S.H., Johnson, J., LaCasce, A., Blum, K.A., Bartlett, N.L., Pitcher, B.N. & Cheson, B.D. (2015) Therapy with bortezomib plus lenalidomide for relapsed/refractory mantle cell lymphoma: final results of a phase II trial (CALGB 50501). *Leuk Lymphoma*, **56**, 958-964.
- Morschhauser, F., Depil, S., Jourdan, E., Wetterwald, M., Bouabdallah, R., Marit, G., Solal-Celigny, P., Sebban, C., Coiffier, B., Chouaki, N., Bauters, F. & Dumontet, C. (2007) Phase II study of gemcitabine-dexamethasone with or without cisplatin in relapsed or refractory mantle cell lymphoma. *Ann Oncol*, **18**, 370-375.
- Morschhauser, F., Seymour, J.F., Kluin-Nelemans, H.C., Grigg, A., Wolf, M., Pfreundschuh, M., Tilly, H., Raemaekers, J., van 't Veer, M.B., Milpied, N., Cartron, G., Pezzutto, A., Spencer, A., Reyes, F. & Dreyling, M. (2008) A phase II study of enzastaurin, a protein kinase C beta inhibitor, in patients with relapsed or refractory mantle cell lymphoma. *Ann Oncol*, **19**, 247-253.
- Morschhauser, F.A., Cartron, G., Thieblemont, C., Solal-Celigny, P., Haioun, C., Bouabdallah, R., Feugier, P., Bouabdallah, K., Asikanius, E., Lei, G., Wenger, M., Wassner-Fritsch, E. & Salles, G.A. (2013) Obinutuzumab (GA101) monotherapy in relapsed/refractory diffuse large b-cell lymphoma or mantle-cell lymphoma: results from the phase II GAUGUIN study. *J Clin Oncol*, **31**, 2912-2919.
- Nabhan, C., Smith, S.M., Helenowski, I., Ramsdale, E., Parsons, B., Karmali, R., Feliciano, J., Hanson, B., Smith, S., McKoy, J., Larsen, A., Hantel, A., Gregory, S. & Evens, A.M. (2012) Analysis of very elderly (≥ 80 years) non-hodgkin lymphoma: impact of functional status and co-morbidities on outcome. *Br J Haematol*, **156**, 196-204.
- Nickenig, C., Dreyling, M., Hoster, E., Pfreundschuh, M., Trumper, L., Reiser, M., Wandt, H., Lengfelder, E., Unterhalt, M., Hiddemann, W. & German Low-Grade Lymphoma Study, G. (2006) Combined cyclophosphamide, vincristine, doxorubicin, and prednisone (CHOP) improves response rates but not survival and has lower hematologic toxicity compared with combined mitoxantrone, chlorambucil, and prednisone (MCP) in follicular and mantle cell lymphomas: results of a prospective randomized trial of the German Low-Grade Lymphoma Study Group. *Cancer*, **107**, 1014-1022.
- Nodit, L., Bahler, D.W., Jacobs, S.A., Locker, J. & Swerdlow, S.H. (2003) Indolent mantle cell lymphoma with nodal involvement and mutated immunoglobulin heavy chain genes. *Hum Pathol*, **34**, 1030-1034.
- Orchard, J., Garand, R., Davis, Z., Babbage, G., Sahota, S., Matutes, E., Catovsky, D., Thomas, P.W., Avet-Loiseau, H. & Oscier, D. (2003) A subset of t(11;14) lymphoma with mantle cell features displays mutated IgVH genes and includes patients with good prognosis, nonnodal disease. *Blood*, **101**, 4975-4981.
- Peggs, K. (2016) Reduced intensity conditioning transplantation (RIC) as part of first line therapy for mantle cell lymphoma: results from the phase II Mini Allo trial (CRUK:C7627/A9080). In: *BMT*, Vol. 51.
- Ritchie, D.S., Seymour, J.F., Grigg, A.P., Roberts, A.W., Hoyt, R., Thompson, S., Szer, J. & Prince, H.M. (2007) The hyper-CVAD-rituximab chemotherapy programme followed by high-dose busulfan, melphalan and autologous stem cell transplantation produces

- excellent event-free survival in patients with previously untreated mantle cell lymphoma. *Ann Hematol*, **86**, 101-105.
- Robak, T., Huang, H., Jin, J., Zhu, J., Liu, T., Samoilova, O., Pylypenko, H., Verhoef, G., Siritanaratkul, N., Osmanov, E., Alexeeva, J., Pereira, J., Drach, J., Mayer, J., Hong, X., Okamoto, R., Pei, L., Rooney, B., van de Velde, H., Cavalli, F. & Investigators, L.Y.M. (2015) Bortezomib-based therapy for newly diagnosed mantle-cell lymphoma. *N Engl J Med*, **372**, 944-953.
- Robak, T., Smolewski, P., Cebula, B., Szmigielska-Kaplon, A., Chojnowski, K. & Blonski, J.Z. (2006) Rituximab combined with cladribine or with cladribine and cyclophosphamide in heavily pretreated patients with indolent lymphoproliferative disorders and mantle cell lymphoma. *Cancer*, **107**, 1542-1550.
- Robinson, S., Boumendil, A., Finel, H., Thomson, K., Chevallier, P., Sierra, J., Finke, J., Poiré, X., Maillard, N., Milpied, N., Yakoub-Agha, I., Koh, M., Kröger, N., Nagler, A., Koc, Y., Dietrich, S., Dreger, P. & Montoto, S. (2017) Reduced Intensity Allogeneic Stem Cell Transplantation For Mantle Cell Lymphoma: Long Term Outcomes. *BMT*, **52**, S17–S115
- Romaguera, J.E., Fayad, L.E., Feng, L., Hartig, K., Weaver, P., Rodriguez, M.A., Hagemester, F.B., Pro, B., McLaughlin, P., Younes, A., Samaniego, F., Goy, A., Cabanillas, F., Kantarjian, H., Kwak, L. & Wang, M. (2010) Ten-year follow-up after intense chemoimmunotherapy with Rituximab-HyperCVAD alternating with Rituximab-high dose methotrexate/cytarabine (R-MA) and without stem cell transplantation in patients with untreated aggressive mantle cell lymphoma. *Br J Haematol*, **150**, 200-208.
- Romaguera, J.E., Medeiros, L.J., Hagemester, F.B., Fayad, L.E., Rodriguez, M.A., Pro, B., Younes, A., McLaughlin, P., Goy, A., Sarris, A.H., Dang, N.H., Samaniego, F., Brown, H.M., Gagneja, H.K. & Cabanillas, F. (2003) Frequency of gastrointestinal involvement and its clinical significance in mantle cell lymphoma. *Cancer*, **97**, 586-591.
- Rosenbluth, B.D. & Yahalom, J. (2006) Highly effective local control and palliation of mantle cell lymphoma with involved-field radiation therapy (IFRT). *Int J Radiat Oncol Biol Phys*, **65**, 1185-1191.
- Ruan, J., Martin, P., Coleman, M., Furman, R.R., Cheung, K., Faye, A., Elstrom, R., Lachs, M., Hajjar, K.A. & Leonard, J.P. (2010) Durable responses with the metronomic rituximab and thalidomide plus prednisone, etoposide, procarbazine, and cyclophosphamide regimen in elderly patients with recurrent mantle cell lymphoma. *Cancer*, **116**, 2655-2664.
- Rubio, M. (2010) Is There Still a Place for Total Body Irradiation (TBI) In the Conditioning Regimen of Autologous Stem Cell Transplantation In Mantle Cell Lymphoma?: a Retrospective Study From the Lymphoma Working Party of the EBMT. In: *52nd ASH annual meeting*, Vol. 116. Blood.
- Rule, S., Dreyling, M., Goy, A., Hess, G., Auer, R., Kahl, B., Cavazos, N., Liu, B., Yang, S., Clow, F., Goldberg, J.D., Beaupre, D., Vermeulen, J., Wildgust, M. & Wang, M. (2017) Outcomes in 370 patients with mantle cell lymphoma treated with ibrutinib: a pooled analysis from three open-label studies. *Br J Haematol*, **179**, 430-438.
- Rule, S., Smith, P., Johnson, P.W., Bolam, S., Follows, G., Gambell, J., Hillmen, P., Jack, A., Johnson, S., Kirkwood, A.A., Kruger, A., Pocock, C., Seymour, J.F., Toncheva, M., Walewski, J. & Linch, D. (2016) The addition of rituximab to fludarabine and cyclophosphamide chemotherapy results in a significant improvement in overall survival in patients with newly diagnosed mantle cell lymphoma: results of a

- randomized UK National Cancer Research Institute trial. *Haematologica*, **101**, 235-240.
- Rummel, M.J., Al-Batran, S.E., Kim, S.Z., Welslau, M., Hecker, R., Kofahl-Krause, D., Josten, K.M., Durk, H., Rost, A., Neise, M., von Grunhagen, U., Chow, K.U., Hansmann, M.L., Hoelzer, D. & Mitrou, P.S. (2005) Bendamustine plus rituximab is effective and has a favorable toxicity profile in the treatment of mantle cell and low-grade non-Hodgkin's lymphoma. *J Clin Oncol*, **23**, 3383-3389.
- Rummel, M.J., Niederle, N., Maschmeyer, G., Banat, G.A., von Grunhagen, U., Losem, C., Kofahl-Krause, D., Heil, G., Welslau, M., Balsler, C., Kaiser, U., Weidmann, E., Durk, H., Ballo, H., Stauch, M., Roller, F., Barth, J., Hoelzer, D., Hinke, A., Brugger, W. & Study group indolent, L. (2013) Bendamustine plus rituximab versus CHOP plus rituximab as first-line treatment for patients with indolent and mantle-cell lymphomas: an open-label, multicentre, randomised, phase 3 non-inferiority trial. *Lancet*, **381**, 1203-1210.
- Sachanas, S., Pangalis, G.A., Vassilakopoulos, T.P., Korkolopoulou, P., Kontopidou, F.N., Athanasoulia, M., Yiakoumis, X., Kalpadakis, C., Georgiou, G., Masouridis, S., Moschogiannis, M., Tsirkinidis, P., Pappis, V., Kokoris, S.I., Siakantaris, M.P., Panayiotidis, P. & Angelopoulou, M.K. (2011) Combination of rituximab with chlorambucil as first line treatment in patients with mantle cell lymphoma: a highly effective regimen. *Leuk Lymphoma*, **52**, 387-393.
- Schulz, H., Bohlius, J.F., Trelle, S., Skoetz, N., Reiser, M., Kober, T., Schwarzer, G., Herold, M., Dreyling, M., Hallek, M. & Engert, A. (2007) Immunochemotherapy with rituximab and overall survival in patients with indolent or mantle cell lymphoma: a systematic review and meta-analysis. *J Natl Cancer Inst*, **99**, 706-714.
- Shrestha, R., Bhatt, V.R., Guru Murthy, G.S. & Armitage, J.O. (2015) Clinicopathologic features and management of blastoid variant of mantle cell lymphoma. *Leuk Lymphoma*, **56**, 2759-2767.
- Smith, M.R., Li, H., Gordon, L., Gascoyne, R.D., Paietta, E., Forero-Torres, A., Kahl, B.S., Advani, R., Hong, F. & Horning, S.J. (2012) Phase II study of rituximab plus cyclophosphamide, doxorubicin, vincristine, and prednisone immunochemotherapy followed by yttrium-90-ibritumomab tiuxetan in untreated mantle-cell lymphoma: Eastern Cooperative Oncology Group Study E1499. *J Clin Oncol*, **30**, 3119-3126.
- Sorrer, M.L., Maris, M.B., Storb, R., Baron, F., Sandmaier, B.M., Maloney, D.G. & Storer, B. (2005) Hematopoietic cell transplantation (HCT)-specific comorbidity index: a new tool for risk assessment before allogeneic HCT. *Blood*, **106**, 2912-2919.
- Sorrer, M.L., Storb, R.F., Sandmaier, B.M., Maziarz, R.T., Pulsipher, M.A., Maris, M.B., Bhatia, S., Ostronoff, F., Deeg, H.J., Syrjala, K.L., Estey, E., Maloney, D.G., Appelbaum, F.R., Martin, P.J. & Storer, B.E. (2014) Comorbidity-age index: a clinical measure of biologic age before allogeneic hematopoietic cell transplantation. *J Clin Oncol*, **32**, 3249-3256.
- Spurgeon, S.E., Pindyck, T., Okada, C., Chen, Y., Chen, Z., Mater, E., Abbi, K. & Epner, E.M. (2011) Cladribine plus rituximab is an effective therapy for newly diagnosed mantle cell lymphoma. *Leuk Lymphoma*, **52**, 1488-1494.
- Swerdlow, S.H., Habeshaw, J.A., Murray, L.J., Dhaliwal, H.S., Lister, T.A. & Stansfeld, A.G. (1983) Centrocytic lymphoma: a distinct clinicopathologic and immunologic entity. A multiparameter study of 18 cases at diagnosis and relapse. *Am J Pathol*, **113**, 181-197.
- Tam, C.S., Bassett, R., Ledesma, C., Korbling, M., Alousi, A., Hosing, C., Kebraei, P., Harrell, R., Rondon, G., Giralt, S.A., Anderlini, P., Popat, U., Pro, B., Samuels, B., Hagemester, F., Medeiros, L.J., Champlin, R.E. & Khouri, I.F. (2009) Mature results

- of the M. D. Anderson Cancer Center risk-adapted transplantation strategy in mantle cell lymphoma. *Blood*, **113**, 4144-4152.
- Teodorovic, I., Pittaluga, S., Kluin-Nelemans, J.C., Meerwaldt, J.H., Hagenbeek, A., van Glabbeke, M., Somers, R., Bijmens, L., Noordijk, E.M. & Peeters, C.D. (1995) Efficacy of four different regimens in 64 mantle-cell lymphoma cases: clinicopathologic comparison with 498 other non-Hodgkin's lymphoma subtypes. European Organization for the Research and Treatment of Cancer Lymphoma Cooperative Group. *J Clin Oncol*, **13**, 2819-2826.
- Thomas, D.W., Owen, R.G., Johnson, S.A., Hillmen, P., Seymour, J.F., Wolf, M.M. & Rule, S.A. (2005) Superior quality and duration of responses among patients with mantle-cell lymphoma treated with fludarabine and cyclophosphamide with or without rituximab compared with prior responses to CHOP. *Leuk Lymphoma*, **46**, 549-552.
- Tiemann, M., Schrader, C., Klapper, W., Dreyling, M.H., Campo, E., Norton, A., Berger, F., Kluin, P., Ott, G., Pileri, S., Pedrinis, E., Feller, A.C., Merz, H., Janssen, D., Hansmann, M.L., Krieken, H., Moller, P., Stein, H., Unterhalt, M., Hiddemann, W., Parwaresch, R. & European, M.C.L.N. (2005) Histopathology, cell proliferation indices and clinical outcome in 304 patients with mantle cell lymphoma (MCL): a clinicopathological study from the European MCL Network. *Br J Haematol*, **131**, 29-38.
- Till, B.G., Gooley, T.A., Crawford, N., Gopal, A.K., Maloney, D.G., Petersdorf, S.H., Pagel, J.M., Holmberg, L., Bensinger, W. & Press, O.W. (2008) Effect of remission status and induction chemotherapy regimen on outcome of autologous stem cell transplantation for mantle cell lymphoma. *Leuk Lymphoma*, **49**, 1062-1073.
- Tucker, D.L., Naylor, G., Kruger, A., Hamilton, M.S., Follows, G. & Rule, S.A. (2016) Ibrutinib is a safe and effective therapy for systemic mantle cell lymphoma with central nervous system involvement - a multi-centre case series from the United Kingdom. *Br J Haematol*.
- Unterhalt, M., Herrmann, R., Tiemann, M., Parwaresch, R., Stein, H., Trumper, L., Nahler, M., Reuss-Borst, M., Tirier, C., Neubauer, A., Freund, M., Kreuser, E.D., Dietzfelbinger, H., Bodenstern, H., Engert, A., Stauder, R., Eimermacher, H., Landys, K. & Hiddemann, W. (1996) Prednimustine, mitoxantrone (PmM) vs cyclophosphamide, vincristine, prednisone (COP) for the treatment of advanced low-grade non-Hodgkin's lymphoma. German Low-Grade Lymphoma Study Group. *Leukemia*, **10**, 836-843.
- Vandenberghe, E., De Wolf-Peeters, C., Vaughan Hudson, G., Vaughan Hudson, B., Pittaluga, S., Anderson, L. & Linch, D.C. (1997) The clinical outcome of 65 cases of mantle cell lymphoma initially treated with non-intensive therapy by the British National Lymphoma Investigation Group. *Br J Haematol*, **99**, 842-847.
- Vandenberghe, E., Ruiz de Elvira, C., Loberiza, F.R., Conde, E., Lopez-Guillermo, A., Gisselbrecht, C., Guilhot, F., Vose, J.M., van Biesen, K., Rizzo, J.D., Weisenburger, D.D., Isaacson, P., Horowitz, M.M., Goldstone, A.H., Lazarus, H.M. & Schmitz, N. (2003) Outcome of autologous transplantation for mantle cell lymphoma: a study by the European Blood and Bone Marrow Transplant and Autologous Blood and Marrow Transplant Registries. *Br J Haematol*, **120**, 793-800.
- Vidal, L. (2016) Maintenance therapy for patients with mantle cell lymphoma (MCL) - a systematic review and meta-analysis of randomized controlled trials (RCTs). In: *58th Annual ASH Meeting*, Vol. 128. Blood.
- Visco, C., Finotto, S., Zambello, R., Paolini, R., Menin, A., Zanotti, R., Zaja, F., Semenzato, G., Pizzolo, G., D'Amore, E.S. & Rodeghiero, F. (2013) Combination of rituximab,

- bendamustine, and cytarabine for patients with mantle-cell non-Hodgkin lymphoma ineligible for intensive regimens or autologous transplantation. *J Clin Oncol*, **31**, 1442-1449.
- Visco, C., Chiappella, A., Nassa, L., Patti, C., Ferrero, S., Barbero, D., Evangelista, A., Spina, M., Molinari, A., Rigacci, L., Tani, M., Di Rocco, A., Pinotti, G., Fabbri, A., Zambello, R., Finotto, S., Gotti, M., Carella, A. M., Salvi, F., Pileri, S. A., Ladetto, M., Ciccone, G., Gaidano, G., Ruggeri, M., Martelli, M. & Vitolo, U. (2017) Rituximab, bendamustine and low-dose cytarabine as induction therapy in elderly patients with mantle cell lymphoma: a multicentre, phase 2 trial from Fondazione Italiana Linfomi. *Lancet Haematol*, **4**, e15-23.
- Walter, H.S., Rule, S.A., Dyer, M.J., Karlin, L., Jones, C., Cazin, B., Quittet, P., Shah, N., Hutchinson, C.V., Honda, H., Duffy, K., Birkett, J., Jamieson, V., Courtenay-Luck, N., Yoshizawa, T., Sharpe, J., Ohno, T., Abe, S., Nishimura, A., Cartron, G., Morschhauser, F., Fegan, C. & Salles, G. (2016) A phase 1 clinical trial of the selective BTK inhibitor ONO/GS-4059 in relapsed and refractory mature B-cell malignancies. *Blood*, **127**, 411-419.
- Wang, M., Fayad, L., Wagner-Bartak, N., Zhang, L., Hagemester, F., Neelapu, S.S., Samaniego, F., McLaughlin, P., Fanale, M., Younes, A., Cabanillas, F., Fowler, N., Newberry, K.J., Sun, L., Young, K.H., Champlin, R., Kwak, L., Feng, L., Badillo, M., Bejarano, M., Hartig, K., Chen, W., Chen, Y., Byrne, C., Bell, N., Zeldis, J. & Romaguera, J. (2012) Lenalidomide in combination with rituximab for patients with relapsed or refractory mantle-cell lymphoma: a phase 1/2 clinical trial. *Lancet Oncol*, **13**, 716-723.
- Wang, M., Popplewell, L.L., Collins, R.H., Jr., Winter, J.N., Goy, A., Kaminski, M.S., Bartlett, N.L., Johnston, P.B., Lister, J., Fanning, S.R., Tuscano, J.M., Beck, J.T., Kaya, H., Robeva, A., Fan, J., Klimovsky, J., Cheung, W., Cherfi, A. & O'Connor, O.A. (2014) Everolimus for patients with mantle cell lymphoma refractory to or intolerant of bortezomib: multicentre, single-arm, phase 2 study. *Br J Haematol*, **165**, 510-518.
- Wang, M.L., Lee, H., Chuang, H., Wagner-Bartak, N., Hagemester, F., Westin, J., Fayad, L., Samaniego, F., Turturro, F., Oki, Y., Chen, W., Badillo, M., Nomie, K., DeLa Rosa, M., Zhao, D., Lam, L., Addison, A., Zhang, H., Young, K.H., Li, S., Santos, D., Medeiros, L.J., Champlin, R., Romaguera, J. & Zhang, L. (2016) Ibrutinib in combination with rituximab in relapsed or refractory mantle cell lymphoma: a single-centre, open-label, phase 2 trial. *Lancet Oncol*, **17**, 48-56.
- Wang, M.L., Rule, S., Martin, P., Goy, A., Auer, R., Kahl, B.S., Jurczak, W., Advani, R.H., Romaguera, J.E., Williams, M.E., Barrientos, J.C., Chmielowska, E., Radford, J., Stilgenbauer, S., Dreyling, M., Jdrzejczak, W.W., Johnson, P., Spurgeon, S.E., Li, L., Zhang, L., Newberry, K., Ou, Z., Cheng, N., Fang, B., McGreivy, J., Clow, F., Buggy, J.J., Chang, B.Y., Beaupre, D.M., Kunkel, L.A. & Blum, K.A. (2013) Targeting BTK with ibrutinib in relapsed or refractory mantle-cell lymphoma. *N Engl J Med*, **369**, 507-516.
- Wang, M., Rule, S., Zinzani, P.L., Goy, A., Casasnovas, O., Smith, S.D., Damaj, G., Doorduijn, J., Lamy, T., Morschhauser, F., Panizo, C., Shah, B., Davies, A., Eek, R., Dupuis, J., Jacobsen, E., Kater, A.P., Le Gouill, S., Oberic, L., Robak, T., Covey, T., Dua, R., Hamdy, A., Huang, X., Izumi, R., Patel, P., Rothbaum, W., Slatter, J.G. & Jurczak, W. (2017) Acalabrutinib in relapsed or refractory mantle cell lymphoma (ACE-LY-004): a single-arm, multicentre, phase 2 trial. *Lancet*. Dec 11. pii: S0140-6736(17)33108-2. doi: 10.1016/S0140-6736(17)33108-2.

- Weigert, O., Unterhalt, M., Hiddemann, W. & Dreyling, M. (2007) Current management of mantle cell lymphoma. *Drugs*, **67**, 1689-1702.
- Weigert, O., Weidmann, E., Mueck, R., Bentz, M., von Schilling, C., Rohrberg, R., Jentsch-Ullrich, K., Hiddemann, W. & Dreyling, M. (2009) A novel regimen combining high dose cytarabine and bortezomib has activity in multiply relapsed and refractory mantle cell lymphoma - long-term results of a multicenter observation study. *Leuk Lymphoma*, **50**, 716-722.
- Weisenburger, D.D., Vose, J.M., Greiner, T.C., Lynch, J.C., Chan, W.C., Bierman, P.J., Dave, B.J., Sanger, W.G. & Armitage, J.O. (2000) Mantle cell lymphoma. A clinicopathologic study of 68 cases from the Nebraska Lymphoma Study Group. *Am J Hematol*, **64**, 190-196.
- William, B.M., Allen, M.S., Loberiza, F.R., Jr., Bociek, R.G., Bierman, P.J., Armitage, J.O. & Vose, J.M. (2014) Phase I/II study of bortezomib-BEAM and autologous hematopoietic stem cell transplantation for relapsed indolent non-Hodgkin lymphoma, transformed, or mantle cell lymphoma. *Biol Blood Marrow Transplant*, **20**, 536-542.
- Zaja, F., De Luca, S., Vitolo, U., Orsucci, L., Levis, A., Salvi, F., Rusconi, C., Ravelli, E., Tucci, A., Bottelli, C., Balzarotti, M., Brusamolino, E., Bonfichi, M., Pileri, S.A., Sabbatini, E., Volpetti, S., Monagheddu, C., Vacca, A., Ria, R. & Fanin, R. (2012) Salvage treatment with lenalidomide and dexamethasone in relapsed/refractory mantle cell lymphoma: clinical results and effects on microenvironment and neo-angiogenic biomarkers. *Haematologica*, **97**, 416-422.
- Zinzani, P.L., Magagnoli, M., Moretti, L., Battista, R., Ronconi, F., De Renzo, A., Zaccaria, A., Gentilini, P., Guardigni, L., Gherlinzoni, F., Cellini, C., Fattori, P.P., Bendandi, M., Bocchia, M., Aitini, E. & Tura, S. (1999) Fludarabine-based chemotherapy in untreated mantle cell lymphomas: an encouraging experience in 29 patients. *Haematologica*, **84**, 1002-1006.
- Zinzani, P.L., Magagnoli, M., Moretti, L., De Renzo, A., Battista, R., Zaccaria, A., Guardigni, L., Mazza, P., Marra, R., Ronconi, F., Lauta, V.M., Bendandi, M., Gherlinzoni, F., Gentilini, P., Ciccone, F., Cellini, C., Stefoni, V., Ricciuti, F., Gobbi, M. & Tura, S. (2000) Randomized trial of fludarabine versus fludarabine and idarubicin as frontline treatment in patients with indolent or mantle-cell lymphoma. *J Clin Oncol*, **18**, 773-779.
- Zucca, E., Roggero, E., Pinotti, G., Pedrinis, E., Cappella, C., Venco, A. & Cavalli, F. (1995) Patterns of survival in mantle cell lymphoma. *Ann Oncol*, **6**, 257-262.