



Burdett, H. L., Perna, G., McKay, L., Broomhead, G. and Kamenos, N. A. (2018)
Community-level sensitivity of a calcifying ecosystem to acute in situ CO₂ enrichment.
Marine Ecology Progress Series, 587, pp. 73-80.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/162830/>

Deposited on: 29 May 2018

Enlighten – Research publications by members of the University of Glasgow_
<http://eprints.gla.ac.uk>

1 **Community-level sensitivity of a calcifying ecosystem to acute in situ CO₂ enrichment**

2 Heidi L. Burdett^{1,2*}, Gabriela Perna^{3,4}, Lucy McKay^{5,6}, Gemma Broomhead⁵, Nicholas A.

3 Kamenos⁴

4 ¹ Lyell Centre for Earth and Marine Science and Technology, Edinburgh, UK, EH 14 4AP

5 ² School of Energy, Geoscience, Infrastructure and Society, Heriot-Watt University, UK,

6 EH14 4AS

7 ³ School of Geographical and Earth Science, University of Glasgow, UK, G12 8QQ

8 ⁴ King Abdullah University for Science and Technology, Saudi Arabia, 23955

9 ⁵ School of Earth & Environmental Sciences, University of St Andrews, UK, KY16 9AL

10 ⁶ School of Engineering, University of Glasgow, Glasgow, UK, G12 8QQ

11

12 * Corresponding author: E: h.burdett@hw.ac.uk

13

14 Running head: CO₂ enrichment of calcifying ecosystem

15

16 **Abstract**

17 The rate of change in ocean carbonate chemistry is a vital determinant in the magnitude of
18 impacts observed. Benthic marine ecosystems are facing an increasing risk of acute CO₂
19 exposure, that may be natural or anthropogenically-derived (e.g. engineering and industrial
20 activities). However, our understanding of how acute CO₂ events impact marine life is
21 restricted to individual organisms, with little understanding for how this manifests at the
22 community level. Here, we investigated, in situ, the effect of acute CO₂ enrichment on the
23 coralline algal ecosystem - a globally ubiquitous, ecologically and economically important
24 habitat, but one which is likely to be sensitive to CO₂ enrichment due to its highly calcified
25 reef-like structures engineered by coralline algae. Most notably, we observed a rapid
26 community-level shift to favour net dissolution rather than net calcification. Smaller changes
27 from net respiration to net photosynthesis were also observed. There was no effect on the net
28 flux of dimethylsulphide / dimethylsulphoniopropionate (algal secondary metabolites), nor
29 the nutrients nitrate and phosphate. Following return to ambient CO₂ levels, only a partial
30 recovery was seen within the monitoring timeframe. This study highlights the sensitivity of
31 biogenic carbonate marine communities to acute CO₂ enrichment, and raises concerns over
32 the capacity for the system to ‘bounce back’ if subjected to repeated acute high-CO₂ events.

33

34 **Keywords:** calcification, photosynthesis, community, ecosystem, maerl bed, carbon dioxide,
35 acidification

36 **Introduction**

37 Long-term environmental change as a result of rising atmospheric CO₂ levels are
38 projected to have significant impacts on marine organisms, especially those with calcified
39 body parts (Kroeker et al. 2010). Simultaneously, the risk of exposure to acute periods of
40 high-CO₂ conditions is also increasing, due to coastal / marine processes (e.g. tides (Abril et
41 al.), upwelling (Lachkar 2014)), land runoff (Strong et al. 2014) and the development of
42 engineering activities such as carbon capture and storage (Blackford et al. 2015). Research
43 has shown that the rate of environmental change is critical in determining the extent of
44 organismal damage, and that acute high-CO₂ exposures can have long-lasting effects (Burdett
45 et al. 2012, Kamenos et al. 2013). However, our understanding of how marine ecosystems
46 (rather than individuals) impact, and are impacted by, acute changes in ocean carbon
47 chemistry is poorly understood (Pfister et al. 2014). This is despite the known importance of
48 key biological processes such as calcification, photosynthesis, respiration and nutrient uptake
49 in driving marine ecosystem variability.

50 In the natural environment, an organism's response to environmental change is mediated
51 by community dynamics within the ecosystem. Failure to take these community-level
52 interactions into account prevents macro-scale predictions of future ecosystem change
53 (Queirós et al. 2014). To date, the majority of acute or chronic environmental change
54 experiments have focused on one, or maybe two, environmental factors (e.g. increased CO₂ /
55 temperature), and consider organisms in isolation (Riebesell & Gattuso 2015). However,
56 whilst informing our mechanistic understanding of physiological responses, these types of
57 experiments are not representative of real-world impacts due to laboratory artefacts and the
58 lack of appreciation for community-wide interactions (Cornwall & Hurd 2015, Riebesell &
59 Gattuso 2015). Consequently, efforts in developing methods for in situ experimentation have
60 recently increased.

61 Natural CO₂ vents, where the water column is enriched with CO₂ due to benthic bubbling
62 of volcanic gases, have proven useful for understanding the impacts of long-term exposure to
63 a high CO₂ environment on marine ecosystem structure (Hall-Spencer et al. 2008, Fabricius
64 et al. 2011, Kamenos et al. 2016). However, these study areas are typically characterised by
65 conditions more extreme or more variable than those predicted for the future, due to variation
66 in physical factors such as water currents and venting rates (Hall-Spencer et al. 2008). ‘Free
67 Ocean CO₂ Enrichment’ (FOCE) experimental setups attempt to bridge the gap between the
68 precise control of laboratory experiments and the natural setting of CO₂ vents (Gattuso et al.
69 2014), by artificially exposing organisms or communities to a high CO₂ environment. This
70 also allows the effects of both chronic and acute CO₂ enrichment to be tested. Partially-
71 artificial designs (where organisms are manually placed in the chambers, rather than
72 examining the natural system) have been conducted on tropical reefs (Kline et al. 2012) and
73 in the deep sea (Barry et al. 2014), whilst smaller chambers deployed on tropical seagrass
74 beds have investigated the community-level response of this vegetated habitat to short-term
75 CO₂ enrichment (e.g. Campbell & Fourqurean 2014).

76 One of the potentially most susceptible groups of organisms to both long and short-term
77 CO₂ enrichment are the red coralline algae (Kroeker et al. 2010) – key ecosystem engineers
78 in the coastal zone (Riosmena-Rodríguez 2017). Coralline algal beds – supported by a free-
79 living coralline algal framework – are globally distributed (van der Heijden & Kamenos
80 2015), highly diverse (BIOMAERL 1999, Barbera et al. 2003) and biogeochemically active
81 (Burdett et al. 2015b, van der Heijden & Kamenos). However, the community susceptibility
82 of coralline algal habitats is currently unknown, despite the real-world relevance of this
83 question compared to laboratory-based single organism studies (Gattuso et al. 2014).
84 Coralline algal beds are listed as ‘Vulnerable’ or ‘Endangered’ by the IUCN (Gubbay et al.
85 2016), a status driven by the sensitivity of coralline algae to environmental change, but also

86 due to the paucity of data available on the functioning of these habitats at the community
87 level.

88 Our understanding of coralline algal community functioning remains limited, even under
89 ambient conditions. Despite substantial gross primary production, coralline algal
90 communities exhibit net heterotrophy (i.e. O₂ uptake; Attard et al. 2015), acting as both a
91 CO₂ source (Martin et al. 2007a) and organic carbon sink (Attard et al. 2015). While nutrient
92 availability is not thought to limit the growth of coralline algal ecosystems (Steller et al
93 2009), there is evidence that coralline algal communities act as a nutrient source, at least in
94 the Mediterranean (Martin et al. 2007b). Coralline algae also represent a globally significant
95 stock of dimethylsulphoniopropionate (DMSP; Burdett et al. 2015a) – an algal secondary
96 metabolite that is the major precursor to the climate-gas dimethylsulphide (DMS). DMSP and
97 DMS (DMS/P) drive a range of community interactions (e.g. grazing behaviour; Lyons et al.
98 2007), but it is not yet known if coralline algal communities are a net source or sink of these
99 compounds. At an individual level, we know that CO₂ enrichment can affect the
100 photosynthesis, calcification and DMSP production of coralline algae (Burdett et al. 2012;
101 Kamenos et al. 2013), but it is not currently understood how this is manifest at a community
102 level, despite the significant implications for ecosystem functioning.

103 Here, we investigated the effect of acute in situ CO₂ enrichment on key community-level,
104 biologically-driven processes in a temperate coralline algal bed. Periodic CO₂ enrichment is a
105 risk to marine habitats in this region due to the prevalence of human activities such as
106 aquaculture – a rapidly expanding industry in Scotland and globally (OECD-FAO 2014).
107 Diel-scale pulsed release of CO₂ can occur from aquaculture infrastructures due to periodicity
108 in fish metabolism, e.g. after feeding (Forsberg 1997, Zakeś et al. 2003). In addition, the
109 development of carbon capture and storage facilities may further accentuate the risk of
110 periodic acute CO₂ release in the future (Blackford et al. 2015). *Lithothamnion glaciale*, the

111 coralline algal ecosystem engineer of this system, is known to be highly sensitive to acute
112 CO₂ exposure (Burdett et al. 2012, Kamenos et al. 2013), but sensitivity at a community level
113 remains unclear. Here, we investigated the integrated community-level response of a
114 coralline algal habitat to short-term CO₂ enrichment via in situ experimentation.

115

116 **Materials and Methods**

117 **Study site and experimental set-up**

118 The experiment was performed on a coralline algal bed in Loch Sween, on the west
119 coast of Scotland, UK, at a depth of 6 m. The ecosystem framework is dominated by the free-
120 living non-geniculate red coralline alga *Lithothamnion glaciale*, supporting a highly diverse
121 community across multiple trophic levels. This includes both calcified and non-calcified
122 macroalgae (including Laminariales) and invertebrates, being particularly rich in Mollusca
123 (e.g. *Aequipecten opercularis* - queen scallops [~ 4 per 20 m^2]) and particularly abundant in
124 Ophiuroidea (sea stars & brittle stars, e.g. *Ophiocomina nigra* [up to 10,000 per m^2] and
125 *Asterias rubens* [~ 11 per 20 m^2]) (BIOMAERL 1999, Barbera et al. 2003, Kamenos 2004).
126 Community biodiversity was not further quantified in this study. Four benthic chambers (28
127 litre volume, diameter = 38 cm) were deployed within the coralline algal bed by SCUBA
128 divers pushing them into the seabed. Chambers were left open for 24 hours to allow the water
129 within the chambers to equilibrate with the surrounding environment. Following
130 equilibration, lids were fitted and the experiment begun, which consisted of three phases: (1)
131 before CO_2 enrichment at ambient (control) conditions (15 hours), (2) during CO_2 enrichment
132 (28 hours) and (3) post-enrichment recovery (37 hours).

133 Chambers were individually connected to the surface via a flow-through system,
134 which continually pumped water through the chamber via the surface at a rate of 120 L hr^{-1}
135 (Swell UK Filter pump 5000). Pumps were located perpendicular from the chambers in
136 relation to the tidal current, to prevent the re-pumping of water through the system. CO_2
137 enrichment was achieved by bubbling pure CO_2 directly into a mixing chamber on the
138 surface, prior to the water being directed to the main in situ chambers. pH (total scale) of
139 water in the mixing chamber was monitored using a pH probe (VitalSINE, daily 3 point

140 calibration following the manufacturer's instructions) and the rate of CO₂ bubbling was
141 adjusted as required to maintain a stable ~0.2 pH unit offset from the incoming water supply.
142 Actual pH change in the chambers (reflecting both the CO₂ addition and biogeochemical
143 community processes) was determined by sampling the in-chamber water during the
144 experimental periods and analysing for total alkalinity (A_T) and dissolved inorganic carbon
145 (C_T), from which pH is calculated (details below). Flow-through circulation was maintained
146 for the duration of the experiment, except during 2-hour incubation periods when the water
147 flow was stopped, but within-chamber circulation was maintained by stirring paddles (Attard
148 et al. 2015). Water samples were taken for determination of dissolved oxygen, carbonate
149 chemistry, nutrients and dimethylated sulphur at the beginning and end of a 2-hour incubation
150 periods, which was carried out every ~12 hours during the experiment (i.e. around midday
151 and midnight during the three experimental phases). Measurements from the beginning and
152 end of the incubation were used for the determination of seabed flux measurements of each
153 parameter to gain understanding of the community response to CO₂ enrichment. All water
154 samples were collected in borosilicate glass syringes using SCUBA diving. Immediately after
155 collection, water samples were returned to the shore and prepared for various water chemistry
156 parameters, as detailed below.

157 **Net photosynthesis / respiration (dissolved oxygen)**

158 Winkler reagents (200 µl each of 3M MnSO₄.H₂O solution and 200 µl of 8M NaOH+4M
159 NaI) were added to 12 ml unfiltered water samples for subsequent dissolved oxygen (DO)
160 determination, and stored in the dark at 4°C until analysis. DO concentrations were
161 determined using the Winkler titration method (Grasshoff et al. 2007): The sample was
162 acidified with 200 µl 5M sulphuric acid and titrated against 0.05M sodium thiosulphate
163 solution with potassium iodate as a standard.

164 **Net calcification / dissolution (carbonate chemistry)**

165 Samples for A_T and C_T were stored in borosilicate glass vials (Labco Ltd, UK) and poisoned
166 with mercuric chloride, following Dickson et al. (2007). A_T was measured on a Metrohm 848
167 Titrino Plus using the 2-stage open-cell potentiometric titration method on 10 ml sample
168 volumes with 0.01 M HCl (Dickson et al. 2007). All A_T samples were analysed at $25 \pm 0.1^\circ\text{C}$
169 with temperature regulation using a water-bath (Julabo 19). C_T was determined by infra-red
170 detection of CO_2 from acidified samples on a dissolved inorganic carbon analyser (Marianda
171 Airica). Additional carbonate chemistry parameters (pH_{NBS} , $p\text{CO}_2$, $[\text{HOC}_3^-]$, $[\text{CO}_3^{2-}]$,
172 aragonite saturation state $[\Omega_{\text{Arg}}]$) were calculated from A_T and C_T using CO2SYS (Pierrot et
173 al. 2006) with dissociation constants from Mehrbach et al. (1973), refit by Dickson and
174 Millero (1987) and KSO4 using Dickson (1990). In situ water temperature ($^\circ\text{C}$), salinity and
175 pH was measured hourly throughout the experimental period using an Exo2 multiparameter
176 sonde (YSI Inc). Nitrate and phosphate concentrations were calculated throughout the
177 experimental period (below) and included in carbonate chemistry calculations. Net
178 community calcification rates were calculated using the alkalinity anomaly technique
179 (Chisholm & Gattuso 1991), based on the change in seawater A_T during the incubation
180 period. For each mole of CaCO_3 precipitated (i.e. calcification), A_T is lowered by two molar
181 equivalents. Therefore, the change in alkalinity can be converted to the mass of CaCO_3
182 precipitated. Certified seawater references materials for oceanic CO_2 (Scripps Institution of
183 Oceanography, University of California, San Diego) were used as A_T and C_T standards,
184 following Dickson et al. (2007).

185 **Net DMS+DMSP (DMS/P_T) flux**

186 Samples for total (dissolved+particulate) DMS+DMSP (DMS/P_T) were stored in 50 ml
187 crimp-top serum vials (Wheaton) fitted with Pharma-Fix lids. NaOH was added to a final

188 concentration of 0.03 M to hydrolyse DMSP into DMS. Samples were analysed by purge-
189 and-trap gas chromatography (Turner et al. 1990), using an SRI 8610C GC fitted with a
190 flame photometric detector (nitrogen carrier gas @ 8 psi). Sample concentrations were
191 quantified via comparison to a DMSP standard (Research Plus Inc); sample detection limit
192 was $<1 \text{ nmol L}^{-1}$, precision and accuracy for standards and samples was within 1%.

193 **Net nitrate and phosphate flux**

194 Unfiltered samples for nitrate and phosphate were stored in HDPE bottles (Fisher Scientific)
195 and frozen within 1 hour of collection. 10 ml samples were analysed for nitrate following the
196 cadmium reduction spectrophotometric method (Grasshoff et al. 2007); absorbance was
197 measured at 400 nm, with sodium nitrate used as a standard. 10 ml samples were analysed for
198 phosphate using the ammonium molybdate/ascorbic acid method (Grasshoff et al. 2007);
199 absorbance was measured at 885 nm, with potassium phosphate used as a standard.

200

201 **Statistical analyses**

202 Where parametric assumptions for normality and homogeneity of variance were met,
203 parametric tests were used to interrogate the data. One-way ANOVAs were used to test for
204 differences between ambient, CO₂ enrichment and recovery experimental phases in terms of
205 carbonate chemistry and net fluxes of DO, calcification rate, DMS/P_T, nitrate and phosphate
206 (i.e. experimental phase included as a factor; no data transformation was required).

207 Correlation tests were used to test correlation significance between fluxes of dissolved
208 oxygen, calcification, DMS/P_T, nitrate and phosphate. Kruskal-Wallis tests were used to test
209 for differences in DO fluxes (parametric assumptions could not be met). Analyses were
210 conducted using Minitab V14.1.

211 **Results**

212 **Environmental conditions**

213 Water temperature was $15.3 \pm 0.32^\circ\text{C}$ and salinity was 33.0 ± 0.38 throughout the experimental
214 period (mean \pm SD, n=80). No significant difference in T_A was observed between the three
215 experimental phases ($F_{2,20} = 0.11$, $p = 0.89$; Table 1). In contrast, C_T was significantly higher
216 during the CO_2 enrichment compared to the ambient / recovery phases ($F_{2,20} = 31.6$, $p <$
217 0.001 ; Table 1), resulting in a significant increase in HCO_3^- ($F_{2,20} = 10.45$, $p = 0.001$) and
218 pCO_2 ($F_{2,20} = 4.24$, $p = 0.03$). Mean aragonite saturation state and pH were reduced during
219 CO_2 enrichment compared to the ambient / recovery phases, but not to the extent that
220 significant differences were observed (Ω_{Ar} : $F_{2,20} = 1.47$, $p = 0.26$; pH: $F_{2,20} = 2.76$, $p = 0.09$;
221 Table 1). Average in situ pH at the site in the 38 days before and during the experiment was
222 8.04 ± 0.04 (mean \pm SD) (Figure S1).

223 **Net photosynthesis / respiration (dissolved oxygen)**

224 At ambient conditions, an average net uptake of O_2 (i.e. net respiration) was observed,
225 characterised by a small net release of O_2 during the day (i.e. net photosynthesis) to net
226 respiration during the night (Figure 1). During the CO_2 enrichment average net O_2 release
227 increased compared to the ambient / recovery phases, reducing the difference between day
228 (higher net O_2 release) and night (lower net O_2 release / net uptake) measurements ($F_{2,27} =$
229 2.98 , $p = 0.07$). During the recovery phase, net O_2 uptake decreased towards initial levels, but
230 did not quite reach the magnitude of net photosynthesis originally observed. When compared
231 separately, net oxygen flux was significantly higher in CO_2 -enriched conditions than ambient
232 or recovery periods during the night ($H_1 = 4.20$, $p = 0.040$), but not during the day ($H_1 = 1.70$,
233 $p = 0.192$), reflecting the observed overall trend towards increased O_2 flux under CO_2
234 enrichment (Figure 1).

235 **Net calcification / dissolution (carbonate chemistry)**

236 A significant reduction in net calcification was observed during the CO₂ enrichment
237 compared to the ambient / recovery phases ($F_{2,25} = 5.49$, $p = 0.01$; Figure 1). Under ambient
238 CO₂ conditions, the coralline algal community consistently exhibited a net calcification.
239 During CO₂ enrichment, a significant shift towards net dissolution was observed. The
240 recovery phase was characterised by an intermediate rate of net calcification. A significant
241 negative correlation between DO flux and net calcification rate was observed ($r = -0.40$, $p =$
242 0.05 ; Figure 1).

243 **Net DMS/P_T flux**

244 Under ambient CO₂ conditions, there was a net uptake of DMS/P_T by the coralline
245 algal community of between 11 – 24 $\mu\text{mol m}^{-2} \text{h}^{-1}$ (Table 2). During CO₂ enrichment there
246 was a small reduction in net uptake rates, manifest as a shift towards the occasional net
247 release of DMS/P_T, but this change was not significant between experimental phases ($F_{2,27} =$
248 0.62 , $p = 0.54$; Table 2). DMS/P_T flux was not significantly correlated with any of the other
249 biogeochemical parameters, at $p < 0.05$.

250 **Net nitrate and phosphate flux**

251 Average net nutrient release and uptake rates were balanced (i.e. flux of ~zero), and
252 no significant change was observed during CO₂ enrichment compared to the ambient /
253 recovery phases (nitrate: $F_{2,25} = 0.80$, $p = 0.46$; phosphate: $F_{2,25} = 0.01$, $p = 0.99$; Table 2).
254 Net benthic flux of phosphate, but not nitrate, was significantly correlated with benthic
255 oxygen flux ($r = 0.46$, $p = 0.02$). No other significant correlations between net O₂, nitrate,
256 phosphate and DMS_{Pt} flux and net calcification rate (at $p < 0.05$) were observed.

257

258 Discussion

259 Despite the known issues with investigating the effect of elevated CO₂ in a laboratory
260 setting, only a handful of in situ CO₂ enrichment experiments have been conducted, and even
261 less on the whole natural community. This is the first community-level in situ acute CO₂
262 enrichment study in mid/high latitudes, and the first to consider the rate of recovery following
263 acute CO₂ perturbation. In this study, there was a rapid community level response to acute
264 CO₂ enrichment. This was particularly evident for net calcification, demonstrating the
265 sensitivity of the whole community to acute CO₂ exposure, not just individual species.

266 Unlike single-organism laboratory experiments, this study integrated the response of
267 the whole community. Whilst this means we are unable to assign individual species to
268 specific biogeochemical changes, the results obtained are relevant to real-world challenges
269 such as the designation of marine management strategies, which by necessity incorporate
270 whole communities (even if a particular species is the target focus). At the level of CO₂
271 enrichment used in this study, the skeleton and epithelial cell surface of *Lithothamnion*
272 *glaciale* is compromised (Burdett et al. 2012, Kamenos et al. 2013), allowing for skeletal
273 dissolution (Langdon et al. 2000) – supporting the observed shift towards net community
274 dissolution. This may have also been facilitated by dissolution of carbonate sediment and
275 dead sections of coralline algae, which cannot exert biological control and buffering against
276 changes in carbonate chemistry (Kamenos et al. 2013). Like other reef-based marine
277 ecosystems, this coralline algal community is highly diverse across multiple trophic levels
278 (BIOMAERL 1999, Barbera et al. 2003, Kamenos 2004). Calcifying invertebrates are
279 especially abundant (e.g. *Ophiocomina nigra*, which can make up 47% of total faunal
280 biomass; BIOMAERL 1999), and CO₂ enrichment is known to lead to a reduction in
281 calcification rate / increase in dissolution rate of these organisms (Kroeker et al. 2010). Thus,
282 these organisms are likely to have also contributed to the observed shift towards net

283 dissolution, impacting their contribution to coastal CO₂ flux (Davoult et al. 2009). Due to the
284 high heterotrophic diversity of coralline algal beds (Barbera et al. 2003), only a small net
285 photosynthesis during the day was observed, supporting previous measurements using the
286 Eddy correlation technique (Attard et al. 2015) and providing confidence that results recorded
287 do not represent treatment artefacts. CO₂ enrichment led to a small increase in net O₂ release,
288 suggesting an increased capacity for net photosynthesis – supporting the likely benefits of
289 elevated CO₂ conditions for aquatic photosynthetic organisms (Kroeker et al. 2010).
290 Photosynthetic use of CO₂ can also provide a potential refuge for calcifying species by
291 buffering against the damaging effects of CO₂ enrichment (e.g. crustose coralline algae;
292 Cornwall et al. 2014, Short et al. 2014, Kamenos et al. 2016), although this was not observed
293 in this study. Increased photosynthetic capacity may also increase the carbon sequestration
294 potential of these ecosystems (a key process in blue carbon storage; van der Heijden &
295 Kamenos 2015), but a shift towards net dissolution may impact the stability of coralline algal
296 carbonate deposits. The balance and interaction of photosynthesis and calcification /
297 dissolution, and subsequent impact on carbon sequestration / storage is exemplified by the
298 observed correlation between net O₂ flux and net calcification.

299 Change in the community-level flux of dimethylated sulphur compounds appears to
300 be robust to acute CO₂ enrichment, despite the known sensitivity of coralline algal DMSP
301 dynamics to acute CO₂ exposure (Burdett et al. 2012). Thus, it may be hypothesised that
302 although DMS/P_T concentrations did not change, the proportion of the molecular species (e.g.
303 dissolved vs particulate, DMSP vs DMS) may have been altered, but this was not calculable
304 by the approach employed. Nutrient fluxes were also insensitive to acute CO₂ enrichment, at
305 least at the level used in this study. However, the correlation between phosphate and DO
306 suggests that a larger CO₂ perturbation (in duration and / or magnitude) may impact
307 phosphorus cycling processes.

308 Acute CO₂ enrichment is just one aspect of carbon-chemistry pressures on marine
309 habitats. In addition, the combined effects of acute CO₂ enrichment and chronic, long-term
310 changes in carbonate chemistry may exacerbate biological responses. This has yet to be tested
311 at the community scale, despite the known importance of both acute and chronic CO₂
312 enrichment in driving responses in marine organisms. Surprisingly, even after a recovery
313 phase almost 1.5 times the length of the CO₂ enrichment, a full recovery (i.e. complete return
314 of all parameters to the initial measured rates) was not seen, at least in terms of the
315 parameters measured here, suggesting that, at best, there is considerable lag in community
316 recovery response times. This calls into question the capacity for the system to ‘bounce back’
317 following repeated exposure to acute CO₂ inputs, which would be likely given the sources of
318 short-term CO₂ enrichment (e.g. aquaculture, CCS). Previous studies have shown that
319 damage to the coralline algal skeletal structure under CO₂-enriched conditions can rapidly
320 occur (Burdett et al. 2012, Kamenos et al. 2013). In situ, this effect may manifest through to
321 the community level. Results from this study and others (e.g. Hall-Spencer et al. 2008,
322 Fabricius et al. 2011) collectively suggest that CO₂ enrichment may cause change across
323 biological scales, from the individual to community levels. If these changes persist in the
324 long-term, we may observe permanent transitions in community composition, perhaps one
325 that favours net photosynthesis, thereby tipping the balance in terms of biodiversity, and / or
326 net dissolution. Such transitions would not favour the growth of carbonate-depositing
327 ecosystem engineers such as coralline algae.

328

329 **References**

330 Abril G, Commarieu MV, Maro D, Fontugne M, Guérin F, Etcheber H (2004) A massive dissolved
331 inorganic carbon release at spring tide in a highly turbid estuary. *Geophys Res Lett*
332 31:L09316

333 Attard K, Stahl H, Kamenos N, Turner G, Burdett H, Glud R (2015) Benthic oxygen exchange in a live
334 coralline algal bed and an adjacent sandy habitat: an eddy covariance study. *Mar Ecol Prog*
335 *Ser* 535:99-115

336 Barbera C, Bordehore C, Borg JA, Glemarec M, Grall J, Hall-Spencer JM, De la Huz C, Lanfranco E,
337 Lastra M, Moore PG, Mora J, Pita ME, Ramos-Espla AA, Rizzo M, Sanchez-Mata A, Seva A,
338 Schembri PJ, Valle C (2003) Conservation and management of northeast Atlantic and
339 Mediterranean maerl beds. *Aquat Conserv: Mar Freshwat Ecosyst* 13:S65-S76

340 Barry JP, Lovera C, Buck KR, Peltzer ET, Taylor JR, Walz P, Whaling PJ, Brewer PG (2014) Use of a Free
341 Ocean CO₂ Enrichment (FOCE) system to evaluate the effects of ocean acidification on the
342 foraging behavior of a deep-sea urchin. *Environ Sci Technol* 48:9890-9897

343 BIOMAERL (1999) BIOMAERL project: Final Report (Coordinator: P.G.Moore, University Marine
344 Biological Station Millport, Scotland), EC Contract No. MAS3-CT95-0020.

345 Blackford J, Stahl H, Kita J, Sato T (2015) Preface to the QICS special issue. *Int J Greenhouse Gas*
346 *Control*:1

347 Burdett HL, Aloisio E, Calosi P, Findlay HS, Widdicombe S, Hatton AD, Kamenos NA (2012) The effect
348 of chronic and acute low pH on the intracellular DMSP production and epithelial cell
349 morphology of red coralline algae. *Mar Biol Res* 8:756-763

350 Burdett HL, Hatton AD, Kamenos NA (2015a) Coralline algae as a globally significant pool of marine
351 dimethylated sulfur. *Global Biogeochem Cycles* 29:1845-1853

352 Burdett HL, Hatton AD, Kamenos NA (2015b) Effects of reduced salinity on the photosynthetic
353 characteristics and intracellular DMSP concentrations of the red coralline alga,
354 *Lithothamnion glaciale*. *Mar Biol* 162:1077-1085

355 Campbell JE, Fourqurean JW (2014) Ocean acidification outweighs nutrient effects in structuring
356 seagrass epiphyte communities. *J Ecol* 102:730-737

357 Chisholm JR, Gattuso JP (1991) Validation of the alkalinity anomaly technique for investigating
358 calcification of photosynthesis in coral reef communities. *Limnol Oceanogr* 36:1232-1239

359 Cornwall CE, Boyd PW, McGraw CM, Hepburn CD, Pilditch CA, Morris JN, Smith AM, Hurd CL (2014)
360 Diffusion boundary layers ameliorate the negative effects of ocean acidification on the
361 temperate coralline macroalga *Arthrocardia corymbosa*. *PLoS One* 9:e97235

362 Cornwall CE, Hurd CL (2015) Experimental design in ocean acidification research: problems and
363 solutions. *ICES J Mar Sci* 73:572-581

364 Davoult D, Harlay J, Gentil F (2009) Contribution of a Dense Population of the Brittle Star *Acrocnida*
365 *brachiata* (Montagu) to the Biogeochemical Fluxes of CO₂ in a Temperate Coastal Ecosystem.
366 *Estuar Coasts*:1103-1110

367 Dickson AG (1990) Standard potential of the reaction: $AgCl(s) + 1/2H_2(g) = Ag(s) + HCl(aq)$, and and
368 the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15
369 K. *JChTh* 22:113-127

370 Dickson AG, Millero FJ (1987) A comparison of the equilibrium constants for the dissociation of
371 carbonic acid in seawater media. *Deep-Sea Res Part A Oceanogr Res Pap* 34:1733-1743

372 Dickson AG, Sabine CL, Christian JR (eds) (2007) Guide to best practices for ocean CO₂
373 measurements. *PICES Special Publication* 3

374 Fabricius KE, Langdon C, Uthicke S, Humphrey C, Noonan S, De'ath G, Okazaki R, Muehllehner N, Glas
375 MS, Lough JM (2011) Losers and winners in coral reefs acclimatized to elevated carbon
376 dioxide concentrations. *Nature Climate Change* 1:165

377 Forsberg O (1997) The impact of varying feeding regimes on oxygen consumption and excretion of
378 carbon dioxide and nitrogen in post-smolt Atlantic salmon *Salmo salar* L. *Aquacult Res* 28:29-
379 41

380 Gattuso JP, Kirkwood W, Barry JP, Cox E, Gazeau F, Hansson L, Hendriks I, Kline DI, Mahacek P,
381 Martin S, McElhany P, Peltzer ET, Reeve J, Roberts D, Saderne V, Tait K, Widdicombe S,
382 Brewer PG (2014) Free-ocean CO₂ enrichment (FOCE) systems: present status and future
383 developments. *Biogeosciences* 11:4057-4075

384 Grasshoff K, Kelming K, Ehrhardt M (2007) Methods of seawater analysis. Wiley-VCH, Weinheim
385 Gubbay S, Sanders N, Haynes T, Janssen JAM, Rodwell JR, Nieto A, Criado MG, Beal S, Borg J,
386 Kennedy M, Micu D, Otero M, Saunders G, Calix M (2016) European Red List of Habitats: Part
387 I. Marine habitats. Publications Office of the European Union, Luxembourg
388 Hall-Spencer JM, Rodolfo-Metalpa R, Martin S, Ransome E, Fine M, Turner SM, Rowley SJ, Tedesco D,
389 Buia M-C (2008) Volcanic carbon dioxide vents show ecosystem effects of ocean
390 acidification. *Nature* 454:96-99
391 Kamenos N (2004) The importance of maerl grounds as nursery areas for commercially important
392 species with emphasis on the Queen scallop (*Aequipecten opercularis*). PhD, University of
393 London, 257pp
394 Kamenos N, Perna G, Gambi M, Micheli F, Kroeker K (2016) Coralline algae in a naturally acidified
395 ecosystem persist by maintaining control of skeletal mineralogy and size. *Proc R Soc Biol Sci*
396 *Ser B* 283:20161159
397 Kamenos NA, Burdett HL, Aloisio E, Findlay HS, Martin S, Longbone C, Dunn J, Widdicombe S, Calosi P
398 (2013) Coralline algal structure is more sensitive to rate, rather than the magnitude, of
399 ocean acidification. *Global Change Biol* 19:3621-3628
400 Kline DI, Teneva L, Schneider K, Miard T, Chai A, Marker M, Headley K, Opdyke B, Nash M, Valetich
401 M, Caves JK, Russell BD, Connell SD, Kirkwood BJ, Brewer P, Peltzer E, Silverman J, Caldeira K,
402 Dunbar RB, Koseff JR, Monismith SG, Mitchell BG, Dove S, Hoegh-Guldberg O (2012) A short-
403 term in situ CO₂ enrichment experiment on Heron Island (GBR). *Sci Rep* 2:413
404 Kroeker KJ, Kordas RL, Crim RN, Singh GG (2010) Meta-analysis reveals negative yet variable effects
405 of ocean acidification on marine organisms. *Ecol Lett* 13:1419-1434
406 Lachkar Z (2014) Effects of upwelling increase on ocean acidification in the California and Canary
407 Current systems. *Geophys Res Lett* 41:90-95
408 Langdon C, Takahashi T, Sweeney C, Chipman D, Goddard J, Marubini F, Aceves H, Barnett H,
409 Atkinson MJ (2000) Effect of calcium carbonate saturation state on the calcification rate of
410 an experimental coral reef. *Global Biogeochem Cycles* 14:639-654
411 Lyons D, Van Alstyne K, Scheibling R (2007) Anti-grazing activity and seasonal variation of
412 dimethylsulfoniopropionate-associated compounds in the invasive alga *Codium fragile* ssp.
413 *tomentosoides*. *Marine Biology* 153:179-188
414 Martin, S, Clavier, J, Chauvaud, L, Thouzeau, G (2007a) Community metabolism in temperate maerl
415 beds. I. Carbon and carbonate fluxes. *Mar Ecol Prog Ser* 335:19-29
416 Martin, S, Clavier, J, Chauvaud, L, Thouzeau, G (2007b) Community metabolism in temperate maerl beds. II.
417 Nutrient fluxes. *Mar Ecol Prog Ser* 335:31-41
418 Mehrbach C, Culbertson CH, Hawley JE, Pytkowicz RM (1973) Measurement of the Apparent
419 Dissociation Constants of Carbonic Acid in Seawater at Atmospheric Pressure. *Limnol*
420 *Oceanogr* 18:897-907
421 OECD-FAO (2014) OECD-FAO Agricultural Outlook 2014-2023 : FISHERIES. Accessed 4th September
422 2017. <http://stats.oecd.org/index.aspx?queryid=58653>
423 Pfister CA, Esbaugh AJ, Frieder CA, Baumann H, Bockmon EE, White MM, Carter BR, Benway HM,
424 Blanchette CA, Carrington E (2014) Detecting the unexpected: a research framework for
425 ocean acidification. *Environ Sci Technol* 48:9982-9994
426 Pierrot D, Lewis E, Wallace DWR (2006) CO₂SYS DOS Program developed for CO₂ system calculations.
427 ORNL/CDIAC-105. Carbon Dioxide Information analysis Center. Oak Ridge National
428 Laboratory, US Department of Energy, Oak Ridge, TN.
429 Queirós AM, Fernandes JA, Faulwetter S, Nunes J, Rastrick SPS, Mieszkowska N, Artioli Y, Yool A,
430 Calosi P, Arvanitidis C, Findlay HS, Barange M, Cheung WWL, Widdicombe S (2014) Scaling
431 up experimental ocean acidification and warming research: from individuals to the
432 ecosystem. *Global Change Biol* 21:130-143
433 Riebesell U, Gattuso J-P (2015) Lessons learned from ocean acidification research. *Nature Clim*
434 *Change* 5:12-14

- 435 Riosmena-Rodríguez R (2017) Natural History of Rhodolith/Maërl Beds: Their Role in Near-Shore
436 Biodiversity and Management. In: Riosmena-Rodríguez R, Nelson W, Aguirre J (eds)
437 Rhodolith/Maërl Beds: A Global Perspective. Springer International Publishing
438 Short J, Kendrick G, Falter J, McCulloch M (2014) Interactions between filamentous turf algae and
439 coralline algae are modified under ocean acidification. *J Exp Mar Biol Ecol* 456:70-77
440 Steller DL, Riosmena-Rodríguez R, Foster MS (2009) Living rhodolith bed ecosystems in the Gulf of
441 California, in Johnson M. E. & Ledesma-Vasquez J. (eds), *Atlas of Coastal Ecosystems in the*
442 *Western Gulf of California*. University of Arizona Press, Tucson: 72–82
443 Strong AL, Kroeker KJ, Teneva LT, Mease LA, Kelly RP (2014) Ocean Acidification 2.0: Managing our
444 Changing Coastal Ocean Chemistry. *Bioscience* 64:581-592
445 Turner SM, Malin G, Bagander LE, Leck C (1990) Interlaboratory Calibration and Sample Analysis of
446 Dimethyl Sulfide in Water. *Mar Chem* 29:47-62
447 van der Heijden LH, Kamenos NA (2015) Calculating the global contribution of coralline algae to
448 carbon burial. *Biogeosciences* 12:7845-7877
449 Zakęś Z, Demska-zakęś K, Kata K (2003) Rates of oxygen consumption and ammonia excretion of
450 juvenile Eurasian perch *Perca fluviatilis* L. *Aquacult Int* 11:277-288

451

452

453

454 **Acknowledgements**

455 This work was supported by a Marine Alliance for Science and Technology for Scotland
 456 (MASTS) Research Fellowship awarded to HB. MASTS is funded by the Scottish Funding
 457 Council (grant reference HR09011) and contributing institutions.

458

459 **Table 1.** System parameters under ambient, CO₂ enrichment and recovery phase conditions,
 460 in benthic chambers deployed on a coralline algal bed in Loch Sween, Scotland. Water
 461 temperature, salinity, photosynthetically active radiation (PAR), A_T (total alkalinity) and C_T
 462 (dissolved inorganic carbon) were directly measured; all other carbonate parameters were
 463 calculated as detailed in the methods (pH is on NBS scale; Ω_{Arg} = aragonite saturation state).
 464 Data presented as mean±SD (n=18, except for temperature and salinity, where n=80). Bold
 465 text denotes parameters that were significantly different during the CO₂ enrichment phase (at
 466 p < 0.05).

	Ambient conditions	CO ₂ enrichment	Recovery period
Temperature (°C)	15.3±0.32	15.3±0.32	15.3±0.32
Salinity	33.0±0.38	33.0±0.38	33.0±0.38
Max PAR (μmol photons m ⁻² s ⁻¹)	158	158	158
A _T (μmol kg ⁻¹)	2190.7±87.2	2202.0±123.28	2210.8±68.2
C_T (μmol kg⁻¹)	2084.8±12.8	2168.9±31.20	2066.2±23.2
pH _{NBS}	7.9±0.2	7.7±0.39	8.0±0.2
pCO₂ (μatm)	821.6±343.4	1747.7±1403.33	646.7±320.6
HCO₃⁻ (μmol kg⁻¹)	1961.1±27.5	2033.5±20.35	1927.6±49.2
CO ₃ ²⁻ (μmol kg ⁻¹)	92.0±45.9	67.8±50.77	113.5±45.5
Ω _{Arg}	1.4±0.7	1.0±0.78	1.7±0.7

467

468

469 **Table 2.** Community response of acute in-situ CO₂ enrichment in terms of net DMSPt, nitrate
470 and phosphate flux, under initial ambient CO₂ conditions, during CO₂ enrichment and during
471 the recovery phase at ambient CO₂. Data presented as mean±SD.


	Ambient conditions	CO ₂ enrichment	Recovery period
Net DMSPt flux (μmol m ⁻² h ⁻¹)	-23.13±27.12	-13.46±28.12	-11.47±11.39
Net nitrate flux (mg m ⁻² h ⁻¹)	-11.40±36.11	-0.55±19.90	7.71±27.37
Net phosphate flux (mg m ⁻² h ⁻¹)	0.04±0.44	0.02±0.24	0.05±0.29

472

473

474

475 **Figure 1.** Community response of acute in-situ CO₂ enrichment in terms of net dissolved
476 oxygen flux and net calcification rate, under initial ambient CO₂ conditions (black circle),
477 during CO₂ enrichment (white circle) or during the recovery phase at ambient CO₂ (grey
478 circle). Data presented as mean±SD.



479

480

481