

Tissue metabolic changes drive cytokine responses to *Mycobacterium tuberculosis*

Authors

Ekta Lachmandas¹, Ana B Rios-Miguel¹, Valerie A C M Koeken¹, Eva van der Pasch¹, Vinod Kumar^{1,2}, Vicky Matzaraki², Yang Li², Marije Oosting¹, Leo A B Joosten¹, Richard A Netea^{1,4}, Mahdad Noursadeghi³, Mihai G Netea¹, Reinout van Crevel¹ and Gabriele Pollara^{3*}

¹Department of Internal Medicine and Radboud Center for Infectious Diseases, Radboud University Medical Center, PO Box 9102, Nijmegen, The Netherlands

²University of Groningen, UMC Groningen, P.O. Box 30001, Groningen, Netherlands

³Division of Infection and Immunity, University College London, London, WC1E 6BT, United Kingdom

⁴Laboratory of Food Microbiology, Wageningen University and Research, Wageningen, the Netherlands

Corresponding author: Dr Gabriele Pollara, <mailto:g.pollara@ucl.ac.uk>

Conflict of interests

The authors do not have a commercial or other association that might pose a conflict of interest relating to the content of the manuscript.

Funding

This work was supported by the Wellcome Trust Research Fellowship to GP (WT101766/Z/13/Z), Wellcome Trust Investigator Award to MN (207511/Z/17/Z), the National Institute for Health Research Biomedical Research Centre at University College London Hospital, and by an ERC Consolidator Grant to MGN (#310372) and a Spinoza grant of the Netherlands Organization for

© The Author(s) 2018. Published by Oxford University Press for the Infectious Diseases Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Scientific Research (to MGN). RvC and EL were supported by The European Union's Seventh Framework Programme (EU FP7) project TANDEM (HEALTH-F3-2012-305279).

Presentations

The information in this manuscript has not been presented at any meetings.

Accepted Manuscript

Abstract

Cellular metabolism can influence host immune responses to *Mycobacterium tuberculosis* (Mtb). Using a systems biology approach, differential expression of 292 metabolic genes involved in glycolysis, glutathione, pyrimidine and inositol phosphate pathways was evident at the site of a human tuberculin skin test challenge in patients with active tuberculosis infection. For 28 metabolic genes, we identified single nucleotide polymorphisms (SNPs) that were trans-acting for in vitro cytokine responses to Mtb stimulation, including glutathione and pyrimidine metabolism genes that alter production of Th1 and Th17 cytokines. Our findings identify novel therapeutic targets in host metabolism that may shape protective immunity to tuberculosis.

Accepted Manuscript

Background

The activity of many cellular metabolic pathways can impact the host immune response to infections [1]. Individual metabolic pathways have been implicated in anti-mycobacterial responses: glutathione enhances IL-12 and IFN γ secretion following Mtb stimulation [2]; tryptophan catabolism is involved in Mtb-induced production of IL-1 β and IL-23 [3] and control of Mtb growth [4]; and a shift towards aerobic glycolysis in Mtb infected macrophages regulates IL-1 β production [5,6]. However, as metabolic reactions are intrinsically inter-dependent, the challenge lies in determining the relative roles of these and other pathways during in vivo Mtb infection.

Our group has made use of the human tuberculin skin test (TST) challenge model to faithfully reflect the inflammatory changes that occur at the site of tuberculosis (TB) disease, characterising the tissue immunological pathways induced early after mycobacterial antigen exposure [7]. However, to date, no studies have explored the metabolic changes and their functional consequences on downstream cytokine responses in such a model. Quantitative cytokine production in response to mycobacterial stimulation has been associated with genetic polymorphisms [8]. These cytokine quantitative trait loci (cQTLs) provide a functional insight into how genetics influences an inflammatory response, and in turn identify critical pathways that may be amenable to host-directed therapy. In this study, we use the TST model to test the hypothesis that differential tissue expression of genes involved in regulating metabolic pathways can directly influence cytokine production following Mtb stimulation. In turn, our findings provide putative mechanistic links between the activity of cellular metabolic pathways and immune effector functions.

Methods

Transcriptomic data analysis

Transcriptomes were derived from TST and blood of patients with active TB, and human Mtb-infected and healthy lymph nodes from a separate cohort (table S1). All patients were HIV seronegative and none had diabetes mellitus.

We used the KEGG pathway database (<http://www.genome.jp/kegg/pathway.html>) to derive 33 pathways associated with human metabolism, yielding a list of 1422 metabolic genes containing no duplicate genes and no annotation to the original pathways (Tables S2 & S3). Bioinformatic analyses were performed as previously described [7] (see Supplementary Methods).

SNP extraction and cQTL mapping

Single nucleotide polymorphisms (SNPs) within metabolic genes and cQTLs were identified as previously described [8] (see Supplementary Methods).

Metabolite reporter analysis

Reporter metabolite analysis [9] was performed in Matlab using the RAVEN Toolbox (<http://biomet-toolbox.org/index.php?page=downtools-raven>) and the human genome-scale metabolic reconstruction network HMR 2.00 provided in Human Protein Atlas (<http://www.metabolicatlas.org/downloads/hmr>) (see Supplementary Methods).

Metabolite depletion experiments

Peripheral blood mononuclear cells (PBMC) were isolated from 9 healthy Dutch adult volunteers and stimulated with Mtb lysate in the presence or absence of pharmacological manipulators of the glutathione pathway (see Supplementary Methods).

Results

We have previously demonstrated that the transcriptional response at the site of TST is characterised by upregulation of 1725 genes that closely reflect changes seen in dissected human TB granuloma relative to healthy lung tissue [7]. We now show that the TST signature is also enriched within Mtb-infected relative to healthy lymph nodes (Figure 1A), confirming that the TST transcriptional responses mirrors pathology seen at established sites of human TB disease.

Out of 1422 metabolic genes, 292 were differentially expressed at the site of TST versus saline injection in patients with active TB disease (Figure S1 and Table S4). Expression of these genes at tissue level showed a remarkable lack of concordance with the blood transcriptome of the same patients, as only 9 metabolic genes were differentially expressed in the blood (Figure 1B, Figure S2 & Table S5). This demonstrates that the TST challenge model provides greater molecular resolution to identify host differential metabolic gene expression to mycobacterial infection than the blood compartment [7].

Given the interconnected relationship between multiple metabolic pathways, we generated a network depicting the ten most enriched KEGG annotated metabolic pathways based on the 292 differentially expressed metabolic genes (Figure 1C). As expected, the TST induced changes in multiple inter-linked metabolic pathways. These included not only previously described pathways such as glycolysis and glutathione metabolism, but also others such as inositol phosphate metabolism and specific amino acid metabolic pathways not previously associated with TB [1,2]. Since differentially expressed genes within the same metabolic pathway were both up- or down-regulated (Figure S3), we evaluated which metabolites in these pathways were most affected using Reporter Metabolite analysis. This identified metabolites in the human metabolic network around which the most transcriptional changes occur [10]. The model predicted changes in several key metabolites between TST and saline, including 1,3-bisphospho-D-glycerate (glycolysis), L-formylkynurenine (tryptophan metabolism), 1-phosphatidyl-1D-myo-inositol-3,4-bisphosphate and glutathione (Figure 1D). Therefore, the model predicts that gene expression changes in the TST alter the concentration of several bioactive metabolites in multiple pathways, and that this has the potential to impact the nature of the host immune response to Mtb.

Next, we sought to test the hypothesis that metabolic changes in the tissue environment influence Mtb-induced cytokine production. We used natural genetic variation to identify putative cis-acting SNPs for metabolic gene expression and tested whether they might be trans-acting SNPs for cytokine responses. We identified metabolic genes differentially expressed in the TST and assessed their impact on cytokine secretion following Mtb lysate-stimulation in a cohort of 500 healthy individuals [8]. First, we used genotypes extracted from the 1000 Genomes Project to identify 16061 SNPs from the 109 metabolic genes that comprised the 10 most enriched metabolic pathways in the TST (Figure 1C). We then assessed which of these SNPs were associated with variable cytokine secretion, generating 2376 putative cQTLs. To reduce multiple testing false positives, we focused on SNPs in the mRNA coding region of the gene of interest that influence the same gene's transcription (i.e. metabolic gene SNPs that were cis-expression QTLs [eQTLs]).

This analysis generated 47 cQTL SNPs from 28 metabolic genes (Figures 2A-B & Tables S6 & S7). Based on these genes, the most over-represented metabolic pathways included the glutathione, glycolysis, inositol phosphate metabolism and pyrimidine pathways (Figure S4). Many amino acid metabolism pathways were also observed, and ALDH3A2, ALDH3A1, LDHA and IL4I1 were the most frequent constituent genes from these pathways (Figure S4). Glutathione and pyrimidine metabolism predominantly influenced the secretion of IFN γ and IL-17: 7 of 15 (47%) cQTLs that regulated IFN γ secretion and 4 of 7 (57%) cQTLs that regulated IL-17 secretion were derived from genes assigned to glutathione or pyrimidine metabolic pathways, whereas no cQTLs from these pathways influenced cytokine secretion by macrophages (Figure 2B). In contrast, genes involved in glycolysis, amino acid and inositol phosphate metabolism acted as cQTLs more ubiquitously, influencing the secretion of both T cell and myeloid cell derived cytokines (IL-1 β , IL-6, IL-22 and TNF α) (Figure 2B).

We validated some of the relationships between metabolic changes and cytokine secretion in vitro using buthionine sulfoximine (BSO), an inhibitor of glutathione synthesis, and diethylmaleate (DEM) that depletes cells of glutathione [11]. Both resulted in increased IL-6 and decreased IL-17 secretion

by Mtb stimulated PBMC (fig 2C), consistent with the cQTL data demonstrating that variable expression of glutathione genes impacted the secretion of both these cytokines (fig 2B).

Finally, to explore the physiological relevance of the 28 metabolic genes that act as cQTLs, we showed that their expression in Mtb-infected lymph nodes strongly correlated with expression in the TST (Figure 2D), indicating that these metabolic genes have the potential to exert cQTL activity at the site of human TB disease, thus shaping the local inflammatory response to Mtb.

Accepted Manuscript

Discussion

Individual metabolic pathways have been shown to affect the host response to Mtb [2,3,5], but their relative contribution in a multicellular tissue infection setting has not been investigated. We took a systems approach using the human in vivo TST challenge model, revealing gene expression changes in multiple metabolic pathways that in turn predict enrichment of several bioactive metabolites. Genetic polymorphisms in these differentially expressed metabolic genes, as well as pharmacological inhibition, were found to control Mtb-induced cytokine production. Finally, expression of these same genes at the site of TB disease closely correlated that observed in the TST, underlying the functional relevance of our findings to the pathogenesis of human TB infection.

The inflammatory response to TST challenge reflects the immunopathological changes in human TB disease [7]. Interestingly, in terms of metabolic gene expression the TST site also closely mirrors Mtb-infected lymph nodes, unlike the peripheral blood of TB patients, which showed quantitatively smaller changes. Although this may reflect the TST response occurring after acute antigenic challenge in contrast to steady state assessment in blood, in these same patients the blood transcriptome shows significant deviation from health [7,12], illustrating that tissue may be the more appropriate context when studying changes in metabolism.

We observed differential gene expression in the TST of multiple metabolic pathways, including glycolysis, glutathione, pyrimidine and inositol phosphate metabolism. The relevance of changes in the metabolic environment after Mtb infection was supported by the fact that we identified several SNPs in these genes that both act as cis-eQTLs and also impact Mtb-induced cytokine secretion by PBMC and macrophages [8], cells that are enriched at the site of TST [7]. For functional validation, we selected one of the pathways identified through gene expression and cQTL-analysis, the glutathione pathway, whereby pharmacological inhibition affected Mtb-induced production of IL-6 and IL-17. Importantly, the 500FG cohort data demonstrated that variable expression of genes involved in glutathione metabolism (GSTM2, ANPEP, LAP3 and GSTA3) impacted the secretion of these same cytokines. Furthermore, we have previously demonstrated that inhibiting glycolysis using 2-deoxyglucose impaired IL-22 secretion by Mtb-stimulated PBMC, in turn validating the observation from the 500FG data that SNPs in the glycolysis gene ADPGK are cQTLs for IL-22 secretion [6].

Our analyses also identified new roles for pyrimidine metabolism, which has recently been associated with regulating inflammasome activity and cytokine secretion [13]. We demonstrate that pyrimidine and glutathione metabolism genes predominantly impact the secretion of T cell-derived cytokines IFN γ and IL-17. As such, therapeutic manipulation of pyrimidine and glutathione pathways may influence Th1 and Th17 polarisation, and thus the balance between protection and pathology in Mtb-infected tissues [14]. In contrast, genes involved in amino acid and inositol phosphate metabolism impact a wider array of cytokines, including those secreted from myeloid cells, which may in part relate to the central role of the metabolite phosphatidylinositol as the backbone of signal transduction components IP3 and Akt [6].

Our study has some limitations. Firstly, we restricted our cQTL analyses to SNPs that were also cis-eQTLs to limit multiple testing errors, thus likely missing other functionally relevant cQTLs that may act in trans via other genes. Equally, identification of SNPs that were eQTLs was limited to databases that probed blood and tissues not infected with Mtb, possibly missing other functionally relevant

SNPs in the context of Mtb infection. Furthermore, the effector cytokines studied for cQTL analysis comprise only one facet of the host immune response to Mtb infection and were restricted to the manually selected panel available in the 500FG database, introducing bias into the breadth of immunological effector functions exerted by metabolic gene SNPs. Finally, although Reporter Metabolite analysis predicted computationally that the gene expression changes impacted on the concentration of several key metabolites, we were not able to assess this directly. The study was conceived retrospectively, and thus tissue samples had already been used wholly for transcriptomic analysis. The relationship between tissue gene expression changes and metabolite concentrations will need to be verified experimentally in future studies.

In summary, this study made a comprehensive assessment of the human tissue metabolic transcriptional response to in vivo mycobacterial antigenic stimulation. A number of known and novel metabolic pathways were differentially expressed, and genetic variation in identified genes affected cytokine responses to Mtb. Therefore, our systems approach provided a novel list of putative metabolic gene cQTLs that are amenable to further experimental validation, and revealed a new layer of complexity to the host anti-mycobacterial response, supporting the use of host-directed strategies that target cellular metabolism, such as the regulation of glycolysis by metformin [15]. Furthermore, our approach combining transcriptomics and functional genomics illustrates a pipeline that can be used to identify novel and clinically relevant pathways in the context of other infectious diseases.

References

1. Lachmandas E, Boutens L, Ratter JM, et al. Microbial stimulation of different Toll-like receptor signalling pathways induces diverse metabolic programmes in human monocytes. *Nat Microbiol.* **2016**; 2:16246.
2. Guerra C, Morris D, Sipin A, et al. Glutathione and Adaptive Immune Responses against *Mycobacterium tuberculosis* Infection in Healthy and HIV Infected Individuals. *PLOS ONE.* **2011**; 6(12):e28378.
3. Memari B, Bouttier M, Dimitrov V, et al. Engagement of the Aryl Hydrocarbon Receptor in *Mycobacterium tuberculosis*-Infected Macrophages Has Pleiotropic Effects on Innate Immune Signaling. *J Immunol.* **2015**; 195(9):4479–4491.
4. Zhang YJ, Reddy MC, Ioerger TR, et al. Tryptophan Biosynthesis Protects *Mycobacteria* from CD4 T-Cell-Mediated Killing. *Cell.* **2013**; 155(6):1296–1308.
5. Gleeson LE, Sheedy FJ, Palsson-McDermott EM, et al. Cutting Edge: *Mycobacterium tuberculosis* Induces Aerobic Glycolysis in Human Alveolar Macrophages That Is Required for Control of Intracellular Bacillary Replication. *J Immunol.* **2016**; 196(6):2444–2449.
6. Lachmandas E, Beigier-Bompadre M, Cheng S-C, et al. Rewiring cellular metabolism via the AKT/mTOR pathway contributes to host defence against *Mycobacterium tuberculosis* in human and murine cells. *Eur J Immunol.* **2016**; 46(11):2574–2586.
7. Bell LCK, Pollara G, Pascoe M, et al. In Vivo Molecular Dissection of the Effects of HIV-1 in Active Tuberculosis. *PLOS Pathog.* **2016**; 12(3):e1005469.
8. Li Y, Oosting M, Smeekens SP, et al. A Functional Genomics Approach to Understand Variation in Cytokine Production in Humans. *Cell.* **2016**; 167(4):1099–1110.e14.
9. Patil KR, Nielsen J. Uncovering transcriptional regulation of metabolism by using metabolic network topology. *Proc Natl Acad Sci U S A.* **2005**; 102(8):2685–2689.
10. Agren R, Liu L, Shoaie S, Vongsangnak W, Nookaew I, Nielsen J. The RAVEN Toolbox and Its Use for Generating a Genome-scale Metabolic Model for *Penicillium chrysogenum*. Maranas CD, editor. *PLoS Comput Biol.* **2013**; 9(3):e1002980.
11. Mitchell JB, Russo A, Biaglow JE, McPherson S. Cellular Glutathione Depletion by Diethyl Maleate or Buthionine Sulfoximine: No Effect of Glutathione Depletion on the Oxygen Enhancement Ratio. *Radiat Res.* **1983**; 96(2):422–428.
12. Roe JK, Thomas N, Gil E, et al. Blood transcriptomic diagnosis of pulmonary and extrapulmonary tuberculosis. *JCI Insight.* **2016**; 1(16):e87238.
13. Furman D, Chang J, Lartigue L, et al. Expression of specific inflammasome gene modules stratifies older individuals into two extreme clinical and immunological states. *Nat Med.* **2017**; 23(2):174–184.

14. Mourik BC, Lubberts E, Steenwinkel JEM de, Ottenhoff THM, Leenen PJM. Interactions between Type 1 Interferons and the Th17 Response in Tuberculosis: Lessons Learned from Autoimmune Diseases. *Front Immunol* [Internet]. **2017** [cited 2017 Jul 3]; 8. Available from: <http://journal.frontiersin.org/article/10.3389/fimmu.2017.00294/full>
15. Singhal A, Jie L, Kumar P, et al. Metformin as adjunct antituberculosis therapy. *Sci Transl Med*. **2014**; 6(263):263ra159.

Accepted Manuscript

Figure legends



Figure 1: Metabolic pathways enriched in TST responses relative to saline injection.

(A) Expression of TST signature in Mtb infected lymph nodes (LN) relative to healthy LN. Each dot represents one sample. Horizontal lines represent median value expression. * $p < 0.0001$ by Mann-Whitney test. (B) Pairwise comparison of 292 genes differentially expressed in TST relative to saline injection compared to the expression difference between the blood of patients with active TB and healthy volunteers (HV). r^2 = Spearman's rank correlation coefficient, CoV = covariance. (C) The top ten most statistically enriched KEGG metabolic pathways are represented in a network plot, in which the edges indicate associations between genes (yellow nodes) and named pathways (blue nodes), and the node size is proportional to the respective pathway $-\log_{10}$ p value enrichment statistic. (D) Reporter metabolites predicted to be differentially expressed in TST compared to saline injection. Metabolites selected for known association with metabolic pathways and ranked by increasing p value.

Figure 2: Identification of cQTLs within metabolic genes differentially expressed in a TST.

(A) Representative box plots of association between SNP genotypes and Mtb-induced cytokine levels. Length of the box is the interquartile range and whiskers indicate the range of $1.5 \times$ the length of the box from either end of the box. P values were obtained using linear regression analysis of cytokine on genotype data. (B) Heatmap of all 47 cQTL SNPs and their relationship to cytokine secretion following PBMC or macrophage (Mfs) stimulation with Mtb lysate. (C) IL-6 & IL-17 secretion from Mtb lysate-stimulated PBMC in the presence or absence of BSO or DEM. Horizontal lines represent median value expression. * $p < 0.01$ Wilcoxon signed-rank test. (D) Gene expression of 28 metabolic genes with cQTL SNPs in TST relative to saline compared to the expression in Mtb infected lymph nodes relative to healthy lymph nodes. r^2 = Spearman's rank correlation coefficient.

Figure 1



D

Metabolite	p value Strength	Associated Metabolic Pathway
1,3-bisphospho-D-glycerate	0.0057	Glycolysis
L-formylkynurenine	0.0057	Tryptophan metabolism
1-phosphatidyl-1D-myo-inositol-3,4-bisphosphatidyl-1D-myo-inositol-3-phosphate	0.0072	Inositol phosphate metabolism
glycine	0.0075	Glycine, serine and threonine metabolism
3-phospho-D-glycerate	0.0077	Glycolysis
serotonin	0.0089	Tryptophan metabolism
Glutathione	0.0097	Glutathione metabolism
(R)-methylmalonyl-CoA	0.0118	Valine, leucine and isoleucine degradation
1-phosphatidyl-1D-myo-inositol-3-phosphate	0.0128	Inositol phosphate metabolism
1-phosphatidyl-1D-myo-inositol-5-phosphate	0.0128	Inositol phosphate metabolism
1-phosphatidyl-1D-myo-inositol-5-phosphate	0.0128	Inositol phosphate metabolism

Figure 2

