

1

2

3 **Lactation modifies stress-induced immune changes in laboratory rats**

4

5

6

7 **Katrin M Jaedicke^{1,2}, Marco D Fuhrmann¹, Volker Stefanski^{1,3} ***

8

9

10 ¹ *Department of Animal Physiology, University of Bayreuth, Bayreuth, Germany*

11 ² *Institute of Cellular Medicine, Newcastle University, Newcastle upon Tyne, UK*

12 ³ *Leibniz Institute of Zoo and Wildlife Research, Berlin, Germany*

13

14

15

16 ***Corresponding author:** Volker Stefanski, Leibniz Institute of Zoo and Wildlife Research,

17 Alfred-Kowalke-Str.17, D-10315 Berlin, Germany,

18 E-mail: stefanski@izw-berlin.de

19 Phone ++49 (0)30 5168214, Fax ++49 (0)30 5168110

20

21 Abstract

22

23 Lactation and stressor exposure both influence the activity of the immune system, but the
24 interaction of both factors on the immune defense is poorly understood. The aim was
25 therefore to investigate in lactating Long-Evans rats the effect of social stress on aspects of
26 cellular immunity in the blood and mesenteric lymph nodes (MLN). Acute social stress (2 h)
27 was induced in lactating and non-lactating female intruders using a confrontation model that
28 yielded into social defeat and increased plasma corticosterone concentrations. Stress as well
29 as lactation had marked effects on the immune system. Acute social stress caused
30 granulocytosis, reduced lymphocyte proliferation, and cytokine production in the blood, but
31 had no significant effects in MLN. In the blood of lactating rats, increased numbers of
32 granulocytes and enhanced phagocytosis, but decreased B cell numbers and reduced IL-2
33 production was observed. However, in MLN both lymphocyte proliferation and monocyte
34 numbers were increased in lactating rats. The effect of stress on the immune measures was
35 often similar in lactating and non-lactating females, but a few important differences were
36 evident: Only non-lactating animals showed an increase in blood granulocyte numbers and a
37 decrease in IL-2 production in response to stressor exposure. Thus, during lactation, a
38 neuroendocrine status may exist which impedes stress-induced modulations at least of some
39 immune parameters.

40

41

42

43 **Key words**

44

45 Females, gender, lactation, social stress, blood cellular immunity, mesenteric lymph nodes,
46 cytokines, lymphocytes, phagocytes, corticosterone

47 Introduction

48 The period of lactation takes up a large part of life in many mammalian species, with
49 profound hormonal, morphological, and behavioral changes occurring during that time. Apart
50 from changes related to milk production, alterations in certain brain regions such as the
51 supraoptic nucleus (Mann and Bridges, 2002; Theodosios et al., 1981) and in the HPA system
52 (Carter et al., 2001; Magiakou et al., 1997; Shanks et al., 1999) occur. In addition, the
53 behavior of lactating females is often characterized by reduced anxiety (Hard and Hansen,
54 1985; Neumann et al., 2000; Toufexis et al., 1999) and increased aggression towards
55 unfamiliar conspecifics (Gammie et al., 2005; Wise, 1974). Whereas in animals lactation is
56 associated with an increased susceptibility to parasitic infections (Barger, 1993; Festa-
57 Bianchet, 1989; Ngwenya, 1977; Shubber et al., 1981), studies in humans show a postpartum
58 relapse of established autoimmune disorders such as rheumatoid arthritis and multiple
59 sclerosis, often described as a flare-up due to the rebound of the immune system after
60 pregnancy (Buchel et al., 2002; Nelson and Ostensen, 1997; van Walderveen et al., 1994).
61 Furthermore, the postpartum period in humans is also associated with the onset of the
62 postpartum autoimmune thyroid syndrome (Amino et al., 1982; Muller et al., 2001).

63 The immune system of lactating animals was the focus of only a small number of studies
64 with mostly different experimental designs. It appears, however, that some aspects of
65 immune function become suppressed, while others remain unaffected or become enhanced.
66 For example, antibody production after immunization (Jäckel, 2003; Ngwenya, 1977) and IL-
67 2 production in the spleen (Shanks et al., 1997) was found to be suppressed during lactation
68 in rodents. Conversely, increased concentrations of plasma IL-6 or an enhanced proliferative
69 response of lymphocytes from mesenteric lymph nodes (MLN) suggest activation of other
70 immune responses (Shanks et al., 1997). Similarly, in humans, an increase in serum IL-6,
71 TNF- α , IFN- γ and IL-10 concentrations and higher numbers of blood cytotoxic T cells and B

72 cells were observed during lactation (Groer et al., 2005). For NK cell numbers, conflicting
73 data exist: Groer (2005) reported a decreased NK cell number during lactation, whereas
74 Watanabe and colleagues (1997) show increased or unaltered values.

75 Stressors have a major impact on the immune system in male (Avitsur et al., 2002; Shurin et
76 al., 1994; Stefanski and Engler, 1998; Stefanski et al., 1996) and female rats (Stefanski and
77 Grüner, 2006). In females, the stress responses can vary considerably depending on the
78 reproductive status. For example, susceptibility to NK cell-dependent tumor metastasis after
79 surgery stress differs among oestrous phases (Ben-Eliyahu et al., 2000; Page and Ben-
80 Eliyahu, 1997). So far, however, only a few studies have investigated the effects of stress on
81 the immune system during lactation. In humans, a psychological stressor (public speaking
82 and mental arithmetic) was effective to decrease blood lymphocyte proliferation in response
83 to PWM in the non-postpartum group only. In contrast, PHA-induced lymphocyte
84 proliferation increased in the group of women who were recently parturient and
85 breastfeeding, whereas the non-postpartum group remained unaffected (Redwine et al.,
86 2001). Shanks and colleagues reported differential effects of conditioned stress (electrical
87 shock) on the immune system in lactating and non-lactating Sprague-Dawley rats (Shanks et
88 al., 1997). Stressed lactating rats showed an enhanced proliferation in MLN and higher
89 plasma IL-6 concentrations, but a reduced splenocyte proliferation compared to the stressed
90 non-lactating group. No difference however, was observed with respect to blood lymphocyte
91 proliferation.

92 The aim of the present study is to investigate the impact of stress on the immune system
93 during lactation using a social confrontation model with high face validity (Stefanski and
94 Engler, 1998; Stefanski and Grüner, 2006; Stefanski et al., 2005). Here, we focus on the
95 effects of an acute (2h) social stressor on several aspects of cellular immunity in the blood
96 and in MLN. This lymph node region plays a key role during lactation because it provides

97 protection for the offspring against potentially harmful gut pathogens (Head and Seelig,
98 1983; Lamm et al., 1978; Roux et al., 1977).

99

100 2 Methods

101 2.1 Animals

102 Adult Long-Evans rats (about 4.5 month old) were housed in male/female pairs in
103 polycarbonate rat cages (26 x 42 x 15 cm) under standard laboratory conditions ($20 \pm 1^\circ\text{C}$, 40
104 $\pm 5\%$ humidity). Animals were maintained on a 12:12 h light–dark cycle and had ad libitum
105 access to rat standard diet and water. All experiments were conducted in the dark phase of the
106 cycle (the active period of the animals).

107

108 2.2 Social stress procedure

109 Social stress was induced using an adapted form of the resident-intruder confrontation
110 paradigm as described previously (Stefanski et al., 2005). Briefly, lactating and non-lactating
111 female rats (“intruders”) were introduced to confrontation-experienced resident female rats
112 (Long-Evans) housed in a larger enclosure with chipboard walls (height: 75 cm) and 0.5 m²
113 of tiled floor. Intruders were attacked by residents within 5 minutes. Pre-tests with resident
114 females ensured reliability of attacking intruders. The confrontation lasted 2 h. No evidence
115 of wounding was observed after the confrontation. Non-stressed control animals remained
116 undisturbed in their home cages.

117

118 2.3 Synchronisation of lactation

119 All confrontations and samplings were conducted on mid-lactation day 9 ± 1 , the time of
120 lactation where milk production is fully established (Morag et al. 1975, Russell 1980). This
121 sampling time well after end of pregnancy also assured that lactation-associated immune
122 changes were measured, as pregnancy-associated immune changes return to pre-pregnancy
123 values within five days after offspring removal (Stefanski et al., 2005). To create comparable
124 experimental conditions in each set of experiments (see below) the time of lactation was
125 synchronized. For synchronization, females were separated from their partners and housed
126 alone for 25 d to exclude pregnancy. Males were re-introduced for three days and then
127 removed. For the following periods of pregnancy and lactation, females and their litter were
128 housed alone again. This mating paradigm resulted in an about 50 % chance of pregnancy.
129 The non-pregnant rats later served as non-lactating female control group. To exclude possible
130 health abnormalities negatively affecting conception and pregnancy, only females with a
131 previous successful pregnancy were included. (Under the housing in our laboratory, > 95 %
132 of females became repeatedly pregnant). By taking advantage of the 5 day oestrus cycle, this
133 study design therefore did not require sterilized males as mating partners, improving animal
134 welfare.

135 **2.4 Collection of blood and mesenteric lymph nodes**

136 Blood and mesenteric lymph nodes from stressed females were collected immediately after
137 the end of the 2 h confrontation; samples from non-stressed controls were taken at
138 corresponding time points. Up to 2 ml of blood was collected from the lateral tail vein within
139 5 min as described previously (Stefanski, 2000). Blood collection was conducted without
140 requirement for anesthesia. For corticosterone measurement, time was restricted to 3 min.
141 For functional immune assays, heparinized blood was processed immediately. K₂-EDTA-
142 treated blood was stored at room temperature and analyzed with flow cytometry within 3 h.
143 Untreated blood was centrifuged and serum was stored at $-20\text{ }^{\circ}\text{C}$ for future corticosterone

144 measurement. Immediately after blood collection, animals were sacrificed by CO₂ inhalation,
145 mesenteric lymph nodes were collected and transferred to complete RPMI medium (RPMI
146 1640, Biochrom, supplemented with 10 % FCS and 50 µg/ml gentamicin, both Life
147 Technologies). Under sterile conditions, fat tissue was removed and mesenteric lymph nodes
148 were passed through a 40 µm nylon net (BD Biosciences). The cell suspension was
149 centrifuged at 300 x g for 5 min, resuspended in 0.9 % isotonic NaCl, and then treated like
150 blood samples for further analysis.

151 For technical reasons and in order to keep processing time for each sample to a minimum, the
152 entire study was conducted as a series of experimental sets (repeats) with 4 sets (12-15
153 animals each) for MLN analysis and corticosterone measurements and additional 4 sets (20-
154 25 animals each) for blood immune measures. Limitations due to blood volume (2 ml)
155 did not allow analysis of all blood immune parameters simultaneously. Thus, analysis of
156 blood phagocytic capacity, lymphocyte proliferation (ConA only), and leukocyte subsets was
157 conducted in two sets, blood cytokine concentrations were analyzed in other two sets. Each
158 set consisted of about equal numbers of stress and control as well as lactating and non-
159 lactating animals.

160

161 **2.5 Behavioral observation**

162 The resident-intruder confrontations were videotaped under infrared light conditions.
163 Agonistic behavior of both intruder and resident female rats was evaluated for the first 30
164 min of confrontation using “continuous recording“ (Martin and Bateson, 1993). The
165 following behavioral elements (Engler and Stefanski, 2003; Stefanski et al., 2005) were
166 analyzed: *attack* (jump at the opponent with physical contact), *submissive posture* (standing
167 with the front legs on the motionless opponent), *chase* (following the opponent at running
168 pace), *sideway* (“lateral threat“, curved broadside orientation in close proximity to the

169 opponent), *approach* (directed movement towards the opponent at walking speed), *upright*
170 (standing on hind legs with ventral body directed towards the opponent), *full defensive*
171 *posture* (lying motionless on the back with ventral surface exposed to the opponent), *retreat*
172 (directed movement away from the opponent at walking speed), *flight* (like *retreat* but at
173 running pace). A dominance index, DI, ranging from 1 (completely dominant) to 0
174 (completely defeated) was calculated for each female in a confrontation dyad with the
175 behavioral elements *flight* and *retreat* (Stefanski and Grüner, 2006). Only intruder females
176 with a $DI < 0.2$ were included in the study.

177

178 **2.6 Corticosterone concentration**

179 Serum corticosterone concentrations were determined with a standard radioimmunoassay
180 (Foster and Dunn, 1974). The specific antibody (Ab 24/6) was kindly provided by Dr. Vescei,
181 Institute of Pharmacology, University of Heidelberg, Germany. [³H]-labeled corticosterone
182 was obtained from Amersham Biosciences. Cross-reactivity of the corticosterone antibody
183 with other relevant steroids was as follows: cortisol 4.4 %, cortisone 0.65 %,
184 deoxycorticosterone 30.0 %, progesterone 35 %, deoxycortisol < 0.4 %, 17 α -OH
185 progesterone < 0.4 %, testosterone 5.5 %, androstendione 3.2 %, aldosterone 0.8 %. The
186 detection limit of the assay was 5 ng/ml. Intra- and inter-assay variance was < 5 % and
187 < 10 %, respectively.

188

189 **2.7 Leukocyte counts and subpopulations**

190 Total leukocytes were counted with an automatic cell counter (Coulter Counter Z2, Coulter
191 Electronics Ltd.). Cells were then incubated for 20 min at room temperature in buffer (2 %
192 FBS, 0.1 % NaN₃ in PBS) with anti-rat CD45LCA: PE (clone OX-1, 20 μ g/ml), anti-rat

193 CD172a: FITC (clone ED9, 10 µg/ml, Serotec Ltd., Düsseldorf, Germany), anti-rat CD3: PE
194 (clone G4.18, 12,5 µg/ml), anti-rat CD4: FITC (clone OX-38, 50 µg/ml), anti-rat CD8b:
195 FITC (clone 341, 50 µg/ml), or anti-rat NKR-P1A: FITC (clone 10/78, 50 µg/ml). Unless
196 otherwise stated, all antibodies were obtained from BD PharMingen (Heidelberg, Germany).
197 Following incubation, erythrocytes were lysed with FACS lysing solution (BD
198 Immunocytometry Systems) and the percentage of leukocyte and lymphocyte subpopulations
199 was determined on a flow cytometer (FACSCalibur, BD Immunocytometry Systems). Ten
200 thousand cells were analyzed for each sample. Leukocyte subpopulations were identified by
201 forward and sidescatter characteristics and by differences in CD45 and ED9 expression.
202 Lymphocyte subpopulations were identified by characteristic expression of the surface
203 markers CD3⁺/CD4⁺ (T helper cells), CD3⁺/CD8b⁺ (cytotoxic T cells) and CD3⁻/NKR-P1A⁺
204 (NK cells). The percentage of B cells was obtained by subtracting all other subpopulations
205 from the total lymphocytes.

206

207 **2.8 Proliferative response**

208 Peripheral blood mononuclear cells (PBMC) or mesenteric lymph node cells (MLNC) were
209 isolated by Nycoprep density gradient (NycoprepTM 1.077A, Axis-Shield). After washing,
210 cells were adjusted to 1.5 x 10⁶ (PBMC) or 2 x 10⁶ (MLNC) / ml in complete RPMI medium.
211 One hundred µl of this cell suspension were then transferred to each well of a 96-well round-
212 bottomed tissue culture plate. For mitogenic stimulation of lymphocytes, 100 µl of ConA or
213 PWM (both Sigma-Aldrich) in complete RPMI medium were added yielding final
214 concentrations of 0.625 µg/ml (ConA or PWM, MLNC) and 5 µg/ml (ConA, PBMC). Due
215 to limited amounts of blood, no stimulation of PBMC with PWM could be conducted. ConA
216 primarily stimulates T cells and PWM primarily stimulates B cells. Unstimulated controls
217 were treated with complete RPMI medium. All stimulations were set up in triplicates. Cells

218 were incubated at 37 °C, 5 % CO₂ for 48 h, pulsed with 0.5 µCi Methyl-[³H]-thymidine
219 (NEN, Boston) and incubated for an additional 24 h. Cells were harvested on glass filters
220 (Filtermats W/Binding, Molecular Devices) and radioactivity was measured as counts per
221 minute (cpm).

222

223 **2.9 Phagocytosis**

224 Phagocytic activity was determined in whole blood samples. 20 µl of heparinized blood were
225 incubated in 460 µl medium with 10 µl of lucigenin solution (5.1 mg/ml PBS, Sigma) at 37
226 °C, 5 % CO₂ for 30 min. After addition of 10 µl Zymosan A (12.5 mg/ml PBS, Sigma),
227 chemiluminescence was measured for 30 min at 37 °C on a luminometer (Multi Bioluminat
228 LB 9505 C, Berthold, Germany). Results are expressed as total counts per 30 min.

229 Preliminary tests revealed no detectable phagocytic activity for isolates from MLN.

230

231 **2.10 Cytokine production**

232 For the determination of cytokine production in cell culture supernatants, cells were isolated
233 by Nycoprep density gradient, adjusted to 2.5 x 10⁶ (PBMC) or 3 x 10⁶ cells/ml (MLNC)
234 and cultured in 96-well plates as described above. Cells were stimulated with ConA at final
235 concentrations of 5 µg/ml (PBMC) or 1.25 µg/ml (MLNC) for 24 h. Supernatants were
236 centrifuged and stored at -80 °C for later analysis.

237 IL-2 and IFN-γ concentrations in supernatants were determined with a standard ELISA
238 procedure. All antibodies and standards were obtained from Biolegend, San Diego. A 96-well
239 high affinity-protein binding ELISA plate was coated with 2 µg/ml of anti-rat IL-2 (clone:
240 BL-7015) or anti-rat IFN-γ (clone: DB-1) in sodium carbonate buffer and incubated at 4 °C
241 overnight. Nonspecific binding was blocked (10 % FCS in PBS) for 1 h at room temperature.

242 One hundred μl of samples or standard (recombinant rat IL-2 or recombinant rat IFN- γ ,
243 source *E. coli*, in blocking solution) were applied and incubated for 3 h at room temperature.
244 The standard curve (20 ng/ml to 40 pg/ml) was analyzed in triplicates, each sample in
245 duplicates. Then, 0.5 $\mu\text{g/ml}$ biotinylated anti-rat IL-2 (clone BL-7030) or anti-rat IFN- γ
246 (clone Poly5109) in blocking solution were applied and incubated for 1 h at room
247 temperature, followed by incubation with streptavidin-horseradish peroxidase (Southern
248 Biotech, 1:6000 in blocking solution) for 30 min at room temperature. Finally, 50 μl of TMB
249 substrate solution (Biomeda) were applied for 8-12 min, the color reaction was stopped with
250 0.5 M H_2SO_4 , and the absorption measured at 450 nm. Cytokine concentrations were
251 calculated from standard curves created in Origin 7G SR4 for Windows. In the present study,
252 IL-4 concentrations were measured in supernatants from stimulated cell cultures as well, with
253 an IL-4 ELISA using anti-rat IL-4 (clone BL-7045) for capture and biotinylated anti-rat IL-4
254 (clone BL-7060) for detection. In all supernatants, IL-4 concentrations were below the
255 detection limit of the assay (40 pg/ml).

256

257 **2.11 Statistics**

258 Behavioral data were analyzed using the Mann-Whitney *U* test. Immunological and hormone
259 data were analyzed with two-way ANOVA for the factors *stress* and *lactation*. Shapiro Wilk
260 testing for normal distribution and Levene testing for homogeneity of variance were
261 performed prior to ANOVA. Standard transformations were applied to achieve normal
262 distribution and homogeneity of variance. Student's *t*-test was used for *post hoc* analysis.
263 Benjamini-Hochberg corrections of *p*-values were applied to adjust for multiple comparisons.
264 A *p*-value of < 0.05 was considered significant. Relations of immune parameters with
265 corticosterone were analyzed with Pearson's correlation. All statistics were calculated using
266 SPSS 12.0 for Windows.

267 **Results**

268 **2.12 Agonistic behavior of resident and intruder rats**

269 Females from both intruder groups were clearly defeated ($DI \leq 0.2$). However, lactating
270 intruders showed less frequently defensive behavior (Fig. 1A) and were less frequently
271 attacked by the residents (Fig. 1B).

272 **2.13 Serum corticosterone concentrations**

273 Serum corticosterone concentrations were about twofold higher in confronted rats as
274 compared to controls (non-lactating females: 455 ± 34 (confrontation) vs. 227 ± 27 ng/ml
275 (control); lactating: 411 ± 29 vs. 258 ± 33 ng/ml) [*stress*: $F_{1,41} = 36.98$; $p < 0.001$]. Neither a
276 difference between lactating and non-lactating rats [*lactation*: $F_{1,41} = 0.041$; n.s.] nor
277 interaction between factors *stress x lactation* [$F_{1,41} = 1.44$; n.s.] was observed.

278 **2.14 Stress- and lactation-induced changes in leukocyte and lymphocyte subsets**

279 Several leukocyte and lymphocyte subsets were affected by either stress or lactation (Tab. 1).
280 In general, more changes were observed in the blood than in the MLN.
281 *Lymphocyte subsets*: Blood lymphocyte numbers after confrontation were about 40 % lower
282 in both lactating and non-lactating rats [*stress*: $F_{1,41} = 16.73$; $p < 0.001$]. This is
283 predominantly due to a reduction in the numbers of B cells [*stress*: $F_{1,40} = 54.04$; $p < 0.001$]
284 and, to a lesser extent, of NK cells [*stress*: $F_{1,40} = 4.17$; $p < 0.05$]. Generally, lactating rats had
285 lower blood B cell numbers [*lactation*: $F_{1,40} = 17.14$; $p < 0.001$]. In addition, in MLN a
286 tendency towards higher NK cell numbers in lactating animals [*lactation*: $F_{1,48} = 3.22$; $p =$
287 0.079] was observed. T cell populations in both blood and MLN were neither affected by
288 stress nor lactation. No significant interactions between the factors *stress* and *lactation* were
289 observed.

290 *Granulocytes*: In non-lactating animals, blood granulocyte numbers were more than 100 %
291 higher in stressed animals than in non-stressed controls [*stress*: $F_{1,41} = 6.04$; $p < 0.05$]. In
292 contrast, lactating control animals had very high granulocyte numbers even before
293 confrontation, but stress had no additional effect. This asymmetry resulted in a strong
294 interaction of the factors *stress* and *lactation* [$F_{1,41} = 9.6$; $p < 0.01$]. The effect of *lactation*
295 [$F_{1,41} = 2.51$; $p = 0.12$] alone did not reach the level of significance. There was no effect of
296 *stress* and *lactation* on MLN granulocytes.

297 *Monocytes*: Stress had no effect on monocyte numbers in both the blood and the MLN.
298 However, an up to twofold higher monocyte cell number was observed in MLN of lactating
299 rats [$F_{1,48} = 31.65$; $p < 0.001$], while blood monocytes remained unaffected. A significant
300 interaction of factors *stress* and *lactation* was not observed.

301 **2.15 Effects of stress and lactation on lymphocyte proliferation in blood and MLN**

302 Stress had a negative impact on ConA-induced proliferation of blood lymphocytes [$F_{1,32} =$
303 9.42 ; $p < 0.01$], resulting in an up to 60 % decrease (Fig. 2A). In MLN, a slightly negative
304 effect of stress on PWM induced lymphocyte proliferation (a decrease of about 10 %) [$F_{1,48}$
305 $= 5.57$; $p < 0.05$] (Fig. 2C) was evident. The effect on ConA-induced proliferation did not
306 reach level of significance [$F_{1,48} = 2.43$; $p = 0.125$] (Fig. 2B, C).

307 Lactation had no effect on mitogen-induced proliferation of blood lymphocytes [$F_{1,32} = 0$;
308 n.s.]. However, proliferation of MLN lymphocytes was higher in lactating animals in
309 response to both PWM [$F_{1,48} = 32.49$; $p < 0.001$] and ConA [$F_{1,48} = 12.12$; $p < 0.01$] (Fig. 2B,
310 C). No significant *stress* x *lactation* interaction was observed.

311 **2.16 Stress- and lactation-induced effects on whole blood phagocytosis**

312 No significant effect of stress on whole blood phagocytosis was observed [$F_{1,38} = 0.15$; $p =$
313 0.7] (Fig. 2D). Both control and stressed lactating animals had a substantially higher

314 phagocytic capacity [$F_{1, 38} = 22.88$; $p < 0.001$], resulting in up to more than twofold increase.

315 No significant *stress x lactation* interaction was observed.

316

317 **2.17 Effects of stress and lactation on cytokine production**

318 IFN- γ production of blood lymphocytes was strongly reduced in stressed animals [*stress*: $F_{1, 30} = 16.08$;

319 $p < 0.001$] (Fig. 3A), reaching only one fourth of control levels. Lactation had no

320 significant effect on IFN- γ production, and no *stress x lactation* interaction was observed.

321 In contrast, an effect of stress on blood lymphocyte IL-2 production [*stress*: $F_{1, 30} = 14.34$; $p <$

322 0.01] (Fig. 3B) could only be detected in non-lactating females [*stress x lactation*: $F_{1, 30} =$

323 4.97 ; $p < 0.05$; *lactation*: $F_{1, 30} = 2.7$; $p = 0.11$]. Lactating females had a very low IL-2

324 production, and no additional reduction in IL-2 production occurred in response to stressor

325 exposure (Fig. 3B). In MLN, neither an effect of stress nor of lactation on cytokine

326 production was noted (Fig. 3C, D).

327

328 **2.18 Relationship between MLN immune parameters and corticosterone**

329

330 Experimental sets involving MLN included both immune and endocrine measurements from

331 the same individuals, allowing correlation analyses between corticosterone and the twelve

332 MLN immune parameters determined in each of the four experimental groups. No significant

333 relationships between serum corticosterone concentrations and MLN immune parameters

334 were detected.

335

336

337

338 **Discussion**

339

340 In this study we demonstrate that lactation strongly affects the immune system in laboratory
341 rats, but the impact on blood and MLN varies. In blood, a reduced mitogen-induced IL-2
342 production and lower B cell numbers were observed, while an enhanced lymphocyte
343 proliferation was found in MLN. The present findings complement a previous study by
344 Shanks et al. (1997) who reported reduced IL-2 production of splenocytes and increased
345 proliferation of MLN lymphocytes during lactation. A low number of B cells and a decreased
346 IL-2 production in blood and spleen correspond with clinical observations such as a reduced
347 ability to mount antibody response or an increased susceptibility to infection (Drazen et al.,
348 2003; Jäckel, 2003). However, based on the present data, one should not draw the conclusion
349 that the immune system is generally suppressed during lactation: first, functionally active
350 immune cells from the blood or spleen might migrate into other body compartments in
351 lactating females. Thus, it cannot be ruled out that a decreased IL-2 production may be
352 assessed in the blood while simultaneously IL-2 production in other body parts such as the
353 intestines, breasts or the reproductive tract is increased. Second, the increased number of
354 phagocytes in the blood suggests an enhancement of at least some aspects of the innate
355 immunity in the circulation. Third, lymphocyte activity is increased in MLN. During
356 lactation, this enhanced activity appears to be especially beneficial in order to assure an
357 adequate maternal-offspring antibody transfer. B cells from MLN travel to the breast,
358 producing antibodies specific to gut pathogens (Head and Seelig, 1983; Roux et al., 1977).
359 These antibodies (particularly IgA) enter the milk and protect the offspring against gut
360 pathogen-associated diseases (Lamm et al., 1978; Shanks et al., 1997).
361 It has been hypothesized in eco-immunological literature that a trade-off may exist between
362 reproduction and immune activity (reviewed in Harshman and Zera, 2007). According to this

363 concept, particularly during the energy demanding process of lactation (Hansen and Ferreira,
364 1986; Papworth and Clubb, 1995), a reallocation of resources from the immune system may
365 be required to meet the costs of lactation (Coop and Kyriazakis, 1999; Demas et al., 1997;
366 Jäckel, 2003). One problem with this concept is that maintenance of activity levels of resting
367 immune cells does not consume much energy (Krauss et al., 2001; Maciver et al., 2008).
368 Moreover, the present data show that rather than being immunosuppressed, the immune
369 system appears to be readjusted during lactation. This involves both reducing and enhancing
370 effects in different body parts. More studies on the migration pattern of immune cells during
371 lactation and the assessment of their activity levels in various body compartments are thus
372 required before more generalized conclusions can be made.

373 Although the physiological mechanisms underlying the effects of lactation on the immune
374 system were not the focus of this study, it should be mentioned that the diverse effects on
375 immune cells in the blood and in the MLN might be caused by local concentration
376 differences of hormones. Prolactin would be a potential candidate, as this hormone is known
377 to inhibit lymphocyte proliferation at high concentrations while having enhancing effects at
378 lower concentrations (Matera et al., 1992). High plasma prolactin concentrations during
379 lactation (Arbogast and Voogt, 1996; Grattan, 2001) might therefore affect lymphocytes in
380 circulation quite strongly. In addition, several studies describe a differential expression of
381 prolactin receptor isoforms during lactation in lymphoid organs (Feng et al., 1998; Gunes and
382 Mastro, 1997), which possibly also contributes to the discrepancies observed here.

383

384 Acute social stress in non-lactating female rats causes a well known pattern of changes in
385 blood immune cells, which is characterized by granulocytosis, reduction of B cell numbers,
386 and decreased lymphocyte proliferation (Stefanski, 1998; Stefanski and Engler, 1998;
387 Stefanski and Grüner, 2006; Stefanski et al., 1996). The present study extends our limited

388 knowledge on stress-induced alterations in female blood cytokine production. Reduced
389 mitogen-induced IL-2 and IFN- γ production in the blood in concert with previous findings
390 from spleen and popliteal lymph nodes (Brenner and Moynihan, 1997; Iwakabe et al., 1998;
391 Shanks et al., 1997) suggest a shift from a T_h1 to a T_h2 cytokine response as observed during
392 pregnancy (Saito et al., 2007). Glucocorticoids and catecholamines have been identified as
393 key players in mediating stress-induced effects on blood immune cells. It is likely that the
394 same basic hormonal mechanisms are also involved in affecting functional capacity, cytokine
395 production, migration and adhesion, as well as apoptosis of immune cells in lactating and
396 non-lactating female rats (Dhabhar, 2002; Elenkov and Chrousos, 2002; Moynihan, 2003).
397 In contrast to blood, MLN were only marginally affected by acute social stress. This relative
398 resistance of MLN lymphocytes to stress is in agreement with the literature, indicating that
399 the effects in MLN are often less intense than the effects in blood or the spleen (Moraska et
400 al., 2002; Nguyen et al., 2000; Shanks et al., 1997). Possibly, blood leukocytes are more
401 directly exposed to glucocorticoids or catecholamines in the circulation, whereas effective
402 concentrations could be lower in MLN. Furthermore, glucocorticoid receptor expression
403 levels are known to differ between tissues (Miller et al., 1998) and may thus also be lower in
404 MLN. However, to the best of our knowledge, there is no study on local hormone
405 concentrations in MLN. Considering the relative resistance of MLN to stressor action it is
406 also not surprising that correlations between serum corticosterone concentrations and
407 measures of MLN immunity were absent in the present study.

408 Many effects of stress on the immune system are similar in non-lactating and lactating
409 animals, but the present study also reveals a few important differences. One is that stress has
410 no further impact on high granulocyte numbers and low blood IL-2 production in lactating
411 females. It appears that there is a mechanism preventing a strong activation or suppression of
412 some key immune functions. Certainly, a minimum capacity of lymphocytes to produce IL-2

413 is essential to maintain basic immune functions. The “ceiling effect” on granulocytes is not
414 exclusive to lactation (Stefanski et al., 2005) and can also be observed in pregnant rats. Here
415 it was argued that a limitation of neutrophil release into circulation prevents a self-
416 destructive activation of innate immunity. One additional difference between non-lactating
417 and lactating animals is a dampened effect of stress on ConA-induced T cell proliferation. A
418 possible mechanism could be a resistance of lymphocytes to glucocorticoids, which has also
419 been described in other contexts before (Sheridan et al., 2000).

420 In a report particularly relevant for the present study, Shanks (1997) investigated the effect of
421 conditioned stress on immune function in lactating Sprague-Dawley rats. Interestingly,
422 although the present and previous studies indicate that lactation alters the effects of stress on
423 immune function, the pattern of change and the parameter affected was not always uniform.
424 In contrast to the aforementioned study we found a stress-buffering effect of lactation on
425 blood lymphocyte proliferation, but no evidence for an increase in MLN lymphocyte
426 proliferation in stressed lactating rats. Further discrepancies also exist with respect to
427 corticosterone secretion. Plasma corticosterone concentrations in response to social stressors
428 were similar in both lactating and non-lactating rats in the present study. This finding does
429 not agree with several previous studies that observed a buffering effect of lactation on HPA
430 responsiveness (Neumann et al., 1998; Shanks et al., 1997; Shanks et al., 1999; Sibolboro
431 Mezzacappa et al., 2003; Stern et al., 1973; Torner et al., 2002; Windle et al., 1997). Since
432 only corticosterone concentrations were assessed in the present report, the possibility cannot
433 be ruled out that other indices of HPA axis activity such as ACTH concentrations differ in
434 lactating females. On the other hand, our findings of a similar corticosterone response in
435 lactating and non-lactating rats do correspond well with a study in humans involving a
436 psychosocial stressor (Trier social stress test), which also does not report an attenuated HPA
437 response (cortisol and ACTH) during lactation (Altemus et al., 2001). The reasons for these

438 discrepancies are not clear, but might be caused by methodological differences such as the
439 nature and duration of stressor, the last time point of suckling, or the phase of lactation. It is
440 also worthy of note that the condition of the pups (especially when stressed) may diminish
441 the buffering of the corticosterone response in lactating rats (Smotherman et al., 1977;
442 Smotherman et al., 1976). However, this sensitizing effect is unlikely to play a role in the
443 present paradigm because mothers were not exposed to their pups during or after the acute
444 social stressor exposure.

445 Lactating females less frequently displayed defensive behavior as compared to non-lactating
446 intruders, which may lead to the assumption that the differential effect of social stress on the
447 immune system in lactating females is related to behavioral differences. The reduced amount
448 of defensive behavior in lactating animals may be due to reduced anxiety levels, as
449 previously reported in other behavioral settings during lactation (Hansen and Ferreira, 1986;
450 Hard and Hansen, 1985; Sibolboro Mezzacappa et al., 2003; Torner et al., 2002; Toufexis et
451 al., 1999; Windle et al., 1997). Alternatively, a dampened resident aggressiveness may be
452 caused by the specific odor of lactating and pregnant females aimed to protect themselves and
453 the litter from conspecifics' aggression (Clegg and Williams, 1983; Kilpatrick et al., 1983;
454 Mennella and Moltz, 1989; Moltz and Lee, 1981). In any case, lactating females may have
455 perceived social conflict in confrontations as less threatening. This, however, was not
456 reflected in reduced CORT secretion, but may have been associated with lowered
457 sympathetic activation. Since catecholamines have major immuno-modulatory properties
458 (Engler et al., 2004; Schedlowski et al., 1996; Stevenson et al., 2001), a specific role of these
459 sympathetic mediators in lactating females should be addressed in the future.

460 In our study we used a social stressor that allows active coping behavior of the intruder and
461 therefore represents a realistic picture of stressor-induced changes also occurring in natural
462 environments. Lactation- and stress-induced changes of the immune system do not only

463 affect the mother but might also directly influence the offspring via an altered transfer of
464 maternal antibodies, which emphasizes the importance of lactation. In future studies,
465 measurements of prolactin and IgA and also the study of other immune compartments such as
466 spleen, lymph node areas other than MLN, or breast and reproductive organs would provide
467 further useful information. In the present study, we investigated a time of lactation during
468 which milk production is already well established and quite energy demanding for the
469 mother. Nevertheless, a time course over the whole period of lactation would be of interest,
470 since lactation-associated effects might differ at the beginning and towards the end of
471 lactation (normal weaning starts at about day 20). Since the effects of acute and prolonged
472 stressor exposure differ considerably, chronic social stress situations should also be taken into
473 account. The present study nevertheless clearly demonstrates that an acute social stressor
474 significantly alters both behavior and immune system during lactation. The findings
475 contribute to an understanding of the complex interactions between the female reproduction
476 system and a stressor.

477

478

479

480 **Acknowledgement**

481 Parts of this study were supported by the DFG (STE 633 /5-1). We gratefully acknowledge
482 the technical assistance of Mrs. Andrea Berger and Inge Zerenner-Fritsche.

483

484

485

486

487 **References**

488

489 Altemus, M., Redwine, L.S., Leong, Y.M., Frye, C.A., Porges, S.W., Carter, C.S., 2001.

490 Responses to laboratory psychosocial stress in postpartum women. *Psychosom Med*

491 63, 814-821.

492 Amino, N., Mori, H., Iwatani, Y., Tanizawa, O., Kawashima, M., Tsuge, I., Ibaragi, K.,

493 Kumahara, Y., Miyai, K., 1982. High prevalence of transient post-partum

494 thyrotoxicosis and hypothyroidism. *N Engl J Med* 306, 849-852.

495 Arbogast, L.A., Voogt, J.L., 1996. The responsiveness of tuberoinfundibular dopaminergic

496 neurons to prolactin feedback is diminished between early lactation and midlactation

497 in the rat. *Endocrinology* 137, 47-54.

498 Avitsur, R., Stark, J.L., Dhabhar, F.S., Sheridan, J.F., 2002. Social stress alters splenocyte

499 phenotype and function. *J Neuroimmunol* 132, 66-71.

500 Barger, I.A., 1993. Influence of sex and reproductive status on susceptibility of ruminants to

501 nematode parasitism. *Int J Parasitol* 23, 463-469.

502 Ben-Eliyahu, S., Shakhar, G., Shakhar, K., Melamed, R., 2000. Timing within the oestrous

503 cycle modulates adrenergic suppression of NK activity and resistance to metastasis:

504 possible clinical implications. *Br J Cancer* 83, 1747-1754.

505 Brenner, G.J., Moynihan, J.A., 1997. Stressor-induced alterations in immune response and

506 viral clearance following infection with herpes simplex virus-type 1 in BALB/c and

507 C57B1/6 mice. *Brain Behav Immun* 11, 9-23.

508 Buchel, E., Van Steenberghe, W., Nevens, F., Fevery, J., 2002. Improvement of autoimmune

509 hepatitis during pregnancy followed by flare-up after delivery. *Am J Gastroenterol* 97,

510 3160-3165.

- 511 Carter, C.S., Altemus, M., Chrousos, G.P., 2001. Neuroendocrine and emotional changes in
512 the post-partum period. *Prog Brain Res* 133, 241-249.
- 513 Clegg, F., Williams, D.I., 1983. Maternal pheromone in *Rattus norvegicus*. *Behav Neural*
514 *Biol* 37, 223-236.
- 515 Coop, R.L., Kyriazakis, I., 1999. Nutrition-parasite interaction. *Vet Parasitol* 84, 187-204.
- 516 Demas, G.E., Chefer, V., Talan, M.I., Nelson, R.J., 1997. Metabolic costs of mounting an
517 antigen-stimulated immune response in adult and aged C57BL/6J mice. *Am J Physiol*
518 273, R1631-1637.
- 519 Dhabhar, F.S., 2002. Stress-induced augmentation of immune function--the role of stress
520 hormones, leukocyte trafficking, and cytokines. *Brain Behav Immun* 16, 785-798.
- 521 Drazen, D.L., Trasy, A., Nelson, R.J., 2003. Photoperiod differentially affects energetics of
522 immunity in pregnant and lactating Siberian hamsters (*Phodopus sungorus*) *Can J Zoo*
523 81, 1406-1413.
- 524 Elenkov, I.J., Chrousos, G.P., 2002. Stress hormones, proinflammatory and antiinflammatory
525 cytokines, and autoimmunity. *Ann N Y Acad Sci* 966, 290-303.
- 526 Engler, H., Dawils, L., Hoves, S., Kurth, S., Stevenson, J.R., Schauenstein, K., Stefanski, V.,
527 2004. Effects of social stress on blood leukocyte distribution: the role of alpha- and
528 beta-adrenergic mechanisms. *J Neuroimmunol* 156, 153-162.
- 529 Engler, H., Stefanski, V., 2003. Social stress and T cell maturation in male rats: transient and
530 persistent alterations in thymic function. *Psychoneuroendocrinology* 28, 951-969.
- 531 Feng, J.C., Loh, T.T., Sheng, H.P., 1998. Lactation increases prolactin receptor expression in
532 spleen and thymus of rats. *Life Sci* 63, 111-119.
- 533 Festa-Bianchet, M.J., 1989. Individual differences, parasites, and the costs of reproduction for
534 Bighorn Ewes (*Ovis Canadensis*). *J Anim Ecol* 58, 785-795.

- 535 Foster, L.B., Dunn, R.T., 1974. Single-antibody technique for radioimmunoassay of cortisol
536 in unextracted serum or plasma. *Clin Chem* 20, 365-368.
- 537 Gammie, S.C., Hasen, N.S., Stevenson, S.A., Bale, T.L., D'Anna, K.L., 2005. Elevated stress
538 sensitivity in corticotropin-releasing factor receptor 2 deficient mice decreases
539 maternal, but not intermale aggression. *Behav Brain Res* 160, 169-177.
- 540 Grattan, D.R., 2001. The actions of prolactin in the brain during pregnancy and lactation.
541 *Prog Brain Res* 133, 153-171.
- 542 Groer, M.W., Davis, M.W., Smith, K., Casey, K., Kramer, V., Bukovsky, E., 2005.
543 Immunity, inflammation and infection in post-partum breast and formula feeders. *Am*
544 *J Reprod Immunol* 54, 222-231.
- 545 Gunes, H., Mastro, A.M., 1997. Prolactin receptor gene expression in rat splenocytes and
546 thymocytes during oestrous cycle, pregnancy and lactation. *Cell Prolif* 30, 219-235.
- 547 Hansen, S., Ferreira, A., 1986. Food intake, aggression, and fear behavior in the mother rat:
548 control by neural systems concerned with milk ejection and maternal behavior. *Behav*
549 *Neurosci* 100, 64-70.
- 550 Hard, E., Hansen, S., 1985. Reduced fearfulness in the lactating rat. *Physiol Behav* 35, 641-
551 643.
- 552 Harshman, L.G., Zera, A.J., 2007. The cost of reproduction: the devil in the details. *Trends*
553 *Ecol Evol* 22, 80-86.
- 554 Head, J.R., Seelig, L.L., Jr., 1983. Autoradiographic analysis of lymphocyte migration into
555 the mammary epithelium and milk of lactating female rats. *J Reprod Immunol* 5, 61-
556 72.
- 557 Iwakabe, K., Shimada, M., Ohta, A., Yahata, T., Ohmi, Y., Habu, S., Nishimura, T., 1998.
558 The restraint stress drives a shift in Th1/Th2 balance toward Th2-dominant immunity
559 in mice. *Immunol Lett* 62, 39-43.

- 560 Jäckel, M., 2003. Die Auswirkungen der Reproduktion auf das Immunsystem weiblicher
561 Hausmeerschweinchen (*Cavia porcellus*). Dissertation, Bielefeld.
- 562 Kilpatrick, S.J., Lee, T.M., Moltz, H., 1983. The maternal pheromone of the rat: testing some
563 assumptions underlying a hypothesis. *Physiol Behav* 30, 539-543.
- 564 Krauss, S., Brand, M.D., Buttgereit, F., 2001. Signaling takes a breath--new quantitative
565 perspectives on bioenergetics and signal transduction. *Immunity* 15, 497-502.
- 566 Lamm, M.E., Weisz-Carrington, P., Roux, M.E., McWilliams, M., Phillips-Quagliata, J.M.,
567 1978. Development of the IgA system in the mammary gland. *Adv Exp Med Biol*
568 107, 35-42.
- 569 Maciver, N.J., Jacobs, S.R., Wieman, H.L., Wofford, J.A., Coloff, J.L., Rathmell, J.C., 2008.
570 Glucose metabolism in lymphocytes is a regulated process with significant effects on
571 immune cell function and survival. *J Leukoc Biol* 84, 949-957.
- 572 Magiakou, M.A., Mastorakos, G., Webster, E., Chrousos, G.P., 1997. The hypothalamic-
573 pituitary-adrenal axis and the female reproductive system. *Ann N Y Acad Sci* 816,
574 42-56.
- 575 Mann, P.E., Bridges, R.S., 2002. Prolactin receptor gene expression in the forebrain of
576 pregnant and lactating rats. *Brain Res Mol Brain Res* 105, 136-145.
- 577 Martin, P., Bateson, P., 1993. *Measuring behaviour: recording methods*. Cambridge
578 University Press.
- 579 Matera, L., Cesano, A., Bellone, G., Oberholtzer, E., 1992. Modulatory effect of prolactin on
580 the resting and mitogen-induced activity of T, B, and NK lymphocytes. *Brain Behav*
581 *Immun* 6, 409-417.
- 582 Mennella, J.A., Moltz, H., 1989. Pheromonal emission by pregnant rats protects against
583 infanticide by nulliparous conspecifics. *Physiol Behav* 46, 591-595.

- 584 Miller, A.H., Spencer, R.L., Pearce, B.D., Pisell, T.L., Azrieli, Y., Tanapat, P., Moday, H.,
585 Rhee, R., McEwen, B.S., 1998. Glucocorticoid receptors are differentially expressed
586 in the cells and tissues of the immune system. *Cell Immunol* 186, 45-54.
- 587 Moltz, H., Lee, T.M., 1981. The maternal pheromone of the rat: identity and functional
588 significance. *Physiol Behav* 26, 301-306.
- 589 Morag, M., Popliker, F., Yagil, R., 1975. Effect of litter size on milk yield in the rat. *Lab*
590 *Anim* 9, 43-47.
- 591 Moraska, A., Campisi, J., Nguyen, K.T., Maier, S.F., Watkins, L.R., Fleshner, M., 2002.
592 Elevated IL-1beta contributes to antibody suppression produced by stress. *J Appl*
593 *Physiol* 93, 207-215.
- 594 Moynihan, J.A., 2003. Mechanisms of stress-induced modulation of immunity. *Brain Behav*
595 *Immun* 17 Suppl 1, S11-16.
- 596 Muller, A.F., Drexhage, H.A., Berghout, A., 2001. Postpartum thyroiditis and autoimmune
597 thyroiditis in women of childbearing age: recent insights and consequences for
598 antenatal and postnatal care. *Endocr Rev* 22, 605-630.
- 599 Nelson, J.L., Ostensen, M., 1997. Pregnancy and rheumatoid arthritis. *Rheum Dis Clin North*
600 *Am* 23, 195-212.
- 601 Neumann, I.D., Johnstone, H.A., Hatzinger, M., Liebsch, G., Shipston, M., Russell, J.A.,
602 Landgraf, R., Douglas, A.J., 1998. Attenuated neuroendocrine responses to emotional
603 and physical stressors in pregnant rats involve adenohipophysial changes. *J Physiol*
604 508, 289-300.
- 605 Neumann, I.D., Torner, L., Wigger, A., 2000. Brain oxytocin: differential inhibition of
606 neuroendocrine stress responses and anxiety-related behaviour in virgin, pregnant and
607 lactating rats. *Neuroscience* 95, 567-575.

- 608 Nguyen, K.T., Deak, T., Will, M.J., Hansen, M.K., Hunsaker, B.N., Fleshner, M., Watkins,
609 L.R., Maier, S.F., 2000. Timecourse and corticosterone sensitivity of the brain,
610 pituitary, and serum interleukin-1beta protein response to acute stress. *Brain Res* 859,
611 193-201.
- 612 Ngwenya, B.Z., 1977. Response of nonsensitized and sensitized lactating mice to infection
613 with trichinella spiralis. *Int J Parasitol* 7, 41-45.
- 614 Page, G.G., Ben-Eliyahu, S., 1997. Increased surgery-induced metastasis and suppressed
615 natural killer cell activity during proestrus/estrus in rats. *Breast Cancer Res Treat* 45,
616 159-167.
- 617 Papworth, T.A., Clubb, S.K., 1995. Clinical pathology in the female rat during the pre- and
618 postnatal period. *Comp Haematol Int* 5, 13-24.
- 619 Redwine, L.S., Altemus, M., Leong, Y.M., Carter, C.S., 2001. Lymphocyte responses to
620 stress in postpartum women: relationship to vagal tone. *Psychoneuroendocrinology*
621 26, 241-251.
- 622 Roux, M.E., McWilliams, M., Phillips-Quagliata, J.M., Weisz-Carrington, P., Lamm, M.E.,
623 1977. Origin of IgA-secreting plasma cells in the mammary gland. *J Exp Med* 146,
624 1311-1322.
- 625 Russell, J.A., 1980. Milk yield, suckling behaviour and milk ejection in the lactating rat
626 nursing litters of different sizes. *J Physiol* 303, 403-415.
- 627 Saito, S., Shiozaki, A., Nakashima, A., Sakai, M., Sasaki, Y., 2007. The role of the immune
628 system in preeclampsia. *Mol Aspects Med* 28, 192-209.
- 629 Schedlowski, M., Hosch, W., Oberbeck, R., Benschop, R.J., Jacobs, R., Raab, H.R., Schmidt,
630 R.E., 1996. Catecholamines modulate human NK cell circulation and function via
631 spleen-independent beta 2-adrenergic mechanisms. *J Immunol* 156, 93-99.

- 632 Shanks, N., Kusnecov, A., Pezzone, M., Berkun, J., Rabin, B.S., 1997. Lactation alters the
633 effects of conditioned stress on immune function. *Am J Physiol* 272, R16-25.
- 634 Shanks, N., Windle, R.J., Perks, P., Wood, S., Ingram, C.D., Lightman, S.L., 1999. The
635 hypothalamic-pituitary-adrenal axis response to endotoxin is attenuated during
636 lactation. *J Neuroendocrinol* 11, 857-865.
- 637 Sheridan, J.F., Stark, J.L., Avitsur, R., Padgett, D.A., 2000. Social disruption, immunity, and
638 susceptibility to viral infection. Role of glucocorticoid insensitivity and NGF. *Ann N*
639 *Y Acad Sci* 917, 894-905.
- 640 Shubber, A.H., Lloyd, S., Soulsby, E.J., 1981. Infection with gastrointestinal helminths.
641 Effect of lactation and maternal transfer of immunity. *Z Parasitenkd* 65, 181-189.
- 642 Shurin, M.R., Kusnecov, A., Hamill, E., Kaplan, S., Rabin, B.S., 1994. Stress-induced
643 alteration of polymorphonuclear leukocyte function in rats. *Brain Behav Immun* 8,
644 163-169.
- 645 Sibolboro Mezzacappa, E., Tu, A.Y., Myers, M.M., 2003. Lactation and weaning effects on
646 physiological and behavioral response to stressors. *Physiol Behav* 78, 1-9.
- 647 Smotherman, W.P., Mendoza, S.P., Levine, S., 1977. Ontogenetic changes in pup-elicited
648 maternal pituitary-adrenal activity: pup age and stage of lactation effects. *Dev*
649 *Psychobiol* 10, 365-371.
- 650 Smotherman, W.P., Wiener, S.G., Mendoza, S.P., Levine, S., 1976. Pituitary--adrenal
651 responsiveness of rat mothers to noxious stimuli and stimuli produced by pups. *Ciba*
652 *Found Symp*, 5-25.
- 653 Stefanski, V., 1998. Social stress in loser rats: opposite immunological effects in submissive
654 and subdominant males. *Physiol Behav* 63, 605-613.
- 655 Stefanski, V., 2000. Social stress in laboratory rats: hormonal responses and immune cell
656 distribution. *Psychoneuroendocrinology* 25, 389-406.

- 657 Stefanski, V., Engler, H., 1998. Effects of acute and chronic social stress on blood cellular
658 immunity in rats. *Physiol Behav* 64, 733-741.
- 659 Stefanski, V., Grüner, S., 2006. Gender difference in basal and stress levels of peripheral
660 blood leukocytes in laboratory rats. *Brain Behav Immun* 20, 369-377.
- 661 Stefanski, V., Raabe, C., Schulte, M., 2005. Pregnancy and social stress in female rats:
662 influences on blood leukocytes and corticosterone concentrations. *J Neuroimmunol*
663 162, 81-88.
- 664 Stefanski, V., Solomon, G.F., Kling, A.S., Thomas, J., Plaeger, S., 1996. Impact of social
665 confrontation on rat CD4 T cells bearing different CD45R isoforms. *Brain Behav*
666 *Immun* 10, 364-379.
- 667 Stern, J.M., Goldman, L., Levine, S., 1973. Pituitary-adrenal responsiveness during lactation
668 in rats. *Neuroendocrinology* 12, 179-191.
- 669 Stevenson, J.R., Westermann, J., Liebmann, P.M., Hortner, M., Rinner, I., Felsner, P.,
670 Wolfler, A., Schauenstein, K., 2001. Prolonged alpha-adrenergic stimulation causes
671 changes in leukocyte distribution and lymphocyte apoptosis in the rat. *J*
672 *Neuroimmunol* 120, 50-57.
- 673 Theodosis, D.T., Poulain, D.A., Vincent, J.D., 1981. Possible morphological bases for
674 synchronisation of neuronal firing in the rat supraoptic nucleus during lactation.
675 *Neuroscience* 6, 919-929.
- 676 Torner, L., Toschi, N., Nava, G., Clapp, C., Neumann, I.D., 2002. Increased hypothalamic
677 expression of prolactin in lactation: involvement in behavioural and neuroendocrine
678 stress responses. *Eur J Neurosci* 15, 1381-1389.
- 679 Toufexis, D.J., Rochford, J., Walker, C.D., 1999. Lactation-induced reduction in rats'
680 acoustic startle is associated with changes in noradrenergic neurotransmission. *Behav*
681 *Neurosci* 113, 176-184.

- 682 van Walderveen, M.A., Tas, M.W., Barkhof, F., Polman, C.H., Frequin, S.T., Hommes, O.R.,
683 Valk, J., 1994. Magnetic resonance evaluation of disease activity during pregnancy in
684 multiple sclerosis. *Neurology* 44, 327-329.
- 685 Watanabe, M., Iwatani, Y., Kaneda, T., Hidaka, Y., Mitsuda, N., Morimoto, Y., Amino, N.,
686 1997. Changes in T, B, and NK lymphocyte subsets during and after normal
687 pregnancy. *Am J Reprod Immunol* 37, 368-377.
- 688 Windle, R.J., Wood, S., Shanks, N., Perks, P., Conde, G.L., da Costa, A.P., Ingram, C.D.,
689 Lightman, S.L., 1997. Endocrine and behavioural responses to noise stress:
690 comparison of virgin and lactating female rats during non-disrupted maternal activity.
691 *J Neuroendocrinol* 9, 407-414.
- 692 Wise, D.A., 1974. Aggression in the female golden hamster: effects of reproductive state and
693 social isolation. *Horm Behav* 5, 235-250.
- 694

695

696 **Fig. 1**

697 Agonistic behaviour of female resident and intruder rats. Non-lactating and lactating rats
698 were exposed to resident-intruder confrontations for 2 h. Behaviour was monitored for the
699 first 30 min of confrontation. Results are given as median \pm interquartile range. *subm.*:
700 submissive; *sidew.*: sideway; *appr*: approach; *full def.*: full defense; n = 34 for each group;
701 Statistics: Mann-Whitney *U*-test; * $p < 0.05$; *** $p < 0.001$.

702

703

704 **Fig. 2**

705 Effects of lactation and confrontation on lymphocyte proliferation in blood and MLN and
706 phagocytosis in whole blood. White bars: no lactation; grey bars: lactation. Data are given as
707 mean \pm SEM. Numbers of animals are indicated at the bottom of the bars. Statistics: *t*-test
708 with Benjamini-Hochberg correction; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

709

710

711 **Fig. 3**

712 Effects of lactation and confrontation on ConA-induced cytokine production in lymphocytes
713 from blood and MLN. IFN- γ and IL-2 concentrations in cell culture supernatants. White bars:
714 no lactation; grey bars: lactation. Data are given as mean \pm SEM. Numbers of animals are
715 indicated at the bottom of the bars. Statistics: *t*-test with Benjamini-Hochberg correction; * $p <$
716 0.05 ; ** $p < 0.01$; *int.*: interaction *stress x lactation* ($p < 0.05$)

717

718

719

720