

SURFICIAL GEOLOGY OF MADISON COUNTY, ILLINOIS

David A. Grimley and Andrew C. Phillips
2006

Illinois Department of Natural Resources
ILLINOIS STATE GEOLOGICAL SURVEY
William W. Shilts, Chief

This surficial geology map is a compilation of 1:24,000-scale quadrangle maps funded by the STATEMAP component of the National Cooperative Geologic Mapping Program and undertaken from 1997–2006. Eleven published 7.5-minute quadrangle maps (listed below), 4 unpublished maps (contract reports cited below) as well as surficial geology mapping of remaining portions of Madison County were merged digitally. Slight modifications were made to previously mapped quadrangles where more recent work has improved our understanding of the geology or for consistency of unit mapping. Detailed descriptions of map units as well as extensive text and cross sections for each quadrangle are available with the published quadrangles. Digital products are available on the ISGS website (http://www.isgs.uiuc.edu/isgshome/online_maps_data.htm). Data used to compose the surficial map include the county soil report (Goddard and Sabata 1982), outcrops, and subsurface boring information from stratigraphic tests, engineering tests, coal tests, and water wells.

The surficial geology of the county varies widely from thick alluvium in the broad Mississippi River valley (known as the American Bottoms) in the southwest to thin drift (usually < 50 feet) and abundantly exposed Paleozoic bedrock in the northwest (west of Alton) to thick loess cover in south-central areas (Edwardsville to Collinsville) to ice-contact deposits in the "ridge-drift" in eastern areas. Madison County was covered by continental glaciers twice by ice advancing from the northeast during the Quaternary Period, first, during the pre-Illinois episode and second, during the Illinois Episode (Willman and Frye 1970; Hansel and Johnson 1996; Grimley et al. 2001). During the Wisconsin Episode (last regional glaciation), the area was not covered by ice but did receive glacial meltwater from the north that deposited silt, sand, and gravel (outwash) in the Mississippi Valley. Deposition of silty waterlain deposits in the American Bottoms by prevailing westerly winds during the last glaciation resulted in a significant cover of loess deposits (dashed-line contours on map). The loess is thickest (up to 95 feet) at the bluffs immediately east of the broad Mississippi Valley and thins to the east and northeast. Mississippian bedrock is exposed in areas west of Alton along the bluffs of the Mississippi River and along nearby tributaries. To the east of Alton, Pennsylvanian bedrock is exposed sporadically near bedrock topographic highs where drift has been eroded along ravines and creeks.

Drift thickness (Quaternary deposits) is typically 25 to 150 feet in the county, but can be up to 200 feet over infilled preglacial bedrock valleys (north-central areas). Pre-Illinois episode deposits, up to 125 feet thick, include preglacial Quaternary alluvium (Canteen member, Banner Formation), till and ice-marginal deposits (Banner Formation), clayey and sandy deposits (Banner Formation) up to 145 feet thick in terraces near Wood River, and lake sediment (Equality Formation), up to 100 feet thick. The lake sediment was deposited in backwater areas, now terraces, along Cahokia Creek, Indian Creek, and Wood River valleys during last-glacial aggradation of the Mississippi River. Postglacial deposits (up to 60 feet thick) include various alluvial deposits (facies of the Cahokia Formation); clayey and sandy deposits are distinguished only in the Mississippi Valley) as well as colluvial deposits (Peyton Formation) along steep slopes in northwestern areas. Alluvial fans (Cahokia Formation, fan facies) are mapped at the base of the eastern bluffs of the American Bottoms, where thick loess deposits have been mobilized and redeposited on the eastern edge of the valley. Areas of anthropogenic fill (disturbed ground) are mapped in industrial areas, landfills, and the many interstate interchanges, particularly in the American Bottoms.

References


- Goddard, T.M. and L.R. Sabata, 1982. Soil survey: Madison County, Illinois: University of Illinois Agricultural Experiment Station and United States Dept. of Agriculture.
- Grimley, D.A., A.C. Phillips, L.R. Follmer, H. Wang, and R.S. Nelson, 2001. Quaternary and environmental geology of the St. Louis Metro East area, in David Malone, ed., Guidebook for Field Trip for the 35th Annual Meeting of the North-Central Section of the Geological Society of America: Illinois State Geological Survey Guidebook 33, p. 21–73.
- Hansel, A.K., and W.H. Johnson, 1996. Wedron and Mason Groups: Lithostratigraphic reclassification of deposits of the Wisconsin Episode, Lake Michigan Lobe. Illinois State Geological Survey, Bulletin 104, 116 p.
- Willman, H.B., and J.C. Frye, 1970. Pleistocene stratigraphy of Illinois: Illinois State Geological Survey, Bulletin 94, 204 p.

Published Maps

- Grimley, D.A., 1999. Surficial geology map, Alton Quadrangle (Illinois portion), Madison County, IL: Illinois State Geological Survey, Illinois Geologic Quadrangle Map, IGQ Alton SG, 1:24,000.
- Grimley, D.A., 2002. Surficial geology map, Elshah Quadrangle, Jersey and Madison Counties, IL: Illinois State Geological Survey, Illinois Geologic Quadrangle Map, IGQ Elshah-SG, scale 1:24,000.
- Grimley, D.A., 2004. Surficial geology of Worden Quadrangle, Madison County, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Worden-SG, 1:24,000.
- Grimley, D.A., 2005. Surficial geology of Bethalto Quadrangle, Madison and Macoupin Counties, Illinois: Illinois State Geological Survey, Illinois Geologic Quadrangle Map, IGQ Bethalto-SG, 1:24,000.
- Grimley, D.A., 2005. Surficial geology of New Douglas Quadrangle, Madison and Macoupin Counties, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM New Douglas-SG, 1:24,000.
- Grimley, D.A. and S.W. Lepley, 2005. Surficial geology of Wood River Quadrangle, Madison County, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Wood River-SG, 1:24,000.
- Grimley, D.A. and A.C. Phillips, 2005. Surficial geology of Grantfork Quadrangle, Madison County, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Grantfork-SG, 1:24,000.
- Phillips, A.C., 2004. Surficial geology of Collinsville Quadrangle, Madison and St. Clair Counties, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Collinsville-SG, 1:24,000.
- Phillips, A.C., 2004. Surficial geology of St. Jacob Quadrangle, Madison and St. Clair Counties, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM St. Jacob-SG, 1:24,000.
- Phillips, A.C., 2005. Surficial geology of Highland Quadrangle, Madison and St. Clair Counties, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Highland-SG, 1:24,000.
- Phillips, A.C., and D.A. Grimley, 2004. Surficial geology of Marine Quadrangle, Madison County, Illinois: Illinois State Geological Survey, Illinois Preliminary Geologic Map, IPGM Marine-SG, 1:24,000.


Contract Reports

- Grimley, D.A., A.C. Phillips, and S.W. Lepley, 2000. Surficial geology of Monks Mound Quadrangle: Contract report, U.S. Geological Survey, National Cooperative Geologic Mapping Program 00HQAG0151: Illinois State Geological Survey, Champaign, IL.
- Phillips, A.C., 2003. Surficial geology of Edwardsville Quadrangle: Contract report, U.S. Geological Survey, National Cooperative Geologic Mapping Program 02HQAG0042: Illinois State Geological Survey, Champaign, IL.
- Phillips, A.C., 2003. Surficial geology of Prairieville Quadrangle: Contract report, U.S. Geological Survey, National Cooperative Geologic Mapping Program 02HQAG0042: Illinois State Geological Survey, Champaign, IL.
- Phillips, A.C., D.A. Grimley, and S.W. Lepley, 2000. Surficial geology of Granite City Quadrangle: Contract report, U.S. Geological Survey, National Cooperative Geologic Mapping Program 00HQAG0151: Illinois State Geological Survey, Champaign, IL.





Base map compiled by Illinois State Geological Survey from 1:100,000-scale Digital Line Graph data provided by the United States Geological Survey.

North American Datum of 1983 (NAD 83)
Lambert Conformal Conic Projection



Released by the authority of the State of Illinois: 2006



Geology based on field work and data compilation by D. Grimley and A. Phillips, 1997–2006.

Digital cartography by J. Carrell and J. Domier, Illinois State Geological Survey.

This Illinois Preliminary Geologic Map (IPGM) is a lightly edited product, subject to less scientific and cartographic review than our Illinois Geological Quadrangle (IGQ) series. It will not necessarily correspond to the format of IGQ series maps, or to those of other IPGM series maps. Whether or when this map will be upgraded depends on the resources and priorities of the ISGS.

This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

The Illinois State Geological Survey, the Illinois Department of Natural Resources, and the State of Illinois make no guarantee, expressed or implied, regarding the correctness of the interpretations presented in this document and accept no liability for the consequences of decisions made by others on the basis of the information presented here. The geologic interpretations are based on data that may vary with respect to accuracy of geographic location, the type and quantity of data available at each location, and the scientific and technical qualifications of the data sources. Maps or cross sections in this document are not meant to be enlarged.

ROAD CLASSIFICATION




Illinois Preliminary Geologic Map IPGM Madison County-SG

QUATERNARY DEPOSITS

Description	Unit	Interpretation
HUDSON EPISODE (~12,000 years before present (B.P.) to today)		
Disturbed ground	dg	Man-made fill or excavations
Fill or removed earth; various sediment types		
Mainly silt, silty clay and fine sand; weakly to well stratified; includes some coarser beds	Cahokia Formation (undivided) c	River deposits (alluvium); mapped in valleys tributary to the Mississippi Valley
Silt loam with thin fine sand beds; weakly stratified	Cahokia Formation (fan facies) cf	Alluvial fan deposits; mainly reworked loess from bluffs east of American Bottoms
Silty clay loam, silty clay, and silty loam; massive to stratified; some fine sand lenses	Cahokia Formation (clayey facies) cc	Overbank alluvium, abandoned channel and swale fills; mapped only in Mississippi Valley
Very fine, fine and medium sand; stratified; moderately to well sorted	Cahokia Formation (sandy facies) cs	Alluvium; point bar and channel deposits; mapped only in Mississippi Valley
Silt loam, pebbly silt loam or pebbly silty clay diamicton	Peyton Formation p	Sediments moved downslope by gravity (colluvium); creep layers, slumps, or landslides
Silty clay to silt with some fine sand; massive to stratified	Cahokia or Equality Formation (undifferentiated) ce	Overbank alluvium or lake deposits; occurs on or near the Wood River Terrace
WISCONSIN EPISODE (~75,000–12,000 years B.P.)		
Silt loam to silty clay loam with some fine sand; massive to stratified	Equality Formation e	Lake deposits; of slackwater origin during peak Mississippi River aggradation
	(hachures on map where buried)	
Fine, medium and coarse sand; stratified; generally coarsens at depth; some gravelly zones	Henry Formation h	Outwash (glacial meltwater deposits); on Wood River Terrace in northern American Bottoms
Silt loam; massive; upper 3/5 of unit is more tan or gray (Peoria); lower portion has pinkish hue (Roxana)	Peoria and Roxana Silt pr	Loess (windblown silt); blankets all uplands; thickness contours shown on map; thin eastward from Mississippi Valley bluffs
ILLINOIS EPISODE (~200,000–130,000 years B.P.)		
Mixture of loam, sand and gravel, and diamicton; weakly stratified; generally massive; includes some sand and gravel lenses (especially in upper portion)	Hagarstown Member, Pearl Formation p-h	Ice-contact sediments; deposited mainly in ice-marginal, subglacial, or supraglacial channels; locally includes glacio-tectonic faulting and deformation
	(stipples on map where buried)	
Sand with some gravel; stratified; may include silty or clayey zones, especially near surface	Pearl Formation (outwash facies) p	Outwash; common in loess-covered terraces along Silver Creek
	(hachures on map where buried)	
Pebbly loam diamicton (mixture of clay, silt, sand, and gravel); generally massive; includes some sand and gravel lenses (especially in upper portion)	Glasford Formation (< 5 feet of loess cover) g	Till and ice marginal deposits; includes subglacial and supraglacial deposits
Silt loam to silty clay loam; massive to weakly stratified; locally fossiliferous	Petersburg Silt (cross sections only) pb	Lake sediment; deposited under slackwater conditions of ice marginal settings
PRE-ILLINOIS EPISODE (~700,000–400,000 years B.P.)		
Pebbly silty clay loam diamicton; generally massive; includes sand and gravel lenses, zones of stratified silt near base	Banner Formation, (undivided) (cross sections only) b	Till and ice marginal deposits; common in bluff areas west of Alton; includes Pennsylvanian and Mississippian rocks
Silty clay loam, silty clay, and silt loam; weakly stratified; contains some fine sand beds	Banner Formation, Canteen member (cross sections only) b-c	Preglacial alluvium and lake deposits; may include some residuum or colluvium at base; occurs mainly in preglacial bedrock valleys

PRE-QUATERNARY DEPOSITS

Description	Unit	Interpretation
Shale, siltstone, limestone, and sandstone; less commonly beds of coal and underlay	Pennsylvanian or Mississippian bedrock P, M	Bedrock outcrops or bedrock within 5 feet of land surface; most common in bluff areas west of Alton; includes Pennsylvanian and Mississippian rocks
<p>— — — — — Illinois Episode till border (areas to southwest of line are driftless)</p> <p>— 15 — — — — — Loess thickness contour (in feet)</p> <p>A—A' Line of cross section</p> <p>Note: Loess contours show the combined thickness of Peoria and Roxana Silts on uneroded upland areas. The actual thickness at a given spot may be much less, especially along valley slopes where post-depositional erosion of loess has been significant (see cross sections).</p>		



Cross Sections

	Sand, may contain some gravel or silt
	Sand with gravel
	Laminated silt and clay
	Mixture of loam, sand and gravel, diamicton, and/or silty clay
	Diamicton, massive silt, or other fine-grained sediment
	Contact
	Inferred contact

Horizontal scale: 1 inch = ~8,333 feet
Vertical scale: 1 inch = 200 feet
Vertical exaggeration: ~42x