## PROGRESS ON OPTICAL ROTATIONAL COOLING OF SiO+

## PATRICK R STOLLENWERK, YEN-WEI LIN, BRIAN C. ODOM, Department of Physics and Astronomy, Northwestern University, Evanston, IL, USA.

provided by Illinois Digital Environment for Ac

Producing ultracold molecules is the first step in precision molecular spectroscopy. Here we present some of the challenges and advantages of SiO+ as well as some of our progress toward meeting those challenges. To demonstrate ground state SiO+, we first load about 100 SiO+ via 2+1 REMPI into an ion trap. Translational motion of SiO+ is then sympathetically cooled by co-trapped Ba+, which is laser cooled. To prepare the population into the ground state, we optically pump the P-branch (rotational cooling transitions) in the B: $\Sigma(v'=0) \leftarrow X:\Sigma(v=0)$  band with broadband radiation. Because the band is highly diagonal, population can be effectively driven into the rotational ground state before falling into other manifolds. The broadband source, a fs laser, is spectrally filtered using an ultrashort pulse shaping technique to drive only the P-branch. Attention must be paid when aligning the optics to obtain sufficient masking resolution. We have achieved 3 cm<sup>-1</sup> resolution, which is sufficient to modify a broadband source for rotationally cooling SiO+.