WEIGHT DISTRIBUTION FORMULA FOR SOME CLASS OF CYCLIC CODES

Tadao Kasami

UNIVERSITY OF ILLINOIS -URBANA, ILLINOIS

REPORT R-285

APRIL, 1966

Coordinated Science Laboratory



WEIGHT DISTRIBUTION FORMULA FOR SOME CLASS OF CYCLIC CODES

Tadao Kasami

REPORT R-285

J

And Community

APRIL, 1966

This work was supported in part by the Joint Services Electronics Program (U. S. Army, U. S. Navy, and U. S. Air Force) under Contract No. DA 28 043 AMC 00073(E), and in part by the National Science Foundation under Grant NSF GK-690, and in part by the Air Force Cambridge Research Laboratories under Contract AF 19(628)4379.

Reproduction in whole or in part is permitted for any purpose of the United States Government.

Distribution of this report is unlimited. Qualified requesters may obtain copies of this report from DDC.

WEIGHT DISTRIBUTION FORMULA FOR SOME CLASS OF CYCLIC CODES

Tadao Kasami

Abstract

P

Let $h_1(X)$ and $h_2(X)$ be different irreducible polynomials such that $h_1(\alpha^{-2^{h}-1}) = 0$ for some h(0 < h < m) and $h_2(\alpha^{-1}) = 0$, α being a primitive element of $GF(2^m)$. This paper presents the weight distribution formula of the code of length 2^m -1 generated by $(X^{2^m-1} - 1)/(h_1(X)h_2(X))$ for any m and h. Some applications to the cross-correlation problem between two different maximum length sequences are presented.

1. Introduction

W. W. Peterson [1] calculated a number of weight distributions for BCH codes of lengths 63 to 1023 and their dual codes by digital computation. He observed that some BCH codes with large t for a given m ($5 \le m \le 10$) have a very simple structure of weight distribution. The result presented here is a theoretical development of his observation.

1

Let C be a cyclic code of length $2^{m}-1$. The extended code of C is the code with an overall parity check added to C as the first digit. The first symbol in a code vector is numbered 0, and for i > 1 the i-th digit is numbered α^{i-2} , α being a primitive element of $GF(2^{m})$. Now for $a(\neq 0)$ and $beGF(2^{m})$ and for a code vector v of the extended code, permute the symbol in position X to position aX + b. Then, the resulting vector is denoted by $\pi_{ab}v$. W. W. Peterson [1] proved that the extended codes of BCH codes are invariant under doubly transitive group of permutations $\pi = {\pi_{ab} | a(\neq 0), beGF(2^{m})}$. This paper presents the weight distribution formula for a class of cyclic codes of length 2^{m} -1 whose extended codes are invariant under π .

Let $g_1(X)$ and $g_2(X)$ be different irreducible polynomials such that

(1) $g_1(\alpha^{2^{h}+1}) = 0$ for some h (0 < h < m), (2) $g_2(\alpha) = 0$

The degree of $g_1(X)$ is a factor m' of m and the degree of $g_2(X)$ is m. Let

$$h_1(X) = X^{m'}g_1(X^{-1}), h_2(X) = X^{m}g_2(X^{-1})$$

Let C_o , C and C' denote binary cyclic codes with length 2^m-1 generated by $g_1(X)g_2(X), (X^{2^m-1}-1)/(h_1(X)h_2(X))$ and $(X^{2^m-1}-1)/[(X-1)h_1(X)h_2(X)]$ respectively. Then C is the dual code of C_o and a subcode of C'. If h = 1, then C_o is a double error correcting BCH code, and if m is odd and h = (m-1)/2, then C is a BCH code with the second largest t for given m.

In what follows, the weight distribution formula of code C for any m and h will be derived. This problem is closely related to the crosscorrelation problem between two different maximum length sequences.^{*} Some applications to the problem will be presented in section 6.

2. Preliminary Lemmas

<u>Lemma 1</u>: The extended code of C' or C is invariant under π .

This lemma follows from the definition of C' or C and a general theorem [2]. Let

$$m = m'm''.$$
(1)
Since $\alpha^{2^{m-h}+1}$ is a root of $g_1(X)$, it can be assumed that

$$2h \leq m$$
. (2)

Since $(2^{m'}-1)(2^{h}+1)$ is divisible by $(2^{m}-1) = (2^{m'}-1)(2^{m'}(m''-1) + 2^{m'}(m''-2) + \dots + 1)$,

$$h \ge m'(m''-1)$$
.

From (1) and (2),

$$m'm'' \ge 2h \ge 2m'(m''-1)$$
 (3)

Hence,

$$m'' = 1 \text{ or } 2.$$

If m'' = 2, then it follows from (3) that

$$m' = h$$
.

That is, there are only two cases:

 $m^{\dagger} = m$

and

$$m' = m/2 = h.$$

* Dr. B. Elspas pointed out this relation.

The following well-known lemmas will be used later.

<u>Lemma 2</u>: Let u(l) denote the smallest positive integer u such that $2^{u} \equiv 1 \pmod{l}$. Then, $2^{u'} \equiv 1 \pmod{l}$ if and only if $u' \equiv 0 \pmod{u(l)}$. Let (l, l') denote the greatest common divisor of l and l'.

<u>Corollary 3</u>: Let $u = (u_1, u_2)$. Then, $2^{u}-1 = (2^{u}-1, 2^{u}-1)$.

Lemma 4: $2^{u}+1$ (or $2^{u}-1$) is divisible by $2^{u'}+1$, if and only if u is divisible by u' and u/u' is odd (or even).

Let $c^{*} = (m,h)$, c = (m,2h) and $v = (2^{m}-1,2^{h}+1)$. By Corollary 3,

$$2^{c^{*}}-1 = (2^{m}-1, 2^{h}-1), \qquad (4)$$

$$2^{c}-1 = (2^{m}-1, 2^{2h}-1). \qquad (5)$$

Since $(2^{h}+1, 2^{h}-1) = 1$,

$$(2^{m}-1, 2^{2h}-1) = (2^{m}-1, 2^{h}+1) (2^{m}-1, 2^{h}-1).$$

Thus,

 $(2^{c}-1)/v = 2^{c'}-1.$

By definition, c = c' or 2c'. Therefore, we have:

Lemma 5: If c = c', then v = 1. Otherwise,

$$v = 2^{c'} + 1 = 2^{c/2} + 1. \tag{6}$$

The next lemma is due to Pless [3].

Lemma 6: Let a_j and b_j denote the number of code vectors of weight j in a code A and the number of code vectors of weight j in the dual code of A respectively. If $b_1 = b_2 = 0$, then the following power moment identities hold:

$$\Sigma a_{j} = 2^{k}$$

$$\Sigma j a_{j} = 2^{k-1}n$$

$$\Sigma j^{2}a_{j} = 2^{k-2}n(n+1)$$

$$\Sigma j^{3}a_{j} = 2^{k-3}(n^{3}+3n^{2}) - 3!2^{k-3}b_{3}$$

$$\Sigma j^{4}a_{j} = 2^{k-4}(n^{4}+6n^{3}+3n^{2}-2n) - 4!2^{k-4}nb_{3} + 4!2^{k-4}b_{4},$$

where k denotes the number of information digits.

Let C_1 and C_2 denote binary cyclic codes with length 2^m-1 generated by $(x^{2^m-1}-1)/h_1(x)$ and $(x^{2^m-1}-1)/h_2(x)$ respectively. Codes C_1 and C_2 are subcodes of C and C'. If the degree of $g_1(x)$ is m, then the roots of $h_1(x) = x^m g_1(x^{-1})$:are:

$$\alpha^{-(2^{h}+1)} = \alpha^{2^{m}-2^{h}-2}, \ \alpha^{-2(2^{h}+1)} = \alpha^{2^{m}-2^{h+1}-2-1}, \ \dots,$$
$$\alpha^{-2^{m-h-1}(2^{h}+1)} = \alpha^{2^{m-1}-2^{m-h-1}-1}, \ \alpha^{-2^{(m-h)}(2^{h}+1)} = \alpha^{2^{m}-2^{m-h}-2},$$
$$\dots, \ \alpha^{-2^{m-1}(2^{h}+1)} = \alpha^{2^{m-1}-2^{h-1}-1}.$$

There is no i $(0 \le i < 2^{m-1})$ with $h_1(\alpha^i) = 0$ except for

$$i_1 = 2^{m-1} - 2^{m-h-1} - 1$$

and

I

$$i_1' = 2^{m-1} - 2^{h-1} - 1.$$

By (2), $i_1 < i_1^{\circ}$.

If $m^{\prime} = m/2$, then the roots of $h_1(X)$ are:

$$\alpha^{-(2^{m'}+1)} = \alpha^{2^{m}-2^{m'}-2}, \ \alpha^{-2(2^{m'}+1)} = \alpha^{2^{m}-2^{m'}+1}-2^{-1}, \ \dots$$
$$\alpha^{-2^{m'}-1}(2^{m'}+1) = \alpha^{2^{m-1}-2^{m'}-1}.$$

there is no i $(0 \le i < 2^{m-1})$ with $h_1(\alpha^i) = 0$ except for

$$i_1 = 2^{m-1} - 2^{m'-1} - 1.$$

The roots of $h_2(X)$ are $\alpha^{-1} = \alpha^{2^m-2}, \alpha^{-2} = \alpha^{2^m-3}, \dots, \alpha^{-2^{m-1}} = \alpha^{2^{m-1}-1}$. As it is done above, here we let

$$i_2 = 2^{m-1} - 1$$
.

Let X_1, X_2, \ldots, X_w be the location numbers of code vector v(x) of C'. Then,

$$v(\alpha^{i}) = \sum_{f=1}^{W} X_{f}^{i}$$
, $0 \leq i < 2^{m}-1$

For any $\beta_0 \in GF(2)$, any $\beta_1 \in GF(2^m')$ and any $\beta_2 \in GF(2^m)$, there exists a unique code vector v(x) of C' such that $v(1) = \beta_0$, $v(\alpha^{i_1}) = \beta_1$ and $v(\alpha^{i_2}) = \beta_2$ (Mattson, Solomon [5]). Let $v(\beta_0, \beta_1, \beta_2; x)$ denote the code vector specified by β_0, β_1 and β_2 . By definition,

$$\mathbf{x}^{\ell}\mathbf{v}(\boldsymbol{\beta}_{o},\boldsymbol{\beta}_{1},\boldsymbol{\beta}_{2};\mathbf{x}) = \mathbf{v}(\boldsymbol{\beta}_{o},\boldsymbol{\alpha}^{\ell \mathbf{i}}\boldsymbol{\beta}_{1},\boldsymbol{\alpha}^{\ell \mathbf{i}}\boldsymbol{\beta}_{2};\mathbf{x})$$

If and only if $\beta_0 = 0$, $v(\beta_0, \beta_1, \beta_2; x) \in C$. If and only if $\beta_0 = \beta_1 = 0$ (or $\beta_0 = \beta_2 = 0$), then $v(\beta_0, \beta_1, \beta_2; x) \in C_2$ (or C_1). The cyclic permutations on code word symbols induce a permutation group on the code vectors of C', which divides $C-C_2$ into disjoint sets of transitivity. Since $v = (i_1, 2^m - 1)$, each set consists of $(2^m - 1)/v$ code vectors. In case of m' = m, let $v(0, \alpha^i, \beta_2; x)$ ($0 \le i < v, \beta_2 \in GF(2^m)$) represent each set. In case of m' = m/2, let $v(0, 1, \beta_2; x)$ represent each set.

^{*}A polynomial representation will be used for a code vector [4].

Now, consider the extended code C_{ex} of code C'. Let $\overline{v}(\beta_0, \beta_1, \beta_2)$ denote the vector with an overall parity check added to $v(\beta_0, \beta_1, \beta_2; x)$ as the first digit. By definition $C_{ex} = \{\overline{v}(\beta_0, \beta_1, \beta_2) | v(\beta_0, \beta_1, \beta_2; x) \in C'\}$. Let X_1, X_2, \dots, X_w be the location numbers of $\overline{v}(\beta_0, \beta_1, \beta_2)$ and let

$$S_{i} = \sum_{f=1}^{W} X_{f}^{i}$$
, $0 \le i < 2^{m} - 1$ (7)

Then, by definition

$$S_{i_1} = \beta_1 , \qquad (8)$$

$$i_2 = \beta_2 , \qquad (9)$$

$$S_{i} = 0$$
 ($i \neq i_{1}2^{\ell}, i_{2}2^{\ell} \pmod{2^{m}-1}, 0 \le \ell < m$)

Therefore,

$$S_i = 0$$
 ($i \neq i_1, i'_1, 1 \le i < 2^{m-1} - 1$) (10)

By Lemma 1, $\pi_{1b} \overline{v}(\beta_0, \beta_1, \beta_2) \in \mathbb{C}_{ex}$ for any $b \in GF(2^m)$. Let

$$\pi_{1b} \overline{v}(\beta_{0}, \beta_{1}, \beta_{2}) = \overline{v}(\beta_{0}', \beta_{1}', \beta_{2}') .$$
(11)

The weights of $\overline{v}(\beta_0,\beta_1,\beta_2)$ and $\overline{v}(\beta_0',\beta_1',\beta_2')$ are the same. By the definition of π_{1b} ,

$$B_{1}^{*} = \sum_{f=1}^{W} (X_{f}+b)^{i_{1}}$$
$$B_{2}^{*} = \sum_{f=1}^{W} (X_{f}+b)^{i_{2}} .$$

Hence

$$\beta_{1}^{v} = \sum_{f=1}^{w} \sum_{i=0}^{i_{1}} (i_{i}^{i_{1}}) X_{f}^{i_{b}} \sum_{i=0}^{i_{1}-i} (i_{i}^{i_{1}}) X_{i_{b}}^{i_{1}-i_{1}} = \sum_{i=0}^{i_{1}} (i_{i}^{i_{1}}) X_{i_{b}}^{i_{1}-i_{1}}$$

7

From (8) and (10),

I

$$\beta'_1 = \beta_1 \tag{12}$$

Note that $(X_{f}+b)^{i_{2}} = (X_{f}+b)^{2^{m-1}-1} = (X_{f}^{2^{m-1}}+b^{2^{m-1}})/(X_{f}+b) = X_{f}^{2^{m-1}-1} + X_{f}^{2^{m-1}-1}b + \dots + b^{2^{m-1}-1}$. Then, $\beta_{2}^{v} = \sum_{f=1}^{w} \sum_{i=0}^{2^{m-1}-1} X_{f}^{i_{b}}b^{2^{m-1}-1-i} = \sum_{i=0}^{2^{m-1}-1} S_{i_{i}}b^{2^{m-1}-1-i}$ (13)

Consider the case of m' = m. Since $i'_1 \equiv 2^h i_1 \pmod{2^m - 1}$, $S_{i'_1} = S_{i_1}^{2^h}$. From (8),(9), (10) and (13),

$$B_{2}^{i} = \beta_{1}b^{2^{m-1}-1-i}1 + \beta_{1}^{2^{h}}b^{2^{m-1}-1-i}1 + \beta_{2} = \beta_{1}b^{2^{m-h-1}} + \beta_{1}^{2^{h}}b^{2^{h-1}} + \beta_{2} .$$
(14)

For the case of $m^{\circ} = m/2$, it follows from (8), (9), (10) and (13) that

$$\beta_2' = \beta_1 b^{2^{h-1}} + \beta_2 . (15)$$

Hereafter we shall consider the case of m' = m except for section 5. For each i $(0 \le i < v)$, $V_i = \{\alpha^{i}b^{2^{m-h-1}} + \alpha^{i2^{h}b^{2^{h-1}}} | beGF(2^{m})\}$ forms a subspace of $GF(2^{m})$. Let

$$F_{i}(X) = \alpha^{i} X^{2^{m-h-1}} + \alpha^{i2^{h}} X^{2^{h-1}}$$
$$= \alpha^{i} X^{2^{h-1}} (X^{2^{h-1}} (2^{m-2h} - 1)) + \alpha^{i} (2^{h-1})$$

If i = 0, the order of a nonzero root in $GF(2^m)$ of $F_o(X)$ is a factor of $2^{m-2h}-1$. Since c = (m,2h) = (m,m-2h), $2^c-1 = (2^m-1,2^{m-2h}-1)$. This implies

that the roots in $GF(2^m)$ of $F_0(X)$ are in subfield $GF(2^c)$. Conversely, any element in this subfield is a root of $F_0(X)$. Hence, the dimension of V_0 is m-c. Let V_{00} (= V_0), V_{01}, V_{02}, \dots, V be the cosets of $GF(2^m)$ with respect $02^c - 1$ to V_0 . Each coset has 2^{m-c} elements.

For $i \neq 0$, assume that α^{j} is a root of $F_{i}(X)$. Then,

$$2^{h-1}(2^{m-2h}-1) j \equiv i(2^{h}-1) \pmod{2^{m}-1}$$

$$(1-2^{2h}) j \equiv i2^{h+1}(2^{h}-1) \pmod{2^{m}-1}$$

$$-(2^{h}+1) j \equiv i2^{h+1} \pmod{2^{m}-1}$$

Since ν divides both $2^{h}+1$ and $2^{m}-1$, ν must divide i. However, $0 < i < \nu$. Therefore, there is no root in $GF(2^{m})$ of $F_{i}(X)$ except for zero. Consequently, $\nu_{i} = GF(2^{m})$.

Let
$$B_{0j} = \{ (\alpha^{\ell i}, \alpha^{\ell i} \beta) | 0 \le \ell < 2^{m} - 1, \beta \in V_{0j} \} (0 \le j \le 2^{c} - 1) \text{ and}$$

 $B_{i} = \{ (\alpha^{i+\ell i}, \beta) | 0 \le \ell < 2^{m} - 1, \beta \in GF(2^{m}) \} (0 \le i \le \nu).$ Then,

$$|B_{0j}| = (2^{m}-1)2^{m-c}/\nu$$
 $(0 \le j \le 2^{c})$, (16)

$$|B_i| = (2^m - 1) 2^m / \nu$$
 $(0 < i < \nu)^*$. (17)

It follows from the definition of B_{0j} or B_i that for any (β_1,β_2) and $(\beta_1^{\circ},\beta_2^{\circ})$ in the same B_{0j} or B_i and for any $\beta_0 \in GF(2)$, there exists permutation π_{ab} such that $\pi_{ab}\overline{v}(\beta_0,\beta_1,\beta_2) = \overline{v}(\beta_0^{\circ},\beta_1^{\circ},\beta_2^{\circ})$ and $\beta_0^{\circ} \in GF(2)$. Therefore, $\overline{v}(\beta_0,\beta_1,\beta_2)$ and $\overline{v}(\beta_0^{\circ},\beta_1^{\circ},\beta_2^{\circ})$ have the same weight w. If β_0 (or β_0°) is zero,

8

[|]B| means the number of elements of B.

then $v(0,\beta_1,\beta_2;x)$ (or $v(0,\beta_1',\beta_2';x)$) has weight w. If β_0 (or β_0') is one, then $v(1,\beta_1,\beta_2;x)$ (or $v(1,\beta_1',\beta_2';x)$) has weight w-1 by definition. Since C° contains all one vector $e = (1,1,\ldots,1)$ and $e(\alpha^i) = \sum_{f=0}^{2^m-2} \alpha^{if} = [(\alpha^i)^{2^m-1}-1]/(\alpha^{i}-1) = 0$ ($0 < i < 2^m-1$),

$$v(1,\beta_1,\beta_2;x) = v(0,\beta_1,\beta_2;x) + e(x).$$

Hence, if β_0 (or $\beta_0^{"}$) is one, then $v(0,\beta_1,\beta_2;x)$ (or $v(0,\beta_1^{'},\beta_2^{'};x)$) has weight $2^m - w$. Therefore, we have Lemma 8.

<u>Lemma 8</u>: For each j (or i) $(0 \le j \le 2^c, 0 \le i \le v)$, there is w_{oj} (or w_i) such that for any $(\beta_1, \beta_2) \in B_{0j}$ (or B_i) the weight of $v(0, \beta_1, \beta_2; x)$ is either w_{0j} (or w_i) or $2^m \cdot w_{0j}$ (or $2^m \cdot w_i$).

3. <u>Case I</u>: (m,h) = (m,2h)

Hereafter a will denote the number of code vectors of weight w in C and b will denote the number of code vectors of weight w in C_o .

Lemma 9: For even w,

$$wa_{w} = (2^{m}-w)a_{2^{m}-w},$$

 $wb_{w} = (2^{m}-w)b_{2^{m}-w}.$

This lemma follows from a theorem due to Peterson [1] and Lemma 1. Consequently, if the values of w_{0j} 's $(0 \le j \le 2^{c})$ and w_{i} 's $(0 \le i \le v)$ are known, the weight distribution of C-C₂ is completely determined. Furthermore, any nonzero vector of C₂ has weight 2^{m-1} , because C₂ is a maximum length sequence code.

<u>Lemma 10</u>: $b_1 = b_2 = 0$, $b_3 = (2^{c'-1}-1)(2^m-1)/3$.

<u>Proof</u>: Since C_0 is a subcode of Hamming code,

 $b_1 = b_2 = 0.$ Assume that α^{j_1} , α^{j_2} and α^{j_3} are the location numbers of a code vector of weight 3 in C_o. Then,

$$\alpha^{j_{1}} + \alpha^{j_{2}} = \alpha^{j_{3}}$$
(18)
$$\alpha^{j_{1}(2^{h}+1)} + \alpha^{j_{2}(2^{h}+1)} = \alpha^{j_{3}(2^{h}+1)}$$
(19)

From (18),

$$\alpha^{j_{3}(2^{h}+1)} = (\alpha^{j_{1}}+\alpha^{j_{2}})^{2^{h}+1} = (\alpha^{j_{1}2^{h}}+\alpha^{j_{2}2^{h}})(\alpha^{j_{1}}+\alpha^{j_{2}})$$
$$= \alpha^{j_{1}(2^{h}+1)} + \alpha^{j_{1}2^{h}}\alpha^{j_{2}} + \alpha^{j_{1}}\alpha^{j_{2}2^{h}} + \alpha^{j_{2}(2^{h}+1)}$$

By combining with (19),

$$\alpha^{j_{1}2^{h}} \alpha^{j_{2}} + \alpha^{j_{1}} \alpha^{j_{2}2^{h}} = 0,$$

$$\alpha^{(j_{1}-j_{2})(2^{h}-1)} = 1.$$
 (20)

Thus,

$$(j_1 - j_2)(2^h - 1) \equiv 0 \pmod{2^m - 1}$$

If $c^* = (m,h) = 1$, then $(2^{h}-1, 2^{m}-1) = 1$. Therefore

 $j_1 = j_2$.

This is a contradiction, which leads to the conclusion that $b_3 = 0$. If c' $\frac{1}{3}$ 1, then $(2^m-1, 2^h-1) = 2^c-1$. Let

$$\mu = (2^{m} - 1) / (2^{c} - 1) . \tag{21}$$

Then,

$$j_1 \equiv j_2 \pmod{\mu}$$
.

Let $j_1 = \ell_1 \mu + i$ and $j_2 = \ell_2 \mu + i$ ($0 \le i < \mu$). Since $\alpha^{\ell_1 \mu} + \alpha^{\ell_2 \mu} = \alpha^{\ell_3 \mu}$ for some ℓ_3 , it follows from (18) that $j_3 = \ell_3 \mu + i$. Conversely, for any i ($0 \le i < \mu$) and for ℓ_1 , ℓ_2 , and ℓ_3 such that

$$\alpha^{\ell_{1}\mu} + \alpha^{\ell_{2}\mu} = \alpha^{\ell_{3}\mu}$$
(22)
$$0 \leq \ell_{1}, \ell_{2}, \ell_{3} < 2^{c'-1} ,$$
(23)

 ${}^{\ell_1\mu+i}_{\alpha}$, ${}^{\ell_2\mu+i}_{\alpha}$, and ${}^{\ell_3\mu+i}_{\alpha}$ satisfy (18) and (19). The number of unordered triplets $({}^{\ell_1},{}^{\ell_2},{}^{\ell_3})$'s satisfying (22) and (23) is equal to $({}^{2^{c'}-1}_{2})/3$. Consequently,

$$b_3 = \mu \left(\frac{2^{c'-1}}{2} \right)/3 = (2^m - 1) (2^{c'-1} - 1)/3.$$
 Q.E.D.

<u>Lemma 11</u>: Let I₂ and I₄ denote $\sum_{j\neq 0} (j-2^{m-1})^2 a_j$ and $\sum_{j\neq 0} (j-2^{m-1})^4 a_j$ respectively. Then,

$$I_{2} = 2^{2m-2}(2^{m}-1),$$

$$I_{4} = 2^{3m+c'-4}(2^{m}-1)$$

<u>Proof</u>: Note that k = 2m. By using the power moment identities of Lemma 6,

$$I_{2} = \sum j^{2}a_{j} - 2^{m} \sum ja_{j} + 2^{2m-2} \sum_{\substack{j \neq 0 \\ j \neq 0}} a_{j}$$

= $2^{2m-2}n(n+1) - 2^{3m-1}n + 2^{2m-2}(2^{2m}-1)$
= $(2^{m}-1)(2^{3m-2}-2^{3m-1}+2^{3m-2}+2^{2m-2})$
= $(2^{m}-1)2^{2m-2}$

$$I_{4} = \sum j^{4}a_{j} - 2^{m+1} \sum j^{3}a_{j} + 3 \cdot 2^{2m-1} \sum j^{2}a_{j} - 2^{3m-1} \sum ja_{j}$$

$$+ 2^{4m-4} \sum_{j\neq 0} a_{j}$$

$$= 2^{2m-4} (n^{4}+6n^{3}+3n^{2}-2n) - 2^{3m-2} (n^{3}+3n^{2})$$

$$+ 3 \cdot 2^{4m-3}n(n+1) - 2^{5m-2}n + 2^{4m-4} (2^{2m}-1)$$

$$+ 3(2^{3m-1}-2^{2m-1}n)b_{3} + 3 \cdot 2^{2m-2}b_{4}$$

$$= n\{2^{2m-4}[(n+1)^{3}+3(n+1)(n-1)] - 2^{3m-2}[(n+1)^{2} + n-1] + 3 \cdot 2^{4m-3}(n+1) - 2^{5m-2} + 2^{4m-4} (2^{m}+1)\}$$

$$+ 3 \cdot 2^{2m-1} (b_{3}+b_{4})$$

$$= n\{2^{5m-4}-2^{5m-2}+3 \cdot 2^{5m-3}-2^{5m-2}+2^{5m-4} + 3 \cdot 2^{3m-4} (2^{m}-2) - 2^{3m-2} (2^{m}-2) + 2^{4m-4}\}$$

$$+ 3 \cdot 2^{2m-1} (b_{3}+b_{4})$$

$$= (2^{m}-1)2^{3m-3} + 3 \cdot 2^{2m-1} (b_{3}+b_{4})$$
(24)

Since all one vector $(1, 1, \ldots, 1)$ is in C₀,

$$b_{2^{m}-4} = b_{3}$$
.

By Lemmas 9 and 10,

$$b_3 + b_4 = b_{2^m - 4} + b_4 = 2^{m - 2} b_{2^m - 4} = 2^{m - 2} b_3$$

= $2^{m - 2} (2^{c' - 1} - 1) (2^m - 1) / 3$. (25)

By substituting the right hand side of (25) into (24),

$$I_4 = (2^m - 1) 2^{3m + c^* - 4}$$
. Q.E.D.

Let $\boldsymbol{j}_{\boldsymbol{M}}$ be the smallest nonzero integer such that

 $a_{j_{M}} + a_{2^{m}-j_{M}} \neq 0$.

By the definition of I_2 and I_4 ,

$$(j_M^{-2^{m-1}})^2 \ge I_4/I_2 = 2^{m+c'-2}$$
 (26)

Consider the case where c = c'. Then, v = 1 by Lemma 5. Since all nonzero vectors in C_2 are of weight 2^{m-1} , it follows from Lemma 8 and (16) that $a_j + a_{2^m-j}$ ($j \neq 0, 2^{m-1}$) must be divisible by $2^{m-c'}(2^m-1) = 2^{m-c}(2^m-1)$. Therefore, from (26)

$$L_2 \ge (j_M - 2^{m-1})^2 (a_{j_M} + a_{2^m - j_M}) \ge 2^{2m-2} (2^m - 1)$$
 (27)

By Lemma 11 and (27),

$$I_{2} = (j_{M}^{-2^{m-1}})^{2} (a_{j_{M}^{+a}}^{+a}) = 2^{2m-2} (2^{m}^{-1}).$$

Consequently,

$$(j_{M}^{-2^{m-1}})^{2} = 2^{m+c'-2} = 2^{m+c-2}$$
 (28)

$$a_{j_{M}} + a_{2^{m}-j_{M}} = 2^{m-c}(2^{m}-1)$$
 (29)

$$a_{j} = 0 \ (j \neq 0, j_{M}, 2^{m-1}, 2^{m}-j_{M}).$$

Hence,

$$j_{M} = 2^{m-1} - 2^{(m+c)/2-1}$$

$$a_{2^{m-1}} = 2^{2^{m}-1} - (a_{j+a_{m-j}})$$

$$= (2^{m}-2^{m-c}+1)(2^{m}-1)$$

By Lemma 9, (29) and (30),

$$a_{j_{M}} = (2^{m-c-1}+2^{(m-c)/2-1})(2^{m}-1)$$
$$a_{2^{m}-j_{M}} = (2^{m-c-1}-2^{(m-c)/2-1})(2^{m}-1)$$

Thus, we have the following theorem.

<u>Theorem 1</u>: If (m,h) = (m,2h) = c, then $a_0 = 1$ $a_{2^{m-1}-2}(m+c)/2-1 = (2^{m-c-1}+2^{(m-c)}/2-1)(2^{m}-1)$ $a_{2^{m-1}} = (2^{m}-2^{m-c}+1)(2^{m}-1)$ $a_{2^{m-1}+2}(m+c)/2-1 = (2^{m-c-1}-2^{(m-c)}/2-1)(2^{m}-1)$ $a_{j} = 0$ for other j.

4. Case II: 2(m,h) = (m,2h) = m

Consider the case in which 2(m,h) = (m,2h). Then, $v = 2^{c'}+1$ by Lemma 5. Since $v(0,\beta_1,0;x)$ is in C_1 , $w_{00} = w_0, w_1, \dots, w_{v-1}$ can be found from the weight distribution of C_1 . Each code vector of C_1 is a vconcatenation of a code vector in cyclic code C'_1 of length $(2^m-1)/v$ which is generated by $(x^{(2^m-1)/v}-1)/h_1(x)$. Let $v'(\beta;x)$ denote a code vector v'(x)

14

(30)

in C_1' such that $v'(\alpha^{1}) = \beta$. With the same argument as the one of section 3, set $C_{1i}' = \{v(\alpha^{i+\ell\nu};x) \mid 0 \le \ell < (2^m-1)/\nu\} \ (0 \le i < \nu) \text{ consists of } (2^m-1)/\nu$ vectors of the same weight w_i' . Since $v(0,\beta,0;x) = (v'(\beta;x), \beta)$

 $v'(\beta;x),\ldots,v'(\beta;x))$, w_i can be given by the following equation:

$$w_{i} = v w_{i}^{\prime} \cdot (0 \le i < v)$$

Therefore, by applying Lemma 6 to code C'_1 , we have

$$(2^{m}-1) / \nu \sum_{i=0}^{\nu-1} w_i / \nu = 2^{m-1} (2^{m}-1) / \nu ,$$

$$(2^{m}-1) / \nu \sum_{i=0}^{\nu-1} (w_i / \nu)^2 = 2^{m-2} (2^{m}-1) (2^{m}-1+\nu) / \nu^2$$

Thus,

$$\sum_{i=0}^{\nu-1} w_i = \nu 2^{m-1}$$
(31)
$$\sum_{i=0}^{\nu-1} w_i^2 = \nu 2^{m-2} (2^m - 1 + \nu) = \nu 2^{m-2} (2^m + 2^{c'})$$
(32)

Therefore,

$$\sum_{i=0}^{j-1} (w_i - 2^{m-1})^2 = v 2^{m-2} (2^m - 1 + v) - 2^m v 2^{m-1} + 2^{2m-2} v$$
$$= v 2^{m-2} (v-1) = v 2^{m+c'-2}$$
(33)

On the other hand, it follows from Lemma 8, (16), (17) and the definition of $\rm I_2$ that

$$I_{2} = I_{20} + I_{21},$$

$$I_{20} = 2^{m-c} (2^{m}-1) \sum_{j=1}^{2^{c}-1} (w_{0j}-2^{m-1})^{2}/\nu,$$

$$I_{21} = 2^{m-c} (2^{m}-1) (w_{0}-2^{m-1})^{2}/\nu + 2^{m} (2^{m}-1) \sum_{i=1}^{\nu-1} (w_{i}-2^{m-1})^{2}/\nu.$$
(34)

By Lemma 11 and (33),

$$I_{2} = 2^{2m-2}(2^{m}-1)$$

$$\geq I_{21} = 2^{2m+c'-2}(2^{m}-1) - (2^{m}-1)(2^{m}-2^{m-c})(w_{0}-2^{m-1})^{2}/v.$$
(35)

By a simple calculation,

$$(w_0 - 2^{m-1})^2 \ge 2^{2m-2} (2^{c'} - 1) \nu / (2^m - 2^{m-c})$$

= 2^{m+c-2} (by (6)).

Hence,

$$v_{o} = 2^{m-1} \pm (2^{(m+c)/2-1} + \delta), \quad \delta \ge 0$$
 (36)

Now, by (31) and (32)

$$I_{2}^{\prime} = \sum_{i=0}^{\nu-1} (w_{i} - (2^{m-1} + 2^{m/2-1}))^{2}$$

= $\nu 2^{m-2} (2^{m-1} + \nu) - 2(2^{m-1} + 2^{m/2-1}) \nu 2^{m-1} + (2^{m-1} + 2^{m/2-1})^{2} \nu$
= $\nu \{2^{2m-2} + 2^{m+c'-2} - 2^{2m-1} + 2^{3m/2-1} + 2^{2m-2} + 2^{3m/2-1} + 2^{m-2}\}$
= $2^{m-2} (2^{c'} + 1)^{2}$ (37)

On the other hand,

$$I_{2}' \geq (w_{0} - (2^{m-1} + 2^{m/2-1}))^{2} = [2^{m-1} + (2^{(m+c)/2-1} + \delta) - 2^{m-1} + 2^{m/2-1}]^{2}$$
$$= 2^{m-2} (2^{c'} + 1 + \delta 2^{1-m/2})^{2}$$
(38)

From (35), (36), and (37), we have that

$$\delta = 0$$

$$w_0 = 2^{m-1} \pm 2^{(m+c)/2-1}$$
(39)

$$w_i = 2^{m-1} \mp 2^{m/2-1}$$
 (0 < i < v) (40)

16

Since w_i ($0 \le i < v$) is divisible by v, the <u>+</u> sign is determined by Lemma 4. Thus, we have:

Theorem 2: Let a' denote the number of code vectors of weight j in C'_1 . If m/c is odd (or even), then

$$a_{0}^{\prime} = 1$$

$$a_{2}^{\prime}m^{-1}-2^{(m+c)/2-1} \quad (\text{or } a_{2}^{\prime}m^{-1}+2^{(m+c)/2-1}) = (2^{m}-1)/(2^{c/2}+1)$$

$$a_{2}^{\prime}m^{-1}+2^{m/2-1} \quad (\text{or } a_{2}^{\prime}m^{-1}-2^{m/2-1}) = 2^{c/2}(2^{m}-1)/(2^{c/2}+1).$$

$$a_{j}^{\prime} = 0 \quad \text{for other } j.$$

From (35) and (39) it follows that

$$I_{21} = 2^{2m+c'-2}(2^{m}-1) - (2^{m}-1)(2^{m}-2^{m-c})2^{m+c-2}/n$$

= $2^{2m-2}(2^{m}-1)(2^{c'}-(2^{c}-1)/n)$
= $2^{2m-2}(2^{m}-1)$
= I_{2} (by (35)).

By (34),

$$I_{20} = (2^{m}-1)2^{m-c} / \nu \sum_{j=1}^{2^{c}-1} (w_{0j}-2^{m-1})^{2} = 0$$

Hence,

$$w_{0j} = 2^{m-1} \quad (1 \le j < 2^{c}).$$
 (41)

By (16), (17), Lemma 8, (39), (40) and (41), we have:

$${}^{a}_{2^{m-1}-2^{m/2-1}} + {}^{a}_{2^{m-1}+2^{m/2-1}} = (2^{m}-1)2^{m+c/2}/(2^{c/2}+1)$$
$${}^{a}_{2^{m-1}} = (2^{m}-1)2^{m-c}(2^{c}-1)/(2^{c/2}+1) + 2^{m}-1$$

$$a_{2^{m-1}-2^{(m+c)/2-1}}^{a} + a_{2^{m-1}+2^{(m+c)/2-1}}^{a} = (2^{m}-1)2^{m-c}/(2^{c/2}+1)$$

Thus, the next theorem follows from Lemma 9.

<u>Theorem 3</u>: If 2(m,h) = (m,2h) = c and $c \neq m$, then

$$a_{0} = 1$$

$$a_{2}^{m-1} (m+c) / 2 - 1 = (2^{m} - 1) (2^{(m-c)} / 2 + 1) 2^{(m-c)} / 2 - 1 / (2^{c/2} + 1))$$

$$a_{2}^{m-1} - 2^{m/2} = (2^{m} - 1) (2^{m/2} + 1) 2^{(m+c)} / 2 - 1 / (2^{c/2} + 1))$$

$$a_{2}^{m-1} = (2^{m} - 1) ((2^{c/2} - 1) 2^{m-c} + 1))$$

$$a_{2}^{m-1} + 2^{m/2} = (2^{m} - 1) (2^{m/2} - 1) 2^{(m+c)} / 2 - 1 / (2^{c/2} + 1))$$

$$a_{2}^{m-1} + 2^{m/2} = (2^{m} - 1) (2^{(m-c)} / 2 - 1) 2^{(m-c)} / 2 - 1 / (2^{c/2} + 1))$$

$$a_{j} = 0 \quad \text{for other } j.$$

5. Case III: 2(m,h) = (m,2h) = m

Consider the case of m = 2h. For any $\beta_1 \neq 0$, β_2 in $GF(2^m)$, there exists $beGF(2^m)$ such that

$$\beta_1 b^{2^{h-1}} + \beta_2 = 0$$
,

because $(2^{h-1}, 2^{m}-1) = 1$. From (15) and a similar argument to the one for the case of m = m', it follows that there exists w such that the weight of any code vector in C-C₂ is either w or 2^{m} -w. Since C'₁, the cyclic code of length $2^{m/2}$ -1 generated by $(x^{2^{m/2}-1}-1)/g_1(x)$, is a maximum length sequence code, C₁ consists of one zero vector and $2^{m/2}$ -1 vectors of weight $2^{m/2-1}$ $(2^{m/2}+1)$. Therefore,

$$w = (2^{m/2} + 1) 2^{m/2 - 1}.$$

On the other hand, C_2 is a maximum length sequence code of length 2^m -1. Hence,

$$a_{2^{m-1}} = 2^{m} - 1$$
.

According to Lemma 9, we have:

Theorem 4: If m = 2h, then

$$a_{0} = 1$$

$$a_{2^{m-1}-2^{m/2-1}} = (2^{m/2}-1)(2^{m-1}+2^{m/2-1})$$

$$a_{2^{m-1}} = 2^{m}-1$$

$$a_{2^{m-1}+2^{m/2-1}} = (2^{m/2}-1)(2^{m-1}-2^{m/2-1})$$

$$a_{j} = 0 \quad \text{for other j.}$$

6. Crosscorrelation Functions of Two Maximum Length Sequences

It follows from Lemma 5 that $\alpha^{2^{h}+1}$ is a primitive element if and only if c = c'. Assume that c = c'. Let

$$v(0,1,0;x) = \sum_{f=0}^{2^{m}-2} v_{1f}x^{f}$$
$$v(0,0,1;x) = \sum_{f=0}^{2^{m}-2} v_{2f}x^{f}$$

Then, $v_1 = v_{10}, v_{11}, \dots, v_{12^{m-2}}$ and $v_2 = v_{20}, v_{21}, \dots, v_{22^{m-2}}$ are maximum length sequences of length 2^{m} -1. In v_1 and v_2 , replace 0 by -1. Let $u_1 = u_{10}, u_{11}, \dots, u_{12^{m-2}}$ and $u_2 = u_{20}, u_{21}, \dots, u_{22^{m-2}}$ be the resulting 22^{m-2} sequences of real numbers 1 and -1. Correlation function $\theta(j)$ of u_1 and u_2 is defined by

$$\Theta(j) = \sum_{\substack{f=0}}^{2^{m}-2} u_{1f}u_{2f-j},$$

where suffix f - j is to be taken mod $2^m - 1$. Note that

$$v(0,1,0;x) + x^{j}v(0,0,1;x) = v(0,1,\alpha^{ji_2};x)$$

If $v(0,1,\alpha^{ji_2};x)$ has weight w, then

$$\theta(j) = 2^m - 1 - 2w$$
 (42)

Let s_i denote the number of j's $(0 \le j < 2^m - 1)$ with $\theta(j) = i$. Then, s_i is the number of vectors $v(0,1,\beta;x)$ with weight $(2^m - 1 - i)/2$. From sections 2 and 3, we have theorem 5.

Theorem 5:

$$s_{-2}^{(m+c)/2} = 2^{m-c-1} - 2^{(m-c)/2-1} ,$$

$$s_{-1} = 2^{m} - 2^{m-c} ,$$

$$s_{2}^{(m+c)/2} = 2^{m-c-1} + 2^{(m-c)/2-1} ,$$

$$s_{i} = 0 \quad \text{for other i.}$$

For $0 \le j \le 2^m - 1$, let

$$\theta_{j} = 1 \text{ if } \theta(j) = -2^{(m+c)/2} - 1 \text{ or } 2^{(m+c)/2} - 1$$

 $\theta_{j} = 0 \text{ if } \theta(j) = -1.$

Sequence $\theta = \theta_0, \theta_1, \dots, \theta_{2^m-2}$ will be called the correlation sequence of u_1 and u_2 . We shall characterize the correlation sequence below. Recall that

$$V_0 = V_{00} = \{b^{2^{m-h-1}} + b^{2^{h-1}} | b \in GF(2^m) \}$$

Since $(2^{h-1}, 2^{m}-1) = 1$,

$$V_{00} = \{b^{2^{m-2n}} + b | b \in GF(2^{m})\}$$

Since (m-2h,m) = c, the Galois group of $GF(2^m)$ over $GF(2^c)$ is generated by the automorphism $X \to X^2$ (by Theorem 9, p. 127 of [6]). Therefore, from the trace theorem (p. 121 of [6]) it follows that

$$V_{00} = \{b | \sigma(b) = 0, b \in GF(2^m) \},\$$

where $\sigma(b)$ denotes the trace of b in GF(2^m) over GF(2^c), and

$$\sigma(b) = b + b^{2^{c}} + b^{2^{2^{c}}} + \dots + b^{2^{m-c}}$$

Hence, any element of each coset V_{0j} has the same trace $t_j \in GF(2^c)$, and if $j \neq j', t_j \neq t_{j'}$.

Note that $v^2(0,\beta_1,\beta_2;x) = v(0,\beta_1,\beta_2;x^2) = v(0,\beta_1^2,\beta_2^2;x) \in C$ and that $v(0,\beta_1,\beta_2;x)$ and $v(0,\beta_1,\beta_2;x^2)$ have the same weight. Consequently, if $t_{j'} = t_j^2$, then $w_{0j'} = w_{0j}$. From the proof of Theorem 1 it follows that there is only one j_0 such that

$$w_{0j_0} = 2^{m-1} - 2^{(m+c)/2-1}$$

$$w_{0j} = 2^{m-1}, j \neq j_0$$
.

Since $t_{j_0} = t_{j_0}^2$, $t_{j_0} = 0$ or 1. Since C_1 is a maximum length sequence code, w_{00} must be 2^{m-1} . Therefore, $t_{j_0} = 1$. This implies that the weight of $v(0,1,\beta;x)$ is not equal to 2^{m-1} if and only if $\beta + \beta^{2^c} + \beta^{2^{2^c}} + \dots + \beta^{2^{m-c}} = 1$. From (42), $\theta(j) \neq -1$, if and only if $\alpha^{j_1} + \alpha^{j_2} + \alpha^{j_1} + \alpha^{j_2} + \alpha^{j_1} = 1$. $\dots + \alpha^{j_1} 2^{2^{m-c}} = 1$. Since $\alpha^{2j_1} = \alpha^{2^m-2} = \alpha^{-1}$ and $\alpha^{-2^{m-1}} = \alpha^{j_2}$, we have

Theorem 6.

21

<u>Theorem 6</u>: $\theta_j = 1$ if and only if $\alpha^{-j} + \alpha^{-j2} + \alpha^{-j2} + \dots + \alpha^{-j2} = 1$.

Now consider the case of c = 1. By definition, $v(0,0,1;\alpha^{2}) = 1$. Therefore,

 $v(0,0,1;\alpha^{-2^{\ell}}) = 1$ $(0 \le \ell \le m)$.

On the other hand, for any $\alpha^{j} \neq \alpha^{-2^{\ell}}$ ($0 \leq \ell < m$),

 $v(0,0,1;\alpha^{j}) = 0$.

By the formula due to Reed and Solomon [7],

$$v_{2f} = \sum_{j=0}^{2^{m}-2} v(0,0,1;\alpha^{j})\alpha^{-jf}$$
$$= \alpha^{f} + \alpha^{2f} + \alpha^{2^{f}f} + \dots + \alpha^{2^{m-1}f}$$

Hence, by Theorem 6

$$\theta_{j} = v_{22^{m}-1-j}.$$

This implies the following corollary.

<u>Corollary 7</u>: If (m,h) = (m,2h) = 1, then the correlation sequence of the maximum length sequence generated by $h_1(X)$ and the one generated by $h_2(X)$ is the maximum length sequence generated by $g_2(X) = X^m h_2(X^{-1})$.

R. Gold and E. Kopitzka [8] observed that for some pairs of maximum length sequences the correlation sequences are also maximum length sequences and they listed all such pairs of sequences of length 8191 or less. Among 28 listed pairs, 25 cases are covered by Corollary 7.

Acknowledgment

The author is grateful to Professors W. W. Peterson, R. T. Chien and J. S. Lin for many valuable suggestions and to Professors M. E. Van Valkenburg and H. Ozaki for their support.

References

- Peterson, W. W., "On the Weight Structure and Symmetry of BCH Codes," Report of University of Hawaii, Contract No. AF19(628)-4379, No. 1; (July, 1965).
- 2. Kasami, T., "Some Invariant Properties of BCH Codes and RM Codes," Seminar Note of CSL, University of Illinois, (March, 1966). A full paper is being prepared jointly with W. W. Peterson and S. J. Lin.
- Pless, V., "Power Moment Identities on Weight Distributions in Error Correcting Codes," <u>Information and Control</u>, <u>6</u>, 147-152 (1963).
- 4. Peterson, W. W., Error Correcting Codes, Wiley, New York (1961).
- 5. Mattson, H. F. and Solomon, G., "A New Treatment of Bose-Chaudhuri Codes," J. Soc. Indust. Appl. Math., 9, No. 4, (December, 1961).
- Reed, I. S. and Solomon, G., "Decoding Procedure for a Polynomial Code," Group Report 47,24, Lincoln Laboratory, (1959).
- 7. Albert, A. A., <u>Fundamental Concepts of Higher Algebra</u>, The University of Chicago Press, (1956).
- 8. Gold, R. and Kopitzka, E., "Study of Correlation Properties of Binary Sequences Vol. $1 \sim 5$," Reports of Magnovon Research Laboratories, (August, 1965).

Distribution list as of March 1, 1965

1	Dr. Chalmers Sherwin Deputy Director (Research & Technology) DD&RE RM 3E1060 The Pentagon Harbierer D. c. 20201	1	Commanding Officer U. S. Army Security Agency Arlington Hall Arlington, Virginia 22212	1
1	Washington, D. C. 20301 Dr. Edward M. Reilley Asst. Director (Research) Ofc. of Defense Res & Eng Department of Defense	1	Commanding Officer U. S. Army Limited War Laboratory Aberdeen Proving Ground Aberdeen, Maryland 21005 Attn: Technical Director	1
1	Washington, D. C. 20301 Dr. James A. Ward Office of Deputy Director (Research and	1	Attn: rechnical Director Commanding Officer Human Engineering Laboratories Aberdeen Proving Ground, Maryland 21005	1
	Information Rm 3D1037) Department of Defense The Pentagon Washington, D. C. 20301	1	Director U. S. Army EngineerGeodesy. Intelligence	1
1	Director Advanced Research Projects Agency Department of Defense Washington, D. C. 20301	1.	U. S. Army Command and General	1
1	Mr. Charles Yost, Director for Materials Sciences Advanced Research Projects Agency		Staff College Fort Leavenworth, Kansas 66207 Attn: Secretary	1
20	Department of Defense Washington, D. C. 20301 Defense Documentation Center	1	Dr. H. Robl, Deputy Director U. S. Army Research Office (Durham) Box CM, Duke Station Durham, North Carolina 27706	1
	Cameron Station, Bidg. 5 Alexandria, Virginia 22314 Attn: TISIA	1	Commanding Officer U. S. Army Research Office (Durham) P. O. Box CM, Duke Station Durham, North Carolina 27706	1
1	Director National Security Agency Fort George G. Meade, Maryland 20755 Attn: Librarian C-332	1	Attm: CRD-AA-IP (Richard O. Ulsh) Commanding General U. S. Army Electronics Command	
1	Chief of Research and Development Headquarters, Department of the Army Washington, D. C. 20310	1	Fort Monmouth, New Jersey 07703 Attn: AMSEL-SC Director	1
1	Attn: Physical Sciences Division P & E Chief of Research and Development Headquarters, Department of the Army Washington, D. C. 20310		U. S. Army Electronics Laboratories Fort Mommouth, New Jersey 07703 Attn: Dr. S. Benedict Levin, Director Institute for Exploratory Research	1
	Attn: Mr. L. H. Geiger, Rm 34442	1	Director U. S. Army Electronics Laboratories	
1	Research Plans Office U. S. Army Research Office 3045 Columbia Pike Arlington, Virginia 22204		Fort Monmouth, New Jersey 07703 Attn: Mr. Robert O. Parker, Executive Secretary, JSTAC (AMSEL-RD-X)	1
1	Commanding General U. S. Army Materiel Command	1	Superintendent U. S. Military Academy West Point, New York 10996	
1	Attn: AMCRD-RS-PE-E Washington, D. C. 20315 Commanding General	1	The Walter Reed Institute of Research Walter Reed Army Medical Center Washington, D. C. 20012	1
	U. S. Army Strategic Communications Command Washington, D. C. 20315	1	Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703	2
1	Commanding Officer U. S. Army Materials Research Agency Watertown Arsenal Watertown, Massachusetts 02172	1	Attn: AMSEL-RD-DR Director	1
1	Commanding Officer U. S. Army Ballistics Research Lab. Aberdeen Proving Ground	1	U. S. Army Electronics Laboratories Attn: AMSEL-RD-X Fort Mommouth, New Jersey 07703 Director	
1	Aberdeen, Maryland 21005 Attn: V. W. Richards Commanding Officer		U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-XE	1
	U. S. Army Ballistics Research Lab. Aberdeen Proving Ground Aberdeen, Maryland 21005 Attn: Keats A. Pullen, Jr.	1	Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-XC	1
1	Commanding Officer U. S. Army Ballistics Research Lab. Aberdeen Proving Ground Aberdeen, Maryland 21005 Attn: George C. Francis, Computing Lab.	1	Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-XS	1
1	Commandant U. S. Army Air Defense School P. O. Box 9390	1	Director U. S. Army Electronics Laboratories Fort Mommouth, New Jersey 07703 Attm: AMSEL-RD-NR	1
1	Fort Bliss, Texas 79916 Attn: Missile Sciences Div., C&S Dept. Commanding General	1	Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703	1
	U, S, Army Missile Command Redstone Arsenal, Alabama 35809 Attn: Technical Library	1	Attn: AMSEL-RD-NE Director U. S. Army Electronics Laboratories	1
1	Commanding General Frankford Arsenal Philadelphia, Pa. 19137 Attn: SMUFA-1310 (Dr. Sidney Ross)	1	Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-NO Director	6
1	Commanding General Frankford Arsenal Philadelphia, Pa. 19137 Attn: SUUFA-1300	1	U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-NP	
1	U. S. Army Munitions Command Picatinney Arsenal Dover, New Jersey 07801		Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-SA	1
1	Attn: Technical Information Branch Commanding Officer Harry Diamond Laboratories Connecticut Ave & Van Ness St. N.W.	1	Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-SE	1
1	Connecticut Ave. & Van Ness St., N.W. Washington, D. C. 20438 Attn: Mr. Berthold Altman Commanding Officer	1	Director U. S. Army Electronics Laboratories Fort Mommouth, New Jersey 07703 Attm: AMSEL-RD-SR	1
	Harry Diamod Laboratories Attn: Library Connecticut Ave. & Van Ness St., N.W. Washington, D. C. 20438	1	Attn: AMSEL-KU-SK Director U, S. Atmy Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-SS	

Disector		
Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-FU#1	1	
Commanding Officer U. S. Army Electronics R&D Activity Fort Huachuca, Arizona 85163		
Commanding Officer U. S. Army Engineers R&D Laboratory Fort Belvoir, Virginia 22060 Attn: STINFO Branch	1	
Commanding Officer U. S. Army Electronics SwD Activity White Sands Missile Range New Mexico 88002	1	
Director Human Resources Research Office The George Washington University 300 N. Washington Street Alexandria, Virginia	1	
Commanding Officer U. S. Army Personnel Research Office Washington 25, D. C.	1	
Commanding Officer U. S. Army Medical Research Laboratory Fort Knox, Kentucky	1	
Commanding General U. S. Army Signal Center and School Attn: Chief, Office of Academic Operations Fort Monmouth, New Jersey 07703	1	
Dr. Richard H. Wilcox, Code 437 Department of the Navy Washington, D. C. 20360		
Chief, Bureau of Weapons Attn: Technical Library, DL1-3 Department of the Navy Washington, D. C. 20360	1	
Washington, D. C. 20360 Chief, Bureau of Ships Department of the Navy Washington, D. C. 20360 Attn: Code 680	1	
Chief, Bureau of Ships Department of the Navy Washington, D. C. 20360 Attn: Code 732	1	
Commander U. S. Naval Air Development Center Johnsville, Pennsylvania Attn: NADC Library	1	
Commanding Officer Naval Electronics Laboratory San Diego, California 92052 Attn: Code 2222(Library)	1	
Commanding Officer Naval Electronics Laboratory San Diego, California 92052 Attn: Code 2800, C. S. Manning	1	
Commanding Officer and Director (Code 142 Library) David W. Taylor Model Basin Washington, D. C. 20007	1	
Director		
Naval Research Laboratory Washington, D. C. 20390 Attn: Technical Information Office (Code 2000)	1	
Commanding Officer Office of Naval Research Branch Office 219 S. Dearborn Street Chicago, Illinois 60604	1	
Chief of Naval Operations Department of the Navy Washington, D. C. 20350 Attn: OP-07T		
Chief of Naval Operations Department of the Navy Washington, D. C. 20350 Attn: OP-03EC	1	

Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-PE

Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-PF

Director U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-PR

Director U. S. Army Electronics Laboratories Fort Mommouth, New Jersey 07703 Attn: AMSEL-RL-GF

U. S. Army Electronics Laboratories Fort Monmouth, New Jersey 07703 Attn: AMSEL-RD-ADT

Director

- Commanding Officer Office of Naval Research Branch Office 1000 Geary Street San Francisco, California 94109
 - Commanding Officer U. S. Naval Weapons Laboratory Asst. Director for Computation Dahlgren, Virginia 22448 Attn: G. H. Gleissner (Code K-4)
- Inspector of Naval Material Bureau of Ships Technical Representative 1902 West Minnehaha Avenue St. Paul 4, Minnesota
- 5 Lt. Col. E. T. Gaines, SREE Chief, Electronics Division Directorate of Engineering Sciences Air Force Office of Scientific Research Washington, D. C. 20333
 - Director of Science & Technology Deputy Chief of Staff (K & D) USAF Washington, D. C. Attn: AFRST-EL/GU

1

- 1 Director of Science & Technology Deputy Chief of Staff (R & D) USAF Washington, D. C. Attn: AFRST-SC
 - Karl M. Fuechsel Electronics Division Director of Engineering Sciences Air Force Office of Scientific Research Washington, D. C. 20333
 - Lt. Col. Edwin M. Myers Headquarters, USAF (AFRDR) Washington 25, D. C.
- Director, Air University Library Maxwell Air Force Base Alabama 36112 Attn: CR-4803a
 - Commander Research & Technology Division AFSC (Mr. Robert L. Feik) Office of the Scientific Director Bolling AFB 25, D. C.
- Commander Research & Technology Division Office of the Scientific Director Bolling AFB 25, D. C. Attn: RTHR
- Commander Air Force Cambridge Research Laboratories Attm: Research Library CRMML-R L. G. Hanscom Field Bedford, Massachusetts 01731
- Dr. Lloyd Hollingsworth AFCRL L. G. Hanscom Field Bedford, Massachusetts 01731
- Commander Air Force Cambridge Research Laboratories Attn: Data Sciences Lab (Lt. S. J. Kahne, CRB) L. G. Hanscom Field Bedford, Massachusetts 01731
- Commander Air Force Systems Command Office of the Chief Scientist (Mr. A. G. Wimer) Andrews AFB, Maryland 20331

Commander Air Force Missile Development Center Attn: MDSGO/Major Harold Wheeler, Jr. Holloman Air Force Base, New Mexico

Commander Research & Technology Division Attn: MAYT (Mr. Evans) Wright-Patterson Air Force Base Ohio 45433

- Directorate of Systems Dynamics Analysis Aeronautical Systems Division Wright-Patterson AFB, Ohio 45433
- Hqs. Aeronautical Systems Division AF Systems Command Attn: Navigation & Guidance Laboratory Wright-Patterson AFB, Ohio 45433
- Rome Air Development Center Attn: Documents Library, RAALD Griffiss Air Force Base Rome, New York 13442
- Commander Rome Air Development Center Attn: RAWI-Major W. Hultarris Griffiss Air Force Base Rome, New York 13442
- Lincoln Laboratory Massachusetts Institute of Technology P. O. Box 73 Lexington 73, Massachusetts Attn: Library A-082

Continued next page

Distribution list as of March 1, 1965 (Cont'd.)

1

1

2

3

1

1

1

1

1

1

1

1

1

1

1

- Lincoln Laboratory Massachusetts Institute of Technology F. O. Box 73 Lexington 73, Massachusetts Attn: Dr. Robert Kingston
- APGC (PGAPI) Eglin Air Force Base Florida
- 1 Mr. Alan Barnum Rome Air Development Center Griffiss Air Force Base Rome, New York 13442
- Director Research Laboratory of Electronics Massachusetts Institute of Technology Cambridge, Massachusetts 02139
- Polytechnic Institute of Brooklyn 55 Johnson Street Brooklyn, New York 11201 Attn: Mr. Jerome Fox Research Coordinator
- Director Columbia Radiation Laboratory Columbia University 538 West 120th Street New York, New York 10027
- Director Coordinated Science Laboratory University of Illinois Urbana, Illinois 61803
- Director Stanford Electronics Laboratories Stanford University Stanford, California
- Director Electronics Research Laboratory University of California Berkeley 4, California
- Professor A. A. Dougal, Director Laboratories for Electronics and Related Science Research University of Texas Austin, Texas 78712
- Professor J. K. Aggarwal Department of Electrical Engineering University of Texas Austin, Texas 78712

1

- Director of Engineering & Applied Physics 210 Pierce Hall Harvard University Cambridge, Massachusetts 02138
- Capt. Paul Johnson (USN Ret.) National Aeronautics & Space Agency 1520 H. Street, N. W. Washington 25, D. C.
- NASA Headquarters Office of Applications 400 Maryland Avenue, S.Q. Weshington 25, D. C. Attn: Code FC Mr. A. M. Greg Andrus
- National Bureau of Standards Research Information Center and Advisory Serv. on Info. Processing Data Processing Systems Division Washington 25, D. C.
- Dr. Wallace Sinaiko Institute for Defense Analyses Research & Eng. Support Div. 1666 Connecticut Avenue, N. W. Washington 9, D. C.
- Data Processing Systems Division National Bureau of Standards Conn, at Van Ness Room 239, Bidg, 10 Washington 25, D. C. Attn: A. K. Smilow
- 1 Exchange and Gift Division The Library of Congress Washington 25, D. C.
- Dr. Alan T. Waterman, Director National Science Foundation Washington 25, D. C.
- H. E. Cochran
 Oak Ridge National Laboratory
 P. O. Box X
 Oak Ridge, Tennessee
- U. S. Atomic Energy Commission Office of Technical Information Extension P. O. Box 62 Oak Ridge, Tennessee
- Mr. G. D. Watson Defense Research Member Canadian Joint Staff 2450 Massachusetts Avenue, N. W. Washington 8, D. C.

1

1

1

- Martin Company P. O. Box 5837 Orlando, Florida Attn: Engineering Library MP-30
- Laboratories for Applied Sciences University of Chicago 6220 South Drexel Chicago, Illinois 60637

- Librarian School of Electrical Engineering Purdue University Lafayette, Indiana
- Donald L. Epley Dept, of Electrical Engineering State University of Iowa Iowa City, Iowa
- Instrumentation Laboratory Massachusetts Institute of Technology 68 Albany Street Cambridge 39, Massachusetts Attn: Library WI-109
- Sylvania Electric Products, Inc. Electronics System Waltham Labs. Library 100 First Avenue Waltham 54, Massachusetts
- Hughes Aircraft Company Centinela and Teale Streets Calver City, California Attn: K. C. Rosenberg, Supervisor Company Technical Document Center
- Autonetics 9150 East Imperial Highway Downey, California Attn: Tech. Library, 3041-11
- Dr. Arnold T. Nordsieck General Motors Corporation Defense Research Laboratories 6767 Hollister Avenue Goleta, California
- University of California Lawrence Radiation Laboratory P. O. Box 808 Livermore, California
- Mr. Thomas L. Hartwick Aerospace Corporation P. O. Box 95085 Los Angeles 45, California
- Lt. Col. Willard Levin Aerospace Corporation P. O. Box 95085 Los Angeles 45, California
- Sylvania Electronic Systems-West Electronic Defense Laboratories P. O. Box 205 Mountain View, California Attn: Documents Center
- Varian Associates 611 Hansen Way Palo Alto, California 94303 Attn: Tech. Library
- Huston Denslow Library Supervisor Jet Propulsion Laboratory California Institute of Technology Pasadena, California
- Professor Nicholas George California Institute of Technology Electrical Engineering Department Pasadena, California
- Space Technology Labs., Inc. One Space Park Redondo Beach, California Attn: Acquistions Group STL Technical Library
- The Rand Corporation 1700 Main Street Santa Monica, California Attn: Library
- Miss F. Cloak Radio Corp. of America RCA Laboratories David Sarnoff Research Center Princeton, New Jersey
- Mr. A. A. Lundstrom Bell Telephone Laboratories Room 2E-127 Whippany Road Whippany, New Jersey
- Cornell Aeronautical Laboratory, Inc. 4455 Genesee Street Buffalo 21, New York Attn: J. P. Desmond, Librarian
 - Sperry Gyroscope Company Marine Division Library 155 Glenn Cove Road Carle Place, L. I., New York Attn: Miss Barbara Judd
 - Library Light Military Electronics Dept, General Electric Company Armament & Control Products Section Johnson City, New York
 - Dr. E. Howard Holt Director Plasma Research Laboratory Rennselaer Polytechnic Institute Troy, New York
 - Battele-DEFENDER Battelle Memorial Institute 505 King Avenue Columbus 1, Ohio

- Laboratory for Electroscience Research New York University University Heights Bronx 53, New York
- National Physical Laboratory Teddington, Middlesex England Attn: Dr. A. M. Uttley, Superintendent, Autonomics Division
- Dr. Lee Huff Behavioral Sciences Advanced Research Projects Agency The Pentagon (Room 3E175) Washington, D. C. 20301
- Dr. Glenn L. Bryan Head, Personnel and Training Branch Office of Naval Research Navy Department Washington, D. C. 20360
- l Instituto de Fisica Aplicado "L. Torres Quevedo" High Vacuum Laboratory Madrid, Spain Attn: Jose L. de Segovia
 - Stanford Research Institute Attn: G-037 External Reports (for J. Goldberg) Menlo Park, California 94025

REVISED U. S. ARMY DISTRIBUTION LIST

(As received at the Coordinated Science Laboratory 27 July 1965)

- 1 Dr. Chalmers Sherwin Deputy Director (Research & Technology) DD&RE Rm 3E1060 The Pentagon Washington, D. C. 20301
- Dr. Edward M. Reilley 1 Asst. Director (Research) Ofc. of Defense Res. & Eng. Department of Defense Washington, D. C. 20301
- Dr. James A. Ward Office of Deputy Director (Research 1 and Information Rm 3D1037) Department of Defense The Pentagon Washington, D. C. 20301
- 1 Director Advanced Research Projects Agency Department of Defense Washington, D. C. 20301
- Mr. E. I. Salkovitz, Director 1 for Materials Sciences Advanced Research Projects Agency Department of Defense Washington, D. C. 20301
- 1 Colonel Charles C. Mack Headquarters Defense Communications Agency (333) The Pentagon Washington, D. C. 20305
- Defense Documentation Center 20 Attn: TISIA Cameron Station, Building 5 Alexandria, Virginia 22314
- 1 Director National Security Agency Attn: Librarian C-332 Fort George G. Meade, Maryland 20755
- U. S. Army Research Office Attn: Physical Sciences Division 3045 Columbia Pike Arlington, Virginia 22204 1
- Chief of Research and Development Headquarters, Department of the Army Attn: Mr. L. H. Geiger, Rm 3D442 Washington, D. C. 20310 1
- Research Plans Office 1 U. S. Army Research Office 3045 Columbia Pike Arlington, Virginia 22204
- Commanding General U. S. Army Materiel Command Attn: AMCRD-RS-PE-E 1 Washington, D. C. 20315
- Commanding General U. S. Army Strategic Communications Command Washington, D. C. 20315 1
- 1 Commanding Officer U. S. Army Materials Research Agency Watertown Arsenal Watertown, Massachusetts 02172
- Commanding Officer 1 Attn: V. W. Richards Aberdeen Proving Ground Aberdeen, Maryland 21005
- 1 Commanding Officer Attn: Keats A. Pullen, Jr. Aberdeen Proving Ground Aberdeen, Maryland 21005
- Commanding Officer U. S. Army Ballistics Research Laboratory Attn: George C. Francis, Computing Lab. Aberdeen Proving Ground, Maryland 21005
- 1 Commandant U. S. Army Air Defense School Attn: Missile Sciences Division, C&S Dept. P. O. Box 9390 Port Pite Fort Bliss, Texas 79916
- 1 Commanding General U. S. Army Missile Command Attn: Technical Library Redstone Arsenal, Alabama 35809
- 1 Commanding General Frankford Arsenal Frankford Arsenal Attn: SMUFA-1310 (Dr. Sidney Ross) Philadelphia, Pennsylvania 19137

1 Commanding General Frankford Arsenal Attn: SMUFA-1300 Philadelphia, Pennsylvania 19137 1

1

1

1

1

1

1

1

Director

- 1 U. S. Army Munitions Command Attn: Technical Information Branch Picatinney Arsenal Dover, New Jersey 07801
- Commanding Officer Harry Diamond Laboratories Attn: Mr. Berthold Altman Connecticut Avenue and Van Ness St.,N.W. Washington, D. C. 20438 1
- Commanding Officer Harry Diamond Laboratories Attn: Library Connecticut Avenue and Van Ness St.,N.W. Washington, D. C. 20438

1 Commanding Officer U. S. Army Security Agency Arlington Hall Arlington, Virginia 22212

- Commanding Officer U. S. Army Limited War Laboratory Attn: Technical Director Aberdeen Proving Ground Aberdeen, Maryland 21005
- 1 Commanding Officer Human Engineering Laboratories Aberdeen Proving Ground, Maryland 21005
- 1 Director U. S. Army Engineer Geodesy, Intelligence & Mapping Research and Development Agency Fort Belvoir, Virginia 22060
- 1 Commandant U. S. Army Command and General Staff College Attn: Secretary Fort Leavenworth, Kansas 66207
- Dr. H. Robl, Deputy Chief Scientist U. S. Army Research Office (Durham) Box CM, Duke Station 1 Durham, North Carolina 27706
- Commanding Officer 1 U. S. Army Research Office (Durham) Attn: CRD-AA-IP (Richard O. Ulsh) Box CM, Duke Station Durham, North Carolina 27706
 - Superintendent U. S. Army Military Academy West Point, New York 10996

1

1

1

The Walter Reed Institute of Research 1 Walter Reed Army Medical Center Washington, D. C. 20012

Commanding Officer U. S. Army Electronics R&D Activity Fort Huachuca, Arizona 85163

- 1 Commanding Officer ATT STINFO Branch Fort Belvoir, Virginia 22060
- Commanding Officer 1 U. S. Army Electronics R&D Activity White Sands Missile Range, New Mexico 88002
- 1 Director Human Resources Research Office The George Washington University 300 N. Washington Street Alexandria, Virginia 22300
- Commanding Officer U. S. Army Personnel Research Office Washington, D. C. 1
- Commanding Officer U. S. Army Medical Research Laboratory Fort Knox, Kentucky 40120
- Commanding General U. S. Army Signal Center and School Fort Monmouth, New Jarsey 07703 Attm: Chief, Office of Academic Operations
- Dr. S. Benedict Levin, Director Institute for Exploratory Research U. S. Army Electronics Command Fort Monmouth, New Jersey 07703

Institute for Exploratory Research U. S. Army Electronics Command Attn: Mr. Robert O. Parker, Executive Secretary, JSTAC (AMSEL-XL-D) Fort Monmouth, New Jersey 07703 Commanding General U. S. Army Electronics Command Fort Monmouth, New Jersey 07703 Attn: AMSEL-SC RD-D RD-G RD-MAF-I RD-MAT RD-GF RD-MN (Marine Corps LnO) XL-D XL-E XL-C XL-S HL-D HL-L HL-J HL-P HL-O HL-R NL-D NL-A NL-P NL-R NL-S KL-D KL-E KL-S KL-T VL-D WL-D 1 Mr. Charles F. Yost Special Assistant to the Director of Research National Aeronautics & Space Admin. Washington, D. C. 20546 Director Research Laboratory of Electronics Massachusetts Institute of Technology Cambridge, Massachusetts 02139 Polytechnic Institute of Brooklyn 55 Johnson Street Brooklyn, New York 11201 Attn: Mr. Jerome Fox Research Coordinator Director Columbia Radiation Laboratory Columbia University 538 West 120th Street New York, New York 10027 Director Stanford Electronics Laboratories Stanford University Stanford, California 94301 Director Electronics Research Laboratory University of California Berkeley, California 94 94700 Director

- Electronic Sciences Laboratory University of Southern California Los Angeles, California 90007
- Professor A. A. Dougal, Director Laboratories for Electronics and Related Science Research 1 University of Texas Austin, Texas 78712
- Professor J. K. Aggarwal Department of Electrical Engineering University of Texas Austin, Texas 78712 1
- 1 Division of Engineering and Applied Physics 210 Pierce Hall Harvard University Cambridge, Massachusetts 02138

KEY WORDS		LIN			КВ	LIN	ĸ	
		ROLE	WT	ROLE	WT	ROLE	ł	
weight distribution					A ALLA	Sec. 1	l	
cyclic codes	1		a block				l	
cross-correlation	and the second	23.4		- and -			l	
maximum length sequences		2.2.2	10 915		1000	1.11.11.11.11	I	
	Mart Statistical		14-11-20		1.2		I	
	10 mm	1	A-2.2	1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1. 1	Ser. C.	1.1.1	I	
	a surger	Davi Til	100,00	the state	016.31	27.23.6	I	
		1.1				1500	I	
		1. (963)	apprenting of	R. P. Mai	alex?	1014.00	I	
	193 6 4	3/12/	1. 1. 1.		Ne's	1983	I	
	Sugar Charles	1.3.6.1		2.200	1000	1. 1. 1. 1.	I	
	and the second			al market	And	1. 200	I	
			121 1		I Talks	1	l	
INSTRUC	TIONS							
1. ORIGINATING ACTIVITY: Enter the name and address of the						y limitatio		
contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing	imposed b	y securi				than those ndard state		
the report.	ments suc							
2a. REPORT SECURITY CLASSIFICATION: Enter the overall secu- rity classification of the report. Indicate whether "Re-			om DDC."	ers may of	otain copi	es of this		
stricted Data" is included. Marking is to be in accordance with appropriate security regulations.						on of this		
				ot authori agencies m		copies of		
2b. GROUP: Automatic downgrading is specified in DoD Direc- tive 5200.10 and Armed Forces Industrial Manual. Enter the	t	(3) "U. S. Government agencies of this report directly from DD DDC users shall request three		DC. Other qualified				
group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as author-		De users	SHALL LC	quest ento	ugn	."		
ized.				encies may				
3. REPORT TITLE: Enter the complete report title in all capital letters. Titles in all cases should be unclassified.				ly from DL t through	C. Other	qualified		
If a meaningful title cannot be selected without classifica- tion, show title classification in all capitals in parenthe-								
sis immediately following the title.				of this re s shall re		ontrolled. ough		
4. DESCRIPTIVE NOTES: If appropriate, enter the type of re-	_			1. 1. 1.		."		
port, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is						ffice of Te ale to the		
covered.						e, if known		
5. AUTHOR(S): Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial.		EMENTARY	NOTES: U	se for add	litional es	xplanatory		
If military, s w rank and branch of service. The name of the principal author is an absolute minimum requirement.	notes.							
	12. SPONS	ORING MI	LITARY AC	TIVITY: En	ter the na	ame of the oring (pay)	1	
6. REPORT DATE: Enter the date of the report as day, month, year; or month, year. If more than one date appears on the				lopment. I				
report, use date of publication.						f and factu	u	
7a. TOTAL NUMBER OF PAGES: The total page count should fol- low normal pagination procedures, i.e., enter the number of	though it	may als	o appear	dicative of elsewhere	in the boo	dy of the		
pages containing information.	technical report. If additional space is required tinuation sheet shall be attached.						or	
7b. NUMBER OF REFERENCES: Enter the total number of refer- ences cited in the report.	It is	highly	desirable	that the	abstract	of classifi	ie	
	reports b	It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security clas						
8a. CONTRACT OR GRANT NUMBER: If appropriate, enter the applicable number of the contract or grant under which the	sification	n of the	informat			h, represent		
report was written.	as (TS),							
8b, 8c, & 8d. PROJECT NUMBER: Enter the appropriate military department identification, such as project number, subproj-						he abstract 225 words.		
ect number, system numbers, task number, etc.	14. KEY W	ORDS: Ke	y words a	re technic	ally mean	ingful term	ms	
9a. ORIGINATOR'S REPORT NUMBER(S): Enter the official report	short phr.	ases tha	t charact	erize a re	port and m	may be used words must	d	
number by which the document will be identified and con- trolled by the originating activity. This number must be	selected	so that	no securi	ty classif	ication is	s required.		
unique to this report.	name, mil	itary pr	oject cod	e name, ge	ographic	ion, trade location, r	ma	
9b. OTHER REPORT NUMBER(S): If the report has been assigned any other report numbers (either by the originator or by the						an indicati s, roles, a		
sponsor), also enter this number(s).	weights i							
	a day of the second		Kalmere	Antin Plan	abarta	man minimum		

Security Classification						
DOCUMENT (Security classification of title, body of abstract and ind	CONTROL DATA					
 ORIGINATING ACTIVITY (Corporate author) University of Illinois Coordinated Science Laborat Urbana, Illinois 61801 REPORT TILE 	2	Unclassification Unclassified				
4. DESCRIPTIVE NOTES (Type of report and inclusive dat		OF CYCLIC CODES				
5. AUTHOR(S) (Last name, first name, initial) Kasami, Tadao						
6. REPORT DATE. April, 1966	7a TOTAL NO. OF PAGES	7b. NO. OF REFS.				
Ba. CONTRACT OR GRANT NO.	24 9a. ORIGINATOR'S REPORT	NUMBER (S)				
DA 28 043 AMC 00073(E) b. project no. 20014501B31F	R-285					
• Also National Science Foundation Grant NSF GK-690 •.	9b.OTHER REPORT NO(S) (Any other numbers that may be assigned this					
10. AVAILABILITY/ LIMITATION NOTICES						
Distribution of this report	is unlimited.					
11. SUPPLEMENTARY NOTES	thru U. S. A	ARY ACTIVITY ces Electronics Program Army Electronics Command A, New Jersey 07703				
13. ABSTRACT	FC. Hormouth	, New Jersey 07705				
Let $h_1(X)$ and $h_2(X)$ be different $h_1(\alpha^{-2^h-1}) = 0$ for some $h(0 < h < element of GF(2^m). This paper proof the code of length 2^m-1 generates m and h. Some applications to the different maximum length sequence$	< m) and $h_2(\alpha^{-1})$ resents the weigh ated by $(x^{2^m-1} -$ he cross-correlat	= 0, α being a primitive at distribution formula 1)/(h ₁ (X)h ₂ (X)) for any tion problem between two				
DD FORM 1473		Security Classification				