

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/309282792>

New identities for ratios of Ramanujan's theta-function

Article in Advanced Studies in Contemporary Mathematics (Kyungshang) · January 2017

DOI: 10.23001/ascm2017.27.1.131

CITATIONS
2

READS
113

3 authors:



M. S. Mahadeva Naika
Bangalore University, Central College Campus
115 PUBLICATIONS 546 CITATIONS

[SEE PROFILE](#)



Chandankumar S.
M. S. Ramaiah University of Applied Sciences
30 PUBLICATIONS 128 CITATIONS

[SEE PROFILE](#)



K. Sushan Bairy
Vijaya College, Bangalore
27 PUBLICATIONS 148 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Continued Fractions [View project](#)



Andrews' singular overpartitions [View project](#)

NEW IDENTITIES FOR RATIOS OF RAMANUJAN'S THETA FUNCTION

M. S. MAHADEVA NAIKA, S. CHANDANKUMAR AND K. SUSHAN BAIRY

ABSTRACT. Ramanujan in his notebooks, has established several new modular equation which he denoted as P and Q . In this paper, we establish several new identities for ratios of Ramanujan's theta function involving $\varphi(q)$. We establish some new explicit evaluations for the ratios of Ramanujan's theta function. We also establish some new modular relations for a continued fraction of order twelve $H(q)$ with $H(q^n)$ for $n = 2, 4, 6, 8, 10, 12, 14$ and 16.

1. INTRODUCTION

In Chapter 16, of his second notebook [12], [3, pp.257-262], Ramanujan develops the theory of theta function and his theta-function is defined by

$$\begin{aligned} f(a, b) &:= \sum_{n=-\infty}^{\infty} a^{n(n+1)/2} b^{n(n-1)/2}, \quad |ab| < 1, \\ &= (-a; ab)_{\infty} (-b; ab)_{\infty} (ab; ab)_{\infty}, \end{aligned}$$

where $(a; q)_{\infty} := \prod_{n=1}^{\infty} (1 - aq^{n-1})$, $|q| < 1$.

Following Ramanujan, we define

$$\varphi(q) := f(q, q) = \sum_{n=-\infty}^{\infty} q^{n^2} = (-q; q^2)_{\infty}^2 (q^2; q^2)_{\infty}, \quad (1.1)$$

$$\psi(q) := f(q, q^3) = \sum_{n=0}^{\infty} q^{n(n+1)/2} = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}}, \quad (1.2)$$

$$f(-q) := f(-q, -q^2) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = (q; q)_{\infty}. \quad (1.3)$$

Now we define a modular equation in brief. The ordinary hypergeometric series ${}_2F_1(a, b; c; x)$ is defined by

$${}_2F_1(a, b; c; x) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} x^n, \quad |x| < 1,$$

2010 *Mathematics Subject Classification.* 11A55, 11B65, 33D15, Secondary 11F27.

Key words and phrases. Theta-function, modular equations.

Research supported by UGC under major research project F.No.34-140\ 2008(SR).

where

$$(a)_0 = 1, (a)_n = a(a+1)(a+2)\dots(a+n-1), \text{ for } n \geq 1.$$

Let

$$z := z(x) := {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)$$

and

$$q := q(x) := \exp\left(-\pi \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-x\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x\right)}\right),$$

where $0 < x < 1$.

Let n denote a fixed natural number, and assume that

$$n \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-\alpha\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; \alpha\right)} = \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; 1-\beta\right)}{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; \beta\right)}. \quad (1.4)$$

Then a modular equation of degree n in the classical theory is a relation between α and β induced by (1.4).

In [2], N. D. Baruah has established the relation between $P := \frac{\varphi(q)}{\varphi(q^3)}$ with $Q := \frac{\varphi(q^n)}{\varphi(q^{3n})}$, for $n = 5$ and 7. M. S. Mahadeva Naika, S. Chandankumar and K. Sushan Bairy [7], have established the relation connecting P and Q for $n = 9, 17$ and 19. Motivated by these works, we establish several new modular identities connecting P and Q for $n = 2, 4, 6, 8, 10, 12, 14$ and 16.

In Section 2, we collect some results which are useful to prove our main results. In Section 3, we establish several new modular equations of degree 3 for the ratios of Ramanujan's theta function. In Section 4, we establish some explicit evaluations for the ratios of Ramanujan's theta function. In section 5, we establish some new modular relations for a continued fraction of order twelve and its evaluations. In Section 6, we explicitly evaluate Ramanujan's remarkable product of theta functions.

2. PRELIMINARIES

In this section, we collect some identities which are useful in establishing our main results.

Lemma 2.1. [3, Entry 27(i), p. 43] *If $\alpha\beta = \pi$, then*

$$\sqrt{\alpha}\varphi(e^{-\alpha^2}) = \sqrt{\beta}\varphi(e^{-\beta^2}). \quad (2.1)$$

Lemma 2.2. [3, Entry 24(iv), p. 39] We have

$$f^3(-q^2) = \varphi(-q)\psi^2(q). \quad (2.2)$$

Lemma 2.3. [3, Entry 10(i), 11(ii), pp. 122–123] For $0 < \alpha < 1$,

$$\varphi(q) = \sqrt{z}, \quad (2.3)$$

$$\sqrt{2}q^{1/8}\psi(-q) = \sqrt{z}\{\alpha(1-\alpha)\}^{1/8}. \quad (2.4)$$

Lemma 2.4. [3, Entry 3 (xii), pp. 352–353] Let α, β and γ be of the first, third and ninth degrees respectively. Let m denote the multiplier connecting α, β and m' be the multiplier relating γ, δ , then

$$\left(\frac{\beta^2}{\alpha\gamma}\right)^{1/4} + \left(\frac{(1-\beta)^2}{(1-\alpha)(1-\gamma)}\right)^{1/4} - \left(\frac{\beta^2(1-\beta)^2}{\alpha\gamma(1-\alpha)(1-\gamma)}\right)^{1/4} = -3\frac{m}{m'}, \quad (2.5)$$

$$\left(\frac{\alpha\gamma}{\beta^2}\right)^{1/4} + \left(\frac{(1-\alpha)(1-\gamma)}{(1-\beta)^2}\right)^{1/4} - \left(\frac{\alpha\gamma(1-\alpha)(1-\gamma)}{\beta^2(1-\beta)^2}\right)^{1/4} = \frac{m'}{m}. \quad (2.6)$$

Lemma 2.5. [3, Entry 11(viii), (ix), p. 384] Let α, β, γ and δ be of the first, third, fifth and fifteenth degrees respectively. Let m denote the multiplier connecting α and β and m' be the multiplier relating γ and δ . Then

$$\left(\frac{\alpha\delta}{\beta\gamma}\right)^{1/8} + \left(\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}\right)^{1/8} - \left(\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}\right)^{1/8} = \sqrt{\frac{m'}{m}}, \quad (2.7)$$

$$\left(\frac{\beta\gamma}{\alpha\delta}\right)^{1/8} + \left(\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}\right)^{1/8} - \left(\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}\right)^{1/8} = -\sqrt{\frac{m}{m'}}. \quad (2.8)$$

Lemma 2.6. [3, Entry 13(i), (ii), p. 401] Let α, β, γ and δ be of the first, third, seventh and twenty first degrees respectively. Let m denote the multiplier connecting α and β and m' be the multiplier relating γ and δ . Then

$$\begin{aligned} & \left(\frac{\beta\gamma}{\alpha\delta}\right)^{1/4} + \left(\frac{(1-\beta)(1-\gamma)}{(1-\alpha)(1-\delta)}\right)^{1/4} - \left(\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}\right)^{1/4} \\ & + 4\left(\frac{\beta\gamma(1-\beta)(1-\gamma)}{\alpha\delta(1-\alpha)(1-\delta)}\right)^{1/6} = \frac{m}{m'}, \end{aligned} \quad (2.9)$$

$$\begin{aligned} & \left(\frac{\alpha\delta}{\beta\gamma}\right)^{1/4} + \left(\frac{(1-\alpha)(1-\delta)}{(1-\beta)(1-\gamma)}\right)^{1/4} - \left(\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}\right)^{1/4} \\ & + 4\left(\frac{\alpha\delta(1-\alpha)(1-\delta)}{\beta\gamma(1-\beta)(1-\gamma)}\right)^{1/6} = \frac{m'}{m}. \end{aligned} \quad (2.10)$$

Lemma 2.7. [4, Entry 51, p. 204] Let $X = \frac{f^2(-q)}{q^{1/6}f^2(-q^3)}$ and $Y = \frac{f^2(-q^2)}{q^{1/3}f^2(-q^6)}$, then

$$XY + \frac{9}{XY} = \left(\frac{Y}{X}\right)^3 + \left(\frac{X}{Y}\right)^3. \quad (2.11)$$

Lemma 2.8. [1, Theorem 5.1] If $M = \frac{\psi(-q)}{q^{1/4}\psi(-q^3)}$ and $N = \frac{\varphi(q)}{\varphi(q^3)}$, then

$$N^4 + M^4N^4 = 9 + M^4. \quad (2.12)$$

3. NEW MODULAR EQUATIONS OF DEGREE 3

In this section, we establish several new modular equations of degree 3 for the ratios of Ramanujan's theta function involving $\varphi(q)$.

Theorem 3.1. If $P := \frac{\varphi(q)\varphi(q^2)}{\varphi(q^3)\varphi(q^6)}$ and $Q := \frac{\varphi(q)\varphi(q^6)}{\varphi(q^2)\varphi(q^3)}$, then

$$P + \frac{3}{P} + Q - \frac{1}{Q} = 4. \quad (3.1)$$

Proof. Equations (2.11) can be rewritten as

$$(UV)^4 + 9(UV)^2 = U^6 + U^6, \quad (3.2)$$

where $U := \frac{f^2(-q)}{q^{1/6}f^2(-q^3)}$ and $V := \frac{f^2(-q^2)}{q^{1/3}f^2(-q^6)}$. Solving the above equation (3.2) for UV and squaring both sides, we find that

$$2U^6V^6 + 729 - 81M + 27U^6 + 27V^6 = MU^6 + MV^6, \quad (3.3)$$

where $M := \pm 81 + 4U^6 + 4V^6$.

$$\begin{aligned} & (X^2Y^2 + X^2 + 3 - 4XY - Y^2)(X^4Y^2 + X^4 - 3Y^2 + Y^4)(X^4Y^4 - X^4 + 9 - Y^4) \\ & (X^4Y^2 - X^4 - 3Y^2 - Y^4)(X^4 + 3X^2 - X^2Y^4 + Y^4)(X^4 - 3X^2 + X^2Y^4 + Y^4) \\ & (X^2Y^2 + X^2 + 3 + 4XY - Y^2)(X^2Y^2 - X^2 + 3 + 4XY + Y^2)(X^2Y^2 - X^2 + 3 \\ & - 4XY + Y^2)(X^4Y^4 + 2X^4Y^2 + X^4 - 6X^2 + 9 + 12X^2Y^2 - 6Y^2 + 2X^2Y^4 + Y^4) \\ & (X^4Y^4 - 2X^4Y^2 + X^4 + 6X^2 + 9 + 12X^2Y^2 + 6Y^2 - 2X^2Y^4 + Y^4) = 0, \end{aligned} \quad (3.4)$$

where $X := \frac{\varphi(q)}{\varphi(q^3)}$ and $Y := \frac{\varphi(q^2)}{\varphi(q^6)}$.

As $q \rightarrow 0$, the first factor vanishes faster than the other factors of the equation (3.4). Setting

$P := XY$ and $Q := \frac{X}{Y}$ in the first factor, we arrive at the equation (3.1). This completes the proof. \square

Theorem 3.2. If $P := \frac{\varphi(q)\varphi(q^4)}{\varphi(q^3)\varphi(q^{12})}$ and $Q := \frac{\varphi(q)\varphi(q^{12})}{\varphi(q^4)\varphi(q^3)}$, then

$$Q^2 + \frac{1}{Q^2} - 16 \left[Q + \frac{1}{Q} \right] + P^2 + \frac{3^2}{P^2} + 2 \left[Q - \frac{1}{Q} \right] \left[P + \frac{3}{P} \right] + 20 = 0. \quad (3.5)$$

Proof. Using the equation (3.1), we arrive at (3.5). \square

Theorem 3.3. If $P := \frac{\varphi(q)\varphi(q^6)}{\varphi(q^3)\varphi(q^{18})}$ and $Q := \frac{\varphi(q)\varphi(q^{18})}{\varphi(q^3)\varphi(q^6)}$, then

$$\begin{aligned} & 9Q^3 - \frac{1}{Q^3} - 36 \left[Q^2 + \frac{1}{Q^2} \right] + 9 \left[7Q + \frac{1}{Q} \right] - 48 + P^3 + \frac{3^3}{P^3} \\ & + \left[P^2 + \frac{3^2}{P^2} \right] \left\{ 12 - \left[5Q + \frac{3}{Q} \right] \right\} + \left[P + \frac{3}{P} \right] \left\{ 3 \left[Q^2 + \frac{1}{Q^2} \right] \right. \\ & \left. - 8 \left[3Q + \frac{5}{Q} \right] + 51 \right\} = 0. \end{aligned} \quad (3.6)$$

Proof. Employing the equations (2.3), (2.4), (2.5) and (2.6), we deduce that

$$d^2b^2 + a^2d^2 + 3c^2a^2 - b^2c^2 = 0, \quad (3.7)$$

where

$$c = \frac{\varphi(q)}{\varphi(q^3)}, \quad d = \frac{\varphi(q^3)}{\varphi(q^9)}, \quad a = \frac{\psi(-q)}{q^{1/4}\psi(-q^3)}, \quad b = \frac{\psi(-q^3)}{q^{3/4}\psi(-q^9)}.$$

Now, using the equations (2.12) and (3.1), we find that

$$\begin{aligned} & (X^4Y^4 - 2Y^4X^2 + Y^4 + 6Y^2 + 9 + 12Y^2X^2 + 6X^2 - 2Y^2X^4 + X^4) \\ & (27 + 63Y^2X^4 + 153Y^2X^2 + Y^6X^6 + 51X^4Y^4 - 45X^2 + 9X^6 - 27Y^2 - Y^6 \\ & + 9X^4 + 9Y^4 + 3Y^2X^6 - 3Y^6X^4 + 3Y^6X^2 + 9Y^4X^2 - 5X^6Y^4 + 36Y^5X \\ & + 120Y^3X - 108YX - 12X^5Y^5 + 40Y^5X^3 + 48Y^3X^3 + 72X^3Y + 24Y^3X^5 \\ & + 36X^5Y)(27 + 63Y^2X^4 + 153Y^2X^2 + Y^6X^6 + 51X^4Y^4 - 45X^2 + 9X^6 \\ & - 27Y^2 - Y^6 + 9X^4 + 9Y^4 + 3Y^2X^6 - 3Y^6X^4 + 3Y^6X^2 + 9Y^4X^2 - 5X^6Y^4 \\ & - 36Y^5X - 120Y^3X + 108YX + 12X^5Y^5 - 40Y^5X^3 - 48Y^3X^3 - 72X^3Y \\ & - 24Y^3X^5 - 36X^5Y) = 0, \end{aligned} \quad (3.8)$$

where $X := \frac{\varphi(q)}{\varphi(q^3)}$ and $Y := \frac{\varphi(q^6)}{\varphi(q^{18})}$.

As $q \rightarrow 0$, the last factor vanishes, whereas, the other factors does not vanish. Setting $P := XY$ and $Q := \frac{X}{Y}$ in the last factor, we arrive at the equation (3.6). This completes the proof. \square

Theorem 3.4. If $P := \frac{\varphi(q)\varphi(q^8)}{\varphi(q^3)\varphi(q^{24})}$ and $Q := \frac{\varphi(q)\varphi(q^{24})}{\varphi(q^3)\varphi(q^8)}$, then

$$\begin{aligned} Q^4 + \frac{1}{Q^4} + 80 \left[Q^3 - \frac{1}{Q^3} \right] + 320 \left[Q^2 + \frac{1}{Q^2} \right] - 80 \left[Q - \frac{1}{Q} \right] + 884 \\ + P^4 + \frac{3^4}{P^4} + 4 \left[P^3 + \frac{3^3}{P^3} \right] \left\{ 4 + \left[Q - \frac{1}{Q} \right] \right\} + 2 \left[P^2 + \frac{3^2}{P^2} \right] \\ \times \left\{ 16 - 8 \left[Q - \frac{1}{Q} \right] + 3 \left[Q^2 + \frac{1}{Q^2} \right] \right\} + 4 \left[P + \frac{3}{P} \right] \\ \times \left\{ -52 + 22 \left[Q - \frac{1}{Q} \right] - 52 \left[Q^2 + \frac{1}{Q^2} \right] + \left[Q^3 - \frac{1}{Q^3} \right] \right\} = 0. \end{aligned} \quad (3.9)$$

Proof. Using the equations (3.1) and (3.5), we obtain (3.9). \square

Theorem 3.5. If $P := \frac{\varphi(q)\varphi(q^{10})}{\varphi(q^3)\varphi(q^{30})}$ and $Q := \frac{\varphi(q)\varphi(q^{30})}{\varphi(q^3)\varphi(q^{10})}$, then

$$\begin{aligned} Q^6 + \frac{1}{Q^6} + 160 \left[Q^5 + \frac{1}{Q^5} \right] + 2468 \left[Q^4 + \frac{1}{Q^4} \right] + 160 \left[Q^3 + \frac{1}{Q^3} \right] + 14864 \\ + 1745 \left[Q^2 + \frac{1}{Q^2} \right] + 8320 \left[Q + \frac{1}{Q} \right] + P^6 + \frac{3^6}{P^6} + \left[P + \frac{3}{P} \right] \left(6 \left[Q^5 - \frac{1}{Q^5} \right] \right. \\ \left. + 640 \left\{ \left[Q^2 - \frac{1}{Q^2} \right] - \left[Q^4 - \frac{1}{Q^4} \right] \right\} + 470 \left[Q^3 - \frac{1}{Q^3} \right] - 580 \left[Q - \frac{1}{Q} \right] \right) \\ + \left[P^2 + \frac{3^2}{P^2} \right] \left(15 \left[Q^4 + \frac{1}{Q^4} \right] - 576 \left[Q^3 + \frac{1}{Q^3} \right] - 1960 \left[Q^2 + \frac{1}{Q^2} \right] \right. \\ \left. - 1600 \left[Q + \frac{1}{Q} \right] - 1415 \right) + \left[P^3 + \frac{3^3}{P^3} \right] \left(20 \left[Q^3 - \frac{1}{Q^3} \right] + 384 \left[Q^2 - \frac{1}{Q^2} \right] \right. \\ \left. + 350 \left[Q - \frac{1}{Q} \right] \right) + \left[P^4 + \frac{3^4}{P^4} \right] \left(15 \left[Q^2 + \frac{1}{Q^2} \right] + 160 \left[Q + \frac{1}{Q} \right] + 324 \right) \\ + 6 \left[P^5 + \frac{3^5}{P^5} \right] \left(Q - \frac{1}{Q} \right) = 0. \end{aligned} \quad (3.10)$$

Proof. Employing the equations (2.3), (2.4), (2.7) and (2.8), we deduce that

$$\frac{a}{b} = \frac{u-v}{u+v}, \quad (3.11)$$

where $u = \frac{\varphi(q)}{\varphi(q^3)}$, $v = \frac{\varphi(q^5)}{\varphi(q^{15})}$, $a = \frac{\psi(-q)}{q^{1/4}\psi(-q^3)}$, $b = \frac{\psi(-q^5)}{q^{5/4}\psi(-q^{15})}$.

Using the equations (2.12), (3.1) and (3.11), we find that

$$\begin{aligned}
& (Y^4 X^4 - 2Y^2 X^4 + X^4 + 6X^2 + 9 + 12Y^2 X^2 + 6Y^2 - 2Y^4 X^2 + Y^4) \\
& (729 + 1215X^4 + 1458X^2 + 540X^6 + 135X^8 + 18X^{10} + 135Y^8 - 1458Y^2 \\
& - 540Y^6 + 15Y^8 X^{12} + 350Y^8 X^{10} - 1415Y^8 X^8 + 580Y^8 X^6 + 1745Y^8 X^4 \\
& - 6Y^{12} X^2 - 1410Y^8 X^2 - 1920YX^9 - 5184YX^7 + 160YX^{11} + 10368YX^5 \\
& + 12960YX^3 - 640Y^3 X^{11} + 160Y^3 X^9 + 1920Y^3 X^7 - 14400Y^3 X^5 \\
& + 12960Y^3 X + 2468Y^2 X^{10} + 1410Y^2 X^8 - 17640Y^2 X^6 + 8320Y^5 X^7 \\
& + 640Y^5 X^9 + 384Y^7 X^{11} - 1600Y^7 X^9 + 8320Y^7 X^5 - 576Y^5 X^{11} \\
& - 14400Y^5 X^3 - 1960Y^6 X^{10} - 580Y^6 X^8 + 14864Y^6 X^6 + 1740Y^6 X^4 \\
& - 17640Y^6 X^2 - 1920Y^7 X^3 - 5184Y^7 X - 10368Y^5 X + 9450Y^2 X^4 \\
& + 26244Y^2 X^2 + 1215Y^4 + 15Y^4 X^{12} - 9450Y^4 X^2 - 12735Y^4 X^4 \\
& - 1740Y^4 X^6 + 470Y^4 X^{10} + 160Y^9 X^{11} - 1600Y^9 X^7 - 640Y^9 X^5 \\
& + 1920Y^9 X - 18Y^{10} + 160Y^{11} X^9 - 384Y^{11} X^7 - 576Y^{11} X^5 + 640Y^{11} X^3 \\
& + Y^{12} + 160Y^{11} X + 324Y^{10} X^{10} - 350Y^{10} X^8 - 1960Y^{10} X^6 - 470Y^{10} X^4 \\
& + 2468Y^{10} X^2 + 6Y^{10} X^{12} + Y^{12} X^{12} - 6Y^{12} X^{10} + 15Y^{12} X^8 - 20Y^{12} X^6 \\
& + 15Y^{12} X^4 + 160Y^9 X^3 + X^{12} + 1745Y^4 X^8 + 6Y^2 X^{12} + 20Y^6 X^{12}) = 0.
\end{aligned} \tag{3.12}$$

where $X := \frac{\varphi(q)}{\varphi(q^3)}$ and $Y := \frac{\varphi(q^{10})}{\varphi(q^{30})}$.

As $q \rightarrow 0$, the second factor vanishes, but the first factor does not vanish. Setting $P := XY$ and $Q := \frac{X}{Y}$ in the second factor, we arrive at the equation (3.12). This completes the proof. \square

Theorem 3.6. If $P := \frac{\varphi(q)\varphi(q^{12})}{\varphi(q^3)\varphi(q^{36})}$ and $Q := \frac{\varphi(q)\varphi(q^{36})}{\varphi(q^3)\varphi(q^{12})}$, then

$$\begin{aligned}
 & 81Q^6 + \frac{1}{Q^6} - \frac{288}{Q^5} + 252 \left[3Q^4 + \frac{35}{Q^4} \right] + 96 \left[9Q^3 + \frac{68}{Q^3} \right] + 52560 \\
 & + 9 \left[1993Q^2 - \frac{551}{Q^2} \right] - 1152 \left[6Q - \frac{49}{Q} \right] + P^6 + \frac{3^6}{P^6} - \left[P^5 + \frac{3^5}{P^5} \right] \\
 & \times \left\{ 48 + 2 \left[5Q + \frac{3}{Q} \right] \right\} + \left[P^4 + \frac{3^4}{P^4} \right] \left\{ 852 + 96 \left[2Q + \frac{3}{Q} \right] + \left[31Q^2 + \frac{15}{Q^2} \right] \right\} \\
 & + \left[P^3 + \frac{3^3}{P^3} \right] \left\{ 102 \left[13Q - \frac{5}{Q} \right] + 96 \left[Q^2 + \frac{5}{Q^2} \right] - 4 \left[3Q^3 + \frac{5}{Q^3} \right] - 2320 \right\} \\
 & + \left[P^2 + \frac{3^2}{P^2} \right] \left\{ 4761 - 192 \left[32Q + \frac{5}{Q} \right] + 8 \left[63Q^2 - \frac{881}{Q^2} \right] - 192 \left[3Q^3 + \frac{10}{Q^3} \right] \right. \\
 & \left. - 3 \left[27Q^4 - \frac{5}{Q^4} \right] \right\} + \left[P + \frac{3}{P} \right] \left\{ 36 \left[231Q - \frac{431}{Q} \right] - 96 \left[57Q^2 - \frac{131}{Q^2} \right] \right. \\
 & \left. + 30 \left[21Q^3 + \frac{67}{Q^3} \right] - 16 \left[27Q^4 + \frac{163}{Q^4} \right] + 6 \left[9Q^5 - \frac{1}{Q^5} \right] - 21696 \right\} = 0.
 \end{aligned} \tag{3.13}$$

Proof. Employing the equations (3.1) and (3.6), we arrive at the equation (3.13). \square

Theorem 3.7. If $P := \frac{\varphi(q)\varphi(q^{14})}{\varphi(q^3)\varphi(q^{42})}$ and $Q := \frac{\varphi(q)\varphi(q^{42})}{\varphi(q^3)\varphi(q^{14})}$, then

$$\begin{aligned}
 & Q^8 + \frac{1}{Q^8} - 504 \left[Q^7 - \frac{1}{Q^7} \right] + 33792 \left[Q^6 + \frac{1}{Q^6} \right] - 100296 \left[Q^5 - \frac{1}{Q^5} \right] \\
 & + 649668 + 118384 \left[Q^4 + \frac{1}{Q^4} \right] + 39816 \left[Q^3 - \frac{1}{Q^3} \right] + 338688 \left[Q^2 + \frac{1}{Q^2} \right] \\
 & - 664776 \left[Q - \frac{1}{Q} \right] + P^8 + \frac{3^8}{P^8} - 8 \left[P^7 + \frac{3^7}{P^7} \right] \left\{ 7 - \left[Q - \frac{1}{Q} \right] \right\} \\
 & + \left[P^6 + \frac{3^6}{P^6} \right] \left\{ 832 + 56 \left[Q - \frac{1}{Q} \right] + 28 \left[Q^2 + \frac{1}{Q^2} \right] \right\} + 56 \left[P^5 + \frac{3^5}{P^5} \right] \\
 & \times \left\{ 69 - 8 \left[Q - \frac{1}{Q} \right] + 43 \left[Q^2 + \frac{1}{Q^2} \right] + \left[Q^3 - \frac{1}{Q^3} \right] \right\} + \left[P^4 + \frac{3^4}{P^4} \right] \\
 & \times \left\{ 7728 - 728 \left[Q - \frac{1}{Q} \right] + 896 \left[Q^2 + \frac{1}{Q^2} \right] + 2328 \left[Q^3 - \frac{1}{Q^3} \right] + 70 \left[Q^4 + \frac{1}{Q^4} \right] \right\} \\
 & - 8 \left[P^3 + \frac{3^3}{P^3} \right] \left\{ 2128 \left[Q - \frac{1}{Q} \right] + 1897 \left[Q^2 + \frac{1}{Q^2} \right] - 2870 \left[Q^3 - \frac{1}{Q^3} \right] \right. \\
 & \left. + 1333 \left[Q^4 + \frac{1}{Q^4} \right] - 7 \left[Q^5 - \frac{1}{Q^5} \right] + 6335 \right\} + 28 \left[P^2 + \frac{3^2}{P^2} \right] \\
 & \times \left\{ -3344 - 2014 \left[Q - \frac{1}{Q} \right] - 2850 \left[Q^2 + \frac{1}{Q^2} \right] + 2714 \left[Q^3 - \frac{1}{Q^3} \right] \right. \\
 & \left. - 976 \left[Q^4 + \frac{1}{Q^4} \right] - 122 \left[Q^5 - \frac{1}{Q^5} \right] + \left[Q^6 + \frac{1}{Q^6} \right] \right\} + 8 \left[P + \frac{3}{P} \right] \\
 & \left\{ 8043 - 42126 \left[Q - \frac{1}{Q} \right] + 12999 \left[Q^2 + \frac{1}{Q^2} \right] + 13272 \left[Q^3 - \frac{1}{Q^3} \right] \right. \\
 & \left. - 4263 \left[Q^4 + \frac{1}{Q^4} \right] - 1792 \left[Q^5 - \frac{1}{Q^5} \right] + 847 \left[Q^6 + \frac{1}{Q^6} \right] + \left[Q^7 - \frac{1}{Q^7} \right] \right\} = 0. \tag{3.14}
 \end{aligned}$$

The proof of the identity (3.14) is similar to the proof of the identity (3.10), except that in place of results (2.7) and (2.8), results (2.9) and (2.10) are used.

Theorem 3.8. If $P := \frac{\varphi(q)\varphi(q^{16})}{\varphi(q^3)\varphi(q^{48})}$ and $Q := \frac{\varphi(q)\varphi(q^{48})}{\varphi(q^3)\varphi(q^{16})}$, then

$$\begin{aligned}
 & P^8 + \frac{3^8}{P^8} + Q^8 + \frac{1}{Q^8} - 832 \left[Q^7 + \frac{1}{Q^7} \right] + 99296 \left[Q^6 + \frac{1}{Q^6} \right] - 315456 \left[Q^5 + \frac{1}{Q^5} \right] \\
 & + 577008 \left[Q^4 + \frac{1}{Q^4} \right] - 734272 \left[Q^3 + \frac{1}{Q^3} \right] + 1906912 \left[Q^2 + \frac{1}{Q^2} \right] - 2587456 \\
 & \times \left[Q + \frac{1}{Q} \right] + 4126660 + 16 \left[P + \frac{3}{P} \right] \left\{ -28287 \left[Q - \frac{1}{Q} \right] - 14952 \left[Q^2 - \frac{1}{Q^2} \right] \right. \\
 & \left. - 100296 \left[Q^3 - \frac{1}{Q^3} \right] + 649668 \left[Q^4 - \frac{1}{Q^4} \right] + 338688 \left[Q^5 - \frac{1}{Q^5} \right] \right\} = 0. \tag{3.15}
 \end{aligned}$$

$$\begin{aligned}
& +15456 \left[Q^3 - \frac{1}{Q^3} \right] + 7632 \left[Q^4 - \frac{1}{Q^4} \right] - 4204 \left[Q^5 - \frac{1}{Q^5} \right] + 1160 \left[Q^6 - \frac{1}{Q^6} \right] \\
& + \left[\frac{Q^7}{2} - \frac{1}{2Q^7} \right] \Big\} + 4 \left[P^2 + \frac{3^2}{P^2} \right] \left\{ -28112 \left[Q + \frac{1}{Q} \right] - 33454 \left[Q^2 + \frac{1}{Q^2} \right] + 42056 \right. \\
& + 2128 \left[Q^3 + \frac{1}{Q^3} \right] - 9640 \left[Q^4 + \frac{1}{Q^4} \right] - 11312 \left[Q^5 + \frac{1}{Q^5} \right] + 7 \left[Q^6 + \frac{1}{Q^6} \right] \Big\} \\
& + 8 \left[P^3 + \frac{3^3}{P^3} \right] \left\{ -5768 \left[Q - \frac{1}{Q} \right] - 2512 \left[Q^2 - \frac{1}{Q^2} \right] + 6758 \left[Q^3 - \frac{1}{Q^3} \right] \right. \\
& + 1760 \left[Q^4 - \frac{1}{Q^4} \right] + 7 \left[Q^5 - \frac{1}{Q^5} \right] \Big\} + 2 \left[P^4 + \frac{3^4}{P^4} \right] \left\{ 3360 \left[Q + \frac{1}{Q} \right] + 9432 \right. \\
& - 7472 \left[Q^2 + \frac{1}{Q^2} \right] + 6048 \left[Q^3 + \frac{1}{Q^3} \right] + 35 \left[Q^4 + \frac{1}{Q^4} \right] \Big\} + 8 \left[P^5 + \frac{3^5}{P^5} \right] \\
& \times \left\{ -528 \left[Q - \frac{1}{Q} \right] - 240 \left[Q^2 - \frac{1}{Q^2} \right] + 7 \left[Q^3 - \frac{1}{Q^3} \right] \right\} + 4 \left[P^6 + \frac{3^6}{P^6} \right] \\
& \times \left\{ -208 \left[Q + \frac{1}{Q} \right] + 7 \left[Q^2 + \frac{1}{Q^2} \right] + 584 \right\} + 8 \left[P^7 + \frac{3^7}{P^7} \right] \left[Q - \frac{1}{Q} \right] = 0.
\end{aligned} \tag{3.15}$$

Proof. Using the equations (3.1) and (3.9), we obtain the equation (3.15). \square

4. EVALUATIONS FOR THE RATIOS OF RAMANUJAN'S THETA FUNCTION

In this section, by using the modular equations obtained in the previous section, we establish some explicit evaluations for the ratios of Ramanujan's theta function.

Theorem 4.1. *We have*

$$\frac{\varphi(e^{-2\pi/\sqrt{3}})}{\varphi(e^{-2\pi\sqrt{3}})} = \frac{3^{1/4}(\sqrt{3}+1)(\sqrt{2}-1)}{\sqrt{2}}, \tag{4.1}$$

$$\frac{\varphi(e^{-\pi/2\sqrt{3}})}{\varphi(e^{-\pi\sqrt{3}/2})} = \frac{3^{1/4}(\sqrt{3}-1)(\sqrt{2}+1)}{\sqrt{2}}, \tag{4.2}$$

$$\frac{\varphi^2(e^{-\pi\sqrt{2/3}})}{\varphi^2(e^{-\pi\sqrt{6}})} = 3\sqrt{2} - 2\sqrt{3} + 3 - \sqrt{6}, \tag{4.3}$$

$$\frac{\varphi^2(e^{-\pi/\sqrt{6}})}{\varphi^2(e^{-\pi\sqrt{3/2}})} = 3\sqrt{2} + 2\sqrt{3} - 3 - \sqrt{6}, \tag{4.4}$$

$$\frac{\varphi^2(e^{-\pi\sqrt{2}})}{\varphi^2(e^{-3\pi\sqrt{2}})} = 21 + 9\sqrt{6} - 15\sqrt{2} - 12\sqrt{3}, \tag{4.5}$$

$$\frac{\varphi^2(e^{-\pi/3\sqrt{2}})}{\varphi^2(e^{-\pi/\sqrt{2}})} = \sqrt{6} + \sqrt{2} - 1, \tag{4.6}$$

$$\frac{\varphi^2(e^{-2\pi\sqrt{2/3}})}{\varphi^2(e^{-2\pi\sqrt{6}})} = \sqrt{3} \left[\sqrt{596 + 344\sqrt{3} - 8A} - \sqrt{595 + 344\sqrt{3} - 8A} \right], \quad (4.7)$$

$$\frac{\varphi^2(e^{-\pi/2\sqrt{6}})}{\varphi^2(e^{-\sqrt{3}\pi/2\sqrt{2}})} = \sqrt{3} \left[\sqrt{596 + 344\sqrt{3} - 8A} + \sqrt{595 + 344\sqrt{3} - 8A} \right], \quad (4.8)$$

where $A := (12 + 7\sqrt{3})(\sqrt{19 + 11\sqrt{3}})$.

$$\frac{\varphi(e^{-4\pi/\sqrt{3}})}{\varphi(e^{-4\pi\sqrt{3}})} = 3^{1/4} \left[\sqrt{\frac{103 + 42\sqrt{6} - 2B}{2}} - \sqrt{\frac{101 + 42\sqrt{6} - 2B}{2}} \right], \quad (4.9)$$

$$\frac{\varphi(e^{-\pi/4\sqrt{3}})}{\varphi(e^{-\pi\sqrt{3}/4})} = 3^{1/4} \left[\sqrt{\frac{103 + 42\sqrt{6} - 2B}{2}} + \sqrt{\frac{101 + 42\sqrt{6} - 2B}{2}} \right], \quad (4.10)$$

where $B := \sqrt{5188 + 2118\sqrt{6}}$.

$$\begin{aligned} \frac{\varphi^2(e^{-\pi\sqrt{14/3}})}{\varphi^2(e^{-\pi\sqrt{42}})} &= \sqrt{3} \left[\sqrt{16352 - 9440\sqrt{3} + 3568\sqrt{21}} - 6180\sqrt{7} \right. \\ &\quad \left. - \sqrt{16351 - 9440\sqrt{3} + 3568\sqrt{21}} - 6180\sqrt{7} \right], \end{aligned} \quad (4.11)$$

$$\begin{aligned} \frac{\varphi^2(e^{-\pi/\sqrt{42}})}{\varphi^2(e^{-\pi\sqrt{3/14}})} &= \sqrt{3} \left[\sqrt{16352 - 9440\sqrt{3} + 3568\sqrt{21}} - 6180\sqrt{7} \right. \\ &\quad \left. + \sqrt{16351 - 9440\sqrt{3} + 3568\sqrt{21}} - 6180\sqrt{7} \right], \end{aligned} \quad (4.12)$$

$$\frac{\varphi^2(e^{-\pi/6})}{\varphi^2(e^{-\pi/2})} = (\sqrt{6} - \sqrt{3})(3^{1/4} + 1) + 3^{1/4}, \quad (4.13)$$

$$\frac{\varphi^2(e^{-2\pi})}{\sqrt{3}\varphi^2(e^{-6\pi})} = 39 + 22\sqrt{3} - \sqrt{2a} + \sqrt{2 \left[3009 + 1737\sqrt{3} - \frac{115713\sqrt{2} + 66807\sqrt{6}}{\sqrt{a}} \right]}, \quad (4.14)$$

where $a := 1479 + 854\sqrt{3}$.

$$\frac{\varphi^2(e^{-\pi\sqrt{10/3}})}{\sqrt{3}\varphi^2(e^{-\pi\sqrt{30}})} = 2\sqrt{10b} + (13\sqrt{3} - 22) - \sqrt{1968 - 1136\sqrt{3} - \sqrt{10} \frac{(4486 - 2590\sqrt{3})}{\sqrt{b}}}, \quad (4.15)$$

$$\frac{\varphi^2(e^{-\pi/\sqrt{30}})}{\sqrt{3}\varphi^2(e^{-\pi\sqrt{3/10}})} = 2\sqrt{10b} + (13\sqrt{3} - 22) + \sqrt{1968 - 1136\sqrt{3} - \sqrt{10} \frac{(4486 - 2590\sqrt{3})}{\sqrt{b}}}, \quad (4.16)$$

where $b := 26 - 15\sqrt{3}$.

Proof of (4.1). Putting $\alpha = \pi/\sqrt{3}$ and $\beta = \pi\sqrt{3}$ in (2.1), we find that

$$\frac{\varphi(e^{-\pi/\sqrt{3}})}{\varphi(e^{-\pi\sqrt{3}})} = 3^{1/4}. \quad (4.17)$$

Using the equation (4.17) in the equation (3.1), with $q = e^{-\pi/\sqrt{3}}$, we deduce that

$$3^{1/4}T + \frac{1}{3^{1/4}T} + \frac{3^{1/4}}{T} - \frac{T}{3^{1/4}} = 4, \quad \text{where } T = \frac{\varphi(e^{-2\pi/\sqrt{3}})}{\varphi(e^{-2\pi\sqrt{3}})}. \quad (4.18)$$

Solving the above equation (4.18) for T and $1 < T < 2$, we arrive at the equation (4.1). \square

Proof of (4.2). Setting $q = e^{-\pi/2\sqrt{3}}$ in the equation (3.1), then using the equation (4.17), we deduce that

$$t^2 - 2t^{3/4} + 2t^{1/4} + 2\sqrt{3} - 3 = 0, \quad \text{where } t = \frac{\varphi(e^{-\pi/2\sqrt{3}})}{\varphi(e^{-\pi\sqrt{3}/2})}. \quad (4.19)$$

Solving the above equation (4.19) for t and $1 < t < 2$, we arrive at the equation (4.2). \square

Since the proofs of the identities (4.3)–(4.16) are similar to the proofs of the identities (4.1) and (4.2). Hence we omit the details.

5. MODULAR IDENTITIES FOR A CONTINUED FRACTION OF ORDER TWELVE

In [11], M. S. Mahadeva Naika, B. N. Dharmendra and K. Shivashankara have established a continued fraction of order twelve. They have also established several explicit evaluations, reciprocity theorems and integral representations.

In this section, we establish some new modular relations for a continued fraction of order twelve using the identities established in Section 3. We also establish some explicit evaluations for a continued fraction of order twelve.

Lemma 5.1. [6], [11, Theorem 3.1] *We have*

$$\frac{\varphi(q)}{\varphi(q^3)} = \frac{1 + H(q)}{1 - H(q)}, \quad \text{where } H(q) = \frac{qf(-q, -q^{11})}{f(-q^5, -q^7)}. \quad (5.1)$$

Theorem 5.1. *If $u := H(q)$ and $v := H(q^2)$, then*

$$(u + v)^2 = u^2v + v. \quad (5.2)$$

Proof. Using the equations (3.1) and (5.1), we obtain the equation (5.2). \square

Theorem 5.2. *If $u := H(q)$ and $v := H(q^4)$, then*

$$(1 - v)(1 + v^2)(v - u^4) + (2v^3 + 2v - 8v^2)u^2 + 4uv(v - 1)(1 - vu^2) = 0. \quad (5.3)$$

Proof. Using the equations (3.5) and (5.1), we obtain the equation (5.3). \square

Theorem 5.3. *If $u := H(q)$ and $v := H(q^6)$, then*

$$\begin{aligned} & (v^5 + v^2 - 1 - v^3)(u^6 - v) + 6(-v^4 + v^5 - v + v^2)(u - u^5) \\ & + 9(1 - 3v - v^3 + 3v^2)(u^2v - v^2u^4) + 2(8v^2 + 8v^4 - 18v^3 - v^5 - v)u^3 = 0. \end{aligned} \quad (5.4)$$

Proof. Using the equations (3.6) and (5.1), we obtain the equation (5.4). \square

Theorem 5.4. *If $u := H(q)$ and $v := H(q^8)$, then*

$$\begin{aligned} & (v^7 - 3v^2 + v - v^6 + 3v^5 - 3v^4 - 1 + 3v^3)(u^8 - v) + 8(v - 3v^5 - v^3 - v^7) \\ & + 2v^4 + 2v^6)u^7 + 4(14v^5 + 5v^7 + 3v^2 + 5v^3 - 10v^4 - 13v^6 - 4v)u^6 \\ & + 8(v^2 + v + 12v^4 - 10v^3 + 3v^6 - 5v^5 - 2v^7)u^5 + 2(v + v^7 + 35v^3 + 35v^5) \\ & - 72v^4 - 4v^6 - 4v^2)u^4 + 8(v^7 - 2v + 12v^4 - 5v^3 + 3v^2 - 10v^5 + v^6)u^3 \\ & + 4(3v^6 - 13v^2 + 14v^3 - 10v^4 + 5v + 5v^5 - 4v^7)u^2 + 8(v^7 + 2v^2 + 2v^4) \\ & - v - v^5 - 3v^3)u = 0. \end{aligned} \quad (5.5)$$

Proof. Using the equations (3.9) and (5.1), we obtain the equation (5.5). \square

Theorem 5.5. *If $u := H(q)$ and $v := H(q^{10})$, then*

$$\begin{aligned} & [1 + 15v^2 - 6v^5 + v^6 - 20v^3 + 15v^4 - 6v][u^{12}v^5 + v] + 2v^5[-1 - 55v^4 + 30v^5 \\ & - 35v^2 + 6v + 60v^3 - 5v^6]u^{11} + [-185v^8 - 1 - 50v^2 + 35v^{11} - 190v^4 + 11v \\ & - 190v^{10} - 56v^6 + 125v^3 + 80v^7 + 176v^5 + 245v^9]u^{10} + 2v^2[-140v + 20 + 340v^2 \\ & + 40v^6 - 95v^7 + 110v^8 - 115v^5 - 425v^3 + 290v^4 - 25v^9]u^9 + [-1020v^4 - 160v^8 \\ & + 60v^2 - 20v - 1630v^6 + 865v^7 + 1675v^5 + 230v^3 - 130v^{10} + 25v^{11} + 105v^9]u^8 \\ & + 4v[10 + 5v^8 + 100v^7 + 280v^3 + 678v^5 - 415v^6 - 575v^4 + 2v^{10} - 10v^9 - 40v \\ & - 35v^2]u^7 + 2v[-10v^8 + 95v^9 - 10v^2 - 420v^7 + 95v + 1150v^4 - 20v^{10} + 1150v^6 \\ & - 1622v^5 - 420v^3 - 20]u^6 + 4v[2 - 10v + 280v^7 - 575v^6 + 100v^3 + 10v^{10} + 678v^5 \\ & - 35v^8 - 415v^4 - 40v^9 + 5v^2]u^5 + [230v^9 - 20v^{11} - 160v^4 - 130v^2 + 25v + 60v^{10} \\ & + 105v^3 + 1675v^7 + 865v^5 - 1020v^8 - 1630v^6]u^4 + 2v[110v - 425v^6 - 25 - 95v^2 \end{aligned}$$

$$\begin{aligned}
 & - 115v^4 - 140v^8 + 40v^3 + 340v^7 + 20v^9 + 290v^5]u^3 + [245v^3 - 190v^2 + 35v \\
 & - 190v^8 - 56v^6 + 11v^{11} + 176v^7 + 80v^5 - 185v^4 + 125v^9 - 50v^{10} - v^{12}]u^2 \quad (5.6) \\
 & + [-2v^7 + 60v^2 + 120v^4 - 110v^3 - 70v^5 + 12v^6 - 10v]u = 0.
 \end{aligned}$$

Proof. Using the equations (3.10) and (5.1), we obtain the equation (5.6). \square

Remark 1. Similarly, one can establish modular relation for $H(q)$ with $H(q^{12})$ using the equation (3.13), $H(q)$ with $H(q^{14})$ using the equation (3.14) and $H(q)$ with $H(q^{16})$ using the equation (3.15).

In the following theorem, we establish some new explicit evaluations for a continued fraction of order twelve $H(q)$, by using the evaluations of ratios of theta function established in section 4.

Theorem 5.6. *We have*

$$H(e^{-2\pi/\sqrt{3}}) = \frac{3^{1/4}\sqrt{(\sqrt{3}+1)(\sqrt{2}-1)} - 1}{3^{1/4}\sqrt{(\sqrt{2}+1)(\sqrt{3}-\sqrt{2})} + 1}, \quad (5.7)$$

$$H(e^{-\pi/2\sqrt{3}}) = \frac{3^{1/4}\sqrt{(\sqrt{3}-1)(\sqrt{2}+1)} - 1}{3^{1/4}\sqrt{(\sqrt{3}-1)(\sqrt{2}+1)} + 1}, \quad (5.8)$$

$$H(e^{-\pi\sqrt{2/3}}) = \frac{\sqrt{(\sqrt{6}+\sqrt{3})(\sqrt{3}-\sqrt{2})} - 1}{\sqrt{(\sqrt{6}+\sqrt{3})(\sqrt{3}-\sqrt{2})} + 1}, \quad (5.9)$$

$$H(e^{-\pi/\sqrt{6}}) = \frac{\sqrt{(\sqrt{6}-\sqrt{3})(\sqrt{3}+\sqrt{2})} - 1}{\sqrt{(\sqrt{6}-\sqrt{3})(\sqrt{3}+\sqrt{2})} + 1}, \quad (5.10)$$

$$H(e^{-\pi\sqrt{2}}) = \frac{\sqrt{21+9\sqrt{6}-15\sqrt{2}-12\sqrt{3}-1}}{\sqrt{21+9\sqrt{6}-15\sqrt{2}-12\sqrt{3}+1}}, \quad (5.11)$$

$$H(e^{-\pi/3\sqrt{2}}) = \frac{\sqrt{\sqrt{6}+\sqrt{2}-1} - 1}{\sqrt{\sqrt{6}+\sqrt{2}-1} + 1}, \quad (5.12)$$

$$H(e^{-\pi/6}) = \frac{\sqrt{\sqrt[4]{3}+(\sqrt[4]{3}+1)(\sqrt{6}-\sqrt{3})} - 1}{\sqrt{\sqrt[4]{3}+(\sqrt[4]{3}+1)(\sqrt{6}-\sqrt{3})} + 1}. \quad (5.13)$$

Remark 2. One can explicitly evaluate $H(e^{-\pi\sqrt{n/3}})$ for $n=8, 1/8, 10, 1/10, 12, 14, 1/14, 16$ and $1/16$.

6. RAMANUJAN'S REMARKABLE PRODUCT OF THETA FUNCTIONS

On page 338 in his first notebook, Ramanujan defines

$$a_{m,n} := \frac{ne^{\frac{-(n-1)\pi}{4}\sqrt{\frac{m}{n}}}\psi^2(e^{-\pi\sqrt{mn}})\varphi^2(-e^{-2\pi\sqrt{mn}})}{\psi^2(e^{-\pi\sqrt{\frac{m}{n}}})\varphi^2(-e^{-2\pi\sqrt{\frac{m}{n}}})}, \quad (6.1)$$

and offers a list of eighteen particular values. All these eighteen values have been established by B. C. Berndt, H. H. Chan and L. C. Zhang [5]. In [10], Mahadeva Naika and B. N. Dharmendra have established some new general theorems for the explicit evaluations of Ramanujan's remarkable product of theta-functions $a_{m,n}$. For more details one can see [8] and [9].

In this section, we compute some new explicit evaluations of Ramanujan's remarkable product of theta functions $a_{3,m}$ for $m = 2, 4, 6, 8, 10, 12, 14$ and 16 .

Lemma 6.1. [10, Theorem 2.4] If $Q = \frac{\varphi(e^{-\pi\sqrt{\frac{m}{3}}})}{\varphi(e^{-\pi\sqrt{3m}})}$, then

$$a_{3,m}^2 = \frac{9(1 - Q^4)}{Q^4(Q^4 - 9)}, \quad Q^4 \neq 9, \quad (6.2)$$

where m is any positive rational.

Theorem 6.1. We have

$$a_{3,2}^2 = (\sqrt{3} + \sqrt{2})(3 - 2\sqrt{2}), \quad (6.3)$$

$$a_{3,4}^2 = \frac{(11\sqrt{3} - 19)(3 + 2\sqrt{2})}{\sqrt{2}}, \quad (6.4)$$

$$a_{3,6}^2 = \frac{4\sqrt{2} + \sqrt{6} - 2 - \sqrt{3}}{9}. \quad (6.5)$$

Remark 3. One can explicitly evaluate $a_{3,m}$ for $m = 8, 10, 12, 14$ and 16 .

REFERENCES

- [1] C. Adiga, Taekyun Kim, M. S. Mahadeva Naika and H. S. Madhusudhan, On Ramanujan's cubic continued fraction and explicit evaluations of theta-functions, Indian J. Pure Appl. Math., 35(9) (2004), 1047–1062.
- [2] N. D. Baruah, Modular equations for Ramanujan's cubic continued fraction, J. Math. Anal. Appl., 268 (2002), 244–255.
- [3] B. C. Berndt, Ramanujan's Notebooks, Part III, Springer-Verlag, New York, 1991.
- [4] B. C. Berndt, Ramanujan's Notebooks, Part IV, Springer-Verlag, New York, 1994.
- [5] B. C. Berndt, H. H. Chan and L.-C. Zhang, Ramanujan's remarkable product of the theta-function, Proc. Edinburgh Math. Soc., 40 (1997), 583–612.

[6]

- [7] M. S. Mahadeva Naika, S. Chandankumar and K. Sushan Bairy, New identities for Ramanujan's cubic continued fraction, *Funct. Approx. Comment. Math.*, 46(1) (2012), 29–44.
- [8] M. S. Mahadeva Naika, S. Chandankumar and M. Harish, On some new P - Q mixed modular equations, *Ann. Univ. Ferrara Sez. VII (N.S.)*, 59(2) (2013), 353–374.
- [9] M. S. Mahadeva Naika, S. Chandankumar and B. Hemanthkumar, On some new modular relations for Ramanujan's remarkable product of theta functions, *Adv. Stud. Contemp. Math.* 23(3) (2013), 431–449.
- [10] M. S. Mahadeva Naika and B. N. Dharmendra, On some new theorems for the explicit evaluations of Ramanujan's remarkable product of theta-function, *Ramanujan J.*, 15(3) (2008), 349–366.
- [11] M. S. Mahadeva Naika, B. N. Dharmendra and K. Shivashankara, A continued fraction of order twelve, *Cent. Eur. J. Math* 6(3) (2008), 393–404.
- [12] S. Ramanujan, Notebooks (2 volumes), Tata Institute of Fundamental Research, Bombay, 1957.
- [13] S. Ramanujan, The lost notebook and other unpublished papers, Narosa, New Delhi, 1988.

(M. S. Mahadeva Naika) DEPARTMENT OF MATHEMATICS, BANGALORE UNIVERSITY, CENTRAL COLLEGE CAMPUS, BANGALORE-560 001, KARNATAKA, INDIA

E-mail address: msmnaika@rediffmail.com

(S. Chandankumar) DEPARTMENT OF MATHEMATICS, M. S. RAMAIAH UNIVERSITY OF APPLIED SCIENCES, PEENYA CAMPUS, PEENYA 4TH PHASE, BENGALURU-560 058, KARNATAKA, INDIA

E-mail address: chandan.s17@gmail.com

(K. Sushan Bairy) PG DEPARTMENT OF MATHEMATICS, VIJAYA COLLEGE, R. V. ROAD, BASAVANAGUDI,, BANGALORE-560 004, INDIA.

E-mail address: ksbairy@rediffmail.com