

## Original Paper

# Role of Dicer Enzyme in the Regulation of Store Operated Calcium Entry (SOCE) in CD4<sup>+</sup> T Cells

Shaqiu Zhang<sup>a,b</sup> Tamer al-Maghout<sup>b</sup> Yuetao Zhou<sup>b</sup> Rosi Bissinger<sup>b</sup>  
 Abeer Abousaab<sup>b</sup> Madhuri S. Salker<sup>b,c</sup> Lisann Pelzl<sup>b</sup> Bradley S. Cobb<sup>d</sup>  
 Anchun Cheng<sup>a</sup> Yogesh Singh<sup>b</sup> Florian Lang<sup>b,e</sup>

<sup>a</sup>Institute of Preventive Veterinary Medicine, Sichuan Agricultural University, Wenjiang, Chengdu city, China; <sup>b</sup>Departments of Cardiology, Vascular Medicine and Physiology, Tuebingen University, <sup>c</sup>Institute of Women's Health, Tuebingen University, Tuebingen, Germany; <sup>d</sup>Department of Comparative Biomedical Sciences, The Royal Veterinary College, London, United Kingdom; <sup>e</sup>Department of Molecular Medicine II, Heinrich Heine University Düsseldorf, Düsseldorf, Germany

**Key Words**

CD4<sup>+</sup> T cells • Dicer • SOCE and miRNAs

**Abstract**

**Background/Aims:** Activation of T cell receptors (TCRs) in CD4<sup>+</sup> T cells leads to a cascade of signalling reactions including increase of intracellular calcium (Ca<sup>2+</sup>) levels with subsequent Ca<sup>2+</sup> dependent stimulation of gene expression, proliferation, cell motility and cytokine release. The increase of cytosolic Ca<sup>2+</sup> results from intracellular Ca<sup>2+</sup> release with subsequent activation of store-operated Ca<sup>2+</sup> entry (SOCE). Previous studies suggested miRNAs are required for the development and functions of CD4<sup>+</sup> T cells. An enzyme called Dicer is required during the process of manufacturing mature miRNAs from the precursor miRNAs. In this study, we explored whether loss of Dicer in CD4<sup>+</sup> T cells affects SOCE and thus Ca<sup>2+</sup> dependent regulation of cellular functions. **Methods:** We tested the expression of Orai1 by q-RT-PCR and flow cytometry. Further, we measured SOCE by an inverted phase-contrast microscope with the Incident-light fluorescence illumination system using Fura-2. Intracellular Ca<sup>2+</sup> was also measured by flow cytometry using Ca<sup>2+</sup> sensitive dye Fluo-4. **Results:** We found that in Dicer deficient (*Dicer*<sup>Δ/Δ</sup>) mice Orai1 was downregulated at mRNA and protein level in CD4<sup>+</sup> T cells. Further, SOCE was significantly smaller in *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells than in CD4<sup>+</sup> T cells isolated from wild-type (*Dicer*<sup>fl/fl</sup>) mice. **Conclusion:** Our data suggest that miRNAs are required for adequate Ca<sup>2+</sup> entry into CD4<sup>+</sup> T cells and thus triggering of Ca<sup>2+</sup> sensitive immune functions.

**Introduction**

After antigen-specific T cell activation, an increase of intracellular Ca<sup>2+</sup> levels is required to induce gene expression, proliferation, cell motility and cytokine expression [1-3]. In resting T cells, Ca<sup>2+</sup> is stored in the endoplasmic reticulum (ER) where it is sensed by stromal

© 2016 The Author(s)  
Published by S. Karger AG, Basel

Yogesh Singh  
Florian Lang

Institute of Physiology I, Tuebingen University, Tuebingen, Gmelinstraße5, D-72076  
Tuebingen (Germany)  
Tel. +49(0)7071 29-72194, E-Mail [ysinghbt@gmail.com](mailto:ysinghbt@gmail.com) / [florian.lang@uni-tuebingen.de](mailto:florian.lang@uni-tuebingen.de)

**KARGER**

cell-interaction molecules (STIM) 1 and 2. In T cells, signalling through the T cell receptor results in the production of inositol trisphosphate (IP<sub>3</sub>), which binds to IP<sub>3</sub> receptors on the ER and results in the release of Ca<sup>2+</sup> into the cytosol. The depletion of ER Ca<sup>2+</sup> results in Ca<sup>2+</sup> influx across the plasma membrane by store-operated Ca<sup>2+</sup> entry (SOCE) [2, 4-7]. It entails activation of the calcium release-activated calcium (CRAC) channel protein 1 (encoded by Orai1 gene) through the binding of the ER Ca<sup>2+</sup> sensors STIM1 and 2. Ca<sup>2+</sup> influx through Orai1 in T cells depends on a negative membrane potential that provides the electrical driving force for Ca<sup>2+</sup> entry [2, 6, 8, 9]. The importance of ion channel function in T cells comes largely from genetic studies in mice through knockout or siRNA mediated knock down of specific ion channel genes [2, 6]. Mice genetically defective for STIM1/2 or Orai1 have impaired T cell development, which is not surprising considering the important role these molecules play in Ca<sup>2+</sup> signalling [5, 8-10]. The regulation of the function of Orai1 channels could affect the signalling in T cells and play an important role in CD4<sup>+</sup> T cells development and function [1-3].

MicroRNAs (miRNAs) are transcribed in the genome as a part of Pol II transcribed messages and function as post-transcriptional gene regulators [11]. Biogenesis of miRNAs involves the RNase Dicer [11, 12]. In the cell nucleus, primary miRNAs are transcribed and processed into precursor miRNAs by the enzyme complex containing the RNase Drosha. These precursor miRNAs then move out from the nucleus to cytoplasm by exportin-5, where these miRNAs are processed into mature miRNAs by Dicer. Thus in the absence of Dicer, miRNAs are not produced. Generally, miRNAs bind to the 3' untranslated region (3'UTR) of mRNAs and inhibit translation and induce message degradation [13-21]. Bioinformatics studies have predicted that almost one third of the genome is targeted by miRNAs [13, 21]. Therefore, miRNAs play decisive roles in the regulation of gene expression. Dicer is involved in the regulation of diverse biological processes including development as well as organogenesis and Dicer contributes to the development of several pathologies including cancer, infection susceptibility as well as autoimmunity [22].

T cell specific deletion of Dicer results in impaired T cell development and severely reduced numbers of regulatory T cells [23, 24]. Dicer deficient mice develop inflammatory bowel disease by age of 6 months [23, 25, 26], which is a similar phenotype as seen in Orai1 and STIM1/2 deficient mice [2, 8, 27]. Dicer impacts on the development of autoimmunity by controlling the immune functions of regulatory T cells and conventional T cells. Therefore, we explored whether Dicer may influence Ca<sup>2+</sup> signalling in CD4<sup>+</sup> T cells.

In this study we found that Dicer deficient (*Dicer*<sup>Δ/Δ</sup>) CD4<sup>+</sup> T cells have reduced expression of Orai1 at mRNA and protein levels. Further studies suggested that Dicer deficient CD4<sup>+</sup> T cells have less Ca<sup>2+</sup> influx after activation with anti-CD3 and anti-CD28 compared with control (*Dicer*<sup>fl/fl</sup>) CD4<sup>+</sup> T cells. Thus, our data suggest that miRNAs are required for proper influx of Ca<sup>2+</sup> after activation of T cells.

## Material and Methods

### Mice

*Dicer*<sup>fl/fl</sup> mice (mixed C57BL/6/129 background) were bred with CD4<sup>Cre</sup> mice to generate CD4 specific *Dicer*<sup>Δ/Δ</sup> mice described earlier [23, 24] and kept in specific pathogen free conditions. Mice used for the experiments were in between 8-16 weeks of age. All the experiments were performed according to the EU Animals Scientific Procedures Act and the German law for the welfare of animals. All the procedures for the experiments were approved by the authorities of the state of Baden-Württemberg.

### CD4<sup>+</sup> T cell isolation and culture

CD4<sup>+</sup> naïve T cells were isolated from *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> mice using the MagniSort<sup>®</sup> Mouse naïve T cell Enrichment kit as described by the manufacturer (eBioscience, Frankfurt, Germany). Purified T cells were cultured in plate-bound anti-CD3: and anti-CD28 (1:2 dilution; 1 μg/ml anti-CD3 and 2 μg/ml anti-CD28) for 2-3 days and then subjected to the measurement of intracellular Ca<sup>2+</sup> by flow cytometry (Fluo-4) and fluorescence microscopy (Fura-2/AM).

## Calcium measurements

Fluorescence measurements were performed using an inverted phase-contrast microscope with the incident-light fluorescence illumination system (Axiovert 100, Zeiss, Germany). Cells were excited alternatively at  $\lambda = 340$  or  $380$  nm and the light deflected by a dichroic mirror into either the objective (Fluar 40 $\times$ /1.30 oil, Zeiss, Germany) or a camera (Proxitronic, Germany). Emitted fluorescence intensity was recorded at  $\lambda = 505$  nm and data were acquired by using specialized computer software (Metafluor, Universal Imaging, USA) [28].

Activated cells (anti-CD3/anti-CD28) for 3 days from both mice strains were loaded with Fura-2-AM (2  $\mu$ M, Molecular Probes, Germany) for 30 min at 37°C in a CO<sub>2</sub> incubator. To measure SOCE, changes in intracellular Ca<sup>2+</sup> [Ca<sup>2+</sup>]<sub>i</sub> were monitored on depletion of the intracellular Ca<sup>2+</sup> stores. In brief, [Ca<sup>2+</sup>]<sub>i</sub> was measured using Ca<sup>2+</sup> containing standard HEPES buffer [125mM/L NaCl, 5mM KCl, 1.2 mM MgSO<sub>4</sub>\*7H<sub>2</sub>O, 32.2 mM HEPES, 2mM Na<sub>2</sub>HPO<sub>4</sub>\*2H<sub>2</sub>O, 5mM Glucose, 1mM CaCl<sub>2</sub>\*2H<sub>2</sub>O; pH=7.4] for 2 minutes and then changed to Ca<sup>2+</sup> free HEPES buffer [125mM NaCl, 5mM KCl, 1.2 mM MgSO<sub>4</sub>\*7H<sub>2</sub>O, 32.2 mM HEPES, 2mM Na<sub>2</sub>HPO<sub>4</sub>\*2H<sub>2</sub>O, 5mM Glucose, 0.5 mM EGTA; pH=7.4] for 3 minutes. In the absence of Ca<sup>2+</sup>, the intracellular Ca<sup>2+</sup> stores were depleted by inhibition of the vesicular Ca<sup>2+</sup> pump by thapsigargin (1  $\mu$ M, Sigma, Germany) and [Ca<sup>2+</sup>]<sub>i</sub> activity was measured for another 5 minutes. Furthermore, Ca<sup>2+</sup> containing HEPES buffer was added for 5 minutes, which allowed assessing SOCE.

## q-RT-PCR

Total mRNA was isolated from *Dicer<sup>f/f</sup>* and *Dicer<sup>Δ/Δ</sup>* mice using the mRNAeasy isolation kit (QIAGEN, Germany) as described by the manufacturer. 1 $\mu$ g mRNA was converted into cDNA using the cDNA synthesis kit (Invitrogen, Germany). Briefly, in 10  $\mu$ l reactions, 10 ng cDNA, 2X SYBR green mastermix (KAPA SYBR® FAST q-PCR kit Master Mix (2x) Bio-Rad iCycler™; Peqlab, Erlangen, Germany) and 250 nM primers were used for q-RT-PCR reactions. q-RT-PCR run and data analysis was performed as described previously [29]. Universal cycling conditions were used (95° C for 10 minutes, 95° C for 15 seconds and 60° C for 1 minute for 40 cycles followed by melting curve analysis) for q-RT-PCR [30]. The following primers were used:

Orai1-F 5'- CCTGGCGCAAGCTCTACTTA-3'  
Orai1-R 5'- CATCGCTACCATGGCGAAGC-3'  
GAPDH-F 5'-TCTGACCACAGTGAGGAATGTCCAC-3'  
GAPDH-R 5'-TTGATGGCAACAATCTCCAC-3'

## Fluo-4 Calcium measurement by flow cytometry

Activated CD4<sup>+</sup> T cells from *Dicer<sup>f/f</sup>* and *Dicer<sup>Δ/Δ</sup>* mice were washed once with PBS and then incubated in Ringer solution with calcium containing Fluo4 and Fluo3 (Invitrogen, Germany) for 30 minutes in a 37 °C incubator as described earlier [31]. After incubation, cells were washed 3 times with Ringer solution. 200  $\mu$ l of Ringer solution was added, and intracellular calcium levels were measured by flow cytometry analysing 20,000 cells. Data were analysed by FlowJo (Treestar, USA)

## Statistics

Data are provided as means  $\pm$  SEM. n represents the number of independent experiments. Data were tested for statistical significance using unpaired Student's t-test. Data were analysed by Excel 2010 or GraphPad Prism Software, USA. A p value of  $\leq 0.05$  was considered statistically significant.


## Results

### Orai1 expression by q-RT-PCR and flow cytometry

Previous studies suggested that various ion channels such as Ca<sup>2+</sup> release activated Ca<sup>2+</sup> (CRAC) channels, K<sup>+</sup> channels, TRPM4 channels, TRPM7 channels and chloride channels contribute to the pathophysiology of asthma, allergy and inflammatory bowel disease [2, 6, 10]. According to published microarray data from *Dicer<sup>Δ/Δ</sup>* mice potassium channel K<sup>+</sup> two pore domain channel subfamily K (KCNK) 1, KCNK6 and Mg<sup>2+</sup> permeable channel (TRPM7) are upregulated in regulatory T cells compared to *Dicer<sup>f/f</sup>* regulatory T cells [32]. Thus, these data suggested that ion channels could be dysregulated in miRNAs deficient immune T cells.


Given the important role of both Ca<sup>2+</sup> signalling and miRNAs in key physiological processes, the role of miRNAs in the regulation of calcium channels in T cell differentiation

and effective function was addressed. Channels accomplishing Ca<sup>2+</sup> entry into lymphocytes include CRAC channels composed of the pore-forming units Orai1, 2 and 3 and their regulators STIM1 and 2, which can be activated by emptying of the intracellular Ca<sup>2+</sup> stores [2, 33]. SOCE following activation of T cells is mediated by the Orai1 channel protein [2]. Therefore, we measured Orai1 expression at mRNA transcript and protein levels. We observed significantly lower mRNA expression in *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells than in *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (Fig. 1A). Similarly, measurement of Orai1 expression by flow cytometry revealed that Orai1 protein abundance was significantly lower in *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells than in *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (Fig. 1B).




**Fig. 1.** *Dicer*<sup>Δ/Δ</sup> mice express reduced Orai1 at transcript and protein levels in CD4<sup>+</sup> T cells. A. mRNA was isolated from *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells and equal amount of mRNA was converted into cDNA. q-RT-PCR was performed for quantification of Orai1 transcript levels. B. Protein expression of Orai1 in *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells by flow cytometry. Left hand side shows the representative FACS histogram of Orai1 expression and right hand side shows arithmetic means ± SEM of n=3 independent experiments. \* indicates statistical significance difference (p<0.05).

**Fig. 2.** *Dicer*<sup>Δ/Δ</sup> mice have decreased calcium influx (SOCE) in ex vivo isolated CD4<sup>+</sup> T cells. CD4<sup>+</sup> T cells were isolated from *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> mice spleen and lymph nodes and Ca<sup>2+</sup> entry into CD4<sup>+</sup> T cells measured using Fura-2 fluorescence under an inverted phase-contrast microscope with the incident-light fluorescence illumination system. A. The average of 100-120 cells was used for measuring the Ca<sup>2+</sup> entry. Representative tracings showing the 340/380 nm fluorescence ratio in Fura-2/AM loaded T cells from *Dicer*<sup>Δ/Δ</sup> and *Dicer*<sup>fl/fl</sup> mice upon removal of extracellular Ca<sup>2+</sup> followed by exposure to thapsigargin (1μM) and Ca<sup>2+</sup> readmission. B. Arithmetic means ± SEM of the slope (left) and peak (right) of the fluorescence ratio change between *Dicer*<sup>Δ/Δ</sup> and *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (n=3; independent experiments). \* indicates statistical significance difference (p<0.05).



**Fig. 3.** SOCE is decreased in activated CD4<sup>+</sup> T cells of *Dicer*<sup>Δ/Δ</sup> mice. CD4<sup>+</sup> T cells were isolated from *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> mice spleen and lymph nodes and activated in the presence of anti-CD3 and anti-CD28 (plate bound) and Ca<sup>2+</sup> entry into CD4<sup>+</sup> T cells measured using Fura-2 fluorescence. A. Representative tracings showing the 340/380 nm fluorescence ratio in Fura-2/AM loaded activated CD4<sup>+</sup> T cells from *Dicer*<sup>Δ/Δ</sup> and *Dicer*<sup>fl/fl</sup> mice upon removal of extracellular Ca<sup>2+</sup> followed by exposure to thapsigargin (1 μM) and Ca<sup>2+</sup> readmission. B. Arithmetic means ± SEM of the slope (left) and peak (right) of the fluorescence ratio change between *Dicer*<sup>Δ/Δ</sup> and *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (n=3; independent experiments). \* indicates statistical significance difference (p<0.05).




#### Ca<sup>2+</sup> entry in ex vivo isolated *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells


We explored whether SOCE into *ex vivo* CD4<sup>+</sup> T cells is modified by *Dicer* deficiency [3, 34]. Thus, we measured the intracellular Ca<sup>2+</sup> activity ([Ca<sup>2+</sup>]<sub>i</sub>) and SOCE from *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells using the Fura-2 AM dye. *Ex vivo* isolated CD4<sup>+</sup> T cells from both *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> mice were loaded with Fura-2 for 30 minutes in standard HEPES buffer containing Ca<sup>2+</sup> and washed once with standard HEPES buffer containing Ca<sup>2+</sup> and then with Ca<sup>2+</sup> free HEPES buffer. Ca<sup>2+</sup> stores were then depleted by addition of sarco/endoplasmic reticulum Ca<sup>2+</sup> ATPase (SERCA) inhibitor thapsigargin (1 μM) in the nominal absence of extracellular Ca<sup>2+</sup>. The intracellular Ca<sup>2+</sup> ([Ca<sup>2+</sup>]<sub>i</sub>) manoeuvre was similar in between *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells (Fig. 2A, B). Addition of extracellular Ca<sup>2+</sup> (1 μM) in the continued presence of thapsigargin triggered SOCE. Peak of SOCE was significantly lower whereas slope of SOCE was not significantly different between *Dicer*<sup>Δ/Δ</sup> and *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (Fig. 2A, B).

#### Ca<sup>2+</sup> entry in activated *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells

The increase of [Ca<sup>2+</sup>]<sub>i</sub> plays a decisive role during the initial phase of T cell activation, particularly for the production of cytokines, cell proliferation and cell death [2, 28, 34-37]. Therefore, we also measured the SOCE from activated CD4<sup>+</sup> T cells. CD4<sup>+</sup> T cells from both *Dicer*<sup>fl/fl</sup> and *Dicer*<sup>Δ/Δ</sup> mice were activated for 2 days in the presence of plate-bound anti-CD3 and anti-CD28, after which Ca<sup>2+</sup> entry was measured. Cells were loaded with Fura-2 for 30 minutes in standard HEPES buffer and washed once with standard HEPES buffer and further with Ca<sup>2+</sup> free HEPES buffer as described in Fig. 2. First, depletion of Ca<sup>2+</sup> stores by the SERCA inhibitor thapsigargin (1 μM) in the nominal absence of extracellular Ca<sup>2+</sup> was performed. Then, [Ca<sup>2+</sup>]<sub>i</sub> measurements were performed prior to and following addition of extracellular Ca<sup>2+</sup> (1 μM) in the continued presence of thapsigargin resulting in SOCE. Slope and peak of SOCE were significantly lower in *Dicer*<sup>Δ/Δ</sup> than in *Dicer*<sup>fl/fl</sup> CD4<sup>+</sup> T cells (Fig. 3A, B). In addition to this, when we compared the Ca<sup>2+</sup> entry in between unactivated and activated T cells from



**Fig. 4.** Intracellular Ca<sup>2+</sup> is increased after activation of T cells in CD4<sup>+</sup> T cells of Dicer<sup>Δ/Δ</sup> mice. Measurement of intracellular Ca<sup>2+</sup> measurement by Fluo-2/AM dye fluorescence in unactivated and activated Dicer<sup>Δ/Δ</sup> and Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells. Activated Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells have significantly higher levels of SOCE compared with unactivated cells, whereas no difference was observed in Dicer<sup>Δ/Δ</sup> T cells. \* indicates statistical significance difference (p<0.05).



**Fig. 5.** Intracellular Ca<sup>2+</sup> is decreased in anti-CD3 and anti-CD28 activated CD4<sup>+</sup> T cells of Dicer<sup>Δ/Δ</sup> mice. Measurement of intracellular Ca<sup>2+</sup> by Fluo-4/AM dye fluorescence in activated Dicer<sup>Δ/Δ</sup> and Dicer<sup>fl/fl</sup> T cells. Dicer<sup>Δ/Δ</sup> T cells have significantly less intracellular Ca<sup>2+</sup> in comparison with Dicer<sup>fl/fl</sup> T cells. \* indicates statistical significance difference (p<0.05).

Dicer<sup>Δ/Δ</sup> and Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells. We found that Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells significantly upregulated the SOCE in activated T cells compared with unactivated T cells whereas no significant difference was observed in Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells (Fig. 4)

#### Ca<sup>2+</sup> measurement by Fluo-4 in activated Dicer<sup>fl/fl</sup> and Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells

To test the effect of Dicer deletion on Ca<sup>2+</sup> signalling by a second, independent method, we employed flow cytometry and measured [Ca<sup>2+</sup>]<sub>i</sub> by Fluo-4 dye. Similar to the experiments with fluorescence microscopy, Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells had significantly lower [Ca<sup>2+</sup>]<sub>i</sub> than Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells (Fig. 5).

## Discussion

The present study revealed a role of miRNAs in Ca<sup>2+</sup> homeostasis as a positive regulator of SOCE into T cells. SOCE was lower in the Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells compared with Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells. Orai1 expression was lower in Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells compared with Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells. Therefore, we speculate that miRNAs are required to suppress the expression of a repressor of Orai1.

Dicer is essential for maturation of miRNAs and is involved in the pathophysiology of various diseases such as cancer, infection and autoimmunity [23, 24]. In this study, we found that activation of T cells with anti-CD3 and anti-CD28 leads to enhanced Ca<sup>2+</sup> entry. This was significantly reduced in Dicer<sup>Δ/Δ</sup> CD4<sup>+</sup> T cells even though most miRNAs show decreased expression in activated T cells due to the degradation of Argonaute proteins, which are the key component of the effector complex that binds the miRNA and mediate the effects on gene regulation [12, 38]. Therefore, there are presumably still sufficient levels of key compensatory miRNAs to regulate Ca<sup>2+</sup> homeostasis in activated Dicer<sup>fl/fl</sup> CD4<sup>+</sup> T cells.

Stimulation of T cells with anti-CD3 and anti-CD28 initiates a cascade of signalling events resulting in activation of various downstream pathways such as PI3K/mTOR, JAK/Stat etc [39]. The influence of miRNAs on SOCE and thus activation of T lymphocytes may play a role in enhancing T lymphocyte activation and function. Limiting the extent and duration of TCR signalling ensures a tightly constrained response, protecting cells from harmful effects of chronic activation [28]. MiRNAs play an additional role in T cell activation through their regulation of mTOR, which is important for regulating the strength of TCR signalling leading to activation or anergy [40]. In contrast to *Dicer<sup>f/f</sup>* CD4<sup>+</sup> T cells, *Dicer<sup>Δ/Δ</sup>* CD4<sup>+</sup> T cells fail to adequately discriminate between activating and anergy-inducing stimuli as the TCR signal can elicit full activation with effective functions or state of anergy [40]. We speculate that Ca<sup>2+</sup> signalling could also play a decisive role in this process because in the absence of miRNAs there was no significant increase in SOCE upon activation in between activated and unactivated CD4<sup>+</sup> T cells from *Dicer<sup>Δ/Δ</sup>* mice. Further studies are required to find out which key miRNAs are involved in Ca<sup>2+</sup> homeostasis.

## Conclusion

The present observations uncovered a novel role of miRNAs in the regulation of Ca<sup>2+</sup> entry into CD4<sup>+</sup> T cells. Further studies are required to define the specific miRNAs and target genes necessary for the regulation of Ca<sup>2+</sup> entry into T cells. In any case, our results suggest that miRNAs are essential for the maintenance of Ca<sup>2+</sup> homeostasis. Our study could have an important implication in infection, autoimmunity and cancer progression.

## Acknowledgements

The authors gratefully acknowledge the meticulous preparation of the manuscript by Tanja Loch and Lejla Subasic and Elfried Farber for technical help. The authors are also thankful to Dr. Jakob Völkl for writing the permission for animal breeding. The Research is supported by DFG (F.L.) and Open Access Publishing Fund of Tuebingen University. MSS is supported by EMBO Long-term Fellowship (ATLF20-2013). Both SZ and YZ are supported by China Scholarship Commission.

## Disclosure Statement

The authors of this manuscript state that they do not have any financial conflict of interests and nothing to disclose.

## References

- 1 Alewine C, Kim BY, Hegde V, Welling PA: Lin-7 targets the Kir 2.3 channel on the basolateral membrane via a L27 domain interaction with CASK. *Am J Physiol Cell Physiol* 2007;293:C1733-1741.
- 2 Feske S, Skolnik EY, Prakriya M: Ion channels and transporters in lymphocyte function and immunity. *Nat Rev Immunol* 2012;12:532-547.
- 3 Fracchia KM, Pai CY, Walsh CM: Modulation of T Cell Metabolism and Function through Calcium Signaling. *Front Immunol* 2013;4:324.
- 4 Oh-Hora M, Komatsu N, Pishyareh M, Feske S, Hori S, Taniguchi M, Rao A, Takayanagi H: Agonist-selected T cell development requires strong T cell receptor signaling and store-operated calcium entry. *Immunity* 2013;38:881-895.

- 5 Oh-Hora M, Yamashita M, Hogan PG, Sharma S, Lamperti E, Chung W, Prakriya M, Feske S, Rao A: Dual functions for the endoplasmic reticulum calcium sensors STIM1 and STIM2 in T cell activation and tolerance. *Nat Immunol* 2008;9:432-443.
- 6 Shaw PJ, Feske S: Regulation of lymphocyte function by ORAI and STIM proteins in infection and autoimmunity. *J Physiol* 2012;590:4157-4167.
- 7 Waite JC, Vardhana S, Shaw PJ, Jang JE, McCarl CA, Cameron TO, Feske S, Dustin ML: Interference with Ca(2+) release activated Ca(2+) (CRAC) channel function delays T-cell arrest in vivo. *Eur J Immunol* 2013;43:3343-3354.
- 8 Feske S, Gwack Y, Prakriya M, Srikanth S, Puppel SH, Tanasa B, Hogan PG, Lewis RS, Daly M, Rao A: A mutation in Orai1 causes immune deficiency by abrogating CRAC channel function. *Nature* 2006;441:179-185.
- 9 Ma J, McCarl CA, Khalil S, Luthy K, Feske S: T-cell-specific deletion of STIM1 and STIM2 protects mice from EAE by impairing the effector functions of Th1 and Th17 cells. *Eur J Immunol* 2010;40:3028-3042.
- 10 McCarl CA, Khalil S, Ma J, Oh-hora M, Yamashita M, Roether J, Kawasaki T, Jairaman A, Sasaki Y, Prakriya M, Feske S: Store-operated Ca2+ entry through ORAI1 is critical for T cell-mediated autoimmunity and allograft rejection. *J Immunol* 2010;185:5845-5858.
- 11 Bartel DP: MicroRNAs: target recognition and regulatory functions. *Cell* 2009;136:215-233.
- 12 Bronevetsky Y, Ansel KM: Regulation of miRNA biogenesis and turnover in the immune system. *Immunol Rev* 2013;253:304-316.
- 13 Carthew RW, Sontheimer EJ: Origins and Mechanisms of miRNAs and siRNAs. *Cell* 2009;136:642-655.
- 14 Filipowicz W, Bhattacharyya SN, Sonenberg N: Mechanisms of post-transcriptional regulation by microRNAs: are the answers in sight? *Nat Rev Genet* 2008;9:102-114.
- 15 Inui M, Martello G, Piccolo S: MicroRNA control of signal transduction. *Nat Rev Mol Cell Biol* 2010;11:252-263.
- 16 Lindsay MA: microRNAs and the immune response. *Trends Immunol* 2008;29:343-351.
- 17 Liu J, Wu CP, Lu BF, Jiang JT: Mechanism of T cell regulation by microRNAs. *Cancer Biol Med* 2013;10:131-137.
- 18 Lodish HF, Zhou B, Liu G, Chen CZ: Micromanagement of the immune system by microRNAs. *Nat Rev Immunol* 2008;8:120-130.
- 19 Lu LF, Liston A: MicroRNA in the immune system, microRNA as an immune system. *Immunology* 2009;127:291-298.
- 20 O'Connell RM, Rao DS, Chaudhuri AA, Baltimore D: Physiological and pathological roles for microRNAs in the immune system. *Nat Rev Immunol* 2010;10:111-122.
- 21 Agarwal V, Bell GW, Nam JW, Bartel DP: Predicting effective microRNA target sites in mammalian mRNAs. *Elife* 2015;4:doi: 10.7554/eLife.05005.
- 22 Rupp LJ, Brady BL, Carpenter AC, De Obaldia ME, Bhandoola A, Bosselut R, Muljo SA, Bassing CH: The microRNA biogenesis machinery modulates lineage commitment during alphabeta T cell development. *J Immunol* 2014;193:4032-4042.
- 23 Cobb BS, Hertweck A, Smith J, O'Connor E, Graf D, Cook T, Smale ST, Sakaguchi S, Livesey FJ, Fisher AG, Merkenschlager M: A role for Dicer in immune regulation. *J Exp Med* 2006;203:2519-2527.
- 24 Cobb BS, Nesterova TB, Thompson E, Hertweck A, O'Connor E, Godwin J, Wilson CB, Brockdorff N, Fisher AG, Smale ST, Merkenschlager M: T cell lineage choice and differentiation in the absence of the RNase III enzyme Dicer. *J Exp Med* 2005;201:1367-1373.
- 25 Chong MM, Rasmussen JP, Rudensky AY, Littman DR: The RNaseIII enzyme Drosha is critical in T cells for preventing lethal inflammatory disease. *J Exp Med* 2008;205:2005-2017.
- 26 Liston A, Lu LF, O'Carroll D, Tarakhovskiy A, Rudensky AY: Dicer-dependent microRNA pathway safeguards regulatory T cell function. *J Exp Med* 2008;205:1993-2004.
- 27 Feske S, Prakriya M, Rao A, Lewis RS: A severe defect in CRAC Ca2+ channel activation and altered K+ channel gating in T cells from immunodeficient patients. *J Exp Med* 2005;202:651-662.
- 28 Bhavsar SK, Schmidt S, Bobbala D, Nurbaeva MK, Hosseinzadeh Z, Merches K, Fajol A, Wilmes J, Lang F: AMPKalpha1-sensitivity of Orai1 and Ca(2+) entry in T - lymphocytes. *Cell Physiol Biochem* 2013;32:687-698.
- 29 Singh Y, Garden OA, Lang F, Cobb BS: MicroRNA-15b/16 Enhances the Induction of Regulatory T Cells by Regulating the Expression of Rictor and mTOR. *J Immunol* 2015;10.4049/jimmunol.1401875.



- 30 Singh Y, Kaul V, Mehra A, Chatterjee S, Tousif S, Dwivedi VP, Suar M, Van Kaer L, Bishai WR, Das G: Mycobacterium tuberculosis controls microRNA-99b (miR-99b) expression in infected murine dendritic cells to modulate host immunity. *J Biol Chem* 2013;288:5056-5061.
- 31 Bissinger R, Modicano P, Alzoubi K, Honisch S, Faggio C, Abed M, Lang F: Effect of saponin on erythrocytes. *Int J Hematol* 2014;100:51-59.
- 32 Zhou X, Jeker LT, Fife BT, Zhu S, Anderson MS, McManus MT, Bluestone JA: Selective miRNA disruption in T reg cells leads to uncontrolled autoimmunity. *J Exp Med* 2008;205:1983-1991.
- 33 Oh-hora M, Rao A: Calcium signaling in lymphocytes. *Curr Opin Immunol* 2008;20:250-258.
- 34 Christo SN, Diener KR, Hayball JD: The functional contribution of calcium ion flux heterogeneity in T cells. *Immunol Cell Biol* 2015;93:694-704.
- 35 Greenberg ML, Yu Y, Leverrier S, Zhang SL, Parker I, Cahalan MD: Orai1 function is essential for T cell homing to lymph nodes. *J Immunol* 2013;190:3197-3206.
- 36 Kim KD, Srikanth S, Yee MK, Mock DC, Lawson GW, Gwack Y: ORAI1 deficiency impairs activated T cell death and enhances T cell survival. *J Immunol* 2011;187:3620-3630.
- 37 Christo SN, Diener KR, Nordon RE, Brown MP, Griesser HJ, Vasilev K, Christo FC, Hayball JD: Scrutinizing calcium flux oscillations in T lymphocytes to deduce the strength of stimulus. *Sci Rep* 2015;5:7760.
- 38 Bronevetsky Y, Villarino AV, Easley CJ, Barbeau R, Barczak AJ, Heinz GA, Kremmer E, Heissmeyer V, McManus MT, Erle DJ, Rao A, Ansel KM: T cell activation induces proteasomal degradation of Argonaute and rapid remodeling of the microRNA repertoire. *J Exp Med* 2013;210:417-432.
- 39 Merckenschlager M, von Boehmer H: PI3 kinase signalling blocks Foxp3 expression by sequestering Foxo factors. *J Exp Med* 2010;207:1347-1350.
- 40 Marçais A, Blevins R, Graumann J, Feytout A, Dharmalingam G, Carroll T, Amado IF, Bruno L, Lee K, Walzer T, Mann M, Freitas AA, Boothby M, Fisher AG, Merckenschlager M: microRNA-mediated regulation of mTOR complex components facilitates discrimination between activation and anergy in CD4 T cells. *J Exp Med* 2014;211:2281-2295.