



Isolation, purification, characterization and applications of serine protease from *Bacillus megaterium*

T. S. Rajesh*, Deepa, Divya, M. Mahesh and Somshekar

Azyme Biosciences Pvt.Ltd 1188, 26th Main, 9th Block Jayanagar, Bangalore-560 069 (Karnataka), INDIA

*Corresponding author. E-mail: srajesht@gmail.com

Abstract: *Bacillus megaterium* isolated from poultry farm soil was identified by standard biochemical tests and screened for the production of serine protease. Production of serine protease was done using 5 different medias by varying the type of amino acid added. The purification was done by salt precipitation, dialysis and DEAE-cellulose ion exchange chromatography. The proline containing media obtained the highest fold purification out of the five different medias (leucine lysine, proline, tryptophan and methionine containing media). The enzyme showed an optimal activity at the temperature 37°C and the pH 6 which are known as its optimum temperature and pH respectively. The enzyme was proved as a Mn²⁺ dependent serine protease as it was activated by Mn²⁺ ions and inhibited by PMSF. The molecular weight of the enzyme was determined by SDS-PAGE technique as around 30kDa. It showed an excellent detergent activity on the blood stains and a very good stability in presence of locally available detergents. The enzyme acted on the keratin protein of the chicken feather and showed a degrading capacity on the protein. So it was proved that the recently studied serine protease has a keratinase activity also. From these data I conclude that the protease isolated from *Bacillus megaterium* is a Mn²⁺ dependent serine protease which has both keratinase and detergent activity.

Keywords: *Bacillus megaterium*, Keratinase, serine protease, phenylmethanesulfonyl fluoride (PMSF)

INTRODUCTION

Serine proteases are generally the endopeptidases. These are present in the body of almost all the organisms. As the serine proteases have diverse role in human health; from non-specific digestion to highly regulated functions like embryonic development, immune response and blood coagulation, which require highly regulated and specifically limited proteolysis. So in many members of this family the activity and specificity are allosterically regulated by macromolecular ligands or small cations like Naz, Ca²⁺ (Page and Di Cera, 2006).

Subtilisins are the best-known microbial alkaline serine proteases produced by *Bacillus subtilis* (Robert, 1975). The high stability and relatively low substrate specificity made them successful for the industrial use. The alkaline proteases are employed primarily as cleansing additives. Among these proteases, the bacterial proteases are most significant, compared with animal and fungal proteases (Ward, 1985). The proteases used in detergent formulation should have a broad range pH and temperature activities (Kunamneni *et al.*, 2003).

The keratinases have several potential applications as detergent for textile fibers (Bergkvist, 1963), in leather processing (Dayanandan *et al.*, 2003, Riffel *et al.*, 2003), waste chicken feather degradation (Bockle *et al.*, 1997, Cheng *et al.*, 1995, Lin *et al.*, 1996) conversion of waste

feather to feather meal for livestock (Onifade *et al.*, 1998), hydrolysis of protein from keratinous waste materials (Kida *et al.*, 1995) etc. These facilitate the processing keratin containing waste from poultry and feather industries by nonpolluting methods (Onifade *et al.*, 1998). In addition, these proteases are used in production of foodstuffs, leather, pharmaceuticals, diagnostic reagents, waste management, silver recovery, skin care ointments, contact lens cleaners, and for research purposes in synthetic organic chemistry.

In this work, the purification and characterization of an extracellular protease isolated from gram-positive soil bacteria is done. The enzyme is proved as the serine protease by being inhibited specifically by PMSF. The detergent activity and the keratinase activity of the enzyme are also studied.

MATERIALS AND METHODS

Isolation and identification of *Bacillus megaterium* form poultry farm soil: The soil sample was collected from the Poultry farm wastage area in Madiwala, Bangalore. The micro organisms were isolated from the soil by serial dilution method (Alexander, 1965, Jackie Reynolds, 2005). The microorganisms were identified by several biochemical tests and staining techniques.

Screening for the production of enzyme: The micro organisms were screened in the Skim- Milk agar for their

Table 1. Purification of the enzyme.

	Protein Content (g)	Activity (U/ml)	Specific Activity (U/mg)	Fold Purified
Media 1:				
Crude	3.6	0.0161	0.00447	1
After Dialysis	1.7	0.6127	0.3604	80.63
After Ion Exchange	0.12	0.071	0.5917	132.4
Media 2:				
Crude	3.6	0.0161	0.00447	1
After Dialysis	1.3	0.1953	0.1502	33.6
After Ion Exchange	0.6	0.162	0.27	60.4
Media 3:				
Crude	2	0.0059	0.00295	1
After Dialysis	1.8	0.2122	0.1179	37.97
After Ion Exchange	0.16	0.0203	0.1269	43
Media 4:				
Crude	2.6	0.0176	0.00677	1
After Dialysis	1.14	0.1145	0.1004	14.83
After Ion Exchange	0.16	0.0385	0.2406	35.54
Media 5:				
Crude	3	0.0264	0.0088	1
After Dialysis	1.58	0.303	0.1918	21.8
After Ion Exchange	0.08	0.023	0.2875	32.7

protease producing ability (Abdelnasser *et al.*, 2007).

Production of serine protease: The enzyme was produced from *Bacillus megaterium* in 5 different medias which differ only in case of the amino acid they possess. Incubated the inoculated medias for 24 hours at 37°C in the shaker incubator (Huang Guangrong *et al.*, 2006).

Purification and characterisation of serine protease: The enzyme was extracted from the production media by centrifuging at 4°C. These crude enzymes were purified by 3 steps, salt precipitation (80% saturation with ammonium sulphate), dialysis and finally ion exchange chromatography. The enzyme activity of the crude, dialysed and purified enzyme samples was assayed by a standard assay method using casein as the substrate. Where one unit will hydrolyze casein to produce color equivalent to 1.0 µmole (181 mg) of tyrosine per minute at pH 7.5 at 37 °C (Anson, 1938). The protein content was also estimated in each case by Folin- Lowry method and BSA as standard. The optimum activity of the enzyme was determined by assaying these at different temperatures (4°C, 25°C, 37°C, 65°C and 100°C) and pH (2, 4, 6, 8, 10) (HUANG Guangrong *et al.*, 2006). The type of protease enzyme was determined by assaying them in presence of mg/ml of MnCl₂ and mg/ml of PMSF. The molecular weight of the enzyme was determined by SDS-PAGE technique.

Applications of serine protease: Two applications of the enzyme were studied. The keratinase activity of the enzyme was studied by treating the raw keratin source (chicken feather) at 37°C for 12 hours. The detergent activity of the enzyme was also studied by treating it on

blood stain along with a local detergent (Surf excel).

RESULTS AND DISCUSSION

The assay results showed that the proline containing media produced better active protease out of all the five medias. So it is proved that the proline has the most inducible activity among the five amino acids (Table 1). The present study supports the statement of Bjorklind and Arvidson (1978). Proline which was also found to be essential for proteinase synthesis, seems to act in the same way by affecting the intracellular concentrations of glutamic acid, aspartic acid and alanine. Proline was essential for proteinase synthesis in post-exponential growth but not for bacterial growth.

Characterization of serine protease: The effect of temperature on the activity of the purified enzyme was analyzed at 4°C to 100°C. The results showed a bell shaped curve with an optimal activity at 37°C. The enzyme will be active in a range of 27 - 55°C and the activity will be almost blocked in a higher and lower temperatures (Fig. 1). The results were much similar to Usharani and Muthuraj, 2009 where they purified protease from *Bacillus laterosporus* which is active at 40°C. The stability was decreased as the temperature was increased from the optimum due to the breakage in the hydrogen bonding. The optimum temperature was slightly higher than from other reporter's results of *Aspergillus nidulans* (Charles *et al.*, 2008).

Effect of pH on protease activity: The enzyme activity of the purified enzyme was detected at pH 2 to 10. The graph showed that the optimum pH for the enzyme is 6. The




Fig. 1. Effect of temperature on enzyme activity.




Fig. 2. Effect of pH on enzyme activity.




Fig.3. Effect of metal ion on enzyme activity.




Fig. 4. Effect of PMSF on enzyme activity.

enzyme was active in a range of 5-7 (Fig. 2). It is different to Saurabh *et al.*, 2007. In Asokan and Jayanthi, 2010. It was found to be 80°C at 9 pH. Extremely high or low pH values generally result in complete loss of activity for most enzymes. pH is also a factor in the stability of enzymes. As with activity, for each enzyme there is also a region of pH optimal stability (Bennett and Frieden, 1969).

Effect of Mn²⁺ and PMSF on protease activity: The investigation on the metal ion activity on serine protease showed that the enzyme activity increased with Mn²⁺ ions concentrations (Fig. 3). Thus the enzyme has an enhanced activity in presence of Mn²⁺ ions. Inhibition studies primarily give an insight into the nature of an enzyme, its cofactor requirement and the nature of the active center (Sigma and Mooser, 1975). The presence of PMSF inhibited the enzyme activity completely (Fig. 4). These results were similar to those of (Tsuchida *et al.*, 1986, Wang *et al.*, 2006, Yamagata and Ichishima, 1989), the protease was completely inhibited by PMSF. This proved that the enzyme is a Mn²⁺ dependent serine protease.

SDS-PAGE: The molecular weight of the protease was determined by the SDS PAGE technique by comparing with the migration distances of standard marker protein. A mixture of carbonic anhydrase, egg albumin and BSA was used as the marker, which has the molecular weight of 67, 45 and 30 kDa respectively (Kunamneni Adinarayana *et al.*, 2003). Depending on the relative mobility, the molecular weight of the protein was calculated to be around 30kDa (Plate 1.). Generally the molecular weight of the protease comes in a range 15- 30 kDa (Kelly and Fogarty, 1976).

Applications of serine protease: The serine protease enzyme produced by *Bacillus megaterium* showed a well characterized keratinase activity. The activity was shown at 37°C and a pH of 7.5 using the sodium phosphate buffer). The feather keratin has been used as the substrate for the keratinase assay in the recent study, as the feather protein has been showed to be an excellent source of the metabolizable protein (Klemersrud *et al.*, 1998) and the microbial keratinases enhance the digestibility of the feather keratin (Lee *et al.*, 1991 and Odetallah *et al.*, 2003).

Besides pH, a good detergent protease is expected to be stable in the presence of commercial detergents. The protease showed an excellent activity when it was used as an additive in detergent (Surf excel). The protease removed the entire bloodstain from the cloth and expressed a very good detergent activity at room temperature. Mohsen, 2005 reported the usefulness of protease from *Pseudomonas aeruginosa* for removal of blood stains from cotton cloth in the presence and absence of detergent.

Serine Proteases have found a wide range of applications




Plate.1. SDS PAGE of the pure enzyme.

in various industries such as food, detergent, pharmaceutical, cosmetic, *etc.* and have been widely commercialised by various companies throughout the world. Though the production of these enzymes has been improved significantly by the utilisation of hyper-producing strains of fungi and bacteria and genetically modified microbes as well, efforts are still being done to find newer sources of enzymes, better production techniques and novel applications of these enzymes in unexplored fields.

REFERENCES

- Abdelnasser, S. S., Ibrahim, Nefisa MA, El-Shayeb and Sohair, S. Mabrouk (2007). Isolation and Identification of Alkaline Protease Producing Alkaliphilic Bacteria from an Egyptian Soda Lake. *Journal of Applied Sciences Research*, 3(11): 1363-1368.
- Anson, M. L. (1938). The estimation of pepsin, trypsin, papain, and cathepsin with hemoglobin *Journal of Gen. Physiology*, 22: 79-89.
- Asokan, S. and Jayanthi, C. (2010). Alkaline protease production by *Bacillus licheniformis* and *Bacillus coagulans*, *Journal of Cell and Tissue Research*, 10 (1): 2119-2123.
- Bennett, T. P. and Frieden (1969). *Modern Topics in Biochemistry*, pg. 43-45, Macmillan, London.
- Bergkvist, R. (1963). The proteolytic enzymes of *Aspergillus oryzae*. I. Methods for estimation and isolation of the proteolytic enzymes, *Acta Chem. Scand.* 17: 1521-1540.
- Bjorklind and S. Arvidson (1978). Influence of amino acids on the synthesis of an extracellular proteinase from *Staphylococcus aureus*. *Journal of General Microbiology*, 107: 367-375.
- Bockle, B., Galunsky and R. Muller (1995). Characterization of a keratinolytic serine proteinase from *Streptomyces pactum* DSM 40530. *Appl. Environ. and Microbiol.*, 63: 55-61.
- Charles, V. Devanathan, Periasamy Anbu, M. N. Ponnuswamy, P. T. Kalaichelvan and Byung-Ki Hur (2008). Purification, characterization and crystallization of an extracellular alkaline protease from *Aspergillus nidulans* HA-10, *Journal of Basic Microbiology*, 48: 347-352
- Cheng, S. W., H. M., Hu, S. W. Shen, H. Takagi, M. Asano and Y. C. Tsai (1995). Production and characterization of keratinase of feather degrading *Bacillus licheniformis* PWD-1, *Biosci. Biotech. Biochem.*, 59:2239-2243.
- Dayanandan, A., J. Kanagaraj, L. Soundarray, R. Govindaraju and G. S. Rajkumar (2003). Application of alkaline protease in leather processing: an eco-friendly approach, *J. Cleaner Production*, 11: 533-536.
- Huang Guangrong, Ying Tiejing, Huo Po and Jiang Jiaxing (2006). Purification and characterization of a protease from *Thermophilic bacillus* strain HS08, *African Journal of Biotechnology*, 5 (24): 2433-2438.
- Jackie Reynolds (2005). Serial Dilution Protocols, Microbe Library (*Journal of Microbiology & Biology Education*).
- Kelly C.T. and Fogarty W. M. (1976) Microbial alkaline enzymes. *Process Biochem.*, 11:3-9
- Kida K, S. Morimura, J. Noda, Y. Nishida, T. Imai and M. Otagiri (1995). Enzymatic hydrolysis of horn and hoof of cow and buffalo. *J. Ferment. Bioeng.*, 80: 478-484.
- Kim, J. M., W. J. Lim and S. J. Suh (2001). Feather degrading *Bacillus* species from poultry waste. *Process Biochem.*, 37: 287- 291.
- Klemersrud, M. J., Klopfenstein, T. J. and Lewis, A. J. (1998). Complementary responses between feather meal and poultry by-product meal with or without ruminally protected methionine and lysine growing calves. *J. Anim. Sci.*, 76: 1970-1975.
- Kunamneni Adinarayana, Poluri Ellaiah and Davuluri Siva Prasad (2003). Purification and partial characterization of thermostable serine alkaline protease from a newly isolated *Bacillus subtilis* PE-11, *AAPS PharmSciTech*, 4 (4) Article 56.
- Lee, C. G, Forket, P. R. and Shih, J. C. H. (1991). Improvement of feather digestibility by bacterial keratinase as a feed additives, *FASEB. J.*, 59:1312.
- Lin, X., J. C. H. Shih and H. E. Swaisgood (1996). Hydrolysis of feather keratin by immobilized keratinase. *Appl. Environ. Microbiol.*, 62: 4273-4275.
- Mohsen Fathi Najafi, Dileep Deobagkar and Deepti Deobagkar (2005). Potential application of protease isolated from *Pseudomonas aeruginosa* PD100. *Electronic Journal of Biotechnology*, 8:2
- Niu Qihong, Huang Xiaowei, Tian Baoyu, Yang Jinkui, Liu Jiang, Zhang Lin and Zhang Keqin (2006). *Bacillus* sp. B16 kills Nematodes with a serine protease identified as a pathologic factor. *Appl. Microbiol Biotechnol.*, 69: 722-730.
- Odetallah, N. H., Wang, J. J., Garlich, J. D. and Shih, J. C. H. (2003). Keratinase in starter diets improves growth of broiler chicks, *Poultry Sci.*, 82: 664- 670.
- Onifade, A. A., N. A. Al-Sane, A. A. Al- Musallam and S. Zarban (1998). A review: Potentials for biotechnological applications of keratin-degrading microorganisms and their enzymes for nutritional improvement of feathers and other keratins as livestock feed resources. *Bioresource Technol.*, 66: 1-11.
- Page, M. J. and Di Cera, E. (2006). Role of Na and K in enzyme function. *Physiol. Rev.*, 86: 1049-1092.
- Riffel, A., F. S. Lucas, P. Heeb and A. Brandelli (2003). Characterization of new keratinolytic bacterium that

- completely degrades native feather keratin. *Archives Microbiol.*, 179: 258-265.
- Robert S. Boethling (1975). Purification and properties of serine protease from *Pseudomonas maltophilia*. *J. Bacteriol.*, 121(3): 933-941.
- Saurabh, S., Jasmine, I., Pritesh, G. and Rajendra Kumar, S. (2007). Enhanced productivity of serine alkaline protease by *Bacillus* sp. using soybean as substrate. *Malaysian Journal of Microbiology*, 3(1): 1-6.
- Sigma, D. S. and Mooser, G. (1975). Chemical studies of enzyme active sites. *Ann. Rev. Biochem.*, 44: 889-931.
- Tsuchida, O., Yamagota, Y., Ishizuka, J., Arai, J., Yamada, J., Takeuchi, M. and Ichishima, E. (1986). An alkaline protease of an alkalophilic *Bacillus* sp. *Curr. Microbiol.*, 14:7-12.
- Usharani, B. and Muthuraj, M. (2009). Production and characterization of protease enzyme from *Bacillus laterosporus*, *Global Journal of Molecular Sciences*, 4 (2): 180-186.
- Wang, R. B., Yang, J. K., Lin, C., Zhang, Y. and Zhang, K.Q.(2006). Purification and characterization of an extracellular serine protease from the nematode-trapping fungus *Dactylella shizishanna*, *Letters in applied Microbiology*, 42: 589-594.
- Ward, O. P. (1985). Proteolytic enzymes, In:Blanch HW, Drew S, Wang DI, eds. *Comprehensive Biotechnology*, 3; 789-818.
- Yamagata, Y. and Ichishima, E. (1989). A new alkaline proteinase with PI 2,8 from alkalophilic *Bacillus* species, *Curr. Microbiol.*, 19: 259-264.