



Journal of Applied and Natural Science 6 (1): 254-257 (2014)



Effect of fungal and bacterial bioagent application on total phenolic content in rice leaves pre-inoculated with *Xanthomonas oryzae* pv. *oryzae* (Uyeda and Ishiyama) Dowson

Gokil Prasad Gangwar* and A. P. Sinha

Department of Plant Pathology, G. B. Pant University of Agriculture and Technology, Pantnagar- 263145 (Uttarakhand), INDIA

*Corresponding author. E-mail: gokil_prasad@rediffmail.com

Received: January 27, 2014; Revised received: May 25, 2014; Accepted: May 30, 2014

Abstract: Present study was carried out to observe the effect of fungal and bacterial bioagents on total phenolic content in rice leaves pre-inoculated with *Xanthomonas oryzae* pv. *oryzae* and on disease severity of bacterial leaf blight of rice. Two commercial formulations of *Trichoderma harzianum* (PBA-1) and *Pseudomonas fluorescens* (PBA-2) and four formulations of fluorescent pseudomonads and *Trichoderma* spp. viz, *P. fluorescens* (Pf 83, rice leaf isolate), fluorescent pseudomonad (FLP 88, rice leaf isolate), *T. harzianum* (rice leaf isolate), *Trichoderma* spp. (isolate 40, isolated from rice field soil) were evaluated. Significantly higher mean value of total phenolic content of rice leaves was observed with the application of bioagent formulations as compared to check (pre-inoculated with *X. oryzae* pv. *oryzae*), chemical treatment and healthy plant. Maximum mean total phenolic content (342.22 µl/g) in rice leaves was observed with Pf 83, which was followed by PBA-2 (334.44 µl/g) and *T. harzianum* (330.00 µl/g). Decrease in disease severity of bacterial leaf blight was observed with the increase of total phenolic content in rice leaves which resulted in increased grain yield and 1000 grain weight.

Keywords: Disease severity, *Pseudomonas fluorescens*, Rice, Total phenolic content, *Trichoderma harzianum*, *Xanthomonas oryzae* pv. *oryzae*

INTRODUCTION

Bacterial leaf blight disease in rice caused by *Xanthomonas oryzae* pv. *oryzae* is known to occur in epidemic proportions in all rice growing areas of the world, particularly destructive in Asia (Mew *et al.*, 1993 and Anonymous, 2002). Hazardous effects of chemical pesticides have imposed the development of suitable eco-friendly means to manage plant diseases. Attempts have been made to manage the disease by means of bioagents (Vasudevan *et al.*, 2002; Manmeet and Thind, 2002; Nzojiyobiri *et al.*, 2003; Babu and Thind, 2005; Palaniyandi *et al.*, 2006; Gangwar and Sinha, 2010a,b; Gangwar and Sinha, 2012a,b and Gangwar, 2013a,b) to reduce crop losses. Application of some biotic and abiotic inducers has been reported (Van Loon, 1983; Kessman *et al.*, 1990; Kuc, 1995 and Biswas *et al.*, 2003) to induce physical and chemical defenses which may resist infection. Induced resistance in rice plants against *Drechslera oryzae* was reported with the application of various fungal fluids (Trivedi and Sinha, 1976). Induction in systemic resistance by *Pseudomonas syringae* pv. *syringae* in rice against *Pyricularia oryzae* was also reported by Smith and Matraux (1991). Kumawat *et al.*, (2008) investigated biochemical evidence of defense response in paddy and reported induction of resistance against

brown leaf spot disease of rice (*Drechslera oryzae*) with application of fungal bioagents (*T. harzianum* and *T. viride*) was due to increase elevated level of soluble proteins and total phenol content. This information implies that the higher total phenolic content for a certain period of time in the host might be associated with defense mechanism. Therefore present investigation was undertaken to observe the effect of application of fungal and bacterial bioagents on total phenolic content in rice leaves pre-inoculated with *X. oryzae* pv. *oryzae* and on disease severity of bacterial leaf blight.

MATERIALS AND METHODS

The present experiment was conducted in the glasshouse of Department of Plant Pathology, G. B. Pant University of Agriculture and Technology, Pantnagar.

Glasshouse experiment: Experiment was carried out using susceptible rice cultivar Pusa Basmati-1. Nursery was raised in field plots at Crop Research Center. Plastic pots were filled ¾ by height with natural field soil and fertilized with NPK (@ 120:60:40 kg/ha). Pots were irrigated for puddling. After puddling, 21 days old seedlings of rice cultivar Pusa Basmati-1 were transplanted in each pot and two seedlings per pot were maintained. Pots were arranged in Randomized Block Design. Regular watering was done by using

tape water. Three replications were maintained for each treatment.

Mass multiplication of fungal and bacterial bioagents and preparation of formulations: Two commercial formulation of *T. harzianum* (PBA-1) and *P. fluorescens* (PBA-2) obtained from Bio-control Laboratory, Department of Plant Pathology, G.B. Pant University of Agriculture and Technology, Pantnagar and four formulations of *P. fluorescens* (*Pf* 83, rice leaf isolate), fluorescent pseudomonad (FLP 88, rice leaf isolate), *T. harzianum* (rice leaf isolate), *Trichoderma* spp. (isolate 40, isolated from rice field soil) were evaluated for ability to induce accumulation of total phenolics in rice leaf challenged with bacterial leaf blight pathogen. *Trichoderma* spp. was mass multiplied on barnyard millet: *Echinochloa frumentacea* (local name: Jhangora) grains. Jhangora grains colonized by *Trichoderma* were air dried in open shade and ground with the help of Willy Mill to get fine powder. This powder was passed through 50 and 80 mesh sieves, simultaneously to obtain spore powder. Fluorescent pseudomonads were multiplied on King's B broth. The isolates were used to inoculate the flask containing 100 ml KB broth and incubated on incubator shaker at 150 rpm for 48 h at 25±2 °C. Bacterial suspension was then mixed directly with sterilized talc powder (@ 1.2, v/w), air dried and mixed well under sterile conditions. The formulations of fungal and bacterial bioagents were prepared by diluting bacterial cell powder or spore powder with talcum powder (mesh = 350 with 95% whiteness) and 1% carboxyl methyl cellulose (CMC) to get desired concentration (10⁶ cfu/g) of bioagents in the formulation.

Inoculation of pathogen, application of bioagent formulations: Pathogen was inoculated by clipping off the leaf tip @ 10⁶ cell/ml inoculum (Kauffman *et al.* 1973) at maximum tillering stage. All bioagent formulations (@ 10 g formulation/ liter water) and chemical treatment (0.03 g streptomycin + 1 g copper oxychloride /liter water) were applied as foliar spray one week after pathogen inoculation.

Measurement of total phenolic content in rice leaves: Phenols were estimated by the procedure described by Sadasivam and Manickam (1997).

Reagents: (1) Methanol, 80% (2) Folin-Ciocalteu reagent – commercially available reagent was diluted with distilled water in 1:1 ratio and used. (3) Sodium carbonate, 20% – prepared by dissolving 20 g sodium carbonate in 100 ml distilled water.

Extraction of phenols: 1 g leaf tissue was grinded in 5 ml 80% methanol. The extract was agitated at 70°C for 15 minutes. Now this methanolic extract was used for estimation of total phenols.

Procedure: To the 1 ml sample (methanolic extract of rice leaves), 5 ml distilled water was added to make the final volume 6 ml. To this 250 µl Folin's reagent was added and the mixture was incubated for 3 min at room temperature. After incubation, 1 ml 20% sodium carbonate and 1 ml distilled water were added and the

solution was incubated for 1 hr at room temperature. Absorbance was recorded at 725 nm. The amount of total phenols was estimated from the standard curve for tannic acid and expressed as µg phenol g⁻¹ fresh leaf weight.

Data collection: Data was recorded as total phenolic content in rice leaves, at 24 h before bioagent application and 84 and 168 h after bioagent application. Disease severity was recorded after 28 days after treatment application. After harvesting, grain yield and 1000 grain weight were recorded.

Statistical analysis: Statistical analysis of the data obtained from field experiment was done using appropriate programme as per the requirement of the experiment. The critical difference (CD) was calculated at 5% level of significance for comparison of difference between the means of different treatments.

RESULTS

Effect of bioagent application on total phenolic content in rice leaves: Total phenolic content of rice leaves varied significantly with all treatments at different time of interval (Table 1). Mean total phenolic content of rice leaves at 24 h before application of treatments was 206.66 µl/g which increased significantly at 84 h (306.66 µl/g) and 168 h (341.66 µl/g) after application of treatments. Significantly higher mean total phenolic content of rice leaves was observed with the application of bioagent formulations over check and chemical treatment. Rice leaves inoculated with pathogen exhibited significantly higher mean total phenolic content (211.66 µl/g) as compared to healthy plant (186.66 µl/g). Maximum mean total phenolic content (342.22 µl/g) in rice leaves was observed with *Pf* 83, which is followed by PBA-2 (334.44 µl/g) and *T. harzianum* (330.00 µl/g). Interaction of total phenolic content in rice leaves with different treatments and time interval was significant. Total phenolic content in rice leaves with all treatments was statistically similar at 24 h before application of treatments. Significant elevation of total phenolic content in rice leaves was observed at 84 h after application of treatments as compared to 24 h before application of treatments with all fungal and bacterial bioagents but the values obtained with chemical treatment, check and healthy plant was statistically equal. Similarly, a significant increase in total phenolic content in rice leaves was observed at 168 after application of treatments as compared to 84 h after application of treatments with all fungal and bacterial bioagents and chemical treatment however, values obtained with check and healthy plant was statistically equal. Maximum total phenolic content in rice leaves was recorded with *Pf* 83 (418.33 µl/g) which is statistically equal to *T. harzianum* (415 µl/g), PBA-2 (405 µl/g) and PBA-1 (400 µl/g) at 168 h after application of treatments. All the bioagent formulations exhibited significantly higher total phenolic content in rice

Table 1. Effect of application of bioagent formulations on total phenolic content of rice leaves pre-inoculated with *X. oryzae* pv. *oryzae* (*Xoo*), under glasshouse conditions.

Treatments	Total phenolic content ($\mu\text{l/g}$)*			Mean
	T ₁	T ₂	T ₃	
<i>Xoo</i> × FLP 88	205.00	333.33	388.33	308.88
<i>Xoo</i> × <i>Pf</i> 83	215.00	393.33	418.33	342.22
<i>Xoo</i> × PBA-2	218.33	380.00	405.00	334.44
<i>Xoo</i> × Isolate 40	211.66	346.66	386.66	315.00
<i>Xoo</i> × PBA-1	206.66	351.66	400.00	319.44
<i>Xoo</i> × <i>T. harzianum</i>	213.33	361.66	415.00	330.00
<i>Xoo</i> × Chemical Treatment	211.66	195.00	243.33	216.66
Check (<i>Xoo</i>)	210.00	210.00	215.00	211.66
Healthy plant	168.33	188.33	203.33	186.66
Mean	206.66	306.66	341.66	285.00
CD at 5%	Time	= 6.77		
	Treatments	= 11.74		
	Time × Treatments	= 20.33		

*Mean of three replications; T₁ = Total phenolic content of rice leaves; 24 h before treatment application; T₂ = Total phenolic content of rice leaves; 84 h after treatment application; T₃ = Total phenolic content of rice leaves; 168 h after treatment application.

Table 2. Effect of application of bioagent formulations on disease severity of bacterial leaf blight of rice after 28 days after treatment application, grain yield and 1000 grain weight, under glasshouse conditions.

Treatments	Disease severity (%)*	Grain yield /Plant (g)*	1000 grain weight (g)*
<i>Xoo</i> × FLP 88	42.33 (40.57)	26.79	24.92
<i>Xoo</i> × <i>Pf</i> 83	35.67 (36.66)	28.41	25.64
<i>Xoo</i> × PBA-2	39.00 (38.60)	28.08	25.48
<i>Xoo</i> × Isolate 40	42.67 (40.78)	25.64	24.04
<i>Xoo</i> × PBA-1	41.67 (40.18)	27.35	25.18
<i>Xoo</i> × <i>T. harzianum</i>	37.67 (37.86)	28.24	25.61
<i>Xoo</i> × Chemical Treatment	51.50 (45.86)	22.87	22.75
Check (<i>Xoo</i>)	94.33 (76.84)	17.02	20.18
Healthy plant	-	27.68	25.02
CD at 5%	6.72	2.45	1.41

* Mean of three replications; Figures in parenthesis are angular transformed values

leaves as compared to chemical treatment, check and healthy plant at 84 and 168 h after application of treatments.

Effect of bioagent application on disease severity, grain yield and 1000 grain weight : It is evident from table 2 that all bioagent formulations and chemical treatment exhibited significantly reduced disease severity of bacterial leaf blight of rice over check. Minimum disease severity was recorded with *Pf* 83 (35.67 %) which was statistically equal to *T. harzianum* (37.67%), PBA-2 (39.00 %), PBA-1 (41.67 %) and FLP 88 (42.33 %). All bioagent formulations, chemical treatment and healthy plant were resulted significantly higher grain yield per plant and 1000 grain weight over check. Maximum grain yield per plant was exhibited by *Pf* 83 (28.41 g) which was at par with *T. harzianum* (28.24 g), PBA-2 (28.08 g), healthy plant (27.68 g), PBA-1 (27.35 g) and FLP 88 (26.79 g). Maximum 1000 grain weight was recorded with *Pf* 83 (25.64 g) which was statistically similar to *T. harzianum* (25.61 g), PBA-2 (25.48 g), PBA-1 (25.18 g), healthy plant (25.02 g) and FLP 88

(24.92 g).

DISCUSSION

A rapid accumulation of phenols at the infection site is the first stage of defense mechanism which slows the growth of the pathogen (Matern and Kneusal, 1988). In the present study, significantly increased total phenolic content in leaves of rice plant pre-challenged with bacterial leaf blight pathogen *X. oryzae* pv. *oryzae* (210, 210 and 215 $\mu\text{l/g}$) was observed as compared to healthy plant (168.33, 188.33 and 203.33 $\mu\text{l/g}$) at 24 h before and 84 h and 168 after application of treatments, respectively. Further significant increase in total phenolic content (195 - 415 $\mu\text{l/g}$) of rice leaves was observed when *X. oryzae* pv. *oryzae* inoculated plant were treated with different fungal and bacterial bioagent formulations (Table 1). Similarly, Kumawat *et al.* (2008) reported increase level of total phenol content (3.24, 3.45 and 3.12 mg/g after 5, 10, and 15 days of inoculation, respectively) due to application of *T. harzianum* and *T. viride* in rice seedling pre-inoculating with brown leaf spot pathogen,

Drechslera oryzae and induction of resistance against disease was also reported. Minimum disease severity of bacterial leaf blight of rice was observed with the application of Pf 83, *T. harzianum*, PBA-2, PBA-1 and FLP 88 (Table 2). Decrease in disease severity of bacterial leaf blight was in accordance to increase in total phenol content (195 - 415 µl/g) in rice leaves with application of fungal and bacterial bioagents. Increase in grain yield and 1000 grain weight was also observed with increase in total phenolic content (195 - 415 µl/g) of rice leaves. Kumawat *et al.* (2008) reported that disease severity of brown leaf spot rice was negatively correlated with total phenol content due to application of *T. harzianum* and *T. viride*. Sivakumar and Sharma (2003) recorded an increase in phenolic content in maize leaf sheaths inoculated with *R. solani* or plants raised from *P. fluorescens* treated seeds. Karthikeyan *et al.* (2006) also reported induction of phenolics in coconut roots treated with biocontrol agents (*Pseudomonas fluorescens*, *Trichoderma viride* and *T. harzianum* against *Ganoderma* disease and they reported maximum level of phenolics after 9 days of treatment application.

Conclusion

The present study concluded that total phenolic content in rice leaves pre-inoculated with *X. oryzae* pv. *oryzae* was increased due to application of fungal and bacterial bioagents. Decrease in disease severity of bacterial leaf blight was observed with the increase of total phenolic content in rice leaves which resulted in increased grain yield and 1000 grain weight.

REFERENCES

- Anonymous (2002). FAO statistic India. File:///A/FAOSTAT Data Base results.com.
- Babu, A.G.C. and Thind, B.S. (2005). Potential use of combinations of *Pantoea agglomerans*, *Pseudomonas fluorescens* and *Bacillus subtilis* as biocontrol agents for the control of bacterial blight of rice. *Annals of the Sri Lanka, Department of Agriculture*, 7: 23-37.
- Biswas, S.K., Srivastava, K.D., Aggarwal, R., Shelly, P., Singh, D.V. (2003). Biochemical changes in wheat induced by *Chaetomium globosum* against spot blotch pathogen. *Indian Phytopath.*, 54: 374-379.
- Gangwar, G.P. (2013a). Efficacy of different isolates of fluorescent pseudomonads against bacterial leaf blight of rice. *Afr. J. Agric. Res.*, 8(37): 4588-4591.
- Gangwar, G.P. (2013b). Field efficacy of formulation of fungal bioagents against bacterial leaf blight of rice caused by *Xanthomonas oryzae* pv. *oryzae* (Uyeda and Ishiyama) Dowson. *J. Appl. & Nat. Sci.*, 5(2): 423-426.
- Gangwar, G.P. and Sinha, A.P. (2010a). Comparative antagonistic potential of *Trichoderma* spp. against *Xanthomonas oryzae* pv. *oryzae*. *Ann. Pl. Protec. Sci.*, 18: 458-463.
- Gangwar, G.P. and Sinha, A.P. (2010b). Evaluation of fluorescent pseudomonads against *Xanthomonas oryzae* pv. *oryzae* causing bacterial leaf blight of rice. *Ann. Pl. Protec. Sci.*, 18: 532-534.
- Gangwar, G.P. and Sinha, A.P. (2012a). Comparative antagonistic potential of fungal and bacterial bioagents against isolates of *Xanthomonas oryzae* pv. *oryzae*. *Ann. Pl. Protec. Sci.*, 20: 154-159.
- Gangwar, G.P. and Sinha, A.P. (2012b). Evaluation of *Trichoderma* spp. and fluorescent pseudomonads for the management of bacterial leaf blight of rice. *Indian Phytopath.*, 65 (1): 89-91.
- Karthikeyan, M., Radhika, K., Mathiyazhagan, S., Bhaskaran, R., Samiyappan, R. and Velazhahan, R. (2006). Induction of phenolics and defense-related enzymes in coconut (*Cocos nucifera* L.) roots treated with biocontrol agents. *Braz. J. Plant Physiol.*, 18(3): 367-377.
- Kauffman, H.E., Reddy, A.P.K., Heisk, S.P.V. and Maraca, S.D. (1973). An improved technique for evaluating resistance of rice varieties to *Xanthomonas oryzae*. *Plant Dis. Rep.*, 57: 537-541.
- Kessman, H., Edwards, R., Geno, P.W. and Dixon, R.A. (1990). Stress responses in alfalfa (*Medicago sativa* L.) constitutive and elicitor-induced accumulation of isoflavonoid conjugates in cell suspension cultures. *Plant Physiol.*, 94: 227-232.
- Kuc, J. (1995). Induced systemic resistance an overview. In: Hammerschmidt, R., Kuc, J. (ed.), *Induced Resistance to Disease in Plants*. Kluwer Academic Publishers, Dordrecht, the Netherlands. pp.182.
- Kumawat, G.L., Biswas, S.K. and Srivastava, S.S.L. (2008). Biochemical evidence of defense response in paddy induced by bio-agents against brown leaf spot pathogen. *Indian Phytopath.*, 61 (2): 197-203
- Manmeet, M. and Thind, B.S. (2002). Management of bacterial blight of rice with bioagents. *Plant Dis. Res.*, 17(1): 21-28.
- Matern, U. and Kneusal, R.E. (1988). Phenolic compounds in plant disease resistance. *Phytoparasitica*, 16:153-170.
- Mew, T.W., Alvarez, A.M., Leach, J.E. and Swings, J. (1993). Focus on bacterial blight of rice. *Plant Dis.*, 77 (1): 5-12
- Nzobjibiri, J.B., Xu, T., Song, F.M. and Shen, Y. (2003). Resistance induced by *Trichoderma harzianum* NF9 against *Magnaporthe grisea* and *Xanthomonas oryzae* pv. *oryzae* in rice. *Chinese Journal of Biological Control*, 19(3):111-114
- Palaniyandi, V., Immanuel, J.E., Gnanamanickam, S.S. and Thomashow, L. (2006). Biological control of rice bacterial blight by plant associated bacteria producing 2,4-diacetylphloroglucinol. *Canadian Journal of Microbiology*, 52(1): 56-65.
- Sivakumar, G. and Sharma, R.C. (2003). Induced biochemical changes due to seed bacterization by *Pseudomonas fluorescens* in maize plants. *Indian Phytopath.*, 56:134-137.
- Smith, J.A. and Mettraux, J.P. (1991). *Pseudomonas syringae* pv. *syringae* Induces systemic resistance to *Pyricularia oryzae* in rice. *Physiol. Mol. Plant Pathol.*, 39: 451.
- Trivedi, N. and Sinha, A.K. (1976). Resistance induced in rice plants against *Helminthosporium* infection by treatment with various fungal fluids. *J. Phytopathol.*, 86: 335-344.
- VanLoon, L.C. (1983). The induction of pathogenesis related proteins by pathogen and specific chemicals. *Neth. J. Plant Pathol.*, 89: 265-73.
- Vasudevan, P., Kavitha, S., Priyadarisini, V.B., Babujee, L. and Gnanamanicka, S.S. (2002). Biological control of rice disease. *Crop Dis.*, 34: 11-23.