



## Combined effect of biopriming and polymer coating on chemical constituents of root exudation in chilli (*Capsicum annuum* L.) cv. K 2 seedlings

S. Sathya\*, S. Lakshmi and S. Nakkeeran

Department of Seed Science and Technology, Tamil Nadu Agricultural University, Coimbatore-641001 (T.N.), INDIA  
 Department of Plant pathology, Tamil Nadu Agricultural University, Coimbatore - 641001 (T.N.), INDIA

\*Corresponding author. E-mail: sathya.agri03@gmail.com

Received: march 7, 2016; Revised received: August 16, 2016; Accepted: November 16, 2016

**Abstract:** A study was carried out to analyze the different volatile compounds in bioprimed chilli (*Capsicum annuum* L.) seedlings of 15 and 30 day old. A common compound found in two stages of chilli seedlings was hydroxylamine, dimethoxydimethyl silane, hexadecanoic acid, 15-methyl- methyl ester. Majority of the compounds in bacterized seedlings had antimicrobial activity. The results on GCMS analysis revealed that, root exudates collected from 15 and 30 days old bacterized seedlings with *B. amyloliquefaciens* VB7 and polymer coating released more number of volatile compounds (65 and 20 compounds respectively) than control (5 and 15 compounds respectively). The root exudates of 15 day old seedling released more volatile compounds (65 nos) than 30 days (20 nos) old seedling.

**Keywords:** Antimicrobial activity, *B. amyloliquefaciens* VB7, Polymer coating, Volatile compounds

### INTRODUCTION

Chilli (*Capsicum annuum*) is one of the important spice crop cultivated around the world for its pungency and colour. The pungency is due to the active principle capsaicin content in the skin and septa of the fruit. It is also used in beverages and preparation of medicines (Zagade *et al.*, 2012). Owing to the potentiality, intensive cultivation of chilli is attacked by several diseases leading to loss of yield in terms of quality and quantity. Among these diseases, damping off incited by *Pythium* spp. is responsible for 90 per cent of plant death either as pre or post-emergence in nurseries and fields (Sowmini, 1961). *Pythium* species are soil borne plant pathogenic fungi, which cause pre and post emergence damping off (Shah Smith and Burns, 1996).

Though fungicides offer a greater degree of protection against pathogens, accumulation of residues in the fruits and their adverse effect on beneficial soil microorganisms and the environment cannot be ignored. Therefore, biocontrol agents appear to hold promise in disease management. Since, biological control is a key component of integrated disease management; it is active against specific pathogens for wider application (Nakkeeran *et al.*, 2006). For effective management of any soil borne disease, the introduced antagonist should colonize the roots (Weller, 1984). The successful antagonist should colonize the rhizosphere at the time of seed germination itself and the antagonist should move from spermosphere to rhizosphere and establish (Weller and Cook, 1983).

Seedling health is determined by the root health.

Biopriming and polymer coating helps in developing a strong root system by promoting biological control of plant diseases besides improving the root system for the active acquisition of water and nutrients for better quality of seedlings (Dorlodot *et al.*, 2007). Heydecker (1973) defined seed priming as a presowing seed invigouration treatment in which seeds are soaked in osmotic solution that allows them to imbibe water and go through the first phase of germination, but does not permit radicle or plumule protrusion through seed coat. Seed treatment with biocontrol agents along with priming agents may serve as an important means of managing many of the soil and seed borne diseases, the process often known as "biopriming" (Rao *et al.*, 2007). Polymer coating is application of a thin, uniform layer of polymer over seeds without significantly increasing seed size and weight. The film formed around the seed acts as a physical barrier, which has been reported to reduce leaching of inhibitors from the seed coverings and may restrict oxygen diffusion to the embryo (Vanangamudi *et al.*, 2003).

Root exudation is a part of rhizodeposition process, which is a major source of soil organic carbon released by plant roots (Nguyen, 2003). The quantity and quality of root exudates are determined by plant species, age of an individual plant and external factors like biotic and abiotic stresses. Root exudation clearly represents a significant carbon cost to the plant with young seedlings typically exuding about 30-40 per cent of their fixed carbon as root exudates (Whipps, 1990). Root exudates contain released ions (*i.e.* H<sup>+</sup>), inorganic acids, oxygen and water, but mainly consist of carbon-based

compounds (Bais *et al.*, 2006).

Hydroponics is a technology for growing plants in nutrient solutions (water and fertilizers) with or without the use of an artificial medium. Hydroponic culture can significantly increase plant growth and produce uniform, stress-free root and shoot material that can be harvested throughout the life span of the plant (Gibeaut, 1997).

The aim of the study was to i) identify the volatile compounds released from the chilli seedling root exudates and compare the root exudates composition of bacterized and untreated seedlings ii) compare the root exudates composition of 15 and 30 days seedlings grown on hydroponic conditions

## MATERIALS AND METHODS

**Seed treatment:** Chilli seeds were surface sterilized with 80 per cent ethanol for 5 min and rinsed four times with distilled water. The seeds were bioprimed with liquid based formulation of *Bacillus amyloliquefaciens* VB7 by soaking the seeds for a period of 12 h and later the seeds were removed and immediately coated with polymer (10 ml kg<sup>-1</sup> of seed) and then shade dried at room temperature (28 ± 2°C).

**Preparation of root exudates:** Seeds bioprimed with 6 per cent *B. amyloliquefaciens* VB7 and untreated seeds were kept for germination using paper medium (between paper). The 14 days old seedlings of uniform size were transplanted into glass test tubes containing 50 ml Hoagland's nutrient solution (Hoagland and Arnon in 1950) prepared with deionized water. Root exudates were collected on 15 and 30 days. The collected liquid was filtrated through a column (20 mm diameter) containing 100 ml of XAD-4 resin, followed by elution with 50 ml methanol and condensed on rotary evaporator (Model IRA<sup>®</sup> RV 10) at 40°C. The solution, with a total volume of 25 ml, was then refrigerated at -20 °C until use.

**Identification of root exudates:** Concentrated methanol solution (5 ml) was transferred to XAD-4 resin column with 200 ml 80 per cent ether + 20 per cent acetate elution to allow the natural evaporation of methanol. The eluate was concentrated under vacuum to dryness and then dissolved in one ml of HPLC grade methanol (Qun *et al.*, 2012). The main component was used in the identification of the root exudates through gas chromatography-mass spectrometry (GC-MS, GC Agilent - 7890B, MS Agilent - 5977A MSD) analysis. One µL aliquots of the reaction mixture were injected directly into the gas chromatograph, operating under the following conditions:

The initial temperature of 80°C was kept for one min, then raised to 250°C at a rate of 8°C min<sup>-1</sup>, then raised to 300°C at a rate of 12°C min<sup>-1</sup> and held for 5 min, total GC run time was 30 min. Injector temperature was 240°C.

## RESULTS AND DISCUSSION

**Identification of volatile compounds in the root exudates of 15 days old chilli seedlings:** The compounds identified in root exudates of nontreated seedlings were shown in Fig.1 and Table 1. The major chemical constituents were hydroxylamine with peak area percentage (91.98 %), dimethoxydimethylsilane (3.47 %) and phenylephrine (2.68 %). Among the identified compounds 2.7 per cent had antimicrobial activity.

Sixty five chemical constituents (Fig.2 and Table 2) have been identified from biopriming with 6 per cent *B. amyloliquefaciens* VB7 and polymer coating @ 10 ml<sup>-1</sup> kg of seed. Among the identified compounds 62.8 per cent had antimicrobial activity. The major chemical constituents with maximum peak area percentage in bacterized seedling root exudates were identified as hydroxylamine (10.13 %), n-decanoic acid (9.24 %), 1-hexadecanol (7.99 %), Z-8-Methyl-9-tetradecenoic acid (5.99 %), cis-undec-4-enal (5.26 %), 13-Octadecenal, (Z)- (4.46 %), 13-Tetradecenal (4.35 %), 9-Octadecenal (3.35 %), Tetrapentacontane, 1,54-dibromo (2.93 %), trans-undec-4-enal (2.84 %), 1,2-Benzenedicarboxylic acid, mono (2-ethylhexyl) ester (2.06 %).

The common compounds identified in treated and nontreated 15 day seedlings root exudates were hydroxylamine, dimethoxydimethylsilane and hexadecanoic acid -15 methyl- methyl ester.

Among the fatty acids, hexadecanoic acid known to have the antibacterial, antifungal activity (Mahadkar *et al.*, 2013) antioxidant, nematocide, 5-alpha reductase inhibitor (Selvamangai and Anusha, 2012). Tetradecanoic acid is known to have potential antibacterial, antifungal activity (Mahadkar *et al.*, 2013) antioxidant and nematocide (Selvamangai and Anusha, 2012). Octadecanoic acid, pentadecanoic acid and heptadecanoic acid have potential antibacterial and antifungal activity (Mahadkar *et al.*, 2013). Another group of fatty acids with potential antifungal activity is the cyclopropane fatty acids (Carballeira, 2008). Alcohols, such as 1-hexanol have antifungal activity and prevent diseases (Archibold *et al.*, 1997).

Hydroxylamines promote seed germination by inhibition of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) decomposition by catalase (Hendricks and Taylorson, 1974). Hydroxylamine is a strong reductant and a strong chelating agent. It reacts to form oximes with aldehydes and ketones or nitrogen ethers with aldehydes, when mono-N substituted (Taylor and Baker, 1937). The marked chelating capacities of hydroxylamine for the iron atoms of haem proteins and the definite but lower capacities for N-aliphatic substituted hydroxylamines indicate the presence of this type of action in seeds.

The use of fatty acids as antifungal agents offers some advantages. Liu *et al.* (2008) proposed that antifungal fatty acids can replace chemicals in use to control plant diseases worldwide, which negatively impact the envi-





Fig. 1. GC-MS chromatogram for untreated 15 days old chilli seedling root exudates.

Table 1. Volatile compounds identified from root exudates of untreated chilli seedling 15 days after sowing.

Peak No	Retention time (min)	Peak area (%)	Name of compound	Nature of compound	Structure	Activity of compound
1	1.695	91.98	Hydroxylamine	Amine		Antioxidants Promote-seed germination
2	1.979	3.47	Dimethoxydimethylsilane	Ether		Precursor-silicone polymer polydimethylsilox-
3	17.467	1.02	Hexadecanoic acid, 15-methyl, methyl ester	Ester		Antioxidant, nematocide, 5-alpha reductase inhibitor
4	19.574	0.86	Propanamide, 2-methyl	Amide		Root growth modulation
5	29.428	2.68	Phenylephrine	Phenethylamines		Antibacterial



**Fig. 2.** GC-MS chromatogram for 6 % *Bacillus amyloliquefaciens* VB7 and polymer coating treated 15 days old chilli seedling root exudates.


ronment by affecting non-target organisms. The fungal membrane has the fundamental role of maintaining cell order and integrity and hence antifungal treatment mostly target the fungal membrane (Avis, 2007). Avis and Belanger (2001) determined the general mechanism which antifungal fatty acids directly interacts with the fungal cell membrane. The antifungal fatty acids naturally insert themselves into the lipid bi-layer of the fungal membranes and physically disturb the membrane, resulting in increased fluidity of the membrane. These elevations in membrane fluidity will cause a generalized disorganization of the cell membrane that leads to conformational changes in membrane proteins, the release of intracellular components, cytoplasmic disorder and eventually cell disintegration.

**Identification of volatile compounds in the root exudates of 30 days old chilli seedlings:** The compounds identified in root exudates of nontreated seedlings was

shown in Fig. 3 and Table 3. Among the identified compounds 1.3 per cent had antimicrobial activity. The major chemical constituents were hydroxylamine with peak area (95.35 %) and dimethoxydimethylsilane (1.59 %).

Twenty chemical constituents (Fig.4 and Table 4) have been identified from bioprimering with 6 per cent *B. amyloliquefaciens* VB7 and polymer coating. Among the identified compounds 8.8 per cent had antimicrobial activity. The major chemical constituents with maximum peak area percentage in bacterized were identified as hydroxylamine (84.04 %), hexadecanoic acid-15 methyl- methyl ester (4.84 %), n-decanoic acid (1.83 %), acetamide, 2,2,2-trifluoro (2.02 %).

The common compounds identified in treated and non-treated 30 day seedlings root exudates were hydroxylamine, dimethoxydimethylsilane, 2-heptanamine-5-




**Fig. 3.** GC-MS chromatogram for untreated 30 days old chilli seedling root exudates.

methyl, acetamide, 2,2,2-trifluoro, methylpent-4-enylamine, hexadecanoic acid-15 methyl-methyl ester, phenylephrine, cyclobutanol, benzeneethanamine-4-methoxy-alpha-methyl.

In this study, root exudates of 15 days old chilli seedling released more volatile compounds than the 30 days old chilli seedling. This result was closely agreeable with the reports of Rovira (1956). More amino acids and sugars were exuded during the first 10 days of growth than during the second 10 days in peas and oats. Vancura and Hovadik (1965) found that 3-pyrazolylalanine was present in the root exudates of cucumber only at the early stage.

The root exudations of volatile compounds are greatly influenced by root age of seedlings (Shukla *et al.*, 2011). Microorganisms may affect the permeability of root cells, metabolism of roots, absorption and excretion of certain compounds in root exudates. It was reported that filtrates of cultures of some bacteria and fungi and also some antibiotics, increased the exudation by oat roots (Blaylock *et al.*, 1997). Some other plant biotic factors like developmental status, shoot herbivory, photosynthesis, supply of carbon from shoot to root, evaporation, transpiration, nutrient deficiency, root architecture, cytosolic concentration, membrane permeability, membrane electrochemical potential,

















**Fig. 4.** GC-MS chromatogram for 6 % *Bacillus amyloliquefaciens* VB7 and polymer coating treated 30 days old seedling root exudates.


















release of microbial signal, allelochemical release, nodulation and some soil biotic factors are also influenced by the root exudation (Shukla *et al.*, 2011). Our results revealed that volatiles can have an effect on secondary metabolites production by *B. amyloliquefaciens*. When exposed to volatiles emitted by *Colimonas pratensis*, *Pseudomonas fluorescens* produced secondary metabolites that had inhibiting activity against a Gram positive bacterium and a fungus but not

against the Gram negative volatile producer. It is plausible that the volatiles served as energy sources and/or signal inducing secondary metabolite production. The volatile triggered antibiotic production in *P. fluorescens* could point a strategy to combine movement (chemotaxis and motility genes) with increasing competitive strength (antibiotics) to invade in to the nutrient providing rhizosphere zone. It is known that bacterial volatiles can have antimicrobial activity and inhibit

**Table 2.** Volatile compounds identified from root exudates of chilli seedling bioprimered with 6 % *Bacillus amyloliquefaciens* and polymer coating after 15 days sowing.

















Peak No	Retention time(min)	Peak area (%)	Name of compound	Nature of compound	Structure	Activity of compound
1	1.693	10.13	Hydroxylamine	Amine		Antioxidants Promote-seed germination
2	1.982	0.35	Dimethoxydimethyl silane	Ether		Precursor-silicone polymer polydimethylsiloxane
3	12.393	0.14	Hexadecanoic acid, 15-methyl-, methyl ester	Ester		Antioxidant, nematocide, 5-alpha reductase inhibitor
4	14.223	0.42	Imidodicarbonic diamide, N-formyl	Amide		Antimicrobial
5	14.787	0.15	Methyl 8-methyl-decanoate	Ester		Antifungal
6	15.560	0.20	Methyl 8-methyl-nonanoate	Ester		Antifungal
7	16.432	1.67	Xylose	Sugar		Antibacterial, antifungal, Precursor-synthetic polymers
8	16.537	1.02	Cyclopropane, nonyl	Cycloalkane		Antifungal- <i>Pythium</i> spp.
9	17.449	1.07	Decanoic acid, methyl ester	Ester		Antifungal
10	17.532	0.21	11,14-Eicosadienoic acid, methyl ester	Ester		Antibacterial
11	17.778	5.26	cis-Undec-4-enal	Aldehyde		Antimicrobial
12	17.915	9.24	n-Decanoic acid	Fatty acid		Antifungal
13	18.234	0.94	1-Eicosanol	Fatty alcohol		Antifungal, Antioxidant
14	18.415	0.42	1,15-Pentadecanediol	Fatty alcohol		Antifungal, antibacterial




















Contd.....

Contd.....						
15	18.575	2.84	trans-Undec-4-enal	Aldehyde		Antimicrobial
16	18.732	0.61	cis-11-Hexadecenal	Aldehyde		Antimicrobial
17	18.954	0.31	1,8-Nonadiene, 2,8-dimethyl-	Alkene		Unknown
18	19.359	5.99	Z-8-Methyl-9-tetradecenoic acid	Fatty acid		Antifungal
19	19.498	0.61	9-Oxabicyclo[6.1.0]nonane	EpoxyCycloalkane		Antibacterial, Antifungal and nematocidal activity
20	19.564	0.63	1,6-Octadiene, 5,7-dimethyl-, (R)-	Alkene		Antioxidant
21	20.021	7.99	1-Hexadecanol	Fatty alcohol		Antibacterial
22	20.228	1.70	tert-Hexadecanethiol	Thiol		Antioxidant, insecticidal, Antifungal
23	21.232	2.06	1,2-Benzenedicarboxylic acid, mono (2-ethylhexyl) ester	Aromatics		Insecticidal activity
24	21.461	0.22	Amphetamine, N-propoxycarbonyl-	Phenethylamines		Antibacterial
25	21.683	0.19	Acetamide, 2-amino-	Amide		Antifungal
26	21.868	0.91	Methyl 11-oxo-9-undecenoate	Ester		Plant growth regulator, insect attractant
27	22.031	0.75	Hexanoic acid, dodecyl ester	Ester		Root growth
28	22.243	1.14	1-Octadecanesulphonyl chloride	Organo sulphonyl halide		Antimicrobial
29	22.366	1.04	(2,2,6-Trimethylbicyclo[4.1.0]hept-1-yl)-methanol	Cyclic alcohol		Unknown
30	22.533	1.48	Heneicosane	Alkane		Pheromone, Antifungal, Antibacterial
31	22.670	0.92	1-Dodecene	Alkene		Inhibitor of ethylene


















Contd.....












32	22.714	0.65	4-Heptafluorobutyryloxyhexadecane	Halogenated alkane		Antibacterial
33	22.813	1.28	Heptane, 3-ethyl-2-methyl	Alkane		To induce shortening and thickening of the stem in cereals and other crop plants increasing the yields or setting of fruit
34	22.955	0.88	3-[N-Aziridyl]propionyl hydrazide	Hydrazide		Antimicrobial
35	23.097	1.63	Cyclodecanone	Alicyclic Ketone		Insecticidal Fungicidal
36	23.234	0.56	2-Piperidinone, 6-methyl	Cyclic amide		Antifungal
37	23.315	0.85	N-Acryloylsarcosine methyl ester	Ester		Unknown
38	23.432	1.82	Decane, 1-fluoro-	Halogenated alkane		Intra and inter-plant communication Attraction or repulsion of parasites
39	23.567	0.90	Octadecane, 1-chloro-	Halogenated alkane		Antifungal
40	23.790	0.21	Ethanamine, 2-phenoxy	Amine		Root growth and development
41	23.843	0.35	Cyclooctene, 1,2-dimethyl-	Cycloalkene		Insecticidal (shoot fly resistance)
42	23.997	0.41	3-Trifluoroacetoxy-pentadecane	Halogenated alkane		Antimicrobial
43	24.117	1.42	8-Hexadecenal, 14-methyl-, (Z)-	Aldehyde		Pesticide-Pheromone Khapra beetle (Ballion Warehouse beetle)
44	24.312	3.35	9-Octadecenal	Fatty aldehyde		Antifungal
45	24.411	1.91	18-Nonadecen-1-ol	Fatty alcohol		Antibacterial
46	24.531	4.35	13-Tetradecenal	Aldehyde		Antimicrobial, Insecticidal
47	24.595	2.93	Tetrapentacontane, 1,54-dibromo-	Halogenated alkane		Antimicrobial and antifungal

Contd.....						
47	24.595	2.93	Tetrapentacontane, 1,54-dibromo-	Halogenated alkane		Antimicrobial and antifungal
48	24.707	4.46	13-Octadecenal, (Z)-	Aldehyde		Pesticide-Pheromone
49	25.063	0.34	Acetic acid, trichloro-, nonyl ester	Ester		Pesticide
50	25.152	0.50	1-Hexacosanol	Fatty alcohol		Antifungal
51	25.223	0.62	Hexadecane, 1-bromo-	Halogenated alkane		Antifungal
52	25.485	0.26	1-Bromodocosane	Halogenated alkane		Antifungal
53	25.606	0.17	Propanenitrile, 3-amino-2,3-di(hydroxyimino)-	Aliphatic nitrile		Unknown
54	25.916	0.52	Heptanal	Aldehyde		Lipid oxidation
55	26.050	0.20	Undecanal	Aldehyde		Antibacterial
56	26.145	0.38	Piperazine, 2-methyl-	Piperazine		Antifungal antibacterial and Plant growth regulation
57	26.321	0.60	7-Oxabicyclo[4.1.0]heptane	Epoxy cyclo alkane		Antibacterial Antifungal Nematicidal
58	26.401	0.09	Oxalic acid, allyl hexadecyl ester	Fatty acid		Antimicrobial
59	26.567	0.49	3-Hydroxy-N,N-dimethylpropanamide	Amide		Unknown
60	26.982	0.57	Urea, (hexahydro-6-methyl-2-oxo-4-pyrimidinyl)-	Amide		Plant growth regulation
61	27.344	0.50	Bicyclo[2.1.1]hexane-1-carboxylic acid, 5,5-dimethyl-	Carboxylic acid		Antifungal
62	27.596	0.50	Sarcosine, N-valeryl-, ethyl ester	Ester		Increase plant biomass
63	27.967	0.38	7-Oxabicyclo[4.1.0]heptane, 1-methyl-4-(2-methyloxiranyl)	Epoxy cyclo alkane		Antibacterial Antifungal Nematicidal
64	28.223	1.78	2,6,6-Trimethylbicyclo[3.1.1]hept-3-ylamine	Bicyclo amine		Insecticidal
65	29.505	0.26	Benzenemethanol, .alpha. [(methylamino) methyl]-	Aromatic		Unknown










**Table 3.** Volatile compounds identified from root exudates of untreated chilli seedling 30 days after sowing.

Peak No	Retention time (min)	Peak area (%)	Name of compound	Nature of compound	Structure	Activity of compound
1	1.689	95.35	Hydroxylamine	Amine		Antioxidants Promote-seed germination
2	1.993	1.59	Dimethoxydimethylsilane	Ether		Precursor-silicone polymerpolydimethylsiloxane
3	6.849	0.10	Acetic acid, [(aminocarbonyl)amino]oxo	Acid		Disease resistance
4	11.154	0.38	2-Heptanamine, 5-methyl	Amine		Antimicrobial
5	11.241	0.56	1-Hexadecanol	<u>Fatty alcohol</u>		Antibacterial
6	14.420	0.20	Acetamide, 2,2-dichloro	Amide		Root development
7	14.518	0.34	Benzeneethanamine, N-methyl	Amine		Plant resistance, Antioxidant
8	15.271	0.16	Methylpent-4-enylamine	Amine		Antibacterial
9	17.468	0.40	Hexadecanoic acid, 15-methyl-, methyl ester	Ester		Antioxidant, nematocide, 5-alpha reductase inhibitor
10	17.842	0.16	Phenylephrine	Phenethylamines		Antibacterial
11	19.569	0.12	Cyclobutanol	Alcohol		Plant resistance, Antioxidant
12	19.949	0.21	1-Methyl-2-phenoxyethylamine	Amine		Plant growth regulation
13	23.311	0.08	2-Thiophenecarboxylic acid, 5-(1,1-dimethylethoxy)-	Organosulfur		Plant growth regulation
14	24.059	0.11	N-Methyl-2-phenyl-1-propanamine	Amine		Plant resistance, Antioxidant
15	24.758	0.23	Benzeneethanamine, 4-methoxy-alpha-methyl	Amine		Plant resistance, Antioxidant

**Table 4.** Volatile compounds identified from root exudates of chilli seedling bioprimered with 6 % *Bacillus amyloliquefaciens* and polymer coating after 30 days sowing.

Peak No	Retention time (min)	Peak area (%)	Name of compound	Nature of compound	Structure	Activity of compound
1	1.693	84.04	Hydroxylamine	Amine		Antioxidants Promote-seed germination
2	1.955	0.37	1,2,3,4-Butanetetrol, [S-(R*,R*)]	Alcohol (Sugar alcohol)		Antimicrobial, Plant growth regulation
3	1.994	0.78	Dimethoxydimethylsilane	Ether		Precursor-silicone polymer polydimethylsiloxane
4	6.177	0.56	2-Heptanamine, 5-methyl	Amine		Antimicrobial
5	6.225	0.42	Cyclohexan-1,4,5-triol-3-one-1-carboxylic acid	Ester		Antibacterial
6	17.098	0.62	1,3-Dioxolane-4-methanol	Hemiacetal		Root growth
7	17.357	0.78	1-Pentanol, 4-amino	Amino alcohol		Root growth, Stress tolerance
8	17.463	4.84	Hexadecanoic acid, 15-methyl, methyl ester	Ester		Antioxidant, nematocide, 5-alpha reductase inhibitor
9	17.879	1.83	n-Decanoic acid	Fatty acid		Antifungal
10	19.514	0.18	Cyclobutanol	Alcohol		Plant resistance, Antioxidant
11	19.568	1.02	p-Hydroxynorephedrine	Phenethylamines		Antibacterial

Contd.....

12	19.711	0.70	3-Azabicyclo [3.2.2]nonane	Amine (Tropane alkaloids)		Antibacterial
13	19.910	0.26	4-Iodo-3-methoxyamphetamine	Phenethylamine		Antibacterial
14	19.982	2.02	Acetamide, 2,2,2-trifluoro	Amide		Antifungal
15	20.203	0.41	Phenylephrine	Phenethylamines		Antibacterial
16	22.501	0.18	Methylpent-4-enylamine	Amine		Antibacterial
17	23.314	0.31	Benzeneethanamine, 4-methoxy-alpha-methyl	Phenethylamine		Antibacterial
18	23.423	0.21	Amphetamine-3-methyl	Phenethylamines		Antibacterial
19	24.060	0.26	Propanamide	Amide		Antifungal
20	24.759	0.21	Metaraminol	Amine		Antibacterial

the growth of other microorganisms (Kai *et al.*, 2007, 2009; Garbeva *et al.*, 2014a, 2014b).

### Conclusion

It was concluded that, the common compound identified in both 15 and 30 day old chilli seedling were, hydroxylamine, dimethoxydimethyl silane and hexadecanoic acid -15-methyl- methyl ester. The results on GCMS analysis revealed that root exudates collected from 15 and 30 day old bacterized seedlings with *B. amyloliquefaciens* VB7 and polymer coating released more number of volatile compounds than control. Between 15 and 30 day old seedlings, bioprimering with *B. amyloliquefaciens* VB7 and polymer coated 15 day old seedling root exudates released more volatile compounds than 30 day old seedling. Majority of the compounds in bacterized seedlings had antimicrobial activity. Those compounds indicate their potential use for

various diseases in traditional system.

### REFERENCES

- Archibold, D.D., Hamilton-Kemp, T.R., Barth, M.M. and Langlois, B.E. (1997). Identifying natural volatile compounds that control gray mold (*Botrytis cinerea*) during postharvest storage of strawberry, blackberry and grape. *J. Agricultural and Food Chemistry*, 45: 4032-4037
- Avis, T.J. (2007). Antifungal compounds that target fungal membrane: Application in plant disease control. *Can. J. Pl. Path.*, 29: 323-329
- Avis, T.J. and Bélanger, R.R. (2001). Specificity and mode of action of the antifungal fatty acid cis-9-heptadecenoic acid produced by *Pseudozyma flocculosa*. *Appl. Environ. Microbiol.*, 67: 956-960
- Bais, H.P., Weir, T.L., Perry, L.G., Gilroy, S. and Vivanco, J.M. (2006). The role of root exudates in rhizosphere interactions with plants and other organisms. *Annu. Rev. Pl. Biol.*, 57: 233-266
- Blaylock, M.S.D.E., Dushenkiv, S., Zakharavo, O., Gussman,

- C., Kapulnik, Y., Ensley, B. and Raskin, E. (1997). Enhanced accumulation of Pb in Indian mustard by soil-applied chelating agents. *Environ. Sci. Technol.*, 31: 860-865
- Carballeira, N.M. (2008). New advances in fatty acids as antimalarial, antimycobacterial and antifungal agents - A review. *Progress in Lipid Research*, 47:50-61
- Dorlodot, S., Forster, B., Pages, L., Price, A., Tuberosa, R. and Draye, X. (2007). Root system architecture: Opportunities and constraints for genetic improvement of crops. *Trends Plant Sci.*, 12: 474-481
- Garbeva, P., Hordijk, C., Gerards, S. and DeBoer, W. (2014a). Volatile-mediated interactions between phylogenetically different soil bacteria. *Frontiers in Microbiology*, 5: 1-9
- Garbeva, P., Hordijk, C., Gerards, S. and DeBoer, W. (2014b). Volatiles produced by the mycophagous soil bacterium *Collimonas*. *FEMS Microbiol. Ecol.*, 87: 639-649
- Gibeaut, D.M., Hulett, J., Cramer, G.R. and Seemann, J.R. (1997). Maximal biomass of *Arabidopsis thaliana* using a simple, low maintenance hydroponic method and favourable environmental conditions. *Plant Physiol.*, 115: 317-319
- Hendricks, S.B. and Taylorson, R.B. (1974). Promotion of seed germination by nitrate, nitrite, hydroxylamine, and ammonium salts. *Plant physiol.*, 54: 304-309
- Heydecker, W. (1973). Germination of an idea: The priming of seeds. *University of Nottingham School of Agriculture Rep.*, 74
- Hoagland, D.R. and Arnon, D.I. (1950). The water culture method for growing plants without soil. California Agricultural Experimental Station Circular No. 347, pp. 1-32. University of California, Berkeley.
- Kai, M., Effmert, U., Berg, G. and Piechulla, B. (2007). Volatiles of bacterial antagonists inhibit mycelial growth of the plant pathogen *Rhizoctonia solani*. *Arch. Microbiol.*, 187: 351-360
- Kai, M., Hausteiner, M., Molina, F., Petri, A., Scholz, B. and Piechulla, B. (2009). Bacterial volatiles and their action potential. *Appl. Microbiol. Biotechnol.*, 81: 1001-1012
- Liu, S., Weibin, R., Jing, L., Hua, X., Jingan, W., Yubao, G. and Jingguo, W. (2008). Biological control of phytopathogenic fungi by fatty acids. *Mycopathologia*, 66: 93-102
- Mahadkar, S., Valvi, S. and Jadhav, V. (2013). Gas chromatography mass spectroscopic (GCMS) analysis of some bioactive compounds from five medicinally relevant wild edible plants. *Asian J. Pharm. Clin. Res.*, 6(1): 136-139
- Nakkeeran, S., Kavitha, K., Chandrasekar, G., Renukadevi, P. and Fernando, W.G.D. (2006). Induction of plant defence compounds by *Pseudomonas chlororaphis* PA23 and *Bacillus subtilis* BSCBE4 in controlling damping-off of hot pepper caused by *Pythium aphanidermatum*. *Biocontrol Sci. and Technol.*, 16(4): 403-416
- Nguyen, C. (2003). Rhizodeposition of organic C by plants: Mechanisms and controls. *Agronomy*, 23: 375-396
- Qun, H. Z., Junan, Z., HaoRu, T. and Zhi, H. (2012). Different vegetables crops in response to allelopathic of hot pepper root exudates. *World Appl. Sci. J.*, 19 (9): 1289-1294
- Rao, M.S.L., Kulkarni, S., Sagar, S.D. and Kulkarni, V.R. (2007). Biopriming induced changes in the activity of defence related enzymes for conferring resistance against *Alternaria* blight of sunflower. *J. Pl. Dis. Sci.*, 2 (1): 14-17
- Rovira, A.D. (1956). A study of the development of the root surface microflora during the initial stages. *J. Appl. Bacteriol.*, 19:72-79
- Selvamangai, G. and Anusha, B. (2012). GC-MS analysis of phytochemicals in the methanolic extract of *Eupatorium triplinerve*. *Asian Pacific Journal of Tropical Biomedicine*, 329-332
- Shah Smith, D.A. and Burns, R.G. (1996). Biological control of damping off of sugar beet by *Pseudomonas putida* applied to seed pellets. *Plant pathol.*, 45: 572-582
- Shukla, K.V., Sharma, S., Singh, N.K., Singh, V., Tiwari, K. and Singh, S. (2011). Nature and role of root exudates: Efficacy in bioremediation. *African J. Biotechnol.*, 10 (48): 9717-9724
- Sowmini, R. (1961). Studies on *Phycomycetes* in agricultural soils with special reference to Pythiaceae. *M.Sc. (Agri.) Thesis*, University of Madras: 160
- Taylor, T.W.J. and Baker, W. (1937). Sidgwick's organic chemistry of nitrogen. *Oxford University Press, New York*. pp. 166-169
- Vanangamudi, K., Srimathi, P., Natarajan, N. and Bhaskaran, M. (2003). Current scenario of seed coating polymer. *In: Proc. of ICAR short course on seed hardening and pelleting technologies for rainfed/ garden land ecosystems*, New Delhi, pp. 80-100.
- Vancura, V. and Hovadik, A. (1965). Composition of root exudates in the course of plant development. *Plant Microb. Relat.*, pp. 21-25
- Weller, D.M. (1984). Distribution of a take-all suppressive strain of *Pseudomonas fluorescens* on seminal roots of winter wheat. *Appl. Environ. Microbiol.*, 48(4): 897-899
- Weller, D.M. and Cook, R.J. (1983). Suppression of take-all of wheat by seed treatment with *Pseudomonas fluorescens*. *Phytopathol.*, 73: 463-469
- Whipps J.M. (1990). Carbon economy. *In: The rhizosphere* (ed. J.M. Lynch), pp. 59-97. JohnWiley & Sons Ltd, Essex, UK.
- Zagade, S.N., Deshpande, G.D., Gawade, D.B., Atnoorkar, A.A. and Pawar, S.V. (2012). Biocontrol agents and fungicides for management of damping off in chilli. *World J. Agric. Sci.*, 8 (6): 590-597