



Studies of alternaria black spot disease of pomegranate caused by *Alternaria alternata* in Punjab

Adesh Kumar,^{*} Tanjeet Singh Chahal, Mandeep Singh Hunjan¹, Harminder Kaur¹ and Roomi Rawal²

Punjab Agricultural University, Fruit Research Station, Jallowal- Lesriwal, Jalandhar-144303 (Punjab), INDIA

¹Department of Plant Pathology, Punjab Agricultural University, Ludhiana-141004 (Punjab), INDIA

²Department of Entomology, Haryana Agricultural University, Hisar-125001 (Haryana), INDIA

^{*}Corresponding author. E-mail: adeshfrs@gmail.com

Received: May 15, 2016; Revised received: September 26, 2016; Accepted: January 15, 2017

Abstract: *Alternaria* black spot of pomegranate caused by *Alternaria alternata* pose significant economic losses in India as it reduce the crop yield. Farm survey was undertaken at Punjab Agricultural University, Pomegranate Research Block, Fruit Research Station, Jallowal- Lesriwal Jalandhar during 2015. Among all the five cultivars highest disease incidence (70%) and severity (30%) was registered in cultivar Bhagwa. Twenty two isolates of *A. alternata* were recovered from infected fruits and clustered using UPGMA (unweighted pair group method with arithmetic averages) on the basis of disease score produced in five cultivars (Ganesh, Ruby, Bhagwa, Jyoti and Mridula). It was revealed that five isolates namely AL14, AL15, AL20, AL21 and AL22 were highly virulent on almost all the pomegranate cultivars. Two cultivars viz. Bhagwa and Mridula were found to be most susceptible as 45.45 % isolates were found to be highly virulent on them. The *in vitro* antifungal effects of the six fungicides on mycelial growth were investigated. Based on the inhibition of mycelial growth, all the fungicides showed most toxic reaction with 50% effective concentrations (EC₅₀) of < 1 ppm. However, they varied in EC₉₀ values. 10 ppm concentration of Tilt (azole group) showed 90% effective concentration (EC₉₀). Folicur also showed approximately same results as Tilt while EC₉₀ value of Natio and SAAF was <25 ppm and > 20 ppm. However, two fungicides namely Dithane Z-78 and Bavistin showed less toxicity against *Alternaria alternata* as compared to other with EC₉₀ value at concentrations <50 ppm and > 25 ppm. Azole group fungicides (Tilt and Folicur) were found most effective to inhibit the pathogen growth.

Keywords: *Alternaria* black spot, *Alternaria alternata*, Cultivars, Fungicides and Pomegranate

INTRODUCTION

Pomegranate (*Punica granatum* L.) a high value fruit crop that cultivated worldwide with dominancy in Mediterranean countries like Spain, Morocco, Egypt, Iran, Afghanistan and Baluchistan since ancient times. Tropical and sub-tropical regions of the Indian sub-continent are the prominent areas under cultivation of this fruit crop. India plays a leading role in pomegranate production with contribution of nearly 50% of global pool. The total area under pomegranate is 132 thousand hectare with annual production of 1357 thousand MT in 2014-2015 (Anonymous, 2015).

A. alternata is an important plant pathogen that causes black spot disease with symptoms present on leaves, flowers, and young fruit of pomegranate (Pala *et al.*, 2009; Holland *et al.*, 2009). This disease is a major hurdle in quality production and high yield in the crop causing up to 15-80% losses in Karnataka including Bijapur, Bagalkot and Koppal districts (Archana and Jamadar, 2014).

A. alternata responsible for fruit rot was first reported in India by Madhukar and Reddy (1976) and subse-

quently from USA and Mexico (Farr *et al.*, 2007). In currently *Alternaria* black spot disease is reported in different part of the world such as Greece (Tziros *et al.* 2007), Israel (Ezra *et al.*, 2010) and Spain (Bergal *et al.*, 2014). This pathogen belongs to omnipresent necrotrophic fungi group and have at least seven pathogenic variants of this fungus, each producing unique host-selective toxins (HSTs) and causing disease on specific host plants (Ito *et al.*, 2004). In Punjab, this disease invariably appears every year in the pomegranate orchards causing significant yield and quality loss. For management of fungal plant pathogens, fungicides have important role to control plant disease. Fungicides inhibit the growth of the fungi under *in vitro* and *in vivo* condition (Archana and Jamadar, 2014; Nel *et al.*, 2007). They have different mode of action to control the fungal growth. Mesta *et al.* (2011) used sterol demethylation inhibitors fungicides namely propiconazole and hexaconazole with conc 0.1% to control *Alternaria* blight of Sunflower. Azoxystrobin fungicides have potential to inhibit mitochondrial respiration and blocking the cytochrome bc1 complex can be used to control *Alternaria solani* pathogen causing

Table 1. Disease severity response of *Alternaria* black spot disease on pomegranate (*Punica granatum* L.).

Response	Per cent Infection on fruit pericarp
0	0.00
1	Up to 1
2	> 1-10
3	> 10-20
4	>20-40
5	> 50

agent of potato early blight (Pasche *et al.*, 2004). Realizing the importance of this disease, the present studies were initiated to isolate, identify and characterize the pathogen with respect to its aggressiveness and also evaluate different fungicides against *A. alternata* under *in vitro* condition.

MATERIALS AND METHODS

Isolation and identification of the pathogen: Symptomatic fruits were collected from Punjab Agricultural University, Pomegranate Research Block, Fruit Research Station, Jallowal- Lesriwal Jalandhar, Punjab in the month of June, 2015. Disease incidence was recorded by counting the number of infected fruits per plant. For disease severity, 20 fruits were selected randomly and scoring was done with help of 1-5 scale. After surface sterilization, the peel pieces were placed on potato dextrose agar separately and grown at 25 °C for 2-3 days. The growing fungal hyphal tips were transferred to PDA and grown for 4-5 days, and cultures were further purified by single spore isolation method. The morpho-logical characteristics of the conidia were analyzed using Leica microscope (DM 2000) at 100X. Twenty two isolates of the *Alternaria* black spot pathogen were recovered and used for this study.

Pathological characterization and aggressiveness: All the twenty two isolates were grown individually on PDA medium and PDA-plugs, 5 mm in diameter, with actively grown mycelium were transferred on fruit wounds made by a scalpel on previously sterilized fruit surfaces. Full sized pomegranate fruits of all cultivars *viz.* Ganaesh, Ruby, Jyoti, Mridula and Bhagwa were inoculated with individual isolate (Tziros *et al.*, 2007). After inoculation fruits were covered with plastic bags for 7- 10 days and moisture was maintained by hand sprayer (one spray daily up to symptom development) in plastic bags. The fruits inoculated with PDA medium only served as control. After inoculation, the plants




Fig. 1. Conidia of *A. alternata* with longitudinal and transverse septa (a-AL15, b-AL20 and c-AL21).




Fig. 2. Clustering of different isolates of *A. alternata* on the basis of virulence character.

were frequently irrigated to keep up high humidity and soil moisture which is vital for disease development. The disease symptoms were observed after 48-72 hrs of inoculation and final observations were recorded after 10 days. Disease severity was recorded on the basis of pathogen response and was divided into five categories (Table-1). The pathogen was re-isolated from the infected inoculated fruits and reconfirmed as *A. alternata*.

In vitro efficacy of different fungicides against A. alternata: Fungicides (Table 2) were evaluated to determine their 50% effective concentration (EC50) and EC90 values for the inhibition of mycelial

Table 2. Fungicides used in this study to control the growth of *A. alternata*.

Fungicides	Chemical group	Active ingredient	Concentration (%)	Formulation
Folicur	Azole	Tebuconazole	25.9	EC
Tilt	Azole	Propiconazole	25	EC
Natio	Mixture of Azole and Strobilurin	Tebuconazole+ Trifloxystrobin	50+25	WG
SAAF	Mixture of benzimidazole and Carbamate	Carbendazim+ Mancozeb	12+63	WP
Bavistin	benzimidazole	Carbendazim	50	WP
Dithane Z-78	Carbamate	zineb	70	WP

Table 3. Disease incidence (DI) and disease severity (DS) of Alternaria black spot disease of pomegranate (*Punica granatum* L.).

Cultivars	DI%	DS%
Ruby	10	3
Ganesh	50	25
Jyoti	56	27
Bhagwa	70	30
Mridula	40	14

growth. To analyze the inhibition of mycelial growth of *A. alternata*, five replicate PDA plates (90 mm in diameter) containing the fungicides were prepared at concentrations of 1, 10, 20, 25 and 50ppm. The control plates contained only PDA medium. Individual agar disks (6 mm in diameter) were removed from the edge of an actively growing culture (AL 22) and placed at PDA plates that containing fungicides and incubated at 25°C for 4-5 days. The mycelium growth of pathogen was measured by scale and compared with growth observed in control. The concentration of each fungicide causing 50% (EC₅₀) or 90% (EC₉₀) reduction in mycelial growth compared to the absence of the fungicide was estimated referring to Matheron and Porchas (2000) and based on the estimated values.

Statistical Analysis: The pathotypic similarity between the various isolates was generated using unweighted paired group mean averages using software programme PAST ver. 2.1.5.

RESULTS AND DISCUSSION

The Alternaria black spot disease incidence on cultivar Bhagwa, a popular variety was observed to be very high. At Fruit Research Station Jallowal, 70% of the plants showed symptoms of this disease with mean disease severity of 30 % (Table-3). It was followed by Jyoti and Ganesh varieties where disease incidence was observed to be 56% and 50 % respectively. In contrast disease incidence and severity of Alternaria




Fig. 3. Virulence frequency of twenty two isolates of *A. alternata* on five pomegranate cultivars.

black spot was low on Ruby and Mridula varieties (Table-3).

Isolation and identification of the pathogen: A total of twenty two isolates were recovered from infected fruit samples. All the isolates showed typical characters of *A. alternata* with Gray-to-black colonies. The pathogen conidiophores were short, septate, branched or unbranched, green to brown, obpyriform with conical or cylindrical beak (Simmons, 1967) (Fig. 1).

Pathological characterization and aggressiveness: Different response with respect to disease aggressiveness (number of disease spot on fruit) was observed within different isolates. Symptoms appeared 48–72 h later, followed by spreading on all fruit pericarp. The data presented in Fig. 2 shows the presence of significantly diverse virulence spectrum in *A. alternata* populations of Punjab. Disease response with respect to twenty two isolates of *A. alternata* clustered using UP-GMA (Fig. 2). The data plotted in dendrogram (Fig.1) showed five isolates namely AL14, AL15, AL20, AL21 and AL22 to be clustered together and were classified in group I. These isolates were found to be highly virulent on almost all the cultivars tested producing the maximum mean disease score 4.92 (Table 4).

Table 4. Grouping of twenty two isolates of *A. alternata* on the basis of virulence reaction on five pomegranate cultivars.

Group	Name of Isolates	No. of Isolates	Disease score		Remark
			Mean	Range	
I	AL14, AL15, AL20, AL21 and AL22	5	4.92	4.6-5	Highly Virulent
II	AL1, AL2, AL3, AL4, AL6, AL7, AL18, AL9, AL16, AL17, AL18, and AL19	12	2.85	2.6-3.2	Moderately Virulent
III	AL5, AL10, AL11, AL12 and AL13	5	2.08	1.8-2.4	Less Virulent

Table 5. Evaluation of different fungicides against *A. alternata* under *in vitro* condition.

Fungicides	Mycelial growth (mm)					
	Concentration in ppm					
	1	5	10	20	25	50
Natio	20.4 (0.23)	18.4 (0.24)	16.2 (0.53)	11.8 (0.23)	0	0
Folicur	24.8 (0.53)	17.2 (0.26)	9.4 (0.23)	8.2 (0.93)	0	0
Tilt	19.2 (0.13)	11.4 (0.22)	9 (0.12)	0	0	0
Bavistin	41.6 (0.33)	23.6 (0.42)	18.4 (0.43)	13 (0.29)	9.2 (0.53)	4.6 (0.83)
Z-78	28.4 (0.22)	19.6 (0.22)	17.2 (0.73)	15.4 (0.28)	12.8 (0.23)	8.6 (0.33)
SAAF	39 (0.63)	23.4 (0.32)	17.8 (0.53)	12.4 (0.27)	8.6 (0.25)	6.2 (0.93)

Figures in parentheses are standard deviations from mean of five replications.




Fig. 4. Inhibition % of *A. alternata* mycelium by different fungicides at different concentrations (a. 1, b. 5, c. 10, d. 20, e. 25 and f. 50ppm).

Twelve out of all the isolates showed disease score ranging from 2.6 to 3.2 with mean 2.85 and were classified as moderately virulent (Table 4). These isolates were unable to cover all the fruit with external symptoms, disease progress was observed only less than half fruit length from the point of inoculation. The lowest disease score ranged from 1.8-2.4 with mean 2.08, which was recorded in 5 isolates where the disease restricted only to point of inoculation (Table 4). Similarly, Gat *et al.* (2012) observed variability in disease severity among the isolates of *A. alternata* and classified them into four groups. Isolates that did not induce any visible symptoms were classified into virulence group. Isolates that caused mild, moderate, or severe symptoms were classified into virulence groups 1, 2, and 3 respectively. Berbegal *et al.* (2014) inoculated fruit and leaves of pomegranate with *A. alternata* and disease symptom was observed after 10 and 3 days respectively.

Virulence frequency of *A. alternata* isolates on five pomegranate cultivars: Currently meager work might be available regarding screening of pomegranate germplasm against this disease. In the present study




Fig. 5. Inhibition of *A. alternata* mycelium growth at 20ppm by Tilt (a), Folicur (b) and without chemical work as control (c).

five pomegranate cultivars were screened (Fig. 3) which showed different reaction to Punjab population of *A. alternata*. Two cultivars i.e. Mridula and Bhagwa were found to be most susceptible as 45.45 % isolates were found to be highly virulent on both. It was followed by Jyoti and Ganesh which were susceptible to 31.18 and 27.27 % isolates respectively. However, Ruby was found to be comparatively less susceptible as only 18.18% isolates showed virulent reaction to Punjab population of *A. alternata*.

***In vitro* efficacy of different fungicides against *A. alternata*:**

The study assessed the effect of several fungicides *viz.* Folicur (tebuconazole), Tilt (propiconazole), Dithane Z-78, Natio, Bavistin and SAAF on pathogen growth inhibition (Table 5). The EC₅₀/ EC₉₀ values for the six fungi-cides were determined and are shown in Table 6. The EC₅₀ values of *A. alternata* for the six fungicides were <1ppm, all the fungicides reduced the fungi growth more than half at this concentration (Fig. 4 and Table 6). Two azole fungicides (Tilt and Folicur) were inhibited the mycelial growth of *A. alternata* most effectively at concentration of 10ppm compared to the others fungicides (Natio, Bavistin, Dithane Z-78 and SAAF) (Fig. 4 and 5). Azole fungicides are sterol demethylation inhibitors (DMIs) that inhibit the C-14 α -demethylation of 24-methylenedihydrolanosterol, a pre-cursor of ergosterol in fungi (Yin *et al.*, 2009). Several field or *in vitro* studies have shown that DMI fungicides such as tebuconazole and propiconazole can control plant pathogenic fungi in different crops (Koller and Scheinpflug 1987; Islam *et al.* 2007; Ivić *et al.* 2011). The EC₉₀ values of *A. alternata* for Natio and SAAF was 25ppm while for Z-78 and Bavistin fungicides was 50ppm. In

Table 6. EC₅₀ and EC₉₀ values of mycelial growth of *A. alternata* for the six fungicides.

Fungicides	EC ₅₀	EC ₉₀
Natio	<1ppm	<25ppm
Folicur	<1ppm	<20ppm
Tilt	<1ppm	10ppm
Bavistin	<1ppm	<50ppm
Z-78	<1ppm	<50ppm
SAAF	<1ppm	<25ppm

this concentration growth of fungi reduced >90% (Table 6) and (Fig. 4). Natio was the third best fungicide which reduced the mycelia growth efficiently with 25ppm. Our study revealed that azole fungicides (Tilt and Folicur) were more effective at inhibiting the mycelial growth of *A. alternata* than other fungicides based on EC₅₀ and EC₉₀ values (Table 6). Similarly, Reuveni and Sheglov (2002) used azoles fungicides to control *A. alternata* pathogen in apple fruit crop. Archana and Jamadar (2014) also studied different fungicides against *A. alternata* and found that azole (propiconazole) reduced more disease as compared to strobilurin and other fungicides. Similar findings were also reported for other plant pathogenic fungi. Nel et al. (2007) evaluated different fungicides against *Fusarium oxysporum* f.sp. *cubense* causal agent of Fusarium wilt of banana and found two fungicides namely prochloraz and propiconazole was very effective to control disease at conc 1 and 5 µg ml⁻¹, respectively. Similarly, Shin et al. (2014) control maize stalk rot disease pathogen *Fusarium subglutinans* and *F. temperatum* control by tebuconazole at conc 0.9 µg/ml. These studies can help to explore the effective chemicals under field condition to manage *A. alternata*.

Conclusion

The present study suggests that pathotypic diversity exist in Punjab isolates of *Alternaria alternata*, they are differentially aggressive on five pomegranate cultivars. It was observed that two cultivars namely Bhagwa and Mridula were found most susceptible, 45.45% isolates showed virulent reaction on both. However, Ruby cultivar was found less susceptible, only 18.18% isolates showed virulent reaction on this cultivar. The Azole fungicides (Tilt and Folicur) were found most effective to inhibit the 90% growth of the mycelium at 10ppm under *in vitro* condition. This group of chemical (Azole fungicides) will be further used in the field condition to manage *Alternaria* black spot disease of pomegranate in Punjab.

REFERENCES

- Anonymous, (2015). Area and Production of pomegranate fruit in India. National horticulture mission online sources.
- Archana, B.C. and Jamadar, M.M. (2014). Management of leaf spot and fruit spot /rot of pomegranate (*Punica granatum* L.) caused by *Alternaria alternata* (Fr.) Keisler. *Karnataka J. Agric. Sci.*, 27:247-49.
- Berbegal, M., López-Cortés, I., Salazar, D., Gramaje, D., Pérez-Sierra, A., García-Jiménez, J. and Armengol, J. (2014). First Report of *Alternaria* Black Spot of Pomegranate Caused by *Alternaria alternata* Spain. *Pl. Dis.*, 98:689
- Ezra, D., Gat, T., Skovorodnikova, Y., Vardi, Y. and Kosto, I. (2010). First report of *Alternaria* black spot of pomegranate by *Alternaria alternata* in Israel. *Aust. Plant Dis. Notes*, 5:1-2
- Farr, D. F., Rossman, A.Y., Palm, M. E. and McCray, E. B. (2007). Fungal Databases, Systematic Botany & Mycology Laboratory, USDA: ARS. Retrieved February 5, 2007, from <http://nt.ars-grin.gov/fungaldatabases>.
- Gat, T., Liarz, I. O., Skovorodnikova, Y., and Ezra, D. (2012). Characterization of *Alternaria alternata* causing black spot disease of pomegranate in Israel using a molecular marker. *Plant Dis.*, 96:1513-18
- Holland, D., Hatib, K. and Bar-Ya'akov, I. (2009). Pomegranate: botany, horticulture, breeding. *Hortic. Rev.*, 35:127-92
- Islam, S.S., Rahman, M. H., Hasan, M. J., Ashadusjaman, M. and Khatun, M.M. (2007). Efficacy of fungicidal seed treatment in Controlling *Alternaria* spp. in Radish Seed. *Int. J. Sustain Crop Prod.*, 2:46-50
- Ito, K., Tanaka, T., Hatta, R., Tamamoto, M., Akimitsu, K. and Tsuge, T. (2004). Dissection of the host range of the fungal plant pathogen *Alternaria alternata* by modification of secondary metabolism. *Mol. Microbiol.*, 52:399-411
- Ivić, D., Sever, Z. and Kuzmanovska, B. 2011. *In vitro* sensitivity of *Fusarium graminearum*, *F. avenaceum* and *F. verticillioi*-des to carbendazim, tebuconazole, flutriafol, metconazole and prochloraz. *Pestic. Phytomed.*, 26:35-42
- Koller, W. and Scheinpflug, H. (1987). Fungal resistance to sterol biosynthesis inhibitors: a new challenge. *Plant Dis.*, 71:1066-74
- Madhukar, J. and Reddy, S. M. (1976). Some new leaf spot diseases of pomegranate. *Indian J. Mycol. Plant Path.*, 18:171-72
- Matheron, M.E. and Porchas, M. (2000). Impact of azoxystrobin, dimethomorph, fluazinam, fosetyl-al, and metalaxyl on growth, sporulation, and zoospore cyst germination of three *Phytophthora* spp. *Plant Dis.*, 84:454-58
- Mesta, R.K., Benagi V. I., Kulkarni, S. and Basavarajappa, M.P. (2011). Management of *Alternaria* blight of sunflower through fungicides. *Karnataka J. Agric. Sci.*, 24: 149-152
- Nel, B., Steinber, C., Labuschagne, N. and Viljoen, A. (2007). Evaluation of fungicides and sterilants for potential application in the management of Fusarium wilt of banana. *Crop Prot.*, 26:697-705
- Pala, H., Tatli, A., Yilmaz, C. and Özgüven, A.I. (2009). Important disease of pomegranate fruit and control possibilities in Turkey. *Acta Hort.*, 818:285-90
- Pasche, J. S., Wharam, C. M. and Gudmestad, N. C. (2004). Shift in sensitivity of *Alternaria solani* in response to QoI fungicides. *Plant Dis.*, 88:181-187
- Reuveni, M.S.D. (2002). Effects of azoxystrobin, difenoconazole, polyoxin B (polar) and trifloxystrobin on germination and growth of *Alternaria alternata* and decay in red delicious apple fruit. *Crop Protec.*, 21:951-55

- Shin, J.H., Han, J.H., Lee, J.K. and Kim, K.S. (2014). Characterization of the maize stalk rot pathogens *Fusarium subglutinans* and *F. temperatum* and the effect of fungicides on their mycelial growth and colony formation. *Plant Pathol. J.*, 30:397-406
- Simmons, E.G. (1967). Typification of *Alternaria*, *Stemphylium* and *Ulocladium*. *Mycologia*, 59:67-92
- Tziros, G.T., Lagopodi, A.L. and Tzavella-Klonari, K. (2007). *Alternaria alternata* fruit rot of pomegranate (*Punica granatum*) in Greece. *New Dis. Rep.*, 15:14
- Yin, Y., Liu, X., Li, B. and Ma, Z. (2009). Characterization of sterol demethylation inhibitor-resistant isolates of *Fusarium asi-aticum* and *F. graminearum* collected from wheat in China. *Phytopathology*, 99:487-497