



Simon, N., Sawkins, E., & Dodd, A. (2018). Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation. *Plant Signaling and Behavior*, [e1457913 ]. <https://doi.org/10.1080/15592324.2018.1457913>

Publisher's PDF, also known as Version of record

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1080/15592324.2018.1457913](https://doi.org/10.1080/15592324.2018.1457913)


[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the final published version of the article (version of record). It first appeared online via Taylor & Francis at <https://www.tandfonline.com/doi/full/10.1080/15592324.2018.1457913> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/pure/about/ebr-terms>



**Editor-in-Chief**  
Michael Bevan  
Professor of Plant and Molecular Biology  
University of Cambridge  
**Section Editors**  
Cellular Signaling  
Developmental Biology  
Plant Physiology  
Plant Pathology  
Plant Systematics  
Plant Molecular Biology  
Plant Genetics  
Plant Biochemistry  
Plant Biophysics  
Plant Evolution  
Plant Ecology  
Plant Development  
Plant Interactions  
Plant Molecular Systematics  
Plant Molecular Evolution  
Plant Molecular Ecology  
Plant Molecular Systematics  
Plant Molecular Evolution  
Plant Molecular Ecology



## Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation

Noriane M. L. Simon, Ellie Sawkins & Antony N. Dodd

To cite this article: Noriane M. L. Simon, Ellie Sawkins & Antony N. Dodd (2018): Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation, *Plant Signaling & Behavior*, DOI: [10.1080/15592324.2018.1457913](https://doi.org/10.1080/15592324.2018.1457913)

To link to this article: <https://doi.org/10.1080/15592324.2018.1457913>



© 2018 The Author(s). Published with license by Taylor & Francis Group, LLC© Noriane M. L. Simon, Ellie Sawkins, and Antony N. Dodd



Accepted author version posted online: 27 Mar 2018.  
Published online: 30 May 2018.



Submit your article to this journal [↗](#)



Article views: 159



View related articles [↗](#)




View Crossmark data [↗](#)

SHORT COMMUNICATION



## Involvement of the SnRK1 subunit KIN10 in sucrose-induced hypocotyl elongation

Noriane M. L. Simon, Ellie Sawkins, and Antony N. Dodd 

School of Biological Sciences, University of Bristol, Life Sciences Building, 24 Tyndall Avenue, Bristol BS8 1TQ, U.K.

### ABSTRACT

A mechanism participating in energy sensing and signalling in plants involves the regulation of sucrose non-fermenting1 (Snf1)-related protein kinase 1 (SnRK1) activity in response to sugar availability. SnRK1 is thought to regulate the activity of both metabolic enzymes and transcription factors in response to changes in energy availability, with trehalose-6-phosphate functioning as a signalling sugar that suppresses SnRK1 activity under sugar-replete conditions. Sucrose supplementation increases the elongation of hypocotyls of developing *Arabidopsis* seedlings, and this response to sucrose involves both the SnRK1 subunit KIN10 and also TREHALOSE-6-PHOSPHATE SYNTHASE1 (TPS1). Here, we measured sucrose-induced hypocotyl elongation in two insertional mutants of KIN10 (*akin10* and *akin10-2*). Under short photoperiods, sucrose supplementation caused great proportional hypocotyl elongation in these KIN10 mutants compared with the wild type, and these mutants had shorter hypocotyls than the wild type in the absence of sucrose supplementation. One interpretation is that SnRK1 activity might suppress hypocotyl elongation in the presence of sucrose, because KIN10 overexpression inhibits sucrose-induced hypocotyl elongation and *akin10* mutants enhance sucrose-induced hypocotyl elongation.

### ARTICLE HISTORY

Received 22 February 2018  
Revised 20 March 2018  
Accepted 21 March 2018

### KEYWORDS

*Arabidopsis*; signal transduction; metabolism; development


We reported recently the involvement of a sugar-signalling mechanism in a pathway that causes hypocotyl elongation in response to sucrose.<sup>1</sup> Hypocotyl elongation in *Arabidopsis thaliana* (*Arabidopsis*) seedlings is caused by cell expansion within the elongating hypocotyl and represents an informative experimental model to study signalling processes that regulate development. In *Arabidopsis*, hypocotyl length is increased by supplementation of the growth media with sucrose.<sup>2–9</sup> We identified that the sugar- and energy-sensing kinase sucrose non-fermenting1 (Snf1)-related protein kinase 1 (SnRK1) regulates sucrose-induced hypocotyl elongation.<sup>1</sup> Under short photoperiods, hypocotyls did not elongate in response to exogenous sucrose in seedlings overexpressing the catalytic alpha subunit of SnRK1, termed SNF1-RELATED PROTEIN KINASE1.1 (KIN10/AKIN10/SnRK1.1).<sup>1</sup> We also found that TREHALOSE-6-PHOSPHATE SYNTHASE1 (TPS1) is required for sucrose-induced hypocotyl elongation under short photoperiods.<sup>1</sup> TPS1 synthesizes the sugar trehalose-6-phosphate (Tre6P), which is a potent inhibitor of SnRK1 activity.<sup>10</sup> Tre6P is thought to function as a signalling sugar that provides information about cellular energy availability.<sup>10,11</sup>

Hypocotyl elongation in response to sucrose might be suppressed in overexpressors of KIN10 (KIN10-ox) because SnRK1 activity is thought to inhibit growth and catabolism under conditions of starvation,<sup>12–14</sup> preventing seedlings from taking advantage of the additional sugars.<sup>1</sup> We reasoned that the converse might be true when SnRK1 activity is low, as occurs under sugar-

replete conditions.<sup>10</sup> To investigate this, we measured the elongation of hypocotyls in response to sucrose in two T-DNA mutants of the KIN10 catalytic subunit of SnRK1 (GABI\_579E09 or *akin10*<sup>15</sup>, and SALKseq\_093965, a new allele named here *akin10-2* for consistency) (Fig S1A). The full-length KIN10 transcript is absent in these *akin10* and *akin10-2* T-DNA lines (Fig. S1B). In the *akin10* mutant, there is a partial loss of phosphorylation of the SnRK1 target bZIP63, most likely due to reduced SnRK1 activity.<sup>15</sup> The remaining phosphorylation of bZIP63 in *akin10* is likely due to KIN11 activity.<sup>15</sup>


Supplementation of wild type seedlings with 3% (w/v) sucrose increased the hypocotyl length under short photoperiods but not under long photoperiods (Fig. 1A, B), as we reported previously.<sup>1</sup> Sucrose supplementation also increased the hypocotyl length of two *akin10* mutants under both short and long photoperiods (Fig. 1A, B). Under short photoperiods, sucrose caused a greater increase in hypocotyl length in *akin10* (6.51 mm longer, 224% increase) and *akin10-2* (6.90 mm longer, 286% increase) compared with the wild type (3.75 mm longer, 67% increase) (Fig. 1A). This greater fold-change in hypocotyl length in the *akin10* mutants under these conditions is because the mutants had significantly shorter hypocotyls than the wild type in the absence of sucrose (Fig. 1A). Under long photoperiods, sucrose supplementation induced hypocotyl elongation in *akin10* mutants, which contrasted the wild type in which sucrose supplementation did not increase hypocotyl length (Fig. 1B). We found previously that sucrose supplementation can decrease the

**CONTACT** Antony N. Dodd  [antony.dodd@bristol.ac.uk](mailto:antony.dodd@bristol.ac.uk)  School of Biological Sciences, University of Bristol, Life Sciences Building, 24 Tyndall Avenue, Bristol BS8 1TQ, U.K.

 Supplemental data for this article can be accessed on the [publisher's website](#).

© 2018 Noriane M. L. Simon, Ellie Sawkins, and Antony N. Dodd. Published with license by Taylor & Francis Group, LLC

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.



**Figure 1.** Sucrose-induced hypocotyl elongation in wild type and *akin10* seedlings measured under (A) 4 h and (B) 16 h photoperiods. Seedlings were cultivated on control media (half strength Murashige and Skoog medium with 0.8% (w/v) agar; 0.5 MS), an equimolar osmotic control (sorbitol), or 3% (w/v) sucrose. Measurement of hypocotyl elongation was conducted as described by Simon et al. 2018. Statistical significance indicated for comparison between seedlings supplemented with 3% (w/v) sucrose and 87.6 mM sorbitol (osmotic control); analysis by univariate ANOVA followed by post-hoc Tukey analysis. Different letters indicate statistically-significant differences between means ( $p < 0.05$ );  $n = 20 \pm$  s.e.m.

hypocotyl length of the Landsberg *erecta* background under long photoperiods,<sup>1</sup> but this did not occur in the Col-0 background used here (Fig. 1B), suggesting that there is some variation between accessions in this developmental response to sucrose.

Hypocotyls of *akin10* and *akin10-2* mutants were significantly shorter than the wild type when cultivated in the absence of sucrose on 0.5MS media (4 h photoperiods, *akin10*  $p < 0.001$ ; *akin10-2*  $p < 0.001$ ; 16 h photoperiods, *akin10*  $p < 0.006$ ; *akin10-2*  $p < 0.001$ ). In addition to changes in phytohormone signalling, the reduced hypocotyl elongation of *akin10* mutations might derive from altered seed quality,<sup>16</sup> attenuated seedling development as occurs in *tps1* knockouts,<sup>17</sup> altered circadian regulation,<sup>18</sup> or altered carbohydrate utilization.<sup>12,19</sup>

The greater proportional increase in hypocotyl length that was caused by sucrose in *akin10* mutants compared with the wild type suggests that SnRK1 activity might contribute to suppression of hypocotyl elongation in response to sucrose. This is because

KIN10 forms a catalytic subunit of SnRK1, and in the absence of this catalytic subunit there was an increase in the magnitude of sucrose-induced hypocotyl elongation. Although KIN10 and KIN11 are thought to confer kinase activity to the SnRK1 complex,<sup>12, 15</sup> *akin10* single mutants change the response of elongating hypocotyls to sucrose (Fig. 1). This indicates that KIN11 cannot completely replace KIN10 within the mechanisms underlying sucrose-induced hypocotyl elongation. This is consistent with the loss of SnRK1 kinase activity in the *akin10* single mutant.<sup>15</sup> An alternative interpretation is that there is some suppression of hypocotyl elongation in the *akin10* mutants in the absence of sucrose, and that this phenotype is lost in the presence of sucrose supplementation (Fig. 1A). Under long photoperiods, sucrose does not cause hypocotyl elongation in the wild type (Fig. 1B), which appears to be due to a combination of photoperiod and daily light input.<sup>1</sup> In comparison, there was sucrose-induced hypocotyl elongation in two *akin10* mutants under long photoperiods. However, under 16 h photoperiods sucrose induced a smaller increase in hypocotyl length in the *akin10* mutants than in *akin10-2* mutants under 4 h photoperiods. Therefore, as with the wild type,<sup>1</sup> photoperiod and/or daily light input influence the magnitude of sucrose-induced hypocotyl elongation in *akin10* mutants. This suggests that mechanisms additional to KIN10 activity within SnRK1 contribute to the photoperiod/daily light input within the response of elongating hypocotyls to sucrose. Such additional mechanisms could include the circadian oscillator, phototransduction pathways, and additional energy-sensing mechanisms.

## Disclosure of potential conflicts of interest

The authors declare no potential conflicts of interest.

## Acknowledgments

We thank BBSRC (UK) for funding (South-West Doctoral Training Partnership grant BB/J014400/1). We thank Prof. Alistair Hetherington for donating *akin10* mutants and Dr. Jean-Charles Isner for discussion about reference transcripts.

## Funding

This work was supported by the Biotechnology and Biological Sciences Research Council (grant BB/J014400/1).

## ORCID

Antony N. Dodd  <http://orcid.org/0000-0001-6859-0105>

## References

- Simon NML, Kusakina J, Fernandez-Lopez A, Chembath A, Belbin FE, Dodd AN. The energy-signaling hub SnRK1 is important for sucrose-induced hypocotyl elongation. *Plant Physiol.* 2018;176:1299–310. doi:10.1104/pp.17.01395.
- Kurata T, Yamamoto KT. *petit1*, a conditional growth mutant of Arabidopsis defective in sucrose-dependent elongation growth. *Plant Physiol.* 1998;118:793. doi:10.1104/pp.118.3.793.
- Takahashi F, Sato-Nara K, Kobayashi K, Suzuki M, Suzuki H. Sugar-induced adventitious roots in Arabidopsis seedlings. *J Plant Res.* 2003;116:83–91.

4. Zhang Y, Liu Z, Wang L, Zheng S, Xie J, Bi Y. Sucrose-induced hypocotyl elongation of Arabidopsis seedlings in darkness depends on the presence of gibberellins. *J Plant Physiol.* 2010;167:1130–6. doi:10.1016/j.jplph.2010.03.007.
5. Liu Z, Zhang Y, Liu R, Hao H, Wang Z, Bi Y. Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in Arabidopsis. *J Plant Physiol.* 2011;168:1771–9. doi:10.1016/j.jplph.2011.04.009.
6. Stewart JL, Maloof JN, Nemhauser JL. PIF genes mediate the effect of sucrose on seedling growth dynamics. *PLoS One.* 2011;6:e19894. doi:10.1371/journal.pone.0019894.
7. Stewart Lilley JL, Gee CW, Sairanen I, Ljung K, Nemhauser JL. An endogenous carbon-sensing pathway triggers increased auxin flux and hypocotyl elongation. *Plant Physiol.* 2012;160:2261–70. doi:10.1104/pp.112.205575.
8. Zhang Z, Zhu J-Y, Roh J, Marchive C, Kim S-K, Meyer C, Sun Y, Wang W, Wang ZY. TOR signaling promotes accumulation of BZR1 to balance growth with carbon availability in Arabidopsis. *Curr Biol.* 2016;26:1854–60. doi:10.1016/j.cub.2016.05.005.
9. Zhang Y, Liu Z, Wang J, Chen Y, Bi Y, He J. Brassinosteroid is required for sugar promotion of hypocotyl elongation in Arabidopsis in darkness. *Planta.* 2015;242:881–93. doi:10.1007/s00425-015-2328-y.
10. Zhang Y, Primavesi LF, Jhurrea D, Andralojc PJ, Mitchell RAC, Powers SJ, Schlupepmann H, Delatte T, Wingler A, Paul MJ. Inhibition of SNF1-related protein kinase 1 activity and regulation of metabolic pathways by trehalose-6-phosphate. *Plant Physiol.* 2009;149:1860–71. doi:10.1104/pp.108.133934.
11. Yadav UP, Ivakov A, Feil R, Duan GY, Walther D, Giavalisco P, Piques M, Carillo P, Hubberten HM, Stütt M, et al. The sucrose–trehalose 6-phosphate (Tre6P) nexus: specificity and mechanisms of sucrose signalling by Tre6P. *J Exp Bot.* 2014;65:1051–68. doi:10.1093/jxb/ert457.
12. Baena-González E, Rolland F, Thevelein JM, Sheen J. A central integrator of transcription networks in plant stress and energy signalling. *Nature.* 2007;448:938–42. doi:10.1038/nature06069.
13. Baena-González E, Sheen J. Convergent energy and stress signaling. *Trends Plant Sci.* 2008;13:474–82. doi:10.1016/j.tplants.2008.06.006.
14. Delatte TL, Sedijani P, Kondou Y, Matsui M, de Jong GJ, Somsen GW, Wiese-Klinkenberg A, Primavesi LF, Paul MJ, Schlupepmann H. Growth arrest by trehalose-6-phosphate: An astonishing case of primary metabolite control over growth by way of the SnRK1 signaling pathway. *Plant Physiol.* 2011;157:160. doi:10.1104/pp.111.180422.
15. Mair A, Pedrotti L, Wurzing B, Anrather D, Simeunovic A, Weiste C, Valerio C, Dietrich K, Kirchner T, Nägele T, et al. SnRK1-triggered switch of bZIP63 dimerization mediates the low-energy response in plants. *Elife.* 2015;4:e05828. doi:10.7554/eLife.05828.
16. Radchuk R, Radchuk V, Weschke W, Borisjuk L, Weber H. Repressing the expression of the SUCROSE NONFERMENTING-1-RELATED PROTEIN KINASE gene in pea embryo causes pleiotropic defects of maturation similar to an abscisic acid-insensitive phenotype. *Plant Physiol.* 2006;140:263–78. doi:10.1104/pp.105.071167.
17. Gómez LD, Gilday A, Feil R, Lunn JE, Graham IA. AtTPS1-mediated trehalose 6-phosphate synthesis is essential for embryogenic and vegetative growth and responsiveness to ABA in germinating seeds and stomatal guard cells. *Plant J.* 2010;64:1–13.
18. Shin J, Sánchez-Villarreal A, Davis AM, Du S-x, Berendzen KW, Koncz C, Ding Z, Li C, Davis SJ. The metabolic sensor AKIN10 modulates the Arabidopsis circadian clock in a light-dependent manner. *Plant Cell Environ.* 2017;40:997–1008. doi:10.1111/pce.12903.
19. Jossier M, Bouly J-P, Meimoun P, Arjmand A, Lessard P, Hawley S, Grahame Hardie D, Thomas M. SnRK1 (SNF1-related kinase 1) has a central role in sugar and ABA signalling in Arabidopsis thaliana. *Plant J.* 2009;59:316–28. doi:10.1111/j.1365-313X.2009.03871.x.