

SCIENTIFIC REPORTS

OPEN

Effect of dietary advanced glycation end products on inflammation and cardiovascular risks in healthy overweight adults: a randomised crossover trial

Estifanos Baye¹, Maximilian PJ de Courten², Karen Walker³, Sanjeeva Ranasinha¹, Arul Earnest⁴, Josephine M Forbes^{5,6} & Barbora de Courten^{1,7}

Diets high in advanced glycation end products (AGEs) are thought to be detrimental to cardiovascular health. However, there remains uncertainty about the beneficial effect of a low AGE diet on cardiovascular risk factors and inflammatory markers in overweight individuals. We thus performed a randomised, double blind, crossover trial to determine whether consumption of low AGE diets reduce inflammation and cardiovascular risks in overweight and obese otherwise healthy adults. All participants ($n = 20$) consumed low and high AGE diets alternately for two weeks and separated by a four week washout period. Low AGE diets did not change systolic ($p = 0.2$) and diastolic blood pressure ($p = 0.3$), mean arterial pressure ($p = 0.8$) and pulse pressure ($p = 0.2$) compared to high AGE diets. Change in total cholesterol ($p = 0.3$), low-density lipoprotein ($p = 0.7$), high-density lipoprotein ($p = 0.2$), and triglycerides ($p = 0.4$) also did not differ and there was no difference in inflammatory markers: interleukin-6 ($p = 0.6$), monocyte chemoattractant protein-1 ($p = 0.9$), tumour necrosis factor α ($p = 0.2$), C-reactive protein ($p = 0.6$) and nuclear factor kappa beta ($p = 0.2$). These findings indicate that consumption of low AGE diets for two weeks did not improve the inflammatory and cardiovascular profiles of overweight and obese adults.

Advanced glycation end products (AGEs) are formed endogenously from non-enzymatic reactions of amino acids with sugars¹. Consumption of foods high in sugar and/or foods exposed to high temperature cooking methods such as deep-frying, broiling, roasting, baking and grilling can increase the total daily AGE intake by 25% compared to the average adult daily intake^{2,3}. High AGE intake from food accelerates the production of endogenous AGEs and increases the level of circulating AGEs in the blood stream^{4,5}. This intake of high AGE contributes to the progression of type 2 diabetes (T2DM) and cardiovascular diseases (CVD) although the exact mechanisms are not clearly understood⁶⁻¹¹. AGE formation can be reduced by shortening the cooking time and lowering temperature to reduce food browning, or by addition of acidic (low pH) ingredients and by high humidity or food moisture content^{7,12}. Given the ease of such changes, reduction of AGE intake may be a promising intervention to lower cardiovascular risk^{3,13-15}.

In humans, the impact of low AGE diets on inflammation and cardiovascular risk factors is not clearly understood. In some studies involving healthy obese individuals, low AGE diets reduced inflammatory markers^{11,16,17} but did not change plasma lipid levels^{17,18} or blood pressure^{17,19}. Whereas, other studies showed a reduction in

¹Monash Centre for Health Research and Implementation, School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia. ²Centre for Chronic Disease, College of Health and Biomedicine, Victoria University, Melbourne, Australia. ³Department of Nutrition and Dietetics, School of Clinical Sciences, Monash University, Melbourne, Australia. ⁴Biostatistics Unit, School of Public Health & Preventive Medicine, Monash University, Melbourne, Australia. ⁵Mater Clinical School, University of Queensland, Brisbane, Australia. ⁶Glycation and Diabetes, Mater Research Institute–University of Queensland, Translational Research Institute, Brisbane, Australia. ⁷Diabetes and Vascular Medicine Unit, Monash Health, Melbourne, Australia. Correspondence and requests for materials should be addressed to B.d.C. (email: barbora.decourten@monash.edu)

Parameters	Low AGE group	High AGE group	P-value
Weight (kg)	94.2 ± 15.2	94.1 ± 14.3	0.94
Body mass index (kg/m ²)	31.4 ± 3.8	31.3 ± 3.7	0.85*
% body fat	30.7 ± 6.5	30.6 ± 6.6	0.83
Waist-to-hip ratio	0.9 ± 0.07	0.9 ± 0.07	0.35*
Systolic BP (mm Hg)	123.1 ± 10.8	122.3 ± 12.3	0.66
Diastolic BP (mm Hg)	77.9 ± 8.9	76.7 ± 9	0.38
Mean arterial pressure (mm Hg)	92.9 ± 9.1	91.9 ± 9	0.37
Pulse pressure (mm Hg)	45.1 ± 6.4	45.6 ± 10.1	0.90*
Total cholesterol (mmol/l)	4.8 ± 1	4.6 ± 1	0.06
LDL (mmol/l)	3.1 ± 0.8	3.1 ± 0.7	0.61
HDL (mmol/l)	1 ± 0.2	1 ± 0.2	0.35
Triglycerides (mmol/l)	1.5 ± 1	1.3 ± 0.7	0.11*
CRP (mg/l)	2.1 ± 1.9	2.4 ± 2.7	0.69*
TNF α (pg/ml)	2.8 ± 3.7	1.6 ± 0.9	0.18
MCP-1 (pg/ml)	126 ± 99	121 ± 66	0.92*
IL-6 (ng/ml)	1.5 ± 2.2	1.6 ± 2.4	0.64*
NF κ B p65 activity (ng/ μ g protein)	14 ± 10	12 ± 10	0.53

Table 1. Participants' baseline characteristics at the beginning of each study period (n = 20). All values are presented as means \pm SDs. Paired t-tests were conducted to determine the p-values. *Data transformation was done. BP, blood pressure; CRP, C-reactive protein; HDL, high-density lipoprotein; IL-6, interleukin -6; LDL, low-density lipoprotein; MAP, mean arterial pressure; MCP-1, monocyte chemoattractant protein-1; NF κ B; Nuclear factor kappa beta; PP, pulse pressure.

plasma lipid levels after low AGE diets^{19–21} but not of markers of inflammation^{18, 22}. In patients with T2DM, low AGE diets improved inflammatory markers^{6, 8, 9}, but not blood pressure⁹, and lipid levels^{6, 9}. Cai and colleagues however reported a reduction in oxidised low-density lipoprotein after 6 weeks of consumption of low AGE diets (5 fold lower in AGE content) in patients with T2DM²³. In all these studies, test and control diets were either not matched for energy and macronutrient content or it was not stated whether similar macronutrient content was achieved between the diets, which could potentially influence the results²⁴. We have therefore investigated the impact of low and high AGE diets carefully matched for both energy and macronutrient profile. These matched diets were given for two weeks each to healthy obese adults in a randomised cross-over design trial to determine effects on cardiometabolic parameters. We have previously reported the main outcomes of the trial which showed that dietary AGEs improved insulin sensitivity but not insulin secretion in overweight and obese non-diabetic individuals²⁴. In this study, we conducted analyses of secondary outcomes of the trial and investigated if dietary AGEs improve blood pressure, plasma lipid profiles and inflammatory markers.

Results

Baseline characteristics. As previously described²⁴, the mean age, body mass index, waist-to-hip ratio and % body fat of the participants were 34 (10) years, 31.3 (3.8) kg/m², 0.9 (0.1) and 31.1 (6.7), respectively. There was no difference in levels of circulating AGEs and metabolic profiles such as insulin sensitivity and secretion between group 1 and 2 before each test diet²⁴. Anthropometric and blood pressure measurements, plasma lipid levels and inflammatory markers were also not different between the two interventions at the beginning of each dietary period (Table 1). The cumulative dietary AGE intakes were significantly lower in the low AGE dietary period compared to the high AGE dietary period (all p < 0.002), as reported previously²⁴. The mean consumption of carboxymethyllysine, carboxyethyllysine and methylglyoxal-derived hydroimidazolidine were decreased by 27%, 38%, and 21%, respectively, in the low AGE group compared to the high AGE group.

Effect of low AGE diet on inflammatory markers. Inflammatory markers such as tumour necrosis factor α (TNF α), C-reactive protein (CRP), monocyte chemoattractant protein 1 (MCP-1) and interleukin 6 (IL-6) levels and nuclear factor kappa beta (NF κ B) activity in peripheral blood mononuclear cells (PBMCs) did not change after consumption of either low or high AGE diets. Between groups comparison also showed that the changes in these markers were not significantly different between low and high AGE diets (Table 2).

Effect of low AGE diets on cardiovascular parameters. Anthropometric measures such as weight and body mass index were significantly reduced only in the low AGE group compared to high AGE group. Percentage body fat and waist-to-hip ratio did not change after consumption of either diet. Between groups comparison have shown that none of these anthropometric measures were different between the diets (Table 2). Systolic and diastolic blood pressure and mean arterial pressure decreased after consumption of both diets whereas pulse pressure was reduced only after intake of low AGE diets. Between groups analyses (after controlling for period and sequence effect) did not show any significant change in these measurements between the diets (Table 2).

The total plasma cholesterol levels decreased in both groups, however, the change between the two diets was not significant. Both plasma low-density lipoprotein (LDL) and high-density lipoprotein (HDL) cholesterol levels

Parameters	Change from baseline in low AGE group		Change from baseline in high AGE group		Difference between the groups	
	Mean \pm SD	P-value [†]	Mean \pm SD	P-value [†]	Mean \pm SD	P-value ^{††}
Weight (kg)	-0.69 \pm 1.34	0.03	-0.16 \pm 1.17	0.54	-0.52 \pm 1.52	0.17
BMI (kg/m ²)	-0.22 \pm 0.45	0.03	-0.06 \pm 0.38	0.49	-0.16 \pm 0.5	0.20
% body fat	-0.06 \pm 1.2	0.82	-0.08 \pm 1.39	0.80	+0.02 \pm 1.56	0.37
WHR	+0.004 \pm 0.03	0.55	-0.004 \pm 0.032	0.54	+0.008 \pm 0.031	0.49
Systolic BP (mm Hg)	-1.2 \pm 7.7	0.49	-1.3 \pm 8.3	0.47	+0.1 \pm 11.4	0.24
Diastolic BP (mm Hg)	-0.8 \pm 8.4	0.67	-1.9 \pm 6.2	0.18	+1.1 \pm 11.6	0.33
Pulse Pressure (mm Hg)	-0.4 \pm 7.7	0.81	+0.5 \pm 9.7	0.80	-0.9 \pm 9.2	0.16
MAP (mm Hg)	-0.9 \pm 7.3	0.57	-1.7 \pm 5.2	0.16	+0.8 \pm 10.7	0.80
Total cholesterol (mmol/l)	-0.4 \pm 0.5	0.0006	-0.15 \pm 0.3	0.04	-0.3 \pm 0.6	0.26
LDL (mmol/l)	-0.3 \pm 0.3	0.002	-0.01 \pm 0.3	0.86	-0.3 \pm 0.5	0.73
HDL (mmol/l)	-0.06 \pm 0.1	0.03	-0.008 \pm 0.08	0.66	-0.05 \pm 0.1	0.21
Triglycerides (mmol/l)	-0.1 \pm 0.3	0.08	-0.2 \pm 0.3	0.01	+0.04 \pm 0.5	0.44
IL-6 (ng/ml)	-0.4 \pm 1.9	0.71*	-0.2 \pm 1.8	0.94*	-0.2 \pm 2.9	0.62*
MCP-1 (pg/ml)	-5.4 \pm 37.4	0.52	+1.9 \pm 44.6	0.84	+7.41 \pm 65.8	0.94
TNF α (pg/ml)	-0.4 \pm 4.4	0.84*	+1.8 \pm 5.5	0.16*	-2.2 \pm 7.1	0.15*
CRP (mg/l)	+0.08 \pm 0.66	0.57	+0.19 \pm 1.55	0.57	+0.1 \pm 1.6	0.60
NF κ B p65 activity (ng/ μ g protein)	-3.4 \pm 9.9	0.14	-1.3 \pm 11.4	0.61	-2.1 \pm 16.7	0.18

Table 2. Effects of low and high AGE diets on markers of inflammation and cardiovascular risk factors (n = 20). [†]Paired t-tests were conducted to determine the p-values within each groups. ^{††}A 2 \times 2 ANOVA for cross over study was used to determine the overall difference between the diets. *Data transformation was done. BP, blood pressure; CRP, C-reactive protein; HDL, high-density lipoprotein cholesterol; IL-6, interleukin -6; LDL, low-density lipoprotein; MAP, mean arterial pressure; MCP-1, monocyte chemoattractant protein-1; NF κ B; nuclear factor kappa B; SD, standard deviation.

decreased with low AGE diet but not changed with high AGE diet whereas the overall difference between the diets was not significant (Table 2). Plasma triglyceride levels decreased with high AGE diet but there was no difference between low and high AGE diets (Table 2).

Correlation analyses between inflammation markers and anthropometric measures and cardiovascular risk factors.

The correlation analyses indicated that the change in diastolic blood pressure and MCP-1 is positively correlated with % body fat ($r = 0.43$, $p = 0.05$) and waist-to-hip ratio ($r = 0.46$, $p = 0.03$), respectively. Change in systolic blood pressure is also positively correlated with NF κ B ($r = 0.47$, $p = 0.04$) whereas plasma lipid levels negatively correlate with CRP (total cholesterol: $r = -0.53$, $p = 0.01$; HDL: $r = -0.59$, $p = 0.006$; LDL: $r = -0.56$, $p = 0.008$) and IL-6 (total cholesterol: $r = -0.51$, $p = 0.03$; HDL: $r = -0.58$, $p = 0.01$; LDL: $r = -0.54$, $p = 0.01$). After adjusting either for change in weight, body mass index, % body fat or waist-to-hip ratio, neither systolic blood pressure nor lipid parameters correlated with inflammatory markers. Other inflammation parameters did not show any correlation with cardiovascular parameters (all $p > 0.1$). Circulating and urinary AGE levels also did not correlate with any cardiovascular parameters (all $p > 0.09$).

Discussion

The present study aimed to measure the effect of consuming low AGE diets for 2 weeks on markers of inflammation and cardiovascular risk factors in healthy obese and overweight individuals. No significant difference was found in changes in blood pressure, plasma lipid levels (total cholesterol, LDL, HDL, triglycerides) or inflammatory markers (CRP, TNF α , MCP-1, IL-6, NF κ B activity) between low and high AGE diets. Our findings on blood pressure measurements were consistent with other studies conducted in healthy overweight individuals^{17, 19}, prediabetes²¹, obese people with metabolic syndrome^{25, 26}, and patients with T2DM⁹.

Regarding plasma lipid profile, we did not find a significant difference between low and high AGE diets. Some studies conducted in healthy individuals^{17, 18}, obese individuals with metabolic syndrome^{25, 26}, and patients with T2DM^{6, 9} have also reported that consumption of low AGE diets did not change plasma lipid levels compared to high AGE diets. In contrast, two other studies that involved young healthy obese volunteers²⁰ and prediabetic individuals²¹ noted a reduction in plasma lipid levels after a low AGE diet. In the first study, the difference between the 2 diets could have been due to the higher intake of dietary fat and carbohydrate on the high AGE diets²⁰. The patients in the second trial in the low AGE arm of the study had relatively lower intima-media thickness compared to those who were in the high AGE diet²¹. This may exaggerated the effect of high AGE diets on lipid profile. In addition, consumption of low AGE diets combined with aerobic exercise program for 12 weeks showed a decrease in plasma lipid levels in overweight and obese men¹⁹. This effect however could have been due to the synergistic effect of physical activity with low AGE diets as it has been widely known for its beneficial effect on metabolic changes²⁷. Therefore, these studies^{19–21} were not able to differentiate between the true effect of low and high AGE diet as they were confounded by other differences in diet, exercise or disease state of the patients.

In this study, intake of low AGE diets for 2 weeks did not change inflammatory markers. This finding was consistent with some studies conducted in healthy overweight and obese individuals^{18, 22}. Other studies conducted in healthy individuals showed an improvement in inflammatory markers after intake of low AGE diets^{11, 16, 17}. However, in these studies, the diets were not matched for macronutrient content, which might have influenced the findings. Studies in obese people with metabolic syndrome²⁶, prediabetes²¹ and patients with T2DM^{8, 9} showed an improvement in inflammatory markers after intake of low AGE diets compared to high AGE diets. Therefore, it could be suggested that intake of low AGE diets was beneficial for those who had higher baseline values in inflammatory markers such as patients with metabolic syndrome or T2DM.

Compared to the previous studies, our study has several strengths that include standardised diets in energy and macronutrient content were provided, study personnel including the investigators were unaware of the dietary allocation, and baseline characteristics such as anthropometric, circulating AGEs and metabolic parameters were not different at baseline during each dietary period. We have also used a crossover design to control for possible confounding factors in addition to the rigorous randomisation process. The small sample size and relatively short duration could be considered as main limitations of this study as one can argue that the change in physiologic processes after intake of low AGE diets may require longer duration.

In conclusion, we have shown that consumption of low AGE diets for 2 weeks did not have beneficial effects in reducing markers of inflammation and cardiovascular risk factors in healthy overweight and obese adults. Long-term well designed trials with larger sample sizes are needed to confirm our findings.

Methods

Study participants. Twenty overweight and obese but otherwise healthy and normoglycemic adults, aged 18–50 years completed the study²⁴, according to a protocol as previously published elsewhere²⁸. Participants did not have diabetes as indicated by a 75 g oral glucose tolerance test (OGTT) (WHO 1999 criteria). All were non-smokers and had no clinical and laboratory signs of infection and none took supplements/medications during the study period. The participants were recruited from the general community through advertisements between January 2006 and December 2010. Ethical approval was obtained from the Alfred Hospital Ethics Committee, Melbourne, Australia and complied with the Declaration of Helsinki. All participants provided written informed consent prior to participation.

Study design. This study employed a two-period randomised cross-over double-blind design. All participants underwent both diets; one low in AGE content and one high in AGE content (typical of a modern Western diet) and diets were matched in energy and macronutrient content. Participants commenced the study after a two-week run-in on their habitual diet but with limitation of their alcohol, fast food and coffee intake. Test diets were consumed for two weeks separated by a four-week washout period (habitual diet). The primary outcomes of this study that focused on insulin resistance/sensitivity have been published²⁴.

Randomisation and masking. Randomisation occurred for 7 blocks of 4 participants stratified by gender and diet order. Participants were masked to the allocation of diet type and to how the diet might affect glucose metabolism. Clinical and laboratory investigators were also masked to diet allocation.

Study Procedures. All participants underwent medical and laboratory screening including a 75 g oral glucose tolerance test (OGTT). Prior to metabolic testing, participants were asked to abstain from strenuous exercise and caffeine for 3 days.

Dietary Intervention. Prior to starting the ‘run-in’ to the first diet, participants kept a 3-day diet record of their habitual diet (2 weekdays, 1 weekend day) based on household measures. Nutrient content was analysed with SERVE (SERVE Nutrition Systems, St Ives, NSW), based on Australian food composition tables. With the use of Australian food composition data from SERVE as well as data from the United States on the N (ϵ)-carboxymethyllysine (CML – the most common AGEs used in clinical studies as an indicator of dietary AGE intake) content of common foods, a menu of carefully matched alternative food choices (of low AGE vs high AGE content) individualised to suit the preferences and habitual diet of each participant was designed. These alternative food choices were matched for macronutrient content and total energy but greatly differed in calculated AGE content, and were provided for each meal of the day, including snacks and beverages. Food choices on the high AGE diet had higher dietary AGE content, while the isoenergetic low AGE diet, matched for macronutrient content, had reduced AGE content through altered cooking techniques⁴ and use of differently processed foods. All food for the two test diets was provided weekly to the participants as ready-to-eat items or as packed food portions, to assist with dietary compliance. For food that required cooking, detailed instructions for storage and heating (method, temperature, and duration) were provided. Participants were asked to eat to appetite throughout both dietary periods to maintain constant body weight. They were required to keep a detailed dietary record indicating cooking method and number of portions eaten for each food item supplied, unconsumed foods or additional foods eaten. Later foods incorporated in these diets were also chemically analysed for their AGE content. The dietician made telephone contact twice a week to provide support and resolve difficulties as well as to ensure dietary compliance. To analyse the dietary AGE content, food items were obtained from local supermarkets and prepared according to the instructions provided to the participants in the study.

Cardiovascular measures. Systolic and diastolic blood pressure were measured using an automated oscillometric measurement system (Omron) after a 30 minute rest. Pulse pressure was calculated by subtracting the diastolic blood pressure from the systolic blood pressure. Mean arterial pressure was computed as diastolic blood pressure + pulse pressure/3²⁹. Plasma lipid levels such as total cholesterol, LDL, HDL and triglycerides were

measured using a standard commercial enzymatic assay (Beckman Coulter LX20PRO Analyser and Synchron and Systems Lipid and Multi Calibrators (Beckman Coulter Diagnostics).

Measurement of inflammatory markers and AGEs. Markers of inflammation (TNF α , MCP-1 and IL-6 levels) were analysed using a commercial automated chemiluminescent enzyme immunoassay and immulite analyser (Diagnostic Products Corporation). Whereas, highly sensitive near infrared particle immunoassay rate methodology was used to measure plasma CRP levels. NF κ B activity in PBMCs was detected and quantified using TransAM NF- κ B DNA-binding activity assay (Active Motif, Carlsbad, CA, USA). Measurement of urinary and serum AGEs were quantified with the use of ultraperformance liquid chromatography–tandem mass spectrometry as previously described³⁰.

Statistical analysis. Change from baseline values was calculated for each parameter in both dietary periods. Paired t-tests were used to compare the participants' characteristics at the beginning of each dietary period and to determine the change in parameters after each study diet. Repeated measures ANOVA was employed to determine the change difference between the two diets after adjusting for carryover, sequence and period effects. The observed carryover effect in this study was very minimal (29.8%) and thus the data in both study periods were combined. Results are presented using means and standard deviations unless and otherwise stated. Appropriate data transformation was undertaken in the event of skewed data to approximate a normal distribution. Statistical analyses were performed using Stata 14 software. P-value of ≤ 0.05 is used to describe statistical significant difference.

Data Availability. The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

References

- Zhang, Q., Ames, J. M., Smith, R. D., Baynes, J. W. & Metz, T. O. A perspective on the Maillard reaction and the analysis of protein glycation by mass spectrometry: probing the pathogenesis of chronic disease. *J Proteome Res* **8**, 754–769, doi:10.1021/pr800858h (2009).
- Uribarri, J. *et al.* Diet-derived advanced glycation end products are major contributors to the body's AGE pool and induce inflammation in healthy subjects. *Ann N Y Acad Sci* **1043**, 461–466, doi:10.1196/annals.1333.052 (2005).
- Negrea, M. *et al.* Effects of low- and high-advanced glycation endproduct meals on macro- and microvascular endothelial function and oxidative stress in patients with type 2 diabetes mellitus. *Am J Clin Nutr* **85**, 1236–1243 (2007).
- Goldberg, T. *et al.* Advanced glycoxidation end products in commonly consumed foods. *J Am Diet Assoc* **104**, 1287–1291, doi:10.1016/j.jada.2004.05.214 (2004).
- Uribarri, J. *et al.* Advanced glycation end products in foods and a practical guide to their reduction in the diet. *J Am Diet Assoc* **110**, 911–916.e912, doi:10.1016/j.jada.2010.03.018 (2010).
- Luevano-Contreras, C., Garay-Sevilla, M. E., Wrobel, K., Malacara, J. M. & Wrobel, K. Dietary advanced glycation end products restriction diminishes inflammation markers and oxidative stress in patients with type 2 diabetes mellitus. *J Clin Biochem Nutr* **52**, 22–26, doi:10.3164/jcfn.12-40 (2013).
- Poulsen, M. W. *et al.* Advanced glycation endproducts in food and their effects on health. *Food Chem Toxicol* **60**, 10–37, doi:10.1016/j.fct.2013.06.052 (2013).
- Uribarri, J. *et al.* Restriction of advanced glycation end products improves insulin resistance in human type 2 diabetes: potential role of AGER1 and SIRT1. *Diabetes Care* **34**, 1610–1616, doi:10.2337/dc11-0091 (2011).
- Vlassara, H. *et al.* Inflammatory mediators are induced by dietary glyco toxins, a major risk factor for diabetic angiopathy. *Proc Natl Acad Sci USA* **99**, 15596–15601, doi:10.1073/pnas.242407999 (2002).
- Vlassara, H. & Striker, G. E. AGE restriction in diabetes mellitus: a paradigm shift. *Nat Rev Endocrinol* **7**, 526–539, doi:10.1038/nrendo.2011.74 (2011).
- Harcourt, B. E. *et al.* Targeted reduction of advanced glycation improves renal function in obesity. *Kidney Int* **80**, 190–198, doi:10.1038/ki.2011.57 (2011).
- Ottum, M. S. & Mistry, A. M. Advanced glycation end-products: modifiable environmental factors profoundly mediate insulin resistance. *J Clin Biochem Nutr* **57**, 1–12, doi:10.3164/jcfn.15-3 (2015).
- Birlouez-Aragon, I. *et al.* A diet based on high-heat-treated foods promotes risk factors for diabetes mellitus and cardiovascular diseases. *American Journal of Clinical Nutrition* **91**, 1220–1226 (2010).
- Clarke, R. E., Dordevic, A. L., Tan, S. M., Ryan, L. & Coughlan, M. T. Dietary Advanced Glycation End Products and Risk Factors for Chronic Disease: A Systematic Review of Randomised Controlled Trials. *Nutrients* **8**, doi:10.3390/nu8030125 (2016).
- Kellow, N. J. & Savige, G. S. Dietary advanced glycation end-product restriction for the attenuation of insulin resistance, oxidative stress and endothelial dysfunction: a systematic review. *European Journal of Clinical Nutrition* **67**, 239–248 (2013).
- Uribarri, J. *et al.* Suppression of native defense mechanisms, SIRT1 and PPARgamma, by dietary glycoxidants precedes disease in adult humans; relevance to lifestyle-engendered chronic diseases. *Amino Acids* **46**, 301–309, doi:10.1007/s00726-013-1502-4 (2014).
- Vlassara, H. *et al.* Protection against loss of innate defenses in adulthood by low advanced glycation end products (AGE) intake: role of the antiinflammatory AGE receptor-1. *J Clin Endocrinol Metab* **94**, 4483–4491, doi:10.1210/jc.2009-0089 (2009).
- Semba, R. D. *et al.* Dietary intake of advanced glycation end products did not affect endothelial function and inflammation in healthy adults in a randomized controlled trial. *J Nutr* **144**, 1037–1042, doi:10.3945/jn.113.189480 (2014).
- Macias-Cervantes, M. H. *et al.* Effect of an advanced glycation end product-restricted diet and exercise on metabolic parameters in adult overweight men. *Nutrition* **31**, 446–451, doi:10.1016/j.nut.2014.10.004 (2015).
- Birlouez-Aragon, I. *et al.* A diet based on high-heat-treated foods promotes risk factors for diabetes mellitus and cardiovascular diseases. *Am J Clin Nutr* **91**, 1220–1226, doi:10.3945/ajcn.2009.28737 (2010).
- Di Pino, A. *et al.* Low advanced glycation end product diet improves the lipid and inflammatory profiles of prediabetic subjects. *J Clin Lipidol* **10**, 1098–1108, doi:10.1016/j.jacl.2016.07.001 (2016).
- Poulsen, M. W. *et al.* Effect of dietary advanced glycation end products on postprandial appetite, inflammation, and endothelial activation in healthy overweight individuals. *Eur J Nutr* **53**, 661–672, doi:10.1007/s00394-013-0574-y (2014).
- Cai, W. *et al.* High levels of dietary advanced glycation end products transform low-density lipoprotein into a potent redox-sensitive mitogen-activated protein kinase stimulant in diabetic patients. *Circulation* **110**, 285–291, doi:10.1161/01.cir.0000135587.92455.0d (2004).
- de Courten, B. *et al.* Diet low in advanced glycation end products increases insulin sensitivity in healthy overweight individuals: a double-blind, randomized, crossover trial. *Am J Clin Nutr* **103**, 1426–1433, doi:10.3945/ajcn.115.125427 (2016).

25. Uusitupa, M. *et al.* Effects of an isocaloric healthy Nordic diet on insulin sensitivity, lipid profile and inflammation markers in metabolic syndrome—a randomized study (SYSDIET). *J Intern Med* **274**, 52–66, doi:10.1111/joim.12044 (2013).
26. Vlassara, H. *et al.* Oral AGE restriction ameliorates insulin resistance in obese individuals with the metabolic syndrome: a randomised controlled trial. *Diabetologia* **59**, 2181–2192, doi:10.1007/s00125-016-4053-x (2016).
27. Ross, R. *et al.* Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* **133**, 92–103 (2000).
28. de Courten, B., de Courten, M. P., Schalkwijk, C. G., Walker, K. Z. & Forbes, J. Dietary Advanced Glycation End Products Consumption as a Direct Modulator of Insulin Sensitivity in Overweight Humans: A Study Protocol for a Double-Blind, Randomized, Two Period Cross-Over Trial. *JMIR Res Protoc* **4**, e93, doi:10.2196/resprot.4552 (2015).
29. Sourris, K. C. *et al.* Plasma advanced glycation end products (AGEs) and NF-kappaB activity are independent determinants of diastolic and pulse pressure. *Clin Chem Lab Med* **52**, 129–138, doi:10.1515/cclm-2012-0850 (2014).
30. Hanssen, N. M. *et al.* Plasma levels of advanced glycation endproducts Nepsilon-(carboxymethyl)lysine, Nepsilon-(carboxyethyl)lysine, and pentosidine are not independently associated with cardiovascular disease in individuals with or without type 2 diabetes: the Hoorn and CODAM studies. *J Clin Endocrinol Metab* **98**, E1369–1373, doi:10.1210/jc.2013-1068 (2013).

Acknowledgements

This study was supported by grants from the National Health and Medical Research Council of Australia and from a Diabetes Australia Research Trust Millennium Award. EB is a recipient of Monash Graduate Scholarship and Monash International Postgraduate Scholarship. BdC is supported by National Heart Foundation Future Leader Fellowship (100864). JMF is supported by Fellowship from the Australian National Health and Medical Research Council. No funder had any role in the study design, data collection, data analysis or interpretation, or writing of the manuscript.

Author Contributions

E.B. analysed the data, wrote the first draft of the manuscript and revised the subsequent drafts. B.d.C., S.R. & A.E. contributed to the analysis and review of the manuscript. M.P.J.d.C., K.W. & J.M.F. contributed to the review of the manuscript. All authors read and approved the manuscript. The corresponding author had access to all the data and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Additional Information

Competing Interests: The authors declare that they have no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2017