

## A Comparison on FCR in fish ponds of Rainbow Trout (*Oncorhynchus mykiss*) fed by extruder food and pellete food

Pirali Kheyraadi, E.<sup>1</sup>; Salehi Farsani, A.<sup>2\*</sup>; Samadi Kuchaksaraei, B.<sup>2</sup>

Received: September 2012 Accepted: May 2013

1-Department of Fisheries and Environment Science, College of natural Resources and Earth Sciences, Shahrekord University, Chaharmahal-va-BAKHTIYARI, Iran

2-Department of Fisheries Science, Islamic Azad University, Science and Research Branch, Tehran, Iran

\*Corresponding author's email: ammar\_1631@yahoo.com

**Keywords:** FCR, Rainbow trout, Extruder food, Pellete food

Food costs in rainbow trout farms are about 50% of production costs (Barrows *et al.*, 2007). Because of inattention of most aquaculturists in exact calculation of FCR and not using qualified food, sometimes food costs will rise a lot and won't be economical.

The newest method for fish food production is production during the process of extrusion. Extrusion process is processing materials with high temperature in a short time (HTST) and is done by combined operation of Humidity, heat, mechanical energy and pressure (Watanabe and Pongmaneerat, 1993).

Basically the longer food is under heat the more its quality will decrease. Because some amino-acids like lysine, tryptophan, cysteine, histidine,... are very sensitive to heat especially dry heat. Loss of these amino-acids reduce food quality (Barrows *et al.*, 2007). Extruder food has these benefits in addition to pellet food benefits:

Starch of food is completely gelatinized in extrusion process. So its digestion will be accelerated and 90% improved. In pellete food starch digestion capability is almost 20%. Extruder food has soft spongy tissue and floats sustainably in the water. So it decreases food waste. Other benefits of extruder food are better food sterilization and deleting of harmful microorganisms (Pourjafar, 2007). Some studies were done about extruder food in recent years (Hilton *et al.*, 1981; Pfeffer *et al.*, 1991; Watababe and Pongmaneerat, 1993; Sørensen *et al.*, 2002; Cheng and Hardy, 2003; Barrows *et al.*, 2008). But there are a few studies about effects of extruder food on FCR (Boccignore *et al.*, 1989; Al-Ruqaie, 2007; Degaura and San Lucjan, 1997).

In this study, it was tried to compare between extruder and pellete food in addition to introduce a correct method for FCR calculation to determine economical food that

give more profit to culturists. This study was done in rainbow trout fish farm in Chaharmahal-va-Bakhtiari province (Iran) in 30 days from 23<sup>th</sup> of August to 22<sup>th</sup> of October 2010. 10000 fish specimens of one pond with average weight of 100 g and total weight of 1000 kg were collected. These fish were divided into 4 groups each contains 2500 specimens (Table 1). Then fish were introduced into 4 ponds each had a volume of 120 m<sup>3</sup>. Input water to each pond was 60 lit/sec. Ponds were named as A, B, C and D treatments. The ponds A and B were fed by extruder food with amount of 2% of body weight and ponds C and D were fed by pelleted food with the same amount in 60 days. Experiment was done with 4 replicates. Fish

were sampled every 15 days. In each randomized sampling there were 5 replicates and average weight and average length of 200 fish were randomly measured. Finally, the FCR were calculated as follows:  
Food conversion rate (FCR) = Food eaten/Weight gain.

For weighing fish digital scale with accuracy of two decimal places and for length determination biometric board were used. Weight and total length (TL) calculation were done in farm and then 200 harvested fish were released to pond again.


Difference between FCR in two different kinds of diets was determined by one way ANOVA by SPSS version 19 (Table 2).

**Table 1-Rainbow trout biological characteristics and the fish ponds features**

| Pond | No. of fish | Average weight (gr) | Total weight (gr) | Average TL (cm) | Water volume (m <sup>3</sup> ) |
|------|-------------|---------------------|-------------------|-----------------|--------------------------------|
| A    | 2500        | 100±10.3            | 250               | 20±5.5          | 120                            |
| B    | 2500        | 100±7.5             | 250               | 20±4.3          | 120                            |
| C    | 2500        | 100±9.2             | 250               | 20±6.2          | 120                            |
| D    | 2500        | 100±9.8             | 250               | 20±6.9          | 120                            |

**Table 2-Average FCR in ponds fed by extruder food compared with ponds fed by pellete food in study period**

| Pond    | Total fish | Average weight (gr) | Average TL(cm) | Mortalities (%) | Total weight | New average weight (gr) | New average length (cm) | Total weight (kg) | Type of food | Amount of food (kg) | Weight gain (kg) | FCR      |
|---------|------------|---------------------|----------------|-----------------|--------------|-------------------------|-------------------------|-------------------|--------------|---------------------|------------------|----------|
| A and B | 500        | 100±8.6             | 20±5.1         | 46              | 500          | 282±30.3                | 35±7.2                  | 1397              | Ex           | 876                 | 897              | 0.9±0.1  |
| C and D | 500        | 100±9.5             | 20±6.4         | 103             | 500          | 211.5±33.2              | 35±5.8                  | 1035              | Pl           | 767                 | 535              | 1.43±0.5 |

**Figure1-A comparison on FCR in ponds fed by extruder food and pellete food**

Attention to tables, diagram 1 and specified factors show that using extruder food has positive effects from all aspects. FCR in fish fed by extruder food was significantly higher than fish fed by pellete food ( $P < 0.05$ ). Factors like food quality, environmental factors, feeding management and fish health are effective on FCR. Using extruded food has a lot of benefits like fast growing in trout fed extruded food (Watabe and Pongmaneerat, 1993; Watabe and Pongmaneerat, 1993; Barrows *et al.*, 2008) degree of extrusion, not only starch degradability by amyloglucosidase but also performance of trout was improved (Pfeffer *et al.*, 1991). Extrusion significantly improves ADCs of dry matter (Cheng and Hardy, 2003), food digestibility (Sørensen *et al.*, 2002) and causes lower weight but higher feed efficiency (Hilton *et al.*, 1981). According to tables 8 and 9, it is clear that FCR obtained with consumed food had significant difference with each other (FCR of extruder food was 0.98 and FCR of pellete food was 1.43). Growth rate, food digestion and keeping nutritive materials in rainbow trout was studied in different temperatures. The more the temperature of extrusion, the more growth will be for fish (Sørensen *et al.*, 2005). If temperature is  $127^{\circ}\text{C}$ , FCR will be more than the time temperature is  $93^{\circ}\text{C}$  (Barrows *et al.*, 2007). Extrusion process will rise the value of protein materials and causes better digestion of proteins (Sørensen, 2007). In the present study, ponds fed by extruder food, had less mortality. But it is very important to consider this point in rainbow trout farms that these results are obtained in this farm condition and with food

made in mentioned date. So culturists should choose the most qualified and economical food with exact calculation of FCR in 15 day periods (Experience has shown that this period is suitable) to obtain the best economic efficiency.

## References

- Al-Ruqaie, I. M., 2007.** Extruded Leftover Food as Animal Feed: I. Effect of Extruded feed on Growth and Feed Utilization of Tilapia (*Oreochromis niloticus*) in Saudi Arabia. *Pakistan Journal of Biological Sciences*, 10 (19), 3248-3253.
- Barrows, F. T., Stone, D. A. J. and Hardy, R. W., 2007.** The effects of extrusion conditions on the nutritional value of soybean meal for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 265, 244-252.
- Barrows, F. T., Gaylord, T. G., Sealey, W. M., Porter, L. and Smith, C. E., 2008.** The effect of vitamin premix in extruded plant-based and fish meal based diets on growth efficiency and health of rainbow trout, *Oncorhynchus mykiss*. *Aquaculture*, 283(1-4), 148-155.
- Boccignone, M., Forneris, G. and Palmegiano, G. B., 1989.** Use of Extruded Maize in Rainbow Trout Feed Improves Growth Rate and Quality. *Aquacultural Engineering*, 8(2), 139-145.
- Chebbaki, K., Akharbach, H., Talbaoui, E., Abrehouch, A., Ait ali, A., Sedki, S., Ben Bani, A. and Idaomar, M., 2010.** Effect of fish meal replacement by protein sources on the extruded and pressed diet of European sea bass juvenile (*Dicentrarchus labrax*). *Agriculture and Biology Journal of North America*, 1(4), 704-710.

- Cheng, Z. J. and Hardy, R. W., 2003.** Effects of extrusion processing of feed ingredients on apparent digestibility coefficients of nutrients for rainbow trout (*Oncorhynchus mykiss*). *Aquaculture Nutrition*, 9(2), 77-83.
- Degaura, S. and San Lucjan, F., 1997.** Evaluation of different pressed and extruded fish meal based diets on the growth of gilthead sea bream, *Sparus aurata* L. *Feeding Tomorrow's Fish*, 5(3), 123-139.
- Hilton, J. W., Cho, C. Y. and Slinger, S. J., 1981.** Effect of extrusion processing and steam pelleting diets on pellet durability, pellet water absorption, and the physiological response of rainbow trout (*Salmo gairdneri* R.). *Aquaculture*, 25(2-3), 185-194.  
<http://www.instrumentationguide.com/utilities/Conductivity/conductivity.pdf>
- Pfeffer, E., Beckmann-Toussaint, J., Henrichfreise, B. and Jansen, H. D., 1991.** Effect of extrusion on efficiency of utilization of maize starch by rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 96(3-4), 293-303.
- Pourjafar, A., 2007.** A comprehensive guide to grow rainbow trout. Parto Vaghe'eh publications. Tehran. Iran. 106P.
- Sadeghi, N. and Maleki, A., 2002.** A guide to grow rainbow trout. Naghshe Mehr publications. Tehran. Iran. 48P.
- Sørensen, M., Ljøkjel, K., Storebakken, T., Shearer, K. D. and Skrede, A., 2002.** Apparent digestibility of protein, amino acids and energy in rainbow trout (*Oncorhynchus mykiss*) fed a fish meal based diet extruded at different temperatures. *Aquaculture*, 211(1-4), 215-225.
- Sørensen, M., Storebakken, T. and Shearer, K. D., 2005.** Digestibility, growth and nutrient retention in rainbow trout (*Oncorhynchus mykiss*) fed diets at two different temperatures. *Aquaculture Nutrition*, 11, 251-256.
- Sørensen, M., 2007.** Ingredient formulation and extrusion processing parameters interfere with nutritional and physical quality of aqua feeds. Feed Technology Update, Oct-2007.
- Watanabe, T. and Pongmaneerat, J., 1993.** Potential of Soybean Meal as a Protein Source in Extruded Pellets for Rainbow Trout. *Nippon Suisan Gakkaishi*, 59(8), 1415-1423.