
Phylogenetic relationships of Iranian Infectious Pancreatic Necrosis Virus (IPNV) based on deduced amino acid sequences of genome segment A and B cDNA

Dadar M.^{1*}; Peyghan R.¹; Rajabi-Memari H.²; Seifi Abad Shapouri M.R.³;
Hasanzadeh R.⁴; Moazzami Goudarzi L.⁴; Vakharia V. N.⁵

Received: July 2013

Accepted: May 2014

Abstract

Infectious Pancreatic Necrosis Virus (IPNV) is the causal agent of a highly contagious disease that affects many species of fish and shellfish. This virus causes economically important diseases of farmed rainbow trout, *Oncorhynchus mykiss*, in Iran which is often associated with the transmission of pathogens from European resources. In this study, moribund rainbow trout fry were collected during an outbreak of IPNV in three different fish farms in one northern province (Mazandaran), and two west provinces (Chaharmahal and Bakhtiari, and Kohgiluyeh and Boyer Ahmad) of Iran. We investigated full genome sequence of Iranian IPNV and compared it with previously identified IPNV sequences. The sequences of different structural and non-structural protein genes were compared with other aquatic birnaviruses sequenced to date. Our results showed that the Iranian isolate fall within genogroup 5, serotype A2 strain SP, having 99 % identity with the strain 1146 from Spain. These results suggest that the Iranian isolate may have originated from Europe.

Keywords: Molecular characterization, IPNV, Virus, Aquatic birnaviruses, Rainbow trout, Iran

1- Department of Aquatic Animal Health, College of Veterinary Medicine, Shahid Chamran University, Ahvaz, Iran

2- Center of Biotechnology Research, College of Agriculture, Shahid Chamran University, Ahvaz, Iran

3-Center of Virology Research, College of Veterinary Medicine, University of Shahid Chamran University, Ahvaz, Iran

4- Iranian Veterinary Organization, Central Veterinary Laboratory, Tehran, Iran

5- Institute of Marine and Environmental Technology, University of Maryland, Baltimore County, Baltimore, USA

*Corresponding author email: dadar.m77@gmail.com

Introduction

Infectious Pancreatic Necrosis (IPN) is one of the most important viral diseases of farmed salmonid fish caused by Infectious Pancreatic Necrosis Virus (IPNV). IPNV is a small, non-enveloped virus belonging to the family Birnaviridae, genus *Aquabirnavirus* (Dobos, 1995; Song *et al.*, 2005). Serologically, *Aquabirnaviruses* have been classified on the basis of cross-neutralization assays and divided into four serogroups A to D (Ruane *et al.*, 2009; Mutoloki and Evensen, 2011). There are nine distinct IPNVs in the A serogroup and only one serotype within B serogroup (Hill and Way, 1995). Most aquatic birnaviruses belong to serogroup A, which include 9 serotypes (A1 to A9), while the minor serogroup B consist of a single serotype, B1 (Caswell-Reno *et al.*, 1989; Bowers *et al.*, 2008). Serotype A1 include USA isolates, serotypes A6-A9 are mainly in Canada, and serotypes A2-A5 and B1 are found in Europe and Asia (Blake *et al.*, 2001). Moreover, genogroups I–VI correlate with the serotypes A1–A9 and B1 as follows: genogroup I (A9 and A1), genogroup II(A3), genogroup III (A2 and B1), genogroup IV (A5 and A6), genogroup V (A7 and A8) and genogroup VI (A4) (Blake *et al.*, 2001; Nishizawa *et al.*, 2005; Romero-Brey *et al.*, 2009). These viruses show significant antigenic variation (Heppell *et al.*, 1995; Ruane *et al.*, 2009). The genome of virus has two segments of double-stranded RNA that are surrounded with a single shelled, icosahedral capsid with 60 nm in diameter (Dobos and Rowe, 1977; Macdonald and Dobos, 1981; Dobos, 1995). Segment A is

3097 bp long and encodes four viral proteins, namely structural proteins VP2 and VP3, and nonstructural proteins VP4 and VP5 (Dobos, 1995). Segment A contains a large open reading frame (ORF) that encodes a 106-KDa polyprotein which is cotranslationally cleaved by VP4 to produce pre-VP2 (pVP2) and VP3 (Dobos and Roberts, 1983; Galloux *et al.*, 2004). There is a small ORF which overlaps with the amino terminal end of the large ORF and generates a 15-KDa (VP5) nonstructural polypeptide (Dobos, 1995; Saint-Jean *et al.*, 2003). VP5 contains Bcl-2 homology domains which is capable of enhancing cell viability with a notable strategy via VP5 to regulate the host anti-apoptosis pathway (Hong *et al.*, 2002). VP2 is an outer capsid protein which contains major neutralizing epitopes of the virus that gives protection. It also contains the markers for virulence and has a particular taxonomic importance for genotyping (Labus *et al.*, 2001; Moon *et al.*, 2004; Das *et al.*, 2007). VP3 is an internal protein with several roles in organizing the IPNV replication cycle (Pedersen *et al.*, 2007; Chiu *et al.*, 2010; Bahar *et al.*, 2013). Segment B is 2784 bp long and encodes VP1 protein, a minor internal polypeptide (94-KDa), which acts as the virion associated RNA-dependent RNA polymerase (RdRP) of IPNV (Duncan *et al.*, 1987; Dobos, 1995). IPN disease can induce high mortality which can result in huge economic loss in both fry and juveniles of rainbow trout, brook trout and Atlantic salmon (Wolf *et al.*, 1968; Skjesol *et al.*, 2011). This virus is widespread in salmonid hatcheries from

America to Europe, Asia, Australia and South Africa (Crane *et al.*, 2000; Davies *et al.*, 2010). Fish that survive an IPNV infection may become carriers of the virus for long period and sequentially transmit the virus to other susceptible species of fish and shellfish (Munro and Midtlyng, 2011). The mortality of an outbreak alters significantly with species, age, environmental condition, physical situation and virulence of the viral strain (Song *et al.*, 2005; Dadar *et al.*, 2013; Salgado-Miranda *et al.*, 2014). Rainbow trout is one of the most favorable species for rearing and its farming is a promising industry in Iran (Dadar *et al.*, 2013). For the first time, IPNV was detected using RT-PCR method in several provinces of Iran in 2007, followed by other reports (Akhlaghi and Hosseini, 2007; Raissy *et al.*, 2010; Oryan *et al.*, 2012; Ahmadi *et al.*, 2013). Therefore, the aim of the present study was to determine IPNV genotype (s) in Iran and compare it with known genotypes of European and American IPNV isolates. The approach taken was to sequence the coding regions of the VP1, VP2, VP3, VP4 and VP5 genes.

Materials and Methods

Fish sampling

During a period of IPN prevalence from 2010 to 2012, IPNV was isolated from rainbow trouts of Iranian farms. Moribund rainbow trout fry were collected during an outbreak of IPNV in different fish farms in North and West Iran in provinces of Mazandaran, Chaharmahal and Bakhtiari, and Kohgiluyeh and Boyer Ahmad. The

farms were run on flow-through system of fresh water with a temperature range of 12 to 15°C. From each farm, 30 moribund fish were selected and transferred to Central Veterinary Laboratory, Tehran, Iran. Virus isolation from fry samples with disease clinical signs, such as darkening of the skin, abdominal swelling, cast-like pseudofaeces and loss of appetite was performed, according to the procedure described by the OIE with minor modification (Crane *et al.*, 2000; Matvienko *et al.*, 2014).

Virus isolation

Each pool contained material from ten fry fishes. Briefly, 2 g of specimen were homogenized in approximately 2ml of minimum essential medium (MEM, Sigma, St. Louis, Missouri, USA), and centrifuged at 3000x g for 10 min. Supernatant was used directly for cell culture inoculation. Chinook salmon embryo (CHSE-214] cells were cultured in MEM containing Earle's salts, L-glutamine, 25 mM HEPES, 10 % fetal bovine serum (FBS), 100 ng/ml of streptomycin sulphate, 100 IU penicillin G., and incubated at 20 °C, up to 70-90 % confluence in 24-well plates before inoculation. Then they were inoculated with 200 microliters of 1:10 and 1:100 dilution of each prepared sample in parallel wells containing CHSE and incubated at 15°C. At 7 days post inoculation (dpi), the cultures were observed for cytopathic effects (CPE) and if they were negative, the cultures were used for the second passage. After subjecting to 1 freeze/thaw cycle, the cell culture lysates from 2 dilutions of each

sample were pooled and centrifuged at 2000 x g for 5 min. Then the fresh CHSE cells were inoculated with the pooled first passage supernatant. Plates were incubated at 15°C and monitored daily up to 21 dpi, for development of viral CPE. If no CPE was observed after a period of 21 dpi, the sample was recorded as negative for IPNV. When CPE was observed, IPNV was confirmed by using an IPNV antigen (Ag) ELISA kit (BIO-X, Jemelle, Belgium), and the culture medium was removed and stored in -80 °C.

ELISA for IPNV antigen

A sandwich ELISA-based Bio-X diagnostics kit (Belgium) was used to confirm the presence of IPNV in cell cultures exhibiting CPE, according to the manufacturer's protocols. Optical density was read by means of a microplate spectrophotometer using a 450 nm filter. Each plate contained positive and negative controls. Positive sample had an A450 at least twice that of the negative control.

RNA extraction

When a sample was identified as IPNV positive, the media was removed and subjected to RNA extraction, using a Roche high pure viral kit (Roche®, Mannheim, Germany), according to manufacturer's recommendations. Concentration and purity of the RNA was obtained and estimated by measuring absorbance at 260 and 280nm in a spectrophotometer (Nanodrop® spectrophotometer ND-1000, Germany).

RT-PCR and sequencing

Since there was adequate information about the Iranian IPNV isolate genome sequence, the existing IPNV sequences

were extracted from NCBI and aligned with Mega5 software (Kumar *et al.*, 2008; Mutoloki and Evensen, 2011; Tamura *et al.*, 2011). Specific primers were designed according to the conserved region in VP1, VP2, VP3, VP4 and VP5 genes. The extracted RNA was amplified using a one-step reverse transcriptase-polymerase chain reaction (RT-PCR) kit in accordance with the manufacturer's protocol (Qiagen, Germany). The primers, including FVP2 (5' ATGAACACAAACAAGGCAAC 3') and RVP2 (5' GACTATGTCTCTCCAGCCCCATGC3') for VP2; FVP3 (5' GCATCCGGGATGGACGAGGA 3') and RVP3 (5' TTACACCTCAGCGTTGTCTC C3') for VP3; FVP4 (5' GGACCAGAGTCTTCAACGAAATC ACG3') and RVP4 (5' TAGATCTCGGCGTCCTGGACTTC3') for VP4, were used to amplify the genes from segment A. Segment B was amplified as two overlapping fragments using specific primers, including FVP1A (5'- ATGTCGGACATCTTCAAYTCACC-3') and RVP1A (5'- GAGCCGTCCTCGTTTGTCCA-3') for the first half segment of VP1 and FVP1B (5'- CACATGCAGGCAATGATGTACTAC3) and RVP1B (5'CTTAGTTTCTTCTCTGCTTCTC-3') for second half segment of VP1. The thermal RT- PCR steps were 1 cycle 50°C for 30 min, 95°C for 15 min, followed by 35 cycles of denaturation at 94°C for 30 sec, annealing temperature at 55°C for 30 sec and extension at 72°C for 1 min for VP3 and 70 sec for VP2. This was

followed by a final extension of 10 min at 72°C. The RT-PCR products were separated by agarose gel electrophoresis, purified with gel extraction kit (Qiagen), and subjected to nucleotide sequence analysis (Bioneer).

Phylogenetic analysis

The sequence data were analyzed using Blast (NCBI), Mega 5 (Kumar *et al.*, 2008; Mutoloki and Evensen, 2011; Tamura *et al.*, 2011) and Vector NTI (Invitrogen) softwares. Finally, the results were compared with other existing sequences in the data bank and phylogenetic trees were drawn with the UPGMA method.

Results

The Iranian IPNV isolates were separated from samples after the second passage on CHSE-214 cell line. The virus CPE was the formation of spindle-shaped cells and pyknosis of nuclei that typically appear in 5-10 days after inoculation. Isolation of the virus in cell culture was confirmed by IPNV antigen ELISA Kit (Bio-X[©] diagnostics kit, Belgium). The specific primer pair for VP1, VP2, VP3 and VP4 amplified the full length of genes from extracted viral RNA successfully. The reactions amplified fragments of 1347 bp, 852 bp, 2535 bp, 402 bp and 720 bp for

VP2, VP2-VP4, VP1, VP5 and VP3 as expected, respectively (Blake *et al.*, 2001). The amplified fragments were sequenced and deposited in the NCBI database with accession numbers VP1: KC900161, VP2: KC489465, VP3: KC489466, VP4: KC710379, VP5: KC900222 and poly protein: KF279643. BlastP alignment comparison of VP1, VP2, VP3 and poly protein amino acid sequences showed that the Iranian isolate of IPNV was closely similar to Sp strain. Amino acid sequences of VP2, VP3, VP1 and polyprotein of Iranian strain and the sequences retrieved from the GenBank were aligned by Mega 5 software and phylogenetic trees were constructed, as shown in Figs. 1-4. The results of this analysis confirmed that the Iranian isolate was of Sp strain and had the highest similarity to isolate 1146 (Q8JK08). In Fig. 5, one can see the presence of a hyper variable region between amino acid 245-257. In VP2 protein, 11 amino acid substitutions were observed at positions 52, 94, 96, 219, 245, 248, 252, 255, 257, 286 and 321 (Fig. 5). In VP3 protein, 5 changes were noted at positions 32, 79, 122, 218 and 235 (Fig. 6). For VP5 protein, we observed 11 amino acid substitutions at positions 6, 11, 13, 29, 36, 54, 65, 68, 97, 108 and 132 (Fig. 7).




Figure1: Phylogenetic analysis of the VP2 protein of selected IPNV strains.




Figure 2: Phylogenetic analysis of the polyprotein of selected IPNV strains.




Figure 3: Phylogenetic analysis of the VP3 protein of selected IPNV strains.




Figure 4: Phylogenetic analysis of the VP1 protein of selected IPNV strains.

1	MNTNKATATYLKSI MLPETGPASIPDDITERHILKQETSSYNLEVSSESGSGLVCFPGAP	60	E7E176	E7E176_9VIRU
1	MNTNKATATYLKSI MLPETGPASIPDDITERHILKQETSSYNLEVSSESGSGLVCFPGAP	60	Q8JK08	Q8JK08_9VIRU
1	MNTNKATATYLKSI MLPETGPASIPDDITERHILKQETSSYNLEVSSESGSGLVCFPGAP	60	Q703G9	POLS_IPNVS
1	MNTNKATATYLKSI MLPETGPASIPDDITERHILKQETSSYNLEVSSESGSGLVCFPGAP	60	E7E192	E7E192_9VIRU
1	MNTNKATATYLKSI MLPETGPASIPDDITERHILKQETSSYNLEVSSESGSGLVCFPGAP	60	20130519603WJS300U	
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	E7E177	E7E177_9VIRU
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	E7E176	E7E176_9VIRU
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	Q8JK08	Q8JK08_9VIRU
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	Q703G9	POLS_IPNVS
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	E7E192	E7E192_9VIRU
61	GSRIGAHYRWNANQTGLEFDQWLETSQDLKKAFFNYGRLISRKYDIQSSTLPAGLYALNGT	120	20130519603WJS300U	
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	E7E177	E7E177_9VIRU
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	E7E176	E7E176_9VIRU
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	Q8JK08	Q8JK08_9VIRU
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	Q703G9	POLS_IPNVS
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	E7E192	E7E192_9VIRU
121	LNAATFEGSLSEVESLTYNLSLSLTTNPQDKVNNQLVTRKGVTVLNLPTGFDRFPYVRELEDE	180	20130519603WJS300U	
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	E7E177	E7E177_9VIRU
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	E7E176	E7E176_9VIRU
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	Q8JK08	Q8JK08_9VIRU
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	Q703G9	POLS_IPNVS
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	E7E192	E7E192_9VIRU
181	TPOGLQSMNGAKMRCTAAIAPRRYEIDLPSQRLPFPATITITLTYEGNADIVNSTTVTG	240	20130519603WJS300U	
241	DINPFLAGQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	E7E177	E7E177_9VIRU
241	DINPFLAGQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	E7E176	E7E176_9VIRU
241	DINPFLAQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	Q8JK08	Q8JK08_9VIRU
241	DINPFLAEQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	Q703G9	POLS_IPNVS
241	DINPFLAEQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	E7E192	E7E192_9VIRU
241	DINPFLAQPADETFHFQDLDFMGLDNDVPPVTVVSSVLAINDNYRGSVSAKMTQSIPTEN	300	20130519603WJS300U	
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	E7E177	E7E177_9VIRU
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	E7E176	E7E176_9VIRU
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	Q8JK08	Q8JK08_9VIRU
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	Q703G9	POLS_IPNVS
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	E7E192	E7E192_9VIRU
301	ITKPIITRVKLSYKINQQTAINVATLGTMGPASVSFSSGNGNVPGLRPIITLVAYEKMTF	360	20130519603WJS300U	
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	E7E177	E7E177_9VIRU
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	E7E176	E7E176_9VIRU
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	Q8JK08	Q8JK08_9VIRU
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	Q703G9	POLS_IPNVS
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	E7E192	E7E192_9VIRU
361	LSILTVAGVSNYELIPNPELLKNMVTRYGKYDPEGLNYAKMILSHREELDIRTVWRTEEY	420	20130519603WJS300U	
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	480	E7E177	E7E177_9VIRU
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	480	E7E176	E7E176_9VIRU
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	480	Q8JK08	Q8JK08_9VIRU
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	480	Q703G9	POLS_IPNVS
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	480	E7E192	E7E192_9VIRU
421	KERTRVFNEITDFSSDLPTS KAWGWRDIVRGIRKVAAPVLSLTFPMAAPLIGMADQFIGD	449	20130519603WJS300U	

Figure 5: Five high similarity hits for blastP on UNIPROTKB of VP2 sorted by descending score. It shows changes in eleven positions at residues 52, 94, 96, 219, 245, 248, 252, 255, 257, 286 and 321. As it can be seen, there is an hypervariable region between residues 245-257. Arrow shows the VP2 sequence of Iranian IPNV isolate in this study.

1	-----ASGMDEELQRLLNATMARAKEVQDAEYKLLRLMAWTRKNDLTDHMY	47	20130519403VIEY8JF	←
721	KRIKYLGEIMASNASGMDEELQRLLNATMARAKEVQDAEYKLLKLMWTRKNDLTDHMY	780	Q8JK03 Q8JK03_9VIRU	
721	KRIKYLGEIMASNASGMDEELQRLLNATMARAKEVQDAEYKLLKLMWTRKNDLTDHMY	780	G9C3Q6 G9C3Q6_9VIRU	
721	KRIKYLGEIMASNASGMDEELQRLLNATMARAKEVQDAEYKLLRLMAWTRKNDLTDHMY	780	A42XA0 A42XA0_9VIRU	
721	KRIKYLGEIMASNASGMDEELQRLLNATMARAKEVQDAEYKLLRLMAWTRKNDLTDHMY	780	Q8JK08 Q8JK08_9VIRU	
*****;				
48	ENSKEDPDALKFGKLISTPPKHPEKPKGPDQHHAEARATRISLDAVRAGADFATPEWVA	107	20130519403VIEY8JF	
781	ENSKEDPDALKFGKLISTPPKHPEKPKGPDQHHAEARATRISLDAVRAGADFATPEWVA	840	Q8JK03 Q8JK03_9VIRU	
781	ENSKEDPDALKFGKLISTPPKHPEKPKGPDQHHAEARATRISLDAVRAGADFATPEWVA	840	G9C3Q6 G9C3Q6_9VIRU	
781	ENSKEDPDALKFGKLISTPPKHPEKPKGPDQHHAEARATRISLDAVRAGADFATPEWVA	840	A42XA0 A42XA0_9VIRU	
781	ENSKEDPDALKFGKLISTPPKHPEKPKGPDQHHAEARATRISLDAVRAGADFATPEWVA	840	Q8JK08 Q8JK08_9VIRU	
*****;				
108	LNNYRGSPGQFKYYLITGREPEPGDEYDYIKQPIVKPTDMNKIRRLANSVYGLPHQEP	167	20130519403VIEY8JF	
841	LNNYRGSPGQFKYYLITGREPEPGDEYDYIKQPIVKPTDMNKIRRLANSVYGLPHQEP	900	Q8JK03 Q8JK03_9VIRU	
841	LNNYRGSPGQFKYYLITGREPEPGDEYDYIKQPIVKPTDMNKIRRLANSVYGLPHQEP	900	G9C3Q6 G9C3Q6_9VIRU	
841	LNNYRGSPGQFKYHLITGREPEPGDEYDYIKQPIVKPTDMNKIRRLANSVYGLPHQEP	900	A42XA0 A42XA0_9VIRU	
841	LNNYRGSPGQFKYYLITGREPEPGDEYDYIKQPIVKPTDMNKIRRLANSVYGLPHQEP	900	Q8JK08 Q8JK08_9VIRU	
*****;				
168	AFEEFYDAVAAVFAQNGGRGPDQDQMDLRELARQMKRRPRNADAPRTRAPAEPAFPGR	227	20130519403VIEY8JF	
901	AFEEFYDAVAAVFAQNGGRGPDQDQMDLRELARQMKRRPRNADAPRTRAPAEPAFPGR	960	Q8JK03 Q8JK03_9VIRU	
901	AFEEFYDAVAAVFAQNGGRGPDQDQMDLRELARQMKRRPRNADAPRTRAPAEPAFPGR	960	G9C3Q6 G9C3Q6_9VIRU	
901	AFEEFYDAVAAVFAQNGGRGPDQDQMDLRELARQMKRRPRNADAPRTRAPAEPAFPGR	960	A42XA0 A42XA0_9VIRU	
901	AFEEFYDAVAAVFAQNGGRGPDQDQMDLRELARQMKRRPRNADAPRTRAPAEPAFPGR	960	Q8JK08 Q8JK08_9VIRU	
*****;				
228	SRFTPSGDNAEV	239	20130519403VIEY8JF	
961	SRFTPSGDNAEV	972	Q8JK03 Q8JK03_9VIRU	
961	SRFTPSGDNAEV	972	G9C3Q6 G9C3Q6_9VIRU	
961	SRFTPSGDNAEV	972	A42XA0 A42XA0_9VIRU	
961	SRFTPSGDNAEV	972	Q8JK08 Q8JK08_9VIRU	
*****;				

Figure 6: Four high similarity hits for blastP on UNIPROTKB of VP3 sorted by score descending. It shows changes in five positions at residues 32, 79, 122, 218 and 235. Arrow shows the VP3 sequence of Iranian IPNV in this study.

1	-----MQDEHKQGNRNLEIHYASRDWTSKHPGRHNGETHLKTDRDLVIQL	45	20130522276M920EB7	←
1	-----MQDEHKQGNRNLEIHYASRDWTSKHPGRHNGETHLKTDRDLVIQL	45	Q82744 Q82744_9VIRU	
1	-----MQDEHKQGNRNLEIHYASRDWTSKHPGXHNGETHXKTDRDLVIQL	45	G9C3Q5 G9C3Q5_9VIRU	
1	MAKALSNDKQTNLNLYSIQDEHKQGNRNLEIHYASRDWTSKHPGRHNGETHPKTRDLVIQ	60	P22496 VP5_IPNVN	
1	-----MQDEHKQGNRNLEIHYASRDWTSKHPGRHNGETHLKTDRDLVIQL	45	Q6U2P6 VP5_IPNVS	
1	MAKALSNDKQTNLNLYSMQDEHKQGNRNLEIHYASRDWTSKHPGRHNGETHLKTDRDLVIQL	60	Q8JK02 Q8JK02_9VIRU	
1	-----MQDEHKQGNRNLEIHYASRDWTSKHPGXHNGETHXKTDRDLVIQL	45	G9C3Q7 G9C3Q7_9VIRU	
1	-----QTNLNLYSIQDEHTQGNRDLEIHYASRDWTSKHPGRHNGETHPKTRDLVIQL	52	Q8JK00 Q8JK00_9VIRU	
*****;				
46	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	105	20130522276M920EB7	
46	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	105	Q82744 Q82744_9VIRU	
46	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	105	G9C3Q5 G9C3Q5_9VIRU	
61	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	120	P22496 VP5_IPNVN	
46	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	105	Q6U2P6 VP5_IPNVS	
61	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	120	Q8JK02 Q8JK02_9VIRU	
46	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	105	G9C3Q7 G9C3Q7_9VIRU	
53	RGLRIRKWHNSCLFPWGTRRLTDRCTLQMECEPDGAGVRPVAGDVAGPEESLQREADLKEI	112	Q8JK00 Q8JK00_9VIRU	
*****;				
106	RHPKLHTGRSLCSERDAQRCHLRRQSV	133	20130522276M920EB7	
106	RHPKLHTGRSLCSERDAQRCHLRRQSV	133	Q82744 Q82744_9VIRU	
106	RHPKLHTGRSLCSERDAQRCHLRRQSV	133	G9C3Q5 G9C3Q5_9VIRU	
121	RHPKLHTGRSLCSERDAQRCHLRRQSV	148	P22496 VP5_IPNVN	
106	RHPKLHTGRSLCSERDAQRCHLRRQSV	133	Q6U2P6 VP5_IPNVS	
121	RHPKLHTGRSLCSERDAQRCHLRRQSV	148	Q8JK02 Q8JK02_9VIRU	
106	RHPKLHTGRSLCSERDAQRCHLRRQSV	133	G9C3Q7 G9C3Q7_9VIRU	
113	RHPKLHTGRSLCSERDAQRCHLRRQSV	140	Q8JK00 Q8JK00_9VIRU	
*****;				

Figure 7: Seven high similarity hits for blastP on UNIPROTKB of VP5 sorted by descending score. It shows changes in eleven positions at residues 6, 11, 13, 29, 36, 54, 65, 68, 97, 108 and 132. Arrow shows the VP5 sequence of Iranian IPNV in this study.

Discussion

Infectious Pancreatic Necrosis (IPN) can cause an important economic impact on salmonids industry and it is considered as a threat for developing this industry. In Iran, this disease was detected for the first time in Fars Province in south of Iran (Akhlaghi and Hosseini, 2007) and spread in many parts of the country in the last few years (Akhlaghi and Hosseini, 2007; Raissy *et al.*, 2010; Oryan *et al.*, 2012; Ahmadi *et al.*, 2013; Dadar *et al.*, 2013). Investigations of Iranian isolates were limited to partial sequence of the VP2 encoding region in the segment A. VP2 is the major capsid protein, which contains all neutralizing epitopes and cell attachment sites that determines host and cell range (Caswell-Reno *et al.*, 1989; Davies *et al.*, 2010; Salgado-Miranda *et al.*, 2014). VP2 is also responsible for the production of type-specific monoclonal antibodies (Caswell-Reno *et al.*, 1989; Nicholson, 1993; Melby and Christie, 1994). The residues of VP2 domains can alter the properties of this protein. This alterations could influence the antigenic characteristics of VP2 and mortality rates in fish (Shivappa *et al.*, 2004). In conclusion, VP2 carries the determinant factors for IPNV virulence (Song *et al.*, 2005). In virulent strains of IPNV, there is threonine and alanine at positions 217 and 221 of VP2, respectively; whereas, moderate to low virulence strains have a proline and alanine at these positions. Strains with threonine at position 221 have been shown to be almost a virulent (Santi *et al.*, 2005; Ruane *et al.*, 2009; Skjesol *et al.*, 2011). Bain *et al.* (2008) demonstrated

that strains with a proline and alanine in positions 217 and 221 respectively, indicate high virulence in the field and experimental conditions, supporting the suggestion that viral, host and environmental factors as well as specific amino acid residues influence pathogenicity. The residue at position 247 of VP2 is also highly variable and may be linked to virulence. Santi and colleagues (2005) showed that the motif Thr217, Ala221, Thr247 was associated with high virulence and the motif Pro217 Ala221 Ala247 is present in viruses with low and moderate virulence. So far, all of detected Iranian isolates had proline and threonine at positions 217 and 221, respectively. Despite the presence of threonine at the position 221 which is indicative of a non-virulent nature, the mortality of Iranian rainbow trout fry corresponds to a moderate virulence (Raissy *et al.*, 2010; Oryan *et al.*, 2012; Ahmadi *et al.*, 2013). The moderate virulence of Iranian IPNV isolates may be related to the alanine residue present in position 247. The VP2 has also been shown to contain the central variable domain that encodes two hypervariable segments. These hypervariable regions determine the virus specific serotypes (Heppell *et al.*, 1995). High similarity hits for blastP on UNIPROTKB of VP2 shows changes in 11 residues at positions 52, 94, 96, 219, 245, 248, 252, 255, 257, 286 and 321, most of which are present in the second hypervariable region between residues 245-257 of the Iranian isolate (Fig. 5).

The first identified Iranian IPNV was similar to Ab strain (Akhlaghi and

Hosseini, 2007) but the present study reports isolation of a virus closely related to SP strain. To study phenotypic and molecular characterization of IPNV, isolates were separated from field outbreaks of IPN in Iran from 2010 to 2012 and used to determine complete sequence of viral genome, including segments A and B. To the best of our knowledge, this is the first report of the full length nucleotide sequences of IPNV in Iran. The VP1, VP2, VP3 and whole polyprotein amino acid region shared the highest amino acid identity (99%) with the Sp isolate 1146 (Q8JK08_9VIRU) (<http://www.uniprot.org/uniprot/Q8JK08>) from Spain with nucleotide ID: AJ489222 (<http://www.ncbi.nlm.nih.gov/nuccore/AJ489222>). The lowest amino acid identity (90%) was with Jasper serotype P22173 (RDRP_IPNVJ) (<http://www.uniprot.org/uniprot/P22173>) with nucleotide ID: AAQ75356 (<http://www.ebi.ac.uk/ena/data/view/AAQ75356>). A comparison of the amino acid sequences from the Iranian isolates showed that they were all nearly the same and all isolates belonged to SP strain.

The position of the start codon of the VP5 protein may vary (Davies *et al.*, 2010). It has been shown that the start codon of VP5 is located at position 68 (Magyar and Dobos, 1994), although Heppell *et al.* (1995) reported that it could be initiated from position 68 or 112. Also Weber (2001) and Shivappa *et al.* (2004) demonstrated that the second in frame methionine codon is responsible for the initiation of VP5 in VR299 and SP strains, respectively. The only known IPNV which

shows to be free of ORF for VP5 is HE strain (Heppell *et al.*, 1995). Alignment of VP5 sequences indicated that Iranian isolates have start codon in position 112 and the second methionine in frame is responsible for the initiation of Iranian isolate (Fig. 7). Since virulence of IPNV isolates have been connected to segment A (Sano *et al.*, 1981), no specific sequences or motifs have been identified in segment B that are linked to virulence. It is demonstrated that all studied pathogenic isolates encoded a truncated VP5 protein (Hong *et al.*, 2002; Santi *et al.*, 2005). In contrast, it is shown that all Australasian strains also contain a truncated VP5 protein and these isolates were usually isolated from healthy fish (Davies *et al.*, 2010). This study indicated that Iranian isolates contain 133 amino acids that encode a truncated VP5 protein, which is similar to what was reported for pathogenic isolates (Hong *et al.*, 2002; Santi *et al.*, 2005).

As demonstrated by Blake *et al.* (2001), IPNV strains appear to cluster into 6 genogroups based on geographical and serological similarity. It is suggested that Iranian isolates fall within genogroup 5, Serotype A2 with moderate to low virulence and appear to be most closely related to the SP strains. In contrast to genogroup 1, which appear to contain virus isolated only from trout in North America (USA and Canada), genogroup 5; consists of isolates from more diverse geographic and host range (Davies *et al.*, 2010). The phylogenetic tree showed clearly that the viruses representing serotype A2 in Iran are more closely

related to European and Asian isolates (Serotypes A2, A3, A4 and A5) and Canada (Serotypes A6, A7 and A8) than isolates representing serotype A1 in the United States and Jasper strains. It is proposed that IPNV was introduced from European sources. In order to perform control of the disease in Iran, it is important to import eyed eggs and smolts which are IPNV-free. This availability of IPNV genome sequences will be useful in further studies such as diagnosis of disease, molecular epidemiology researches and developing native vaccines.

Acknowledgments

The authors are grateful to veterinary faculty of Chamran University for funding this work. Also would like to thank Iranian Veterinary Organization, especially Central Veterinary Laboratory for their assistance in this project.

References

- Ahmadi, N., Oryan, A., Akhlaghi, M. and Hosseini, A., 2013.** Tissue distribution of infectious pancreatic necrosis virus serotype Sp in naturally infected cultured rainbow trout, *Oncorhynchus mykiss* (Walbaum): an immunohistochemical and nested-PCR study. *Journal of Fish Diseases*, 36, 629–637.
- Akhlaghi, M. and Hosseini, A., 2007.** First report on the detection of infectious pancreatic necrosis virus (IPNV) by RT-PCR in rainbow trout fry cultured in Iran. *Bulletin-european association of fish pathologists*, 27, 205.
- Bahar, M.W., Sarin, L.P., Graham, S. C., Pang, J., Bamford, D.H., Stuart, D.I. and Grimes, J.M., 2013.** Structure of a VP1-VP3 complex suggests how birnaviruses package the VP1 polymerase. *Journal of Virology*, 87, 3229-3236.
- Bain, N., Gregory, A. and Raynard, R., 2008.** Genetic analysis of infectious pancreatic necrosis virus from Scotland. *Journal of Fish Diseases*, 31, 37-47.
- Blake, S., Ma, J., Caporale, D., Jairath, S. and Nicholson, B., 2001.** Phylogenetic relationships of aquatic birnaviruses based on deduced amino acid sequences of genome segment A cDNA. *Diseases of Aquatic Organisms*, 45, 89.
- Bowers, R.M., Lapatra, S.E. and Dhar, A.K., 2008.** Detection and quantitation of infectious pancreatic necrosis virus by real-time reverse transcriptase-polymerase chain reaction using lethal and non-lethal tissue sampling. *Journal of Virological Methods*, 147, 226-234.
- Caswell-Reno, P., Lipipun, V., Reno, P. and Nicholson, B., 1989.** Use of a group-reactive and other monoclonal antibodies in an enzyme immunodot assay for identification and presumptive serotyping of aquatic birnaviruses. *Journal of Clinical Microbiology*, 27, 1924-1929.
- Chiu, C.L., Wu, J.L., Her, G.M., Chou, Y.L. and Hong, J.R., 2010.** Aquatic birnavirus capsid protein, VP3, induces apoptosis via the Bad-mediated mitochondria pathway in

- fish and mouse cells. *Apoptosis*, 15, 653-668.
- Crane, M., Hardy-Smith, P., Williams, L.M., Hyatt, A.D., Eaton, L.M., Gould, A., Handlinger, J., Kattenbelt, J. and Gudkovs, N., 2000.** First isolation of an aquatic birnavirus from farmed and wild fish species in Australia. *Diseases of Aquatic Organism*, 43, 1-14.
- Dadar, M., Peyghan, R., Memari, H. R., Shapouri, M.R.S.A., Hasanzadeh, R., Goudarzi, L.M. and Vakharia, V.N., 2013.** Sequence analysis of infectious pancreatic necrosis virus isolated from Iranian reared rainbow trout (*Oncorhynchus mykiss*) in 2012. *Virus Genes*, 47, 574-578.
- Das, B., Collet, B., Snow, M. and Ellis, A., 2007.** Expression kinetics of ISG15 and viral major capsid protein (VP2) in Atlantic cod (*Gadus morhua* L.) fry following infection with infectious pancreatic necrosis virus (IPNV). *Fish & Shellfish Immunology*, 23, 825-830.
- Davies, K.R., Mccoll, K.A., Wang, L.F., Yu, M., Williams, L.M. and Crane, M.S.J., 2010.** Molecular characterisation of Australasian isolates of aquatic birnaviruses. *Diseases of Aquatic Organisms*, 93, 1-15.
- Dobos, P., 1995.** The molecular biology of infectious pancreatic necrosis virus (IPNV). *Annual Review of Fish Diseases*, 5, 25-54.
- Dobos, P. and Roberts, T., 1983.** The molecular biology of infectious pancreatic necrosis virus: a review. *Canadian Journal of Microbiology*, 29, 377-384.
- Dobos, P. and Rowe, D., 1977.** Peptide map comparison of infectious pancreatic necrosis virus-specific polypeptides. *Journal of Virology*, 24, 805-820.
- Duncan, R., Nagy, E., Krell, P.J. and Dobos, P., 1987.** Synthesis of the infectious pancreatic necrosis virus polyprotein, detection of a virus-encoded protease, and fine structure mapping of genome segment A coding regions. *Journal of Virology*, 61, 3655-3664.
- Galloux, M., Chevalier, C., Henry, C., Huet, J.C., Da Costa, B. and Delmas, B., 2004.** Peptides resulting from the pVP2 C-terminal processing are present in infectious pancreatic necrosis virus particles. *Journal of General Virology*, 85, 2231-2236.
- Heppell, J., Tarrab, E., Berthiaume, L., Lecomte, J. and Arella, M., 1995.** Characterization of the small open reading frame on genome segment A of infectious pancreatic necrosis virus. *Journal of General Virology*, 76, 2091-2096.
- Hill, B. and Way, K., 1995.** Serological classification of infectious pancreatic necrosis (IPN) virus and other aquatic birnaviruses. *Annual Review of Fish Diseases*, 5, 55-77.
- Hong, J.R., Gong, H.Y. and Wu, J.L., 2002.** IPNV VP5, a novel anti-apoptosis gene of the Bcl-2 family, regulates Mcl-1 and viral protein expression. *Virology*, 295, 217-229.

- Kumar, S., Nei, M., Dudley, J. and Tamura, K., 2008.** MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Briefings in Bioinformatic*, 9, 299-306.
- Labus, M.B., Breeman, S., Ellis, A.E., Smail, D.A., Kervick, M. and Melvin, W.T., 2001.** Antigenic comparison of a truncated form of VP2 of infectious pancreatic necrosis (IPN) virus expressed in four different cell types. *Fish & Shellfish Immunology*, 11, 203-216.
- Macdonald, R.D. and Dobos, P., 1981.** Identification of the proteins encoded by each genome segment of infectious pancreatic necrosis virus. *Virology*, 114, 414-422.
- Magyar, G. and Dobos, P., 1994.** Expression of infectious pancreatic necrosis virus polyprotein and VP1 in insect cells and the detection of the polyprotein in purified virus. *Virology*, 198, 437-445.
- Matvienko, N., Rud, Y. and Buchatsky, L., 2014.** Replication of Infectious Pancreatic Necrosis Virus in Different Cell Lines and in Rainbow Trout (*Oncorhynchus mykiss*) Fingerlings. *Archives of Polish Fisheries*, 22, 127-133.
- Melby, H. and Christie, K., 1994.** Antigenic analysis of reference strains and Norwegian field strains of aquatic birnaviruses by the use of six monoclonal antibodies produced against the infectious pancreatic necrosis virus N1 strain. *Journal of Fish Diseases*, 17, 409-415.
- Moon, C., Do, J., Cha, S., Bang, J.D., Park, M., Yoo, D., Lee, J., Kim, H., Chung, D. and Park, J., 2004.** Comparison of the immunogenicity of recombinant VP2 and VP3 of infectious pancreatic necrosis virus and marine birnavirus. *Archives of Virology*, 149, 2059-2068.
- Munro, E.S. and Midtlyng, P.J., 2011.** Infectious Pancreatic Necrosis and Associated Aquatic Birnaviruses. *Fish Diseases and Disorders: Volume 3: Viral, Bacterial and Fungal Infections*, 3, 1.
- Mutoloki, S. and Evensen, Ø., 2011.** Sequence similarities of the capsid gene of Chilean and European isolates of infectious pancreatic necrosis virus point towards a common origin. *Journal of General Virology*, 92, 1721-1726.
- Nicholson, B.L., 1993.** Use of monoclonal antibodies in identification and characterization of fish viruses. *Annual Review of Fish Diseases*, 3, 241-257.
- Nishizawa, T., Kinoshita, S. and Yoshimizu, M., 2005.** An approach for genogrouping of Japanese isolates of aquabirnaviruses in a new genogroup, VII, based on the VP2/NS junction region. *Journal of General Virology*, 86, 1973-1978.
- Oryan, A., Ahmadi, N., Akhlaghi, M. and Hosseini, A., 2012.** A comparative histopathological, immunohistochemical and nested-PCR study for diagnosis of infectious pancreatic necrosis in the naturally infected cultured rainbow trout

- (*Oncorhynchus mykiss*). *Aquaculture International*, 20, 725-734.
- Pedersen, T., Skjesol, A. and Jørgensen, J.B., 2007.** VP3, a structural protein of infectious pancreatic necrosis virus, interacts with RNA-dependent RNA polymerase VP1 and with double-stranded RNA. *Journal of Virology*, 81, 6652-6663.
- Raissy, M., Momtaz, H., Ansari, M., Moumeni, M. and Hosseinifard, M., 2010.** Distribution of Infectious Pancreatic Necrosis Virus (IPNV) in two major rainbow trout fry producing provinces of Iran with respect to clinically infected farms. *Journal of Food, Agriculture & Environment*, 8, 614-615.
- Romero-Brey, I., Bandin, I., Cutrín, J., Vakharia, V. and Dopazo, C., 2009.** Genetic analysis of aquabirnaviruses isolated from wild fish reveals occurrence of natural reassortment of infectious pancreatic necrosis virus. *Journal of Fish Diseases*, 32, 585-595.
- Ruane, N., Mccarthy, L., Swords, D. and Henshilwood, K., 2009.** Molecular differentiation of infectious pancreatic necrosis virus isolates from farmed and wild salmonids in Ireland. *Journal of Fish Diseases*, 32, 979-987.
- Saint-Jean, S.R., Borrego, J.J. and Perez-Prieto, S.I., 2003.** Infectious pancreatic necrosis virus: biology, pathogenesis, and diagnostic methods. *Advances in Virus Research*, 62, 113-165.
- Salgado-Miranda, C., Rojas-Anaya, E., García-Espinosa, G. and Loza-Rubio, E., 2014.** Molecular Characterization of the VP2 Gene of Infectious Pancreatic Necrosis Virus (IPNV) Isolates from Mexico. *Journal of Aquatic Animal Health*, 26, 43-51.
- Sano, T., Okamoto, N. and Nishimura, T., 1981.** A new viral epizootic of *Anguilla japonica* Temminck and Schlegel. *Journal of Fish Diseases*, 4, 127-139.
- Santi, N., Song, H., Vakharia, V.N. and Evensen, Ø., 2005.** Infectious pancreatic necrosis virus VP5 is dispensable for virulence and persistence. *Journal of Virology*, 79, 9206-9216.
- Shivappa, R., Song, H., Yao, K., Aas-Eng, A., Evensen, O. and Vakharia, V., 2004.** Molecular characterization of Sp serotype strains of infectious pancreatic necrosis virus exhibiting differences in virulence. *Diseases of Aquatic Organisms*, 61, 23-32.
- Skjesol, A., Skjæveland, I., Elnæs, M., Timmerhaus, G., Fredriksen, B.N., Jørgensen, S.M., Krasnov, A. and Jørgensen, J.B., 2011.** IPNV with high and low virulence: host immune responses and viral mutations during infection. *Virology Journal*, 8, 396.
- Song, H., Santi, N., Evensen, Ø. and Vakharia, V.N., 2005.** Molecular determinants of infectious pancreatic necrosis virus virulence and cell culture adaptation. *Journal of Virology*, 79, 10289-10299.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S.,**

2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2731-2739.

Weber, S., Fichtner, D., Mettenleiter, T. C. and Mundt, E., 2001. Expression of VP5 of infectious pancreatic necrosis virus strain VR299 is initiated at the second in-frame start codon. *Journal of General Virology*, 82, 805-812.

Wolf, K., Quimby, M., Carlson, C. and Bullock, G., 1968. Infectious pancreatic necrosis: selection of virus-free stock from a population of carrier trout. *Journal of the Fisheries Board of Canada*, 25, 383-391.