

## Effects of modified atmosphere packaging on microbiological load and physico-chemical properties of barramundi (*Lates calcarifer* Bloch) fillets at 8°C

Yassoralipour A.<sup>1,2\*</sup>; Bakar J.<sup>2</sup>; Rahman R.A.<sup>2</sup>;  
Fatimah A. Bakar<sup>2</sup>; Özogul F.<sup>3</sup>

Received: March 2014

Accepted: August 2015

### Abstract

Different modified atmosphere conditions with various CO<sub>2</sub> concentrations on microbial load (psychrophile, mesophile aerobic and anaerobic bacteria) and physico-chemical properties of barramundi (*Lates calcarifer* Bloch) fillets stored at 8°C were compared to determine the best packaging conditions. The gas conditions evaluated were 100% CO<sub>2</sub> (M<sub>1</sub>), 75% CO<sub>2</sub>/25% N<sub>2</sub> (M<sub>2</sub>), 50% CO<sub>2</sub>/50% N<sub>2</sub> (M<sub>3</sub>), 25% CO<sub>2</sub>/75% N<sub>2</sub> (M<sub>4</sub>) and 100% N<sub>2</sub> (control). High CO<sub>2</sub> concentration (M<sub>1</sub> and M<sub>2</sub>) of fish fillets delayed the psychrophilic bacteria growth compared to low CO<sub>2</sub> concentration (50% and 25% CO<sub>2</sub>) as well as the mesophilic, anaerobic and histamine forming bacteria. We concluded that the atmosphere with 75:25% and 100:0 (CO<sub>2</sub>:N<sub>2</sub>) had the most appropriate gas composition to inhibit the microbial growth and prolong the shelf life of barramundi fillets ( $p < 0.05$ ).

**Keywords:** Barramundi, CO<sub>2</sub> concentration, Psychrophile bacteria, Histamine forming bacteria, MAP, *Lates calcarifer*.

---

1- Current address: Faculty of Science, Department of Agricultural and Food Science, Universiti Tunku Abdul Rahman, 31900 Kampar, Perak, Malaysia.

2- Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 Serdang, Selangor D.E., Malaysia.

3- Faculty of Fisheries, Department of Seafood Processing Technology, Cukurova University, 01330 Balcali, Adana, Turkey.

\* Corresponding author's Email: [aliyas@utar.edu.my](mailto:aliyas@utar.edu.my)

## Introduction

Fish and fish products are some of the most important protein sources in human nutrition. An increasing number of seafood products are preserved by modified atmosphere packaging with refrigeration and addition of low levels of preservatives (Gram, 2009). Due to the increasing demand of consumers for fresh refrigerated fish products, a lot of research works have been conducted on preservation techniques to control the bacterial growth and the extending the shelf-life of the seafood products. Modified atmosphere packaging (MAP) and vacuum packaging (VP), along with refrigeration, have become increasingly popular preservation techniques, which have brought major changes in storage, distribution and marketing of raw and processed products to meet the consumers' demands (Brody, 1989).

Seafoods have susceptible chemical composition, which influence subsequent spoilage patterns. The presence of microorganisms on the live animal and contamination of raw materials during processing provides favorable conditions for microbial action and subsequent deterioration and spoilage of the products ( Bagge-Ravn *et al.*, 2003; Gram, 2009;).

The growth of microorganisms makes food organoleptically unacceptable for consumption because of changes in colour, smell, odour and texture. To reduce the rate of these changes, the growth of spoilage microorganisms should be inhibited by

application of MAP technology and increasing the lag phase of facultative and anaerobic microorganisms, which results in extension of the potential shelf life of MAP products. Nevertheless, there is a safety concern with the extended shelf life, because anaerobic and facultative pathogens that might have shorter lag times and faster growth rates are capable of growing during the extended storage period (Arashisar *et al.*, 2004; Ravi Sankar *et al.*, 2008; Gram, 2009).

The effect of MAP on lactic acid bacteria can vary depending on the type of product package. The increased CO<sub>2</sub> and decreased O<sub>2</sub> concentrations used in MAP generally favor the growth of lactic acid bacteria. Although the effect of MAP on yeasts is negligible, molds are aerobic microorganisms and therefore CO<sub>2</sub> can cause growth inhibition at concentrations as low as 10% (Molin, 2000). The most common gases used in MAP are carbon dioxide, oxygen and nitrogen ( Reddy *et al.*, 1994; Gimenez *et al.*, 2002). CO<sub>2</sub> concentration in MAP has been extended the shelf-life of foods by inhibiting the microbial growth of *Enterobacteria* and H<sub>2</sub>S-producing bacteria (López-Caballero *et al.*, 2002), histamine forming bacteria (Özogul and Özogul, 2006), *Pseudomonas spp.* (Erkan *et al.*, 2006) and *Lactobacillus sake* (Devlieghere *et al.*, 1998). Carbon dioxide inhibits the growth of psychrophilic microorganisms during the logarithmic phase and extends the lag phase. Oxygen stimulates the

growth of aerobic bacteria, but inhibits the growth of strictly anaerobic bacteria (Arashisar *et al.*, 2004). Nitrogen delays the oxidative rancidity and inhibits the growth of aerobic microorganisms by displacing the oxygen in packs (Farber, 1991). However, the synergetic effects of all these gases need to be established for the specific food application such as seafood.

The most effective CO<sub>2</sub> mixtures has been applied by various researchers for specific fish species such as pearlspot (Ravi Sankar *et al.*, 2008), sardine (Özogul *et al.*, 2002; Özogul *et al.*, 2004; Özogul and Özogul, 2006), tilapia (Reddy *et al.*, 1994), sole (López-Gálvez *et al.*, 1998), trout (100% CO<sub>2</sub>) (Arashisar *et al.*, 2004) and tuna (Emborg *et al.*, 2005). The CO<sub>2</sub> content is as low as 40% and as high as 75% of the total gas mixtures. This CO<sub>2</sub> composition reduces microbial growth and extends the shelf life of the products. Therefore, the aim of this study was to evaluate the effects of CO<sub>2</sub> concentration from 25-100% in combination with N<sub>2</sub> on microbial growth and physico-chemical properties of packed barramundi fillets. The role of microorganisms particularly total plate count (TPC), psychrophile, histamine forming bacteria and anaerobic bacteria on the spoilage process was also evaluated.

## Materials and methods

### *Sample preparation*

Whole barramundi (*L. calcarifer*) with an average weight of 0.5 kg were

bought from local fish market. The fish was transported to the laboratory on ice and upon arrival decapitated and filleted manually. All samples preparation was done according to Bakar *et al.* (2010) and Yassoralipour *et al.* (2012).

### *Modified atmosphere packaging of sample*

Each fillet was packed individually in a pouch with Multivac packaging (GOOD-And-WELL, Taiwan) and stored at 8±1°C for 20 days. The material (220×300 mm) were made of Nylon (thickness 0.09 mm) and had an O<sub>2</sub>, N<sub>2</sub>, CO<sub>2</sub> and H<sub>2</sub>O (at 23°C) transmission rate of 1.55, 0.465, 6.15 cm<sup>3</sup>/m<sup>2</sup>/day atm and 15 g/m<sup>2</sup>/day atm, respectively. The final gas/product ratio in all packages was adjusted to about 2:1 (v/w) by a gas mixer (PBI Dansensor MAP Mix 9001 ME, USA). The treatments were labelled as control (100% N<sub>2</sub>), M<sub>1</sub> (100% CO<sub>2</sub>), M<sub>2</sub> (75% CO<sub>2</sub>/25% N<sub>2</sub>), M<sub>3</sub> (50% CO<sub>2</sub>/50% N<sub>2</sub>), and M<sub>4</sub> (25% CO<sub>2</sub>/75% N<sub>2</sub>). All treatments were prepared in triplicates.

### *Mesophile, psychrophile, histamine forming and anaerobic bacteria*

The enumerations of mesophile and psychrophile bacteria were carried out according to method of AOAC (2000) while the counting of histamine-forming bacteria were carried out according to the procedure of Niven's (Niven *et al.*, 1981). Anaerobic bacteria enumeration on tryptose sulphite cycloserine (TSC) agar (Oxoid, USA) was performed according to Perez-

Alonso *et al.* (2004) and Ravi Sankar *et al.* (2008) and incubated an aerobically in anaerobic jar, at 30°C for 5 days. All samples were replicated twice.

#### Total volatile base-nitrogen (TVB-N)

A vapour distillation method was used for total volatile bases nitrogen (TVB-N) determination (Arashisar *et al.*, 2004; Goulas and Kontominas, 2005). The results were expressed as mg N/100g of fish muscle.

#### pH determination

The muscle was homogenised in distilled water in the ratio of 1:10 (w/v) and the measurement was done by a digital pH-meter (DELTA 320) at room temperature (Goulas and Kontominas, 2005).

#### Statistical analysis


Data collected were analyzed by analysis of variance (ANOVA). Pearson

correlation for the effect of CO<sub>2</sub> was conducted to determine the relationship between mesophile, psychrotrophile, histamine forming and anaerobe bacteria in all treatments. The significance of results was at  $p < 0.05$ . The software used was SPSS for windows (SPSS Inc. 2008).


## Results

### Microbiological changes


The total plate count (TPC), aerobic psychrophile count (APC), histamine forming bacteria (HFB) and anaerobic bacteria counts in barramundi fillets as a function of atmosphere storage at 8°C are shown in Figs. 1, 2, 3 and 4, respectively. The initial TPC for samples were 3.54 log cfu/g (Fig. 1) and APC 4.28 log cfu/g (Fig. 2).




**Figure 1:** Effect of packaging atmosphere on growth of total plate count on barramundi fillets stored at 8 °C. <sup>A, a</sup> TPC (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p < 0.05$ ).



**Figure 2: Effect of packaging atmosphere on growth of psychrophile aerobic bacteria on barramundi fillets stored at 8 °C.**  
 A, a APC (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p<0.05$ ).



**Figure 3: Effect of packaging atmosphere on growth of histamine forming bacteria on barramundi fillets stored at 8 °C.**  
 A, a HFB (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p<0.05$ ).




**Figure 4:** Effect of packaging atmosphere on growth of anaerobic bacteria on barramundi fillets stored at 8 °C.

A, a Anaerobe (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p<0.05$ ).

#### Total volatile base-nitrogen

Changes in TVB-N concentration for control and MAP conditions of barramundi fillet during the 20 days storage is given (Fig. 5). As expected, TVB-N levels were initially low (13.53 mg N/100g fish muscle) for fresh barramundi fillets. TVB-N increased gradually after 4 days of storage for

control barramundi fillet, reaching final (day 20) value of 111.10 mg N/100g fish muscle. TVB-N values of M<sub>1</sub>, M<sub>2</sub>, M<sub>3</sub> and M<sub>4</sub> compared to control also increased significantly ( $p<0.05$ ) after 8 days of storage; however, the rate of enhancement was lower than control.




**Figure 5:** Effect of packaging atmosphere on TVB-N value of barramundi fillets stored at 8 °C. <sup>A, a</sup> TVB-N (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p<0.05$ ).

### pH

pH values of the MAP samples were significantly ( $p<0.05$ ) different throughout storage period and the initial value which was 6.42, reached to 6.49,

6.55, 6.65, 6.67 and 6.92 for 100% CO<sub>2</sub>, 75% CO<sub>2</sub>, 50% CO<sub>2</sub> 25% CO<sub>2</sub> and control, respectively (Fig. 6).



**Figure 6:** Effect of packaging atmosphere on pH value of barramundi fillets stored at 8 °C. <sup>A, a</sup> pH (means±standard deviation) with different upper case (for the same MAP and different days) and different lower case (for the same days and different MAP) are significantly different ( $p<0.05$ ).

## Discussion

### *Microbiological changes*

The total microbiological limit for human consumption in fresh fish, proposed by ICMSF (2002) is 7 log cfu/g, while other authors recommend 6 log cfu/g (Ababouch *et al.*, 1991; Arashisar *et al.*, 2004; Scherer *et al.*, 2006). Psychrophile counts exceeded mesophilic counts in all the samples after 4 days. Similar observations were reported using the gas composition of 90:2.5:7.5% (CO<sub>2</sub>:O<sub>2</sub>:N<sub>2</sub>) and 40:30:30% (CO<sub>2</sub>:O<sub>2</sub>:N<sub>2</sub>) for rainbow trout by Arashisar *et al.* (2004) and Ravi Sankar *et al.* (2008) stored at 4±1°C.

Erkan *et al.* (2006) reported that mesophilic aerobic counts showed a slight increase for sardines (*Sardina pilchardus*) during storage 70:5:25% (CO<sub>2</sub>:O<sub>2</sub>:N<sub>2</sub>) at 4° C, which reached a value of 4 log cfu/g (day 7). Total mesophilic count exceeded 6 log cfu/g after 12 days in barramundi stored in M<sub>3</sub> and M<sub>4</sub>, and after 16 days for samples stored in M<sub>1</sub> and M<sub>2</sub>. In sample stored in M<sub>4</sub> (100% CO<sub>2</sub> concentration), the growth was slower by 1-2 log cfu/g. These results were similar to research of Arashisar *et al.* (2004), Lalitha *et al.* (2005) and Ravi Sankar *et al.* (2008), who kept their samples in MAP. They mentioned that the CO<sub>2</sub> concentration prolongs lag phase of bacteria growth,

increases generation time and delaying bacteria growth in fish filets. The HFB are usually considered part of the normal microflora of the gut, skin, or gills of fish (Huss, 1994). The histamine-forming bacteria grew faster in barramundi fillets kept in M<sub>3</sub> and M<sub>4</sub> sample, as compared to M<sub>1</sub> and M<sub>2</sub> (Fig. 3). However, the different concentration of CO<sub>2</sub> in the MAP sample might have influenced the growth of HFB (Özogul *et al.*, 2004; Özogul and Özogul, 2006). Results of ANOVA (Table 1) indicated that MAP and storage days significantly ( $p<0.05$ ) changed the pH, TVB-N and microbial load of fish fillet. As illustrated by Montgomery (2008), the corresponding variable will be more significant ( $p<0.05$ ) if the absolute F ratio became larger and P value became smaller.

Reddy *et al.* (1994) found that initial aerobic bacteria counts and anaerobic bacteria count in CO<sub>2</sub>:N<sub>2</sub> (75:25%) for tilapia stored at 4°C was 4.3 and 3.2 log cfu/g, respectively. This count reached a 4.6 log cfu/g after 10 days of storage in packed tilapia. Significant reduction of 1 to 2 log cfu/g was noticed in the anaerobic count in all samples until the 12 day of storage (Fig. 4). Reddy *et al.* (1994) and Ravi Sankar *et al.* (2008) also reported the same reduction for CO<sub>2</sub>:O<sub>2</sub> (60:40%) of pearlspot (*Etroplus suratensis*) stored at 0-2°C and CO<sub>2</sub>:N<sub>2</sub> (75:25%) tilapia stored at 4°C, respectively.



**Table 1: ANOVA results for the effect of MAP and storage days for all parameters evaluated for barramundi fillets.**

Parameters	100% CO <sub>2</sub>	75% CO <sub>2</sub>	50% CO <sub>2</sub>	25%CO <sub>2</sub>	100% N <sub>2</sub>	Adjusted <i>p</i> -value	MAP effect	Day (MAP) effect
<b>pH</b>								
							*	*
100% CO <sub>2</sub>		0.03610	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.03610		0.00000	0.00000	0.00000	F-ratio	3953.44	459.99
50% CO <sub>2</sub>	0.00000	0.00000		0.00000	0.00000			
25% CO <sub>2</sub>	0.00000	0.00000	0.00000		0.00000			
100%N <sub>2</sub>	0.00000	0.00000	0.00000	0.00000				
<b>TVB-N</b>								
							*	*
100% CO <sub>2</sub>		0.99900	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.99900		0.00000	0.00000	0.00000	F-ratio	377.57	233.36
50% CO <sub>2</sub>	0.00000	0.00000		0.00000	0.00000			
25% CO <sub>2</sub>	0.00000	0.00000	0.00000		0.00000			
100%N <sub>2</sub>	0.00000	0.00000	0.00000	0.00000				
<b>TPC</b>								
							*	*
100% CO <sub>2</sub>		0.17560	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.17560		0.00020	0.00000	0.00000	F-ratio	480.27	1362.25
50% CO <sub>2</sub>	0.00000	0.00020		0.00000	0.00000			
25% CO <sub>2</sub>	0.00000	0.00000	0.00000		0.00000			
100%N <sub>2</sub>	0.00000	0.00000	0.00000	0.00000				
<b>APC</b>								
							*	*
100% CO <sub>2</sub>		0.05860	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.05860		0.00040	0.00000	0.00000	F-ratio	270.56	980.18
50% CO <sub>2</sub>	0.00000	0.00040		0.00010				
25% CO <sub>2</sub>	0.00000	0.00000	0.00010		0.00000			
100%N <sub>2</sub>	0.00000	0.00000	0.00000	0.00000				
<b>HFB</b>								
							*	*
100% CO <sub>2</sub>		0.01510	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.01510		0.00000	0.00000	0.00000	F-ratio	403.22	833.82
50% CO <sub>2</sub>	0.00000	0.00000		0.99610	0.00000			
25% CO <sub>2</sub>	0.00000	0.00000	0.99610		0.00000			
100%N <sub>2</sub>								
<b>Anaerobe</b>								
							*	*
100% CO <sub>2</sub>		0.99900	0.00000	0.00000	0.00000	<i>p</i> -value	0.0000	0.0000
75% CO <sub>2</sub>	0.99900		0.00000	0.00000	0.00000	F-ratio	258.84	97.63
50% CO <sub>2</sub>	0.00000	0.00000		0.01460	0.00000			
25% CO <sub>2</sub>	0.00000	0.00000	0.01460		0.00000			
100%N <sub>2</sub>	0.00000	0.00000	0.00000	0.00000				

TVB-N: Total volatile base nitrogen; TPC: Total plate count; APC: Aerobic psychrophilic count; HFB: Histamine forming bacteria. \*  $p < 0.05$  is significant difference

### Total volatile base-nitrogen

The levels of 25 mg N/100g fish muscle is thought to indicate incipient spoilage in fish (Lannelongue *et al.*, 1982) but a higher TVB-N level of 30-35 mg N/100g fish muscle depending on fish species, has been suggested as the limit of acceptability for fish spoilage (Connell, 1990). In the current study, barramundi samples stored as control and M<sub>4</sub> exceeded after 4 and 12 days of storage, respectively.

### pH

A significant increase ( $p < 0.05$ ) of pH was observed in control samples during the storage period, and reached to pH 7.23 after 20 days of air storage. Increases in pH indicate the accumulation of alkaline compounds, such as ammonia mainly derived from microbial action (Hebard *et al.*, 1982).

Pastoriza *et al.* (1998) found similar results for pH values in control samples of ice-stored hake (*Merluccius merluccius*). The pH is lower for MAP-stored fish since carbon dioxide can be absorbed into fish muscle surface, acidifying it via the formation of carbonic acid (Banks *et al.*, 1980).

There were no significant differences ( $p > 0.05$ ) among the total plate count and psychrophile aerobic bacteria in M<sub>1</sub> as compared to M<sub>2</sub>; however, there was significant difference ( $p < 0.05$ ) with the other treatments (M<sub>3</sub>, M<sub>4</sub> and control) in 8 days of storage and after that significantly increased. Among four treatments, M<sub>1</sub> and M<sub>2</sub> had the least

changes in the all parameters evaluated. 75 and 100 % CO<sub>2</sub> concentration caused significant reductions ( $p > 0.05$ ) in the contents of TVB-N, pH and microbial counts in barramundi fillets, which has been stored at 8°C for 20 days. The atmosphere with 75:25% and 100:0 (CO<sub>2</sub>:N<sub>2</sub>) had least significant ( $p < 0.05$ ) changes and the most appropriate gas composition to reduce the microbial load that resulted in prolonging the shelf life of barramundi fillets.

### References

- Ababouch, L., Afilal, M.E., Benabdeljelil, H. and Busta, F.F., 1991.** Quantitative changes in bacteria, amino acids and biogenic amines in sardine (*Sardina pilchardus*) stored at ambient temperature (25–28°C) and in ice. *International Journal of Food Science & Technology*, 26(3), 297–306.
- AOAC., 2000.** Official methods of analysis of AOAC International. (W. Horwitz, Ed.). Gaithersburg Md.: Association of Official Analytical Chemists.
- Arashisar, S., Hisar, O., Kaya, M. and Yanik, T., 2004.** Effects of modified atmosphere and vacuum packaging on microbiological and chemical properties of rainbow trout (*Oncorhynchus mykiss*) fillets. *International Journal of Food Microbiology*, 97(2), 209–214.
- Bagge-Ravn, D., Ng, Y., Hjelm, M., Christiansen, J.N., Johansen, C.**

- and Gram, L., 2003.** The microbial ecology of processing equipment in different fish industries—analysis of the microflora during processing and following cleaning and disinfection. *International Journal of Food Microbiology*, 87(3), 239–250.
- Bakar, J., Yassoralipour, A., Bakar, F.A. and Rahman, R.A. 2010.** Biogenic amine changes in barramundi (*Lates calcarifer*) slices stored at 0°C and 4°C. *Food Chemistry*, 119(2), 467–470.
- Banks, H., Nickelson, R. I. I. and Finne, G. 1980.** Shelf life studies on carbon dioxide packaged finfish from the Gulf of Mexico. *Journal of Food Science*, 45(2), 157–162.
- Brody, A.L., 1989.** Modified atmosphere/vacuum packaging of meat, in: Brody, A.L. (Ed.), *Controlled/modified Atmosphere/vacuum Packaging of Foods*. Food and Nutrition Press, Trumbull, Connecticut, pp. 17–37.
- Connell, J.J., 1990.** Methods of assessing and selecting for quality. In *Control of Fish Quality* (3rd ed.). Springer, Berlin.
- Devlieghere, F., Debevere, J. and Van Impe, J., 1998.** Effect of dissolved carbon dioxide and temperature on the growth of *Lactobacillus sake* in modified atmospheres. *International Journal of Food Microbiology*, 41(3), 231–238.
- Emborg, J., Laursen, B.G. and Dalgaard, P., 2005.** Significant histamine formation in tuna (*Thunnus albacares*) at 2 degrees C-effect of vacuum- and modified atmosphere-packaging on psychrotolerant bacteria. *International Journal of Food Microbiology*, 101(3), 263–79.
- Erkan, N., Özden, Ö., Alakavuk, D., Yildirim, Ş. and İnuğur, M., 2006.** Spoilage and shelf life of sardines (*Sardina pilchardus*) packed in modified atmosphere. *European Food Research and Technology*, 222(5), 667–673.
- Farber, J.M., 1991.** Microbiological aspects of modified-atmosphere packaging technology—a review. *Journal of Food Protection*, 54(1), 58–70.
- Gimenez, B., Roncales, P. and Beltran, J.A., 2002.** Modified atmosphere packaging of filleted rainbow trout. *Journal of the Science of Food and Agriculture*, 82(10), 1154–1159.
- Goulas, A.E. and Kontominas, M.G., 2005.** Effect of salting and smoking-method on the keeping quality of chub mackerel (*Scomber japonicus*): biochemical and sensory attributes. *Food Chemistry*, 93(3), 511–520.
- Gram, L. 2009.** Microbiological Spoilage of Fish and Seafood Products. *Compendium of the Microbiological Spoilage of Foods and Beverages*, 87–119.
- Hebard, C.E., Flick, G.J. and Martin, R.E., 1982.** Occurrence and significance of trimethylamine oxide and its derivatives in fish and shellfish. In R. E. Martin, G. J. Flick, C. E. Hebard, and D. . Ward (Eds.),

- Chemistry & Biochemistry of Marine Food Products* (pp. 149–304). Connecticut: Avi, Westport.
- Huss, H. H. 1994.** Assurance of seafood quality. *FAO Fisheries Technical Paper*, 32–35.
- ICMSF., 2002.** International commission on microbiological specifications for foods. microorganisms in foods 7. Microbiological Testing in Food Safety Management. Kluwer Academic/Plenum Publishers, Springer, London.
- Lalitha, K.V, Sonaji, E.R., Manju, S., Jose, L., Gopal, T.K. S. and Ravisankar, C.N., 2005.** Microbiological and biochemical changes in pearl spot (*Etroplus suratensis* Bloch) stored under modified atmospheres. *Journal of Applied Microbiology*, 99(5), 1222–8.
- Lannelongue, M., Finne, G., Hanna, M.O., Nickelson, R. and Vanderzant, C., 1982.** Microbiological and chemical changes during storage of swordfish (*Xiphias gladius*) steaks in retail packages containing CO<sub>2</sub>-enriched atmospheres. *Journal of Food Protection*, 45(13), 1197–1203.
- López-Caballero, M., Gonçalves, A. and Nunes, M., 2002.** Effect of CO<sub>2</sub>/O<sub>2</sub>-containing modified atmospheres on packed deepwater pink shrimp (*Parapenaeus longirostris*). *European Food Research and Technology*, 214(3), 192–197.
- López-Gálvez, D. E., de la Hoz, L., Blanco, M. and Ordóñez, J.A., 1998.** Refrigerated storage (2°C) of sole (*Solea solea*) fillets under CO<sub>2</sub> -enriched atmospheres. *Journal of Agricultural and Food Chemistry*, 46(3), 1143–1149.
- Molin, G., 2000.** Modified atmospheres. In B. Lund, A. C. Baird-Parker, and G. W. Gould (Eds.), *Microbiological Safety and Quality of Food Vol. 1* (pp. 204–234). Gaithersburg, Maryland: Aspen Publishers, Inc.
- Montgomery, D.C. 2008.** Design and Analysis of Experiments. John Wiley & Sons. 680P.
- Niven, C.F., Jeffrey, M.B. and Corlett, D.A., 1981.** Differential plating medium for quantitative detection of histamine-producing bacteria. *Applied and Environmental Microbiology*, 41(1), 321–2.
- Ozogul, F., Taylor, K. D.A., Quantick, P., and Ozogul, Y., 2002.** Changes in biogenic amines in herring stored under modified atmosphere and vacuum pack. *Journal of Food Science*, 67(7), 2497–2501.
- Özogul, F., Polat, A. and Özogul, Y., 2004.** The effects of modified atmosphere packaging and vacuum packaging on chemical, sensory and microbiological changes of sardines (*Sardina pilchardus*). *Food Chemistry*, 85(1), 49–57.
- Özogul, F. and Özogul, Y., 2006.** Biogenic amine content and biogenic amine quality indices of sardines (*S. pilchardus*) stored in modified atmosphere packaging and vacuum

- packaging. *Food Chemistry*, 99(3), 574–578.
- Pastoriza, L., Sampedro, G., Herrera, J.J. and Cabo, M.L., 1998.** Influence of sodium chloride and modified atmosphere packaging on microbiological, chemical and sensorial properties in ice storage of slices of hake (*Merluccius merluccius*). *Food Chemistry*, 61(1-2), 23–28.
- Perez-Alonso, F., Aubourg, S.P., Rodríguez, Ó. and Barros-Velazquez, J., 2004.** Shelf life extension of Atlantic pomfret (*Brama brama*) fillets by packaging under a vacuum-skin system. *European Food Research and Technology*, 218(4), 313–317.
- Ravi Sankar, C.N., Lalitha, K.V, Jose, L., Manju, S. and Gopal, T.K.S., 2008.** Effect of packaging atmosphere on the microbial attributes of pearlspot (*Etroplus suratensis* Bloch) stored at 0-2°C. *Food Microbiology*, 25(3), 518–528.
- Reddy, N.R., Schreiber, C.L., Buzard, K. S., Skinner, G.E. and Armstrong, D.J., 1994.** Shelf life of fresh tilapia fillets packaged in high barrier film with modified atmospheres. *Journal of Food Science*, 59(2), 260–264.
- Scherer, R., Augusti, P.R., Bochi, V.C., Steffens, C., Fries, L.L. M., Daniel, A.P. and Emanuelli, T., 2006.** Chemical and microbiological quality of grass carp (*Ctenopharyngodon idella*) slaughtered by different methods. *Food Chemistry*, 99(1), 136–142.
- Yassoralipour, A., Bakar, J., Rahman, R. A. and Abu Bakar, F., 2012.** Biogenic amines formation in barramundi (*Lates calcarifer*) fillets at 8°C kept in modified atmosphere packaging with varied CO<sub>2</sub> concentration. *LWT - Food Science and Technology*, 48(1), 142–146.