

38

## Assessment of heavy metals in water and fish from Ibrahim Adamu Lake, Jigawa, Nigeria

Sambo, F. / Haruna, / Idris, S. / Mohammed, S. / Nasir, A.

### Abstract

The study was conducted to assess the levels of heavy metals in water and the flesh of tilapia (*Oreochromis niloticus*). Water and tilapia were sampled for a period of 12 months from October, 2003 – September, 2004. Water and muscle tissue were analyzed for Cu, Cr, Pb and Zn using flame atomic absorption spectrophotometer. The assessment showed Cr (3.35mg/l), Cu (0.75mg/l) Pb (0.81mg/l) in water while Cr (2.27mg/kg) and Pb (0.27mg/kg) were found to be high in the tissue samples and were above the values recommended by WHO and the food Agricultural Organization of the United Nations and therefore not safe for human consumption.

Keywords: Heavy metals, *Oreochromis niloticus*, atomic absorption, spectrophotometer.

### Introduction

Contamination of freshwater bodies with a different variety of pollutants has been a problem over the last decades because of the damage caused to aquatic life (Canli et al., 1998). Aquatic systems are very sensitive to heavy metal pollutants. Pollution of heavy metal in aquatic ecosystem is growing at an alarming rate (Malik et al., 2010). The main source of freshwater pollution can be attributed to discharge of untreated waste, dumping of industrial effluent, atmospheric deposition, mining, erosion and runoff from agricultural fields (Ambedkar and Muniyan, 2011).

Metals are not biodegradable and their presence in the food chain through a number of pathways becomes accumulated in different organs of animals (Stainiskiene et al., 2006; Korai et al., 2008). Increase in pollution, urbanization, industrialization and agricultural practices have further aggravated the situation (Gigure et al., 2004; Gupta et al., 2009). The accumulation could also depend on the concentration of the metals and exposure period (Mathew et al., 2011). Heavy metals can accumulate up to toxic concentration and cause damage. They have a great chance of moving up the food chain and cause public health hazards depending on their relative levels (Mansour and Sidky, 2002). Heavy metals such as Cu, Pb, Zn and Cr have been associated with waste water and sewage (Mucheweti et al., 2005) and are known to be toxic at high concentration (Rana et al., 2008).

The aim of this study is to assess the levels of these metals in water and in the flesh of the fish (*Oreochromis niloticus*) collected from Ibrahim Adamu Lake, Jigawa State, Nigeria.

### Materials and Methods

- **The Study Area:** Ibrahim Adamu Lake is located in Kazaure town of Jigawa state, 735km from the state capital Dutse on 13° along the line of latitude. The lake was constructed in 1974 with the primary aim of supplying water to the town, irrigation, fisheries and secondary aim of recreation and wildlife conservation.
- **Collection of fish sample:** Fish species were caught by artisanal fishermen using different fishing gears viz; traps, cast nets of “2-10” mesh and hooks. The fish caught were iced packed in order to maintain freshness and brought to the Research Laboratory immediately at Bayero University, Kano. The fish (*O. niloticus*) was identified at a glance by the characteristic pattern of dark and light bands crossing the caudal fin.
- **Analysis of metals in water:** Water samples for heavy metals analysis were collected in 1litre bottles pre-rinsed with distilled water. A 0.25 litre of the water sample was transferred into a 500cm<sup>3</sup> beaker. It was filtered and evaporated to dryness. The residue was re-dissolved in 0.1M nitric acid in a 100cm<sup>3</sup> volumetric flask ready for analysis after labeling (Ayodele and Abubakar, 1998). Aqueous stock solutions were prepared for Cr, Cu, Zn and Pb using

the appropriate salts. Five working standard solutions were prepared for each metal by serial dilution of stock solutions. These and blank solutions were aspirated into flame Atomic Absorption spectrophotometer. A calibration curve of absorption versus concentration was prepared for each metal and used for the determination of the metal concentrations in the samples using standard calibration plot.

Analysis of metals in fish tissue: Iced packed fish was dissected to remove the flesh. The flesh was dried to constant weight in an oven at 105°C. A method described by (Ayodele and Abubakar, 1998) was used to digest the sample. Dried tissue was ground in a ceramic mortar and sieved through 20 µm mesh. A 1.0g of the dry powdered sample was digested in 10cm<sup>3</sup> concentrated Nitric Acid. The resulting solution was evaporated to dryness and the residue was re-dissolved in 0.1MHNO<sub>3</sub>. The metal concentrations were determined by flame atomic absorption spectrophotometer (Buckmodel 210).

**Results**

In the water samples, the mean concentration of metal Zn, Cr, Cu, and Pb were 1.64mg/L, 3.35mg/L, 0.75mg/L and 0.81mg/L respectively (Table 1). The concentration of chromium was the highest followed by copper. The mean values of metals in fish tissue were 0.87mg/kg, 2.77mg/kg, 0.30mg/kg and 0.27mg/kg (Table 2). The mean values were compared with permissible limits.

Table 1: Mean heavy metals concentration (mg/l) in water samples from Ibrahim Adamu Lake

| Metals | Mean concentration (mg/l) | FAO/WHO (1983) |
|--------|---------------------------|----------------|
| Zn     | 1.64                      | 5.0            |
| Cr     | *3.35                     | 0.05           |
| Cu     | *0.75                     | 0.05           |
| Pb     | *0.81                     | 0.01           |

\*Levels higher than permissible limits

Table 2: Mean heavy metals concentration in fish tissue (mg/kg) from Ibrahim Adamu Lake

| Metals | Mean concentration (mg/kg) | FAO/WHO (1983) |
|--------|----------------------------|----------------|
| Zn     | 0.87                       | 5.00           |
| Cr     | *2.77                      | 1.00           |
| Cu     | 0.30                       | 3.00           |
| Pb     | *0.27                      | 0.21           |

\*Levels higher than permissible limits

**Discussion**

The research area, Ibrahim Adamu is the centre of human and livestock communities in Kazaure Local Government Area of Jigawa state, therefore, the analysis of the water and fish from the lake becomes very important from public health perspectives. The level of heavy metals in the fish flesh is of interest because of potential risk to humans (Asraf, 2005; Barbieri et al., 2010). The mean concentration of Zn in the water and tissue samples were below the FAO/WHO limits of 5.0mg/L and 5.0mg/kg desirable for Zn in drinking water and fish tissue respectively.

Copper is an essential element that promotes the activity of certain enzymes in human body, however when ingested in large quantity may be toxic to man and may pose danger to the community. The mean concentration of copper recorded in water was higher than in tissue when compared with the permissible limits (Table 1 and 2). This could be attributed to human activities taking place. Lawani and Alowade (1987) reported a mean value of 5.89mg/l for Cu.

The mean concentration of chromium (3.35mg/Land 2.77mg/kg) was above the limit for both water and fish tissue when compared with values of 0.05mg/L and 1.00mg/kg by FAO/WHO. Eneji et al. (2010) reported high concentration of chromium in Tilapia from River Benue which he said was attributed to bioavailability, intrinsic fish processes and trophic structure variation.

Lead poisoning is ranked as the most common environmental health hazard even at low concentrations. The mean concentrations of lead obtained were high considering the acute toxicity of the metal. The values in the study 0.81mg/l and 0.27mg/kg were high compared with the safe limits set by FAO/WHO.

**Conclusion**

The result from the study gave important information on the levels of the four metals and the fish were capable of concentrating these metals in their tissue from the water showing the same trend in humans that feed on the fish depending on the consumption by consumers. It is recommended that resources be put in place by the state government to continuously reduce to the minimum the method of the lake pollution.

## REFERENCES

- Ambedkhar, G. and Muniyan, M. (2011). Bioaccumulation of some Heavy metals in the selected five freshwater fish from Kolhdam River, Tamilnadu, India. *Advances in Applied Sciences Research*, 2(5): 221 – 225.
- Ashraf, W. (2005). Accumulation of Heavy metals in kidney and heart tissues of epinephelus microdon fish from the Arabian Gulf. *Environ Monit Assess*, 101(1-3): 311 – 316.
- Ayodele, J.T. and Abubakar, M. (1998). Trace metals in Tiga Lake, Kano, Nigeria. *Trop. Environ Rev*, 3:230 – 237.
- Barbieri, E., Passos, E., Aragoakas, A., Santos, A.B. and Garcia, C.A.B. (2010). Assessment to trace metal in catfish (catharopis spixi) from Sal River, Estuary, Aracail, State of Sergipe, Brazil. *Water Environ. Res*, 82(2): 2301 – 2307.
- Canli, M., Ayo and Kolay, M. (1998). Levels of metals (Cd,Pb,Cu,and Ni) of Euprinus Carpia, Barbus capito and Choudrostrome regtun from the seyhan River, Turkey, Tubitak, Turkj., 22: 149 – 157.
- Eneje, I.S., Rufussha, A. and Annune, P.A. (2010). Bioaccumulation of Heavy metals in fish (Tilapia zilli and Clarias gariepinus) organs from River Benue, North-Central Nigeria. *Pak. J. Anal Environ Chem.*, Vol. 12, No. 1 and 2 (25-31): 127-133.
- FAO (1983). *Compilation of Legal limits for Hazardous Substances*. Rome, Italy, 5-100.
- Gupta, A., Rai, D.K., Pandey, R.S. and Sharma, B. (2009). Analysis of some heavy metals in the riverine water, sediments and fish from river ganges at Allahbad. *Environ Monit. Assess*, 157: 449-458.
- Gigure, A., Campbell, P.G.C., Hare, L., Mc Donald, D.G. and Rasmussen, J.B. (2004). Influence of lake chemistry and fish age on cadmium, copper and zinc concentrations in various organs of indigeneous yellow perch (*Perca flavescens*). *Canadian J. fish Aquat. Sci*, 61: 702-716.
- Korai, A.L., Sahoto, G.A., Kazi, T.G. and Lashari, K.H. (2008). Lead concentrations in fresh water, muscle, gill and liver of Catla catla (Hamilton) from keenjhar Lake. *Pak. J. Anal. Environ. Chem*. 9(10): 11-9.
- Lawani, S.A. and Alawode (1987). Bioaccumulation of heavy metals in fish organs from River Benue, Nigeria. *Nig. J. of Biotech.*, 4(4).
- Mathew, Precious, D. H., Blessing, M.D., Lebogange, F., Rendani, V. and Francis, R. (2011). Heavy metal levels in water, catfish (*Clarias gariepinus*) and African fish eagle (*Haliaeetus*) specimen from the municipal waste water fed modimola dam outside makifeng city, North-west Province, South Africa. *Life Science Journal*, 8(52).
- Mucheweti, M., Birkett, W.J., Chinyangaa, E., Zuausya, R., Scrimshaw, M.D. and Lester, J.N. (2006). Heavy Metal content of vegetables irrigated with mixtures of waste water and sewage sludge in Zimbabwe. Implications for Human Health Agric. *Ecosystem Environ*, 112(1): 1-14.
- Malik, N., Biswass, A.K., Qureshi, T.A., Borana, K. and Virha, R. (2010). Bioaccumulation of heavy metals in fish tissue of a freshwater Lake of Bhopal. *Environ. Monit. Assess*, 160:267 – 268.
- Mansour, S.A. and Sidk, M.M. (2002). Ecotoxicological studies 3: Heavy metals Contamination water and fish from Kayoom Government of Egypt. *Food Chem.*, 78:15-22.
- Rana, T., Sakars Mandal, T.K. and Batabyal, S. (2008). Haemotobiochemical profiles of affected cattle at arsenic prone zone in Haringhata block of Nadia District of West Bengal in India. *Internet. J. Hamotel*, 4(2).
- Staniskiene, B., Matuskevicius, P., Budreckiene, R. and Sibniewaska, K.A. (2006). Distribution of heavy metals in tissues of freshwater fish in Lithuania, *Polish J. Environ. Stud.*, 15(4): 585 – 591.