

50

Evaluation of protein fractions of indigenous clariid fish species (*Clarias gariepinus* and *Heterobranchus bidorsalis*) and their reciprocal hybrids

Agbebi, O. T. / Sofela, S. O. / Awodiran, M. O.

Abstract

Electrophoresis of Myofibrillar and Sarcoplasmic muscle proteins of African catfish: Clarias gariepinus, Heterobranchus bidorsalis and their reciprocal hybrids in South-West Nigeria was carried out to reveal the similarities and dissimilarities among species in order to aid the selection of suitable strains for aquaculture that could lead to production of new varieties of fishes to alleviate the problem of short supply of fast growing quality fish seeds. The study was aimed at analyzing the muscle protein profiles of C. gariepinus, H. bidorsalis and their reciprocal hybrids. Sixteen juveniles fish samples (comprising four samples from each mating combinations) artificially propagated and reared for sixteen weeks were analyzed electrophoretically. Myofibrillar and sarcoplasmic fractions were prepared by homogenizing 150mg of fish muscle in 1.5ml of rigor buffer containing 10 mM Trisemeleates, 60 mM K Cl, 5 mM MgCl₂, 1 mM EDTA. Extracts were centrifuged in a tube at 10,000g for 5 min at 4°C. The resultant pellets (myofibrilla) and supernatant (sarcoplasmic) were separated using 12.5% Sodium Dodecyl-Sulphate Polyacrylamide Gel Electrophoresis (1D SDS-PAGE). The relative concentration of individual protein bands were analyzed using TotalLab™ 1D software. The individual protein bands in the electrophoregram were identified in relation to their molecular weights. A total of eleven and seven protein bands were resolved in Myofibrilla and Sarcoplasmic fractions respectively. The 5th band with molecular weight (MW) of 52.23 KDa of the myofibrilla electrophoregram distinguished C. gariepinus from H. bidorsalis while the 3rd band with MW 119.04, 4th band with MWs 101.49 & 102.13; 8th band with MWs 29.24 and 29.39 KDa distinct the pure breeds from the hybrids. However, in sarcoplasmic fraction, the 3rd and 5th bands with MWs 92.11 KDa and 54.28 KDa respectively distinguished the hybrids in the while the 7th band with MW 41.67 KDa distinct the pure breeds. Therefore, this research will serve as a bridge between the existing gaps of information available on the muscle protein profile of C. gariepinus, H. bidorsalis and their reciprocal hybrids and the study identifies the proteomic classification of Clariid species with the aim of enlightening fish researchers and aquaculturist on the characterization of broodstock selection for successful breeding exercise.

Keywords: Myofibrillar, sarcoplasmic, rigor buffer, C. gariepinus and H. bidorsalis.

Introduction

Clariids are widely considered as an important tropical catfish species for aquaculture. It is well known in both culture and artisanal environments in Nigeria where it serves both socio-cultural and research purposes in most regions. *Clarias gariepinus* and *Heterobranchus bidorsalis* are species of high aquaculture importance in South-Western Nigeria (Oyebola et al., 2013). They are widely cultured owing to their high market price, fast growth rate and ability to withstand adverse pond conditions especially low oxygen content (Idodo-Umeh, 2003). *C. gariepinus* and its hybrid, which is reproductively viable, are cultured throughout Nigeria and most especially in the South Western Zone. Hybridization between the two species is common and widely reported among farmers in order to improve quality of fish seed. Hybrids of the two species must be properly identified so as not to replace them with the pure of *Heterobranchus* spp. Intergeneric hybrids of *Heterobranchus* X *Clarias* have been observed to be fertile (Aluko 1998). It was stated that aquaculture practitioners in Nigeria have exploited the fertility of the F1 hybrids by using these hybrids as breeders for further propagation. This improper hatchery practice of indiscriminate use of F1 hybrids as breeders should be of immediate concern to conservationist and geneticists as the potential hazards of this process on the catfish gene pool: could be the genetic loss of pure indigenous stock of *Heterobranchus* and *Clarias* species (Yisa and Olufagba, 2005).

In the past, the identification of fish species was carried out mainly by examining the external morphological characteristics. In the present day, electrophoresis of whole muscle protein, sarcoplasmic proteins, serum proteins, liver proteins, salivary proteins and a number of enzymes often have been used by some researchers as an aid in the species identification of fish (Pintero et al. 2001, Yilmaaz et al. 2007, Lamy et al. 2008). Although a few serum electrophoresis research (Diyaware et al. (2011, 2012); Akinwande et al. (2012), Majolagbe et al. (2012) and Oyebola et al. (2013)) on *Clarias* and *Heterobranchus sp* have recently surfaced, but there appears dearth of information on the fish muscle protein of Clariid species in Nigeria (Agbebi et al., 2013).

Muscle is the principal component of meat and contains water, proteins, fat, carbohydrates and inorganic constituents (Forrest et al., 1975). Water, as a major constituent, accounts for 65 to 80% of the total muscle weight- Proteins comprise from 16 to 22% of the total muscle mass and are the principal component of the solid matter. Muscle proteins are generally categorised as described by Huss, (1995) as structural proteins (myofibrillar), sarcoplasmic and collagen. Myofibrillar proteins (actin, myosin, torpormyosin and actomyosin) constitute 70-80 percent of the total protein content, sarcoplasmic proteins (myoalbumin, globulin and enzymes) constitute 25-30 percent of the protein while connective tissue proteins (collagen), constitute approximately 3 percent of the protein in teleostei. Myofibrillar proteins are soluble in neutral salt solutions of fairly high ionic strength (0.5 M) while sarcoplasmic which are soluble in neutral salt solutions of low ionic strength (<0.15 M). Their solubility in the various concentrations of salt solutions makes them independently separable in electrophoresis.

In 1D-Electrophoresis, myofibrillar and sarcoplasmic proteins are separated on polyacrylamide gel. Most of the electrophoretic studies carried out are related to either total proteins or myofibrillar protein or corneal proteins (Myla et al., 2013). However, the proteins in the sarcoplasmic fraction are excellently suited to distinguishing fish species, as each species has a characteristic band pattern when separated by isoelectric methods (Huss, 1995). Therefore this research will serve as a bridge between the existing gaps of information available on the muscle protein profile of *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids and also establish the similarities and dissimilarities between the species and their hybrids for the purpose of identification.

Materials and Methods

Muscle sample collection. The experimental fish was produce through intergeneric hybridization between *C. gariepinus* and *H. bidorsalis* obtained from Hepa Fish Farm, Asero, Abeokuta, Nigeria. The following crosses were carried out for parental cross- *C. gariepinus* x *C. gariepinus* and *H. bidorsalis* x *H. bidorsalis*, and hybrids *C. gariepinus* x *H. bidorsalis* – (*Clariabbranchus*) and *H. bidorsalis* x *C. gariepinus* (Heteroclarias).

Sixteen juvenile fish samples (comprising four samples from each mating combinations) were transported live in well open plastic rubber bowls to the Biotechnology laboratory unit of the Animal Breeding and Genetics of the Federal University of Agriculture, Abeokuta. Muscle were isolated from the epiaxial muscle in the region below the cranial to the dorsal fin, above the lateral line of the experimental fish with the aid of a new surgical blade for each species (El-Serafy et al., 2006 and Reedish et al., 2008) after fish killing by medullar transaction (El-Serafy et al., 2006). Myofibrillar and sarcoplasmic extracts were prepared by homogenizing 150mg of muscle with the aid of a mortar and pestle (on ice) in 1.5ml of rigor buffer containing 10 mM trisemeates, 60 mM K Cl, 5 mM MgCl₂, 1 mM EDTA (Agbebi et al., 2013). The extracts were prepared according to (Agbebi et al., 2013).

Sample preparation for SDS-PAGE analysis and gel staining. Protein fractions were centrifuged for 5 minutes at 10,000 x g at room temperature prior the electrophoretic analysis. Samples were loaded into 1mm x 12 cm x 14 cm 12.5% polyacrylamide gel with 3% stacking gel. First, gel capillary chamber was loaded with a broad range molecular weight standard (BioRad Laboratories, Hercules, CA). Proteins were separated by applying constant voltage of 300V for 2hrs 30mins until the dye front reached the bottom of the gel. After electrophoretic separation, gels were stained overnight with gentle agitation on an orbital shaker. Staining buffer contains (400 ml methanol, 50 ml glacial acetic acid and Coomassie Brilliant Blue G250) and subsequently destained with 10% acetic acid. Images were digitized and acquired employing a CCD camera (Samsung ES25- Samsung Opto, Electronics Co. Ltd). Raw images were subsequently imported into TotalLab 1D software (Nonlinear Dynamics, Newsacastle upon Tyne, UK) and analyzed (band detection and molecular weight calculations). User defined parameters employed are background subtraction-rolling ball, radius = 300; band detection: minimum slope = 30; noise reduction = 3; % maximum peak = 1.0; Gaussian volumes fitted to peaks-no advanced fitting; Rf calibration aligned with no lane, and use curve lines and snap R_f lines to bands were checked (Reddish et al., 2008).




Fig. 1: Raw image of 12.5% SDS myofibrillar muscle protein of artificially propagated *C.gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Fig. 2: Raw image of 12.5% SDS sarcoplasmic muscle protein fraction of artificially propagated *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Fig. 1 & 2 are the raw image of Myofibrilla and Sarcoplasmic muscle proteins of *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids. Results




Fig. 3: Calibrated image of 12.5% SDS myofibrillar muscle protein of artificially propagated *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Fig. 4: Calibrated image of 12.5% SDS sarcoplasmic muscle protein fraction of artificially propagated *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Fig. 3 & 4 shows the result of 12.5% SDS electrophoregram analysis. The blue diamonds denotes bands derived from electrophoregram by the TotalLab software.

The myofibrillar muscle fraction resolved a total of eleven different protein bands across the gel. The average number of protein bands resolved among the pure breeds; *C. gariepinus* and *H. bidorsalis* were ten and nine respectively in all examined species. In the hybrids; Clariobranchus and Heteroclaris ten protein bands each were detected. The 1st, 2nd, 6th, 7th, 9th, 10th and 11th bands were commonly detected across all mating combinations. In *C. gariepinus* the 5th band was present but absent in *H. bidorsalis*. The 4th and 8th bands were also present in the pure breeds but absent in the hybrids. The hybrids also had the 3rd band present but absent in the pure breeds. There were no distinguishing bands between the hybrids in myofibrilla fractions.

In the male pure breeds, eight protein bands were resolved for both *C. gariepinus* and *H. bidorsalis* while the male hybrids Clariobranchus spp. and Heteroclaris spp. both resolved six bands. In the female seven protein bands were resolved in both pure breeds; *C. gariepinus* and *H. bidorsalis* and hybrids; Clariobranchus species and Heteroclaris spp. resolved. Although there were species specific bands among individual species there was no clear sexual dimorphic band across the male or female species of all cross combinations.

The sarcoplasmic muscle fraction resolved a total of nine different protein bands were resolved across the gel. The 1st and 9th bands were only present in BRS (Broad Range Standard) therefore, seven protein bands were detected across the mating combinations. The 2nd, 4th, 6th and 8th bands were commonly detected across all mating combinations. The average number of protein bands resolved among the pure breeds; *C. gariepinus* and *H. bidorsalis* were six and five respectively in all examined species. In *C. gariepinus* the 7th band was present but absent in both *H. bidorsalis*. In the hybrids; Clariobranchus and Heteroclaris, an average number of six and four protein bands each were respectively detected. The 3rd and 5th bands were present in Clariobranchus but absent in Heteroclaris.

In the male pure breeds, five protein bands were resolved for both *C. gariepinus* and *H. bidorsalis* while the male hybrids; Clariobranchus and Heteroclaris species resolved five and four protein bands respectively. In the females, five protein bands were resolved in pure breeds; *C. gariepinus* and *H. bidorsalis* while three protein bands were resolved in hybrids Clariobranchus and Heteroclaris species.

The molecular weights of each Myofibrillar (Table 1) and Sarcoplasmic (Table 2) are presented below for artificially propagated *Clarias gariepinus*, *Heterobranchus bidorsalis* and their reciprocal hybrids in Kilo-Daltons (KDa). Myofibrillar protein bands were within 210 KDa to 10KDa while the sarcoplasmic protein bands were within 140KDa to 30KDa.

Table 1: Molecular weights of bands detected for calibrated image of 12.5% SDS myofibrillar muscle protein of artificially propagated *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Band No.	BRS	CLRS	HETB	CLRB	HETC
1	223.00	-	-	-	-
2	116.20	141.53	143.03	146.93	140.92
3	97.40	-	-	92.11	-
4	66.20	64.27	66.71	66.2	63.53
5	-	53.41	54.62	54.28	-
6	45.00	47.39	47.99	48.4	47.47
7	-	41.67	-	-	-
8	31.00	33.81	34.77	35.39	34.43
9	21.5	-	-	-	-

Table 2: Molecular weights of bands detected for calibrated image of 12.5% SDS sarcoplasmic muscle protein of artificially propagated *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids.

Band No.	BRS	CLRS	HETB	CLRB	HETC
1	223.00	175.25	179.35	201.17	205.50
2	-	137.63	141.60	148.99	154.03
3	116.20	-	-	119.04	119.04
4	97.40	101.49	102.13	-	-
5	66.20	52.23	-	53.78	54.12
6	45.00	48.43	49.72	47.28	46.83
7	-	38.09	37.71	37.25	37.65
8	31.00	29.24	29.39	-	-
9	-	22.39	22.18	23.10	23.48
10	21.50	20.35	20.38	20.80	21.37
11	-	18.36	18.36	18.25	17.93

In the myofibrillar electrophoregram, the 5th band that distinguishes the pure breeds; *C. gariepinus* and *H. bidorsalis* had a molecular weight of 52.23KDa. The distinctions between the pure breeds and the hybrids marked by the 4th band had molecular weights 101.49 and 102.13 KDa while the 8th band had molecular weights 29.24 and 29.39 KDa for *C. gariepinus* and *H. bidorsalis* respectively. The 3rd band had the molecular weights 119.04 in both Clariabranchus and Heteroclaris.

In the Sarcoplasmic electrophoregram, the 7th band that distinguishes the pure breeds; *C. gariepinus* and *H. bidorsalis* had the molecular weight 41.67 KDa. The 3rd and 5th bands that distinct the hybrids; Clariabranchus and Heteroclaris had the molecular weight 92.11 KDa and 54.28 KDa respectively.

Discussion

Electrophoretic separation by SDS-PAGE muscle myofibrilla and sarcoplasmic proteins reveal similarity and dissimilarity among *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids in southwest Nigeria based on presence or absence and molecular weights of proteins bands of individual species and their hybrids. A total of eleven and seven protein bands were resolved in myofibrilla and sarcoplasmic fractions respectively. This is not farfetched from results obtained by Machado and Sgarbieri (1991), in the partial characterization and nutritive value of proteins from Pacu (*Colossoma mitrei*, Berg 1895).

Dissimilarities among species are important for identification of the species within the genus. The 5th band of the myofibrilla electrophoregram distinguished *C. gariepinus* from *H. bidorsalis* while the 3rd, 4th and 8th bands distinct the pure breeds from the hybrids. The bands of the hybrids were similar, hence no distinction could be made for the myofibrilla fraction. However, the 3rd and 5th bands distinguished the hybrids in the sarcoplasmic fraction, while the 7th bands distincts the pure breeds. Band similarity was high in both fractions; seven out of eleven in the myofibrilla fraction and four out of seven in sarcoplasmic fraction. Myofibrillar and sarcoplasmic protein bands were found within molecular weight ranges of 10 KDa – 210 KDa of 30KDa – 140KDa. It is therefore observed that both fractions: sarcoplasmic and myofibrilla effectively characterized the pure breeds but sarcoplasmic identifies hybrids species effectively than myofibrilla fractions. This is in support with Yarmohammadi et al. (2011) using molecular technique such as amplified fragment length polymorphism (AFLP) applied to species identification of better hybrids. Hybrids identification was performed by comparison of electrophoresis profiles with parental species.

Agbebi et al. (2013) identified the individual protein bands in each fraction of yellow perch (*Perca flavescens*) whole muscle, sarcoplasmic and myofibrilla in relation to their molecular weights and identified the sarcoplasmic protein in the whole muscle from myofibrilla fraction. In Myla et al. (2013) muscle albumin proteins of three species of pomfrets - *Pampus argenteus*, *P. chinensis* and *Apolectis niger* observed difference between the sexes of three species using protein fraction, relative mobility and molecular weights. The finding does not corroborate with this result for no distinct classification is derived from the sexes of the two fractions. Even though sexually dimorphic bands are not satisfactorily reported in this study, but male and female fish species are easily distinguished from one another under field examination after they are about 10cm long (Yisa and Olufeagba, 2005).

Conclusion

The challenges of morphological variations in *C. gariepinus*, *H. bidorsalis* and their reciprocal hybrids necessitated the use of molecular tool in establishing their biochemical relationship. This research will enable Aquaculturists to recognize or identify the right species to use in breeding and genetic studies for accurate results and profitability. Consequently solving the problem

of poor growth performance and loss of pure indigenous stock of *Heterobranchus* and *Clarias* species.

REFERENCES

- Agbebi O.T., Wick M.P, Popovski Z. T (2013). Evaluation of three protein fractions in yellow fish perch species (*Perca flavescens*) using two extraction buffers: Trichloroacetic Acid Precipitation and Rigor Buffer. *Ife Journal of Science* Vol. 15(2): 197-208.
- Akinwande A. A., O. A. Fagbenro and O. T. Adebayo (2012). Serum protein pattern in interspecific and intergeneric hybrids of *Heterobranchus longifilis*, *Clarias gariepinus* and *Clarias anguillaris* in sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) *International Journal of Fisheries and Aquaculture* Vol. 4(10):202-208.
- Aluko, P. O. and Shaba, M. (1998). Intra- and interspecific hybridization studies between exotic *Clarias gariepinus* and two indigenous Clariid species. *Nigerian J. genet.* 14:59-63.
- Diyaware, M. Y., Haruna, A. B., Abubakar, K. A., Omitogun, G. O. (2011). Serum protein characterization of intergeneric hybrids between *C. anguillaris* X *H. bidorsalis* and their hybrids from semi-arid zones of Nigeria. *J. Microbiol. Biotech. Res.* 1/4:227-234.
- , Haruna, A. B., Abubakar, K. A., Omitogun, G. O. (2012). Serum protein electrophoretic characterization of *C. anguillaris*, *H. bidorsalis* and their hybrids from Northeast Nigeria. *J. Microbiol. Biotech. Res.* 2/1:70-77.
- Huss, H. H. (1995). Quality and quality changes in fresh fish. *FAO Fisheries Technical Paper* 348.
- Idodo-Umeh, G. (2003). *Freshwater Fishes of Nigeria (Taxonomy, Ecological Notes, Diets and Utilization)*. Benin City: Idodo-Umeh Publishers Ltd. 232.
- Lamy E, Gonçalo da Costa, Fernando, C. S., Potes, J., Coelho, A. V., Baptista, E. S. (2008). Comparison of electrophoretic protein profiles from sheep and goat parotid saliva. *J. Chem. Ecol.* 34:388-397.
- Majolagbe, F. A., Awodiran, M. O., Awopetu, J. (2012). Electrophoretic Studies of *Clarias gariepinus* and *Heterobranchus bidorsalis* and their hybrids. *Ife Journal of Science* 14:1 167.
- Myla, S., Chakravarty, M., Navya, P. R., Ganesh, C. (2013). SDS-polyacrylamide gel electrophoresis of muscle albumin proteins in three species of pomfrets—*Pampus argenteus*, *Pampus chinensis*, and *Apolectis niger*. *International Journal of Advanced Biology.* 3/2: 172-180.
- Oyebola, O. O., Omitoyin, B. O., Salako, A. E., Ajani, E. K., Awodiran, M. O. (2013). Genetic and biochemical differentiation of pectoral spine variants in *Clarias gariepinus*. *Int'l J. of Modern Biol. Res.* 8:14.
- Pineiro, C., Vazquez, J., Marina, A. I., Barros-Velazquez, J., Gallardo, J.M. (2001). Characterization and partial sequencing of species-specific sarcoplasmic polypeptides from commercial hake species by mass spectrometry following two-dimensional electrophoresis. *Electrophoresis.* 22/8: 1545-1552.
- Reddish, J. M., St-Pierre, N., Nichols, A., Green-Church, K. (2008). Proteomic analysis of proteins associated with body mass and length in yellow perch (*Perca flavescens*). *Proteomics* 2008, 8, 2333-2343.
- Yilmaz, M., Yilmaz, R. H., Alas, A. (2007). An electrophoretic taxonomic study on serum proteins of *Acanthobrama marmid*, *Leuciscus cephalus*, and *Chondrostoma regium*. *EurAsian J. of BioSciences* 3, 22-27.
- Yisa, M. Olufeagba, S. O. (2004). An exposition on field identification of Clariid catfishes as an important tool in fish breeding and genetics. *Proceedings, 19th FISON Annual Conference, Nov. 29-Dec. 3, Ilorin.*