

Technical University of Denmark



The response of fluorescent organic matter to ozone treatment in pilot freshwater RAS

Spiliotopoulou, Aikaterini; Andersen, Henrik Rasmus; Pedersen, Lars-Flemming; Stedmon, Colin; Hambly, Adam

Publication date:
2018

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Spiliotopoulou, A., Andersen, H. R., Pedersen, L-F., Stedmon, C., & Hambly, A. (2018). The response of fluorescent organic matter to ozone treatment in pilot freshwater RAS. Abstract from World Aquaculture Society: Asian – Pacific Aquaculture 2018 , Taipei, Taiwan, Province of China.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

THE RESPONSE OF FLUORESCENT ORGANIC MATTER TO OZONE TREATMENT IN FRESHWATER PILOT RECIRCULATING AQUACULTURE SYSTEMS

Aikaterini Spiliotopoulou^{*}, Henrik R. Andersen, Lars-Flemming Pedersen, Colin A. Stedmon & Adam C. Hambly

Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet 115, 2800 Kongens Lyngby, Denmark
OxyGuard International A/S, Farum Gydevej 64, 3520 Farum, Denmark
aispil@env.dtu.dk

Water quality in intense recirculating aquaculture systems (RAS) is characterised by the accumulation of organic and inorganic compounds, potentially deteriorating the water quality. Ozone has been implemented in RAS as a continuous water treatment technology improving water quality by oxidizing organic matter and reducing bacterial abundance. The analytical efforts to detect and monitor organic matter (DOM) in aquatic systems have included absorption spectroscopy of the coloured fraction (CDOM). A part of CDOM also fluoresces (FDOM). FDOM fraction has been widely used in aquatic environments as a quantitative and qualitative measure of DOM. FDOM can be characterised by the fluorescence excitation-emission matrix (EEM) spectroscopy which can be further decomposed mathematically with methods such as parallel factor (PARAFAC) analysis to identify the independent FDOM fractions. This provides both a quantitative and qualitative measure of DOM.

Recently, it was shown that the organic matter fluorescence of RAS water is highly sensitive to ozonation and fluorescence spectroscopy could therefore be used as an indirect method to determine ozone delivery within these systems. Therefore, three ozone dosages, including a control, were injected in pilot freshwater RAS where trouts were farmed. The ozonation trial lasted eight days utilising one RAS per dosage. The test levels ranged from 52-130 mg O₃/h, equivalent to 10-25 g O₃/kg feed.

The DOM consisted of four components (Fig. 1) which differed in their fluorescence characteristics and response to ozonation. A UV wavelength fluorescent fraction (C4) typical of proteinaceous material removed by 13-20% immediately after ozone initiation. The remaining fractions that exhibited visible wavelength fluorescence, at first were unaffected but during the following days, were gradually degraded, reaching a removal of 34-66%. By the end of the experiment the fluorescence intensities of all fractions were diminished up to 60% in all applied dosages.

This study provides the first application of fluorescent EEM spectroscopy and PARAFAC analysis to determine the effect of ozone on FDOM character in aquaculture water, more specifically within continuously treated RAS. The results indicate that the response in organic matter fluorescence can be used to depict the impact of ozone dosage.




Figure 1: Effect of ozone on FDOM over time.