

Keywords: gastrointestinal cancer; aspirin; epidemiology; drug repurposing

Effect of low-dose aspirin use on survival of patients with gastrointestinal malignancies; an observational study

M A Frouws¹, E Bastiaannet¹, R E Langley², W K Chia³, M P P van Herk-Sukel⁴, V E P P Lemmens^{5,6}, H Putter⁷, H H Hartgrink¹, B A Bonsing¹, C J H Van de Velde¹, J E A Portielje⁸ and G J Liefers^{*,1}

¹Department of Surgical Oncology, Leiden University Medical Center, Leiden 2300 RC, The Netherlands; ²MRC Clinical Trials Unit, University College London, Institute of Clinical Trials and Methodology, Aviation House 125 Kingsway, London WC2B 6NH, UK; ³Department of Medical Oncology, National Cancer Centre Singapore, 11 Hospital Drive, Singapore 169610, Singapore; ⁴PHARMO Institute for Drug Outcomes Research, Van Deventerlaan 30/40, Utrecht 3528 AE, The Netherlands; ⁵Comprehensive Cancer Organisation The Netherlands, Eindhoven 5600 AE, The Netherlands; ⁶Department of Public Health, Erasmus MC Medical Centre, PO Box 2040, Rotterdam 3000 CA, The Netherlands; ⁷Department of Medical Statistics and Bioinformatics, Leiden University Medical Center, Leiden 2300 RC, The Netherlands and ⁸Department of Medical Oncology, Haga Hospital, Leyweg 275, The Hague 2545 CH, The Netherlands

Background: Previous studies suggested a relationship between aspirin use and mortality reduction. The mechanism for the effect of aspirin on cancer outcomes remains unclear. The aim of this study was to evaluate aspirin use and survival in patients with gastrointestinal tract cancer.

Methods: Patients with gastrointestinal tract cancer diagnosed between 1998 and 2011 were included. The population-based Eindhoven Cancer Registry was linked to drug-dispensing data from the PHARMO Database Network. The association between aspirin use after diagnosis and overall survival was analysed using Cox regression models.

Results: In total, 13715 patients were diagnosed with gastrointestinal cancer. A total of 1008 patients were identified as aspirin users, and 8278 patients were identified as nonusers. The adjusted hazard ratio for aspirin users vs nonusers was 0.52 (95% CI 0.44–0.63). A significant association between aspirin use and survival was observed for patients with oesophageal, hepatobiliary and colorectal cancer.

Conclusions: Post-diagnosis use of aspirin in patients with gastrointestinal tract malignancies is associated with increased survival in cancers with different sites of origin and biology. This adds weight to the hypothesis that the anti-cancer effects of aspirin are not tumour-site specific and may be modulated through the tumour micro-environment.

The incidence of cancer is increasing, particularly in low-and medium-resource countries; by the end of 2015 there were an estimated 15.2 million new cases globally with a predicted increase to 21.6 million by 2030 (Sullivan *et al*, 2015). The cost of health care is also increasing, and there is a real need for reasonably priced, widely available therapeutics to improve cancer outcomes. Although, the US Food and Drug Administration has approved a

higher percentage of oncology drugs since 2008, many of these are expensive targeted agents with approvals based on surrogate end points, and infrequently improve overall survival (Kim and Prasad, 2015). Aspirin (acetylsalicylic acid) was originally synthesised and used as an analgesic in 1897, with the anti-platelet functions of low-dose aspirin subsequently discovered in the early 1970s. This latter discovery led to many large randomised controlled trials

*Correspondence: Dr GJ Liefers; E-mail: g.j.liefers@lumc.nl

Received 12 September 2016; revised 23 November 2016; accepted 28 November 2016; published online 10 January 2017

© 2017 Cancer Research UK. All rights reserved 0007–0920/17

delineating the role of aspirin in the treatment and prevention of vascular disease. Retrospective long-term analyses of cancer outcomes in these randomised trials have revealed two interesting phenomena. Firstly, there was a 24% reduction in cancer incidence in patients allocated to aspirin, and this effect was seen across tumour types but was most marked in tumours arising from the gastrointestinal tract. Secondly, if cancers did develop they were less likely to have metastasised at presentation or subsequently if the patient received aspirin (Rothwell *et al*, 2011, 2012; Algra and Rothwell, 2012; Cook *et al*, 2013).

Much of the work to date relating to aspirin and cancer has focussed on colorectal cancer. In particular there have been several epidemiological studies showing a reduction in cancer mortality and improved overall survival for patients taking aspirin after a diagnosis of colorectal cancer (Chia *et al*, 2012; Jacobs *et al*, 2012; Paleari *et al*, 2015; Coyle *et al*, 2016; Elwood *et al*, 2016). This has led to several ongoing adjuvant studies in colorectal cancer; the Add-Aspirin trial (Langley and Rothwell, 2014), Adjuvant Aspirin for Colon Cancer (NCT02467582), the ALASCCA trial (NCT02647099), the ASCOLT trial (NCT00565708) and the Aspirin trial (NCT02301286). In addition, two other randomised controlled trials have focussed on primary prevention, and after long-term follow-up showed a beneficial effect on primary prevention in both hereditary and sporadic colorectal cancer (Burn *et al*, 2011; Cook *et al*, 2013). Also, a meta-analysis of four other randomised controlled trials showed an absolute risk reduction of 6.7% for the recurrence of adenoma's in patients with a history of these lesions (Cole *et al*, 2009).

The mechanism(s) underlying the beneficial effects of aspirin on cancer outcomes remains unclear. Several different potential biomarkers have been investigated, but due to the multiple potential cellular pathways and conflicting results of previous studies, the mechanism of action remains unknown; although platelets may have a central role (Langley *et al*, 2011). The aim of this study was to provide epidemiological evidence and further mechanistic insights on the potential beneficial effects of aspirin use after diagnosis of cancer that arises from any part of the gastrointestinal tract. Because many studies have tried to differentiate effects of aspirin use both before and after diagnosis, an additional analysis was performed including the patients that use aspirin both pre- and post diagnosis.

MATERIALS AND METHODS

Study population. Data from the Eindhoven Cancer Registry was used to identify patients diagnosed with cancer of the gastrointestinal tract between January 1998 and December 2011 in the south of the Netherlands. This area is served by 10 hospitals, covers a demographic region of approximately 1.5 million Dutch citizens and is part of the nationwide Netherlands Comprehensive Cancer Organisation. The Eindhoven Cancer Registry is linked to the municipal population registry, which records the vital status (alive/dead) of all inhabitants. Patients are informed about the registration and registered except patients who objected to be registered. The Netherlands Cancer Registry is obliged to work according to the law about protection of privacy data; informed consent of the patients for this specific study was not applicable. Patient selection and data cleaning was performed by the Eindhoven Cancer Registry. Follow-up for this project was until 31 December 2012.

The PHARMO Database Network is a population-based network which combines data from different health-care settings in the Netherlands. For this study the out-patient pharmacy database was used, which contains drug-dispensing records from all community pharmacies. Drugs are coded using the Anatomical

Therapeutic Chemical classification (www.whocc.no/atc_ddd_index) and the records include information on the type of product, date prescribed, dose and regimen, quantity and route of administration. The PHARMO database was linked to the Eindhoven Cancer Registry and thus allows drug use by cancer patients to be analysed (van Herk-Sukel *et al*, 2010). From this linked database, prescriptions for aspirin (only the ones that were actually dispensed) were selected.

Definition of users and nonusers. For this study, patients older than 18 years who used aspirin after a diagnosis of a gastrointestinal cancer were selected. The gastrointestinal tumours were coded according to the International Classification of Disease 10 [ICD-10] C15-C26. This comprises cancer from the following sites: oesophagus; stomach; small intestine; colon; recto-sigmoid and rectum; anus; liver and intra hepatic bile ducts; gallbladder and extra hepatic bile ducts; pancreas; and a group 'gastrointestinal tumours not otherwise specified (nos)'.

Patients who used aspirin before diagnosis were excluded from the analyses. Aspirin users (ATC codes: B01AC06, B01AC08, B01AC56, N02BA01, N02BA15, N02BA51, N02BA65) were defined as those prescribed aspirin for at least 30 days. Nonusers were defined as patients who received for less than 30 days or never used aspirin. Time after diagnoses was defined in periods of use and no use by analysing each single prescription during follow-up. Periods of less than 14 days in between two prescriptions were considered consecutive. Follow-up started 14 days after diagnosis because there was no information about in-hospital use of medication. Immortal time bias is avoided by analysing prescriptions as a time-varying covariate, in periods of use and no use (Stricker and Stijnen, 2010; Suissa and Azoulay, 2012).

Statistical analysis. Information from the ECR-contained information about the presence or absence of the following comorbidities at cancer diagnosis: lung disease; cardiovascular disease; diabetes and disorders of the gastrointestinal tract; urinary tract; nervous system; musculoskeletal system; and a group of other comorbidities. Comorbidity was analysed as 0, 1 or ≥ 2 comorbidities. A χ^2 -test was used to assess baseline characteristics for categorical values.

Survival analysis were performed with the Simon-Makuch method, an alternative for Kaplan-Meier and with the ability to process time-varying covariates in survival curves (Simon and Makuch, 1984). A Cox proportional hazards model was used with aspirin use as a time-varying covariate, as described by Stricker and Stijnen (2010). Schoenfeld residuals were tested to verify the assumption of proportional hazards. Follow-up duration (survival) was recorded in months from diagnosis ($t=0$). Multivariable survival models were built with the following covariates: age at diagnosis (continuous), sex, stage of cancer (categorical), number of comorbidities (0, 1 or ≥ 2), treatment (surgery yes/no, radiotherapy yes/no and chemotherapy yes/no). Missing/unknown values were included in the multivariable model as missing indicator. Analysis were performed using Stata statistical software version 12 (StataCorp, 2011). Statistical tests were two-sided and considered significant at the $P < 0.05$ level.

Relative survival rates were used to take into account the risk of dying from causes other than the disease of interest. The excess mortality reflects the difference between the overall survival of patients and the survival that would be expected in the absence of cancer. The excess mortality was calculated as the ratio of the observed (all-cause) survival proportion to the expected survival proportion (Dickman and Adami, 2006). National life tables were used to estimate background mortality (expected survival) according to sex, year of age and incidence year. Relative excess risks were estimated using a flexible parametric model, implemented in the Stata command `stpm2` (Lambert and Royston, 2009).

Different parts of the gastrointestinal tract were analysed separately if there were as at least 10 aspirin users (therefore small bowel, anal cancer and gastrointestinal tumours NOS were not considered separately). Histological subtypes (adenocarcinoma and squamous cell carcinomas) were also analysed separately in groups with at least 10 aspirin users. Statistical interaction for this subgroup was tested by including an interaction term in the model of aspirin use and histological subtype and significance was assessed using the Wald test. A sensitivity analysis was performed by repeating the analysis and excluding patients with stage IV disease and separately repeating the analysis and excluding the first year of follow-up from the analysis. The main analysis and all subgroup analysis were pre-planned.

Pre- and post-diagnosis use of aspirin. For the analysis in patients that use both aspirin before and after diagnosis, the groups were selected with the same method as described in the 'definition of users and nonusers' heading. The only difference was that patients who started aspirin use before diagnosis and continued this after diagnosis were selected for the group of users of aspirin. Patients using aspirin only after diagnosis were excluded for this analysis. The statistical analysis was also equal to the analysis described above, where aspirin use was analysed as time-varying covariate, and the same factors were used for the multivariable analysis.

RESULTS

In total 13 715 patients were identified with a cancer of the gastrointestinal tract diagnosed between January 1998 and

December 2011. The following were excluded from the analysis (CONSORT diagram Figure 1): 4187 patients who were using aspirin before diagnosis, 239 patients with follow-up of less than 14 days and three patients below the age of 18 years. Thus, 9286 patients were included in the survival analysis, of which 8278 patients (person years: 4375) did not use aspirin and 1008 (person years: 2150) used aspirin after diagnosis. In total, 5138 events (deaths) were recorded. Table 1 shows the characteristics of this population.

The majority of patients were diagnosed with colon cancer (43%), rectal cancer (25%) and oesophageal cancer (10%). Median age at diagnosis was 68 years (interquartile range (IQR) 59–76) in the aspirin group and 69 (IQR 61–74) in the nonusers group. Aspirin users were less often female and more frequently diagnosed with stages I and II disease compared with nonusers. In the nonusers group, 26% of patients had stage IV disease compared with 9% in the aspirin users group. Median survival for all patients was 48 months.

Figure 2 shows survival curves for users of aspirin after diagnosis vs nonusers. In the group of aspirin users, 65% (95% CI 59–71%) of patients was alive after five years, in contrast to nonusers, where 45% (95% CI 44–46%) of patients was alive after five years.

A Cox proportional hazard model was used with use of aspirin as a time-varying covariate. The proportional hazard assumption was fulfilled. For all patients with gastrointestinal cancer, aspirin use was associated with a significant reduction in overall mortality, hazard ratio (HR) 0.57 (95% confidence interval (CI) 0.48–0.69) (Table 2). Adjusted for age at diagnosis, sex, stage of cancer,




Figure 1. Flowchart of patients selected for analysis.

Table 1. Characteristics of the cohort

	All patients	%	No aspirin use	%	Aspirin post diagnosis	%	P-value
Total	9286	100	8278	89	1008	11	
Location tumour							
Oesophageal cancer	946	10.19	886	10.7	60	5.95	<0.001
Gastric cancer	750	8.08	700	8.46	50	4.96	
Small intestine cancer	97	1.04	88	1.06	9	0.89	
Colon cancer	3977	42.83	3434	41.48	543	53.9	
Rectal cancer	2358	25.39	2069	24.99	289	28.7	
Anal cancer	67	0.72	60	0.72	7	0.69	
Hepatobiliary cancer	385	4.15	360	4.35	25	2.5	
Pancreatic cancer	692	7.45	667	8.06	25	2.47	
Cancer of the gastrointestinal tract nos	14	0.15	14	0.17	0	0.0	
Sex							
Male	5140	55.35	4517	54.6	623	61.8	<0.001
Female	4146	44.65	3761	45.43	385	38.19	
Age mean (s.d.). Median (IQR)	67.1 (11)	68 (60–75)	66.7 (12)	68 (59–76)	67.7 (9.9)	69 (61–74)	
18–60 Years	2420	26.06	2219	26.8	201	19.9	<0.001
60–69 Years	2763	29.75	2437	29.4	326	32.3	
70–79 years	2831	30.49	2464	29.8	367	36.4	
80 Years and older	1272	13.7	1158	14	114	11.3	
Stage							
0	204	2.2	176	2.13	28	2.78	<0.001
I	1496	16.11	1258	15.2	238	23.56	
II	2222	23.93	1900	22.95	322	31.94	
III	2058	22.16	1788	21.6	270	26.8	
IV	2249	24.22	2162	26.12	87	8.63	
Unknown	1057	11.4	994	12.01	63	6.25	
Surgery							
No	2693	29	2603	31.44	90	8.93	<0.001
Yes	6593	71	5675	68.56	918	91.07	
Chemotherapy							
No	6544	70.47	5798	70.04	746	74.01	0.009
Yes	2742	29.53	2480	29.96	262	25.99	
Radiotherapy							
No	7042	75.83	6291	76	751	74.5	0.3
Yes	2244	24.17	1987	24	257	25.5	
Comorbidities							
None	3383	36.43	3056	36.92	327	32.44	0.05
One	2664	28.69	2359	28.5	305	30.26	
Two or more	2295	24.71	2027	24.49	268	26.63	
Unknown	944	10.17	836	10.1	108	10.69	
Morphology							
Adenocarcinoma	8343	89.84	7378	89.13	965	95.73	<0.001
Squamous cell carcinoma	298	3.21	280	3.38	18	1.79	
Epithelial	140	1.51	135	1.63	5	0.5	
Gastrointestinal Stromal tumour	58	0.62	50	0.6	8	0.79	
Other (not specified)	447	4.81	435	5.25	12	1.19	
Months survival, median (IQR)	48	(15.4–95.4)	24	(7.5–58.6)	89.4	(54.8–132.6)	

Abbreviations: IQR = interquartile range; nos = not otherwise specified; s.d. = standard deviation.

number of comorbidities, treatment (surgery, radiotherapy, and chemotherapy) the multivariable HR was 0.52 (95% CI 0.44–0.63).

Stratification according to tumour type is shown in Table 2 and Figure 3. In patients with oesophageal, colon, rectal, and hepatobiliary tract cancer a significant association was found between the use of aspirin after diagnosis and overall survival. For patients with gastric and pancreatic cancer using aspirin, there was no statistically significant increase in survival. A survival benefit with aspirin was seen regardless of the stage of cancer at presentation and after all primary treatment modalities including chemotherapy, radiotherapy or surgery (Figure 4).

Table 2 additionally shows the relative survival estimates which are a good estimation of the cancer-specific survival (Sarfati *et al*, 2010). Equal to the overall survival rates, the observed relative excess risks were significant in patients with oesophageal cancer, hepatobiliary cancer, colon and rectal cancer.

Squamous cell cancers accounted for 3% of the total cohort of which, 81% ($n = 242$) arose from the oesophagus and 18% ($n = 53$) were anal cancers. Patients with adenocarcinoma of the oesophagus who used aspirin had an adjusted HR 0.24 (95% CI 0.10–0.59) for overall survival, while those with a squamous cell carcinoma of the oesophagus had a HR for overall survival of 1.02 (95% CI 0.37–2.83) for aspirin users compared with nonusers. The test for heterogeneity of the effect of aspirin in patients with oesophageal squamous cell carcinoma vs patients with oesophageal adenocarcinoma was significant (P for interaction = 0.01).

In 72% of prescriptions, 2435 in total, the dose was reported. Of all prescribed dosages, 98% were 100 mg daily or lower. It was therefore not possible to analyse a dose–effect relationship, because only 31 prescriptions were for higher-dose aspirin.

The sensitivity analysis with the exclusion of the first year follow-up showed a similar effect, with an unadjusted HR of 0.56




Figure 2. Survival comparison for aspirin users vs nonusers with Simon-Makuch method.

(95% C.I. 0.45–0.69) and adjusted HR 0.49 (0.39–0.61). The sensitivity analysis or stages I–III showed an adjusted HR 0.49 (95% CI 0.39–0.61), consistent with the stratified analysis by stage in Figure 4.

The analysis in the patients that use aspirin both pre- and post diagnosis can be found in Table 3. Supplementary Table 1 and Supplementary Figure 1 show the PRISMA flowchart for this cohort and the patient characteristics.

DISCUSSION

Aspirin use after diagnosis of a gastrointestinal malignancy is associated with significantly lower mortality rates and this effect remains after adjusting for potential cofounders. It was most marked for tumours arising from the oesophagus, colon, rectum and hepatobiliary tract. This large cohort study of almost 9300 patients is the first observational cohort study evaluating the association of aspirin and survival in various gastrointestinal malignancies. The statistically significant effect on survival seen in patients with tumours of the oesophagus, colon and rectum is consistent with data from other published studies (Chia *et al*, 2012; Jacobs *et al*, 2012; Macfarlane *et al*, 2015; Coyle *et al*, 2016; Elwood *et al*, 2016). The effect in the tumour types was also present in patients that used aspirin both pre- and post diagnosis.

In a recent prospective cohort study, Cao *et al* (2016) found that the reduced overall reduced cancer risk associated with the use of aspirin was primarily owing to gastrointestinal tract cancers. In addition, in a meta-analysis of randomised trials evaluating aspirin for the prevention of cardiovascular disease, Rothwell *et al* (2011) showed a reduced risk of cancer-specific death with aspirin (HR of 0.79 (95% CI 0.68–0.92)) in all types of cancer. Stratified for tumour location, the largest benefit was found in patients with gastrointestinal tumours, with no significant heterogeneity between different gastrointestinal cancers. Consistent with our study they also showed that patients with adenocarcinomas were most likely to benefit from aspirin HR 0.70 (95% CI 0.54–0.91). However, in contrast, patients in our study only started aspirin after diagnosis of cancer, which is most relevant when considering recommendations for subsequent management after a cancer diagnosis. In our study, 11% of patients started using aspirin after

diagnosis, which is also consistent with previous studies in cancer patients (McCowan *et al*, 2013).

A strength of our study is that the data is derived from linked cancer registry and pharmacy data, eliminating both recall and information bias. Although we cannot verify that patients actually ingested the aspirin, the prescriptions registered by the PHARMO institute are actually handed out to the patients by the pharmacy and this therefore adds weight to the definition of user. In addition, immortal time bias and misclassification of exposure in follow-up is avoided by the use of a Cox proportional hazards model with time-varying covariate (Suissa and Azoulay, 2012). With this technique, accurate risk estimates are provided as each individual prescription is analysed (Stricker and Stijnen, 2010). Moreover, the exclusion of patients already using aspirin before diagnosis and the determination of patient characteristics at diagnosis ($t = 0$) mimics the use of aspirin as adjuvant therapy. In our study, patients are identified at diagnosis but before they are exposed to the treatment of interest and differentiated into groups of users and nonusers. This 'new-user design' eliminates important biases associated with observational studies (Ray, 2003). In addition, it has been suggested that for measuring the side effects of drugs, which the effect of aspirin on cancer could theoretically be considered, observational data could in some cases be considered non-inferior to results from randomised controlled trials (Vandenbroucke, 2004). Our study has limitations. First, since exposure to aspirin depends on a clinician's decision to prescribe aspirin to a certain patient, it is prone to confounding by selective prescribing. For instance, oncologists may withhold aspirin treatment (as secondary prevention for cardiovascular disease) in patients diagnosed with incurable (stage IV) cancer because of the poor prognosis. Thus patients with a particularly poor prognosis may end up in the non-user group. This reverse causation was addressed by the pre-planned sensitivity analysis excluding the first year of follow-up, which restricted the study population to patients alive at one year after diagnosis. By introducing this one year exposure lag, any undiagnosed recurrence at baseline or early recurrence would have been likely to become apparent and therefore baseline prognosis between the two groups is believed to be more similar (Chubak *et al*, 2013). Second, proven cardiovascular disease is the main indication for low-dose aspirin in The Netherlands. This could imply that patients prescribed aspirin have a worse life expectancy at baseline because of lifestyle factors and risks associated with both cardiovascular disease and cancer development. Considering the absence of information on cancer-specific survival and cause of death in our study, hypothetically part of the overall survival gain we observed could be explained by the prevention of cardiovascular mortality. However, in a large meta-analysis of individual participant data, the reduction in vascular-specific mortality from aspirin was only 9%, HR 0.91 (Baigent *et al*, 2009). Therefore a reduction in cardiovascular mortality could only partly explain the reduction in mortality we observed. Several of the studies evaluating the effect of aspirin use after a diagnosis of colorectal cancer have shown a significant reduction in colorectal cancer-specific mortality (Chia *et al*, 2012; Coyle *et al*, 2016). Third, over the counter aspirin use was not included. However, prescription data can give valid estimations of association even though available over the counter (Yood *et al*, 2007). No data were available to adjust for lifestyle factors, health-related behaviour and mutational status. Lastly, Table 1 shows that the groups aspirin users and nonusers are different with respect to baseline characteristics. This is to a large extent the result of the size of the cohort. After adjustment for these factors the association between aspirin use and survival remained significant. Nevertheless, the confounding by indication as described remains, and therefore randomised controlled trials remain inevitable before aspirin can be used as regular anti-cancer treatment.

Table 2. Time-dependent survival analysis (overall survival) stratified according to tumour type

		Overall survival						Relative survival			
No. at risk	No. events	Crude hazard ratio	95% CI	P-value	Adjusted hazard ratio ^a	95% CI	P-value	Crude relative excess risk	95% CI	Adjusted relative excess risk ^a	95% CI
Aspirin users vs nonusers (n = 9286)											
Nonusers	4776	1 (Reference)			1 (Reference)						
Aspirin Users	1008	0.57	0.48–0.69	<0.001	0.52	0.44–0.63	<0.001	0.41	0.30–0.56	0.44	0.33–0.58
Aspirin users vs nonusers per tumour type											
Oesophageal cancer (n = 946)											
Nonusers	722	1 (Reference)			1 (Reference)						
Aspirin Users	30	0.42	0.22–0.81	0.01	0.45	0.23–0.88	0.02	0.34	0.15–0.77	0.31	0.14–0.70
Gastric cancer (n = 750)											
Nonusers	555	1 (Reference)			1 (Reference)						
Aspirin Users	30	0.75	0.41–1.36	0.34	0.87	0.47–1.61	0.66	0.50	0.19–1.29	0.70	0.29–1.70
Pancreatic cancer (n = 692)											
Nonusers	631	1 (Reference)			1 (Reference)						
Aspirin users	21	1.06	0.58–1.93	0.84	0.81	0.44–1.48	0.49	1.03	0.56–1.89	0.77	0.42–1.41
Hepatobiliary cancer (n = 385)											
Nonusers	311	1 (Reference)			1 (Reference)						
Aspirin Users	10	0.39	0.16–0.95	0.04	0.34	0.14–0.84	0.02	0.34	0.18–1.06	0.35	0.13–0.95
Colon cancer (n = 3977)											
Nonusers	3434	1 (Reference)			1 (Reference)						
Aspirin Users	543	0.62	0.48–0.80	<0.001	0.56	0.43–0.72	<0.001	0.41	0.23–0.72	0.44	0.27–0.72
Rectal cancer (n = 2358)											
Nonusers	2069	1 (Reference)			1 (Reference)						
Aspirin Users	289	0.51	0.33–0.77	0.001	0.41	0.27–0.63	<0.001	0.26	0.09–0.78	0.25	0.09–0.68

Abbreviation: CI, confidence interval.

^aAdjusted for stage, sex, age at diagnosis, surgery, radiotherapy, chemotherapy and comorbidities.




Figure 3. Overall survival analysis for aspirin users vs nonusers stratified according to tumour type.




Figure 4. Forest plot of adjusted overall survival analysis in patients with gastrointestinal malignancies grouped according to aspirin users vs nonusers and stratified for stage and treatment.

The mechanism responsible for the effect of aspirin on cancer remains unknown. Aspirin (acetylsalicylic acid) reduces prostanoïd generation by irreversible inhibition of platelet COX-1 (cyclooxygenase-1) and COX-2 isozymes. Activated platelets release several growth factors which impact on tumour progression and metastasis (Bruno *et al*, 2012). Maximum platelet inactivation by COX-1 is thought to be obtained by low-dose aspirin (75–100 mg daily) and over 95% of platelet activity is inhibited for up to 24 h (Patrignani *et al*, 2014). A number of potential biomarkers have been identified as predictors of response to aspirin in terms of cancer outcomes. Chan *et al* (2009) reported that the effect of high-dose (325 mg) aspirin after a colorectal cancer diagnosis was predominantly in patients with COX-2 (also called PTGS2) overexpression. However, to achieve constant inhibition of COX in tissues, the administered daily dose of aspirin would have to be higher than 2000 mg (Thun *et al*, 2012). In some studies mutations in phosphatidylinositol 3-kinase (PIK3CA) have been associated with aspirin response; however, in a previous study we did not find this association but showed that the effect of aspirin was associated with tumours that expressed human leukocyte antigen (HLA) class I molecules (Reimers *et al*, 2014; Paleari *et al*, 2015).

Our observation that aspirin use is similarly associated with good prognosis in various tumour types with clearly different biology makes a nonspecific mode of action plausible. It is possible that aspirin executes its effect by inhibiting platelet aggregation around circulating epithelial tumour cells, irrespective of organ site which then facilitates immune clearance. The coming years will hopefully provide answers. Several randomised clinical trials have commenced in the past years (NCT02647099 (Langley and Rothwell, 2014), NCT02467582, NCT02301286, NCT00565708). Many of these trials are united in the ‘Aspirin Trialist Collaborative Group’ and will pool results regarding clinical outcome and expression of biomarkers.

Table 3. Time-dependent survival analysis (overall survival) stratified according to tumour type with prediagnosis aspirin users

	Overall survival							
	No. at risk	No. events	Crude hazard ratio	95% CI	P-value	Adjusted hazard ratio ^a	95% CI	P-value
Aspirin users vs nonusers (n = 12 109)								
Nonusers	8366	4913	1 (Reference)			1 (Reference)		
Pre and post-diagnosis aspirin users ^b	2736	1647	0.69	0.64–0.75	<0.001	0.61	0.57–0.66	<0.001
Aspirin users vs nonusers per tumour type								
Oesophageal cancer (n = 1180)								
Nonusers	894	741	1 (Reference)			1 (Reference)		
Aspirin Users ^b	286	229	0.64	0.52–0.79	<0.001	0.61	0.49–0.76	<0.001
Gastric cancer (n = 933)								
Nonusers	714	574	1 (Reference)			1 (Reference)		
Aspirin Users ^b	219	184	0.90	0.72–1.13	0.37	0.85	0.67–1.07	0.17
Pancreatic cancer (n = 876)								
Nonusers	681	648	1 (Reference)			1 (Reference)		
Aspirin Users ^b	195	183	0.68	0.54–0.84	<0.001	0.67	0.53–0.84	0.001
Hepatobiliary cancer (n = 477)								
Nonusers	364	317	1 (Reference)			1 (Reference)		
Aspirin Users ^b	113	101	0.81	0.61–1.08	0.16	0.69	0.51–0.93	0.02
Colon cancer (n = 4730)								
Nonusers	3469	1642	1 (Reference)			1 (Reference)		
Aspirin Users ^b	1261	612	0.67	0.59–0.76	<0.001	0.55	0.48–0.63	<0.001
Rectal cancer (n = 2687)								
Nonusers	2080	910	1 (Reference)			1 (Reference)		
Aspirin Users ^b	607	306	0.78	0.65–0.94	0.008	0.63	0.52–0.75	<0.001

Abbreviation: CI = confidence interval.

^aAdjusted for stage, sex, age at diagnosis, surgery, radiotherapy, chemotherapy and comorbidities.

^bOnly patients using aspirin both pre- and post diagnosis were analysed.

CONCLUSION

Aspirin use after diagnosis of gastrointestinal malignancies is associated with improved overall survival. This observation makes a nonspecific mode of action for aspirin on cancer plausible. These results offer direction towards future studies, both in terms of new randomised controlled trials as well as further studies to identify biomarkers that predict response to aspirin.

ACKNOWLEDGEMENTS

MPPVH-S is an employee of the PHARMO Institute for Drug Outcomes Research. This independent research institute performs financially supported studies for government and related health-care authorities and several pharmaceutical companies. However, this study was not financially supported by a pharmaceutical company. R.E.L. has received financial support through grants from Cancer Research UK and the National Institute for Health Research (NIHR) Health Technology Assessment (HTA) as Chief Investigator of the Add-Aspirin Trial, has received compensation from Bayer and Aspirin Foundation for service on scientific advisory boards, and has received a supply of aspirin and placebo from Bayer Pharmaceuticals AG for the Add-Aspirin Trial.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHOR CONTRIBUTIONS

MAF contributed to study design, data preparation, data analysis and interpretation, prepared the first draft of the report, contributed to subsequent versions and the final report. EB contributed to study design, data preparation, data analysis and interpretation, contributed to the first draft of the report and drafting of the final report. REL and WKC contributed to data interpretation, reviewed the first draft of the report, contributed to subsequent drafts and the final report. HP contributed to study design, data preparation, data analysis and interpretation and contributed to the drafting of the final report. MPPVH-S, VEPPL, HHH, BB, CJHVDV and JEAP contributed to data interpretation and drafting of the final report. GJL initiated the study, contributed to data analysis and interpretation, contributed to the first draft of the report, subsequent versions and the final version. G.J.L. had full access to all the data in the study and the final responsibility for the decision to submit the article for publication.

REFERENCES

- Algra AM, Rothwell PM (2012) Effects of regular aspirin on long-term cancer incidence and metastasis: a systematic comparison of evidence from observational studies versus randomised trials. *Lancet Oncol* **13**(5): 518–527.
- Baigent C, Blackwell L, Collins R, Emberson J, Godwin J, Peto R, Buring J, Hennekens C, Kearney P, Meade T, Patrono C, Roncaglioni MC, Zanchetti A (2009) Aspirin in the primary and secondary prevention of vascular disease: collaborative meta-analysis of individual participant data from randomised trials. *Lancet* **373**(9678): 1849–1860.
- Bruno A, Dovizio M, Tacconelli S, Patrignani P (2012) Mechanisms of the antitumoural effects of aspirin in the gastrointestinal tract. *Best Pract Res Clin Gastroenterol* **26**(4): e1–e13.
- Burn J, Gerdes AM, Macrae F, Mecklin JP, Moeslein G, Olschwang S, Eccles D, Evans DG, Maher ER, Bertario L, Bisgaard ML, Dunlop MG, Ho JW, Hodgson SV, Lindblom A, Lubinski J, Morrison PJ, Murday V, Ramesar R, Side L, Scott RJ, Thomas HJ, Vasen HF, Barker G, Crawford G, Elliott F, Movahedi M, Pylvanainen K, Wijnen JT, Fodde R, Lynch HT, Mathers JC, Bishop DT (2011) Long-term effect of aspirin on cancer risk in carriers of hereditary colorectal cancer: an analysis from the CAPP2 randomised controlled trial. *Lancet* **378**(9809): 2081–2087.
- Cao Y, Nishihara R, Wu K, Wang M, Ogino S, Willett WC, Spiegelman D, Fuchs CS, Giovannucci EL, Chan AT (2016) Population-wide impact of long-term use of aspirin and the risk for cancer. *JAMA Oncol* **2**(6): 762–769.
- Chan AT, Ogino S, Fuchs CS (2009) Aspirin use and survival after diagnosis of colorectal cancer. *JAMA* **302**(6): 649–658.
- Chia WK, Ali R, Toh HC (2012) Aspirin as adjuvant therapy for colorectal cancer—reinterpreting paradigms. *Nat Rev Clin Oncol* **9**(10): 561–570.
- Chubak J, Boudreau DM, Wirtz HS, McKnight B, Weiss NS (2013) Threats to validity of nonrandomized studies of postdiagnosis exposures on cancer recurrence and survival. *J Natl Cancer Inst* **105**(19): 1456–1462.
- Cole BF, Logan RF, Halabi S, Benamouzig R, Sandler RS, Grainge MJ, Chaussade S, Baron JA (2009) Aspirin for the chemoprevention of colorectal adenomas: meta-analysis of the randomized trials. *J Natl Cancer Inst* **101**(4): 256–266.
- Cook NR, Lee IM, Zhang SM, Moorthy MV, Buring JE (2013) Alternate-day, low-dose aspirin and cancer risk: long-term observational follow-up of a randomized trial. *Ann Intern Med* **159**(2): 77–85.
- Coyle C, Cafferty FH, Langley RE (2016) Aspirin and colorectal cancer prevention and treatment: is it for everyone? *Curr Colorectal Cancer Rep* **12**: 27–34.
- Dickman PW, Adami HO (2006) Interpreting trends in cancer patient survival. *J Intern Med* **260**(2): 103–117.
- Elwood PC, Morgan G, Pickering JE, Galante J, Weightman AL, Morris D, Kelson M, Dolwani S (2016) Aspirin in the treatment of cancer: reductions in metastatic spread and in mortality: a systematic review and meta-analyses of published studies. *PLoS One* **11**(4): e0152402.
- Jacobs EJ, Newton CC, Gapstur SM, Thun MJ (2012) Daily aspirin use and cancer mortality in a large US cohort. *J Natl Cancer Inst* **104**(16): 1208–1217.
- Kim C, Prasad V (2015) Cancer drugs approved on the basis of a surrogate end point and subsequent overall survival: an analysis of 5 Years of US Food and Drug Administration approvals. *JAMA Intern Med* **175**(12): 1992–1994.
- Lambert PC, Royston P (2009) Further development of flexible parametric models for survival analysis. *Stata J* **9**(2): 265.
- Langley RE, Burdett S, Tierney JF, Cafferty F, Parmar MK, Venning G (2011) Aspirin and cancer: has aspirin been overlooked as an adjuvant therapy? *Br J Cancer* **105**(8): 1107–1113.
- Langley RE, Rothwell PM (2014) Aspirin in gastrointestinal oncology: new data on an old friend. *Curr Opin Oncol* **26**(4): 441–447.
- Macfarlane TV, Murchie P, Watson MC (2015) Aspirin and other non-steroidal anti-inflammatory drug prescriptions and survival after the diagnosis of head and neck and oesophageal cancer. *Cancer Epidemiol* **39**(6): 1015–1022.
- McCowan C, Munro AJ, Donnan PT, Steele RJ (2013) Use of aspirin post-diagnosis in a cohort of patients with colorectal cancer and its association with all-cause and colorectal cancer specific mortality. *Eur J Cancer* **49**(5): 1049–1057.
- Paleari L, Puntoni M, Clavarezza M, DeCensi M, Cuzick J, DeCensi A (2015) PIK3CA mutation, aspirin use after diagnosis and survival of colorectal cancer: a systematic review and meta-analysis of epidemiological studies. *Clin Oncol (R Coll Radiol)* **28**(5): 317–326.
- Patrignani P, Tacconelli S, Piazzuelo E, Di Francesco L, Dovizio M, Sostres C, Marcantoni E, Guillem-Llobat P, Del Boccio P, Zucchelli M, Patrono C, Lanasa A (2014) Reappraisal of the clinical pharmacology of low-dose aspirin by comparing novel direct and traditional indirect biomarkers of drug action. *J Thromb Haemost* **12**(8): 1320–1330.
- Ray WA (2003) Evaluating medication effects outside of clinical trials: new-user designs. *Am J Epidemiol* **158**(9): 915–920.
- Reimers MS, Bastiaannet E, Langley RE, van ER, van Vlierberghe RL, Lemmens VE, van Herk-Sukel MP, van WT, Fodde R, Kuppen PJ, Morreau H, van de Velde CJ, Liefers GJ (2014) Expression of HLA class I antigen, aspirin use, and survival after a diagnosis of colon cancer. *JAMA Intern Med* **174**(5): 732–739.

- Rothwell PM, Fowkes FG, Belch JF, Ogawa H, Warlow CP, Meade TW (2011) Effect of daily aspirin on long-term risk of death due to cancer: analysis of individual patient data from randomised trials. *Lancet* **377**(9759): 31–41.
- Rothwell PM, Price JF, Fowkes FG, Zanchetti A, Roncaglioni MC, Tognoni G, Lee R, Belch JF, Wilson M, Mehta Z, Meade TW (2012) Short-term effects of daily aspirin on cancer incidence, mortality, and non-vascular death: analysis of the time course of risks and benefits in 51 randomised controlled trials. *Lancet* **379**(9826): 1602–1612.
- Sarfati D, Blakely T, Pearce N (2010) Measuring cancer survival in populations: relative survival vs cancer-specific survival. *Int J Epidemiol* **39**(2): 598–610.
- Simon R, Makuch RW (1984) A non-parametric graphical representation of the relationship between survival and the occurrence of an event: application to responder versus non-responder bias. *Stat Med* **3**(1): 35–44.
- StataCorp (2011) *Stata Statistical Software: Release 12*. College Station, TX, USA: StataCorp LP.
- Stricker BH, Stijnen T (2010) Analysis of individual drug use as a time-varying determinant of exposure in prospective population-based cohort studies. *Eur J Epidemiol* **25**(4): 245–251.
- Suissa S, Azoulay L (2012) Metformin and the risk of cancer: time-related biases in observational studies. *Diabetes Care* **35**(12): 2665–2673.
- Sullivan R, Alatisse OI, Anderson BO, Audisio R, Autier P, Aggarwal A, Balch C, Brennan MF, Dare A, D'Cruz A, Eggermont AMM, Fleming K, Gueye SM, Hagander L, Herrera CA, Holmer H, Ilbawi AM, Jarnheimer A, Ji J-f, Kingham TP, Liberman J, Leather AJM, Meara JG, Mukhopadhyay S, Murthy SS, Omar S, Parham GP, Pramesh CS, Riviello R, Rodin D, Santini L, Shrikhande SV, Shrima M, Thomas R, Tsunoda AT, van de Velde C, Veronesi U, Vijaykumar DK, Watters D, Wang S, Wu Y-L, Zeiton M, Purushotham A (2015) Global cancer surgery: delivering safe, affordable, and timely cancer surgery. *Lancet Oncol* **16**(11): 1193–1224.
- Thun MJ, Jacobs EJ, Patrono C (2012) The role of aspirin in cancer prevention. *Nat Rev Clin Oncol* **9**(5): 259–267.
- van Herk-Sukel MP, van de Poll-Franse LV, Lemmens VE, Vreugdenhil G, Puijdt JF, Coebergh JW, Herings RM (2010) New opportunities for drug outcomes research in cancer patients: the linkage of the Eindhoven Cancer Registry and the PHARMO Record Linkage System. *Eur J Cancer* **46**(2): 395–404.
- Vandenbroucke JP (2004) When are observational studies as credible as randomised trials? *Lancet* **363**(9422): 1728–1731.
- Yood MU, Campbell UB, Rothman KJ, Jick SS, Lang J, Wells KE, Jick H, Johnson CC (2007) Using prescription claims data for drugs available over-the-counter (OTC). *Pharmacoepidemiol Drug Saf* **16**(9): 961–968.

This work is published under the standard license to publish agreement. After 12 months the work will become freely available and the license terms will switch to a Creative Commons Attribution-NonCommercial-Share Alike 4.0 Unported License.

Supplementary Information accompanies this paper on British Journal of Cancer website (<http://www.nature.com/bjc>)