

Microbiology and antibiotic susceptibility patterns in spontaneous bacterial peritonitis: A study of two Dutch cohorts at a 10-year interval

United European Gastroenterology Journal
2018, Vol. 6(4) 614–621
© Author(s) 2017
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/2050640617744456
journals.sagepub.com/home/ueg



Rosalie C Oey¹, Robert A de Man¹, Nicole S Erler^{2,3}, Annelies Verbon^{4,5} and Henk R van Buuren¹

Abstract

Background: Recent investigations suggest an increasing prevalence of Gram-positive and antibiotic-resistant bacteria causing spontaneous bacterial peritonitis (SBP), probably related to changes in antibiotic prescription patterns, in particular more widespread and long-term use of antibiotic prophylaxis with quinolones.

Objective: The primary objective of this study was to assess potential changes in the microbiology of SBP in two patient cohorts studied at a 10-year interval. Further aims were to study prognostic factors and outcome of SBP.

Methods: A retrospective double-cohort study, including all ascitic cultures from patients with cirrhosis obtained 2003–2005 and 2013–2014, was conducted.

Results: In total 312 patients were included, 125 patients in the first and 187 patients in the second cohort. SBP was diagnosed in 132 of 840 analyzed ascitic fluid samples; 62 samples were culture positive. An increase of Gram-positive bacterial isolates was noted from 26% to 46% between cohorts ($p = 0.122$). The prevalence of multidrug-antibiotic-resistant pathogens increased from 25% to 32% ($p = 0.350$). Survival after SBP among the two cohorts was comparable.

Conclusion: This single-center study in the Netherlands found a modest but nonsignificant increase in the proportion of patients with SBP caused by Gram-positive bacteria and multidrug-antibiotic-resistant bacteria over a 10-year period. Our findings differ from reported data in other countries and suggest empiric antibiotic prophylaxis and treatment of SBP should be based on national and regional microbiological findings and resistance patterns.

Keywords

Antimicrobial resistance, liver cirrhosis, microbiology, prognosis, spontaneous bacterial peritonitis

Received: 13 July 2017; accepted: 24 October 2017

Key summary

- Spontaneous bacterial peritonitis (SBP) is a common complication in patients with liver cirrhosis with a 30-day mortality of 33%.
- A modest but non-significant increase of Gram-positive bacteria and multidrug-antibiotic-resistant bacteria causing SBP was noted over a 10-year period in the Netherlands.
- There are substantial regional differences in bacterial flora and antibiotic susceptibility causing SBP.
- Recommendations in antibiotic prophylaxis and treatment of SBP should carefully take microbiologic geographical differences into account.

¹Department of Gastroenterology and Hepatology, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

²Department of Biostatistics, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

³Department of Epidemiology, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

⁴Department of Medical Microbiology and Infectious Diseases, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

⁵Department of Internal Medicine, Erasmus MC, University Medical Center, Rotterdam, the Netherlands

Corresponding author:

Rosalie C Oey, Erasmus MC, University Medical Center, Department of Gastroenterology and Hepatology, room Ca-411, PO Box 2040, 3000 CA Rotterdam, the Netherlands.
Email: r.oey@erasmusmc.nl

Introduction

Spontaneous bacterial peritonitis (SBP) is a common infection in patients with cirrhosis and ascites. Reportedly, this infection can be diagnosed in up to 30% of cirrhotic patients with ascites who are admitted to the hospital. SBP is associated with significant morbidity and mortality.^{1–4} Intestinal bacterial translocation, altered immunity and the presence of ascites are key in the development of SBP.^{5–7}

SBP is diagnosed by a polymorphonuclear neutrophil (PMN) count in ascitic fluid equal to or greater than 250/ μ l. Approximately 40% of SBP episodes are culture positive.^{3,8,9} Numerous, in particular older, studies reported that Gram-negative enteric bacteria were involved in the majority of SBP episodes. International guidelines recommend third-generation cephalosporin as empirical treatment for SBP and quinolones for secondary prophylaxis.^{10,11} However, in the last decade Gram-positive bacteria and antibiotic-resistant bacteria have been increasingly found to cause SBP.^{3,8,9,12–14} This change in microbiology has been attributed to long-term and widespread quinolone use and increased prevalence of hospital and intensive care unit admissions. These findings have raised doubts about the currently recommended antibiotic strategy in SBP.

The prevalence of antibiotic-resistant pathogens substantially differs geographically.¹⁵ Antibiotic consumption has been identified as the main cause for increasing rates of antibiotic resistance. The Netherlands is known for a restrictive antibiotic policy and has had the lowest antibiotic use in Europe for years.^{15–17} Consequently, microbiological study results in SBP in our country could differ from those observed in other countries—i.e. Spain, Greece, Germany and the United States—over time. This would mean that international guidelines for prophylaxis and treatment of SBP would need to differentiate between countries based on antibiotic resistance rates. Therefore, we investigated causative microorganisms in two patient cohorts who were hospitalized with a 10-year interval in a tertiary referral hospital in the Netherlands. In addition, we aimed to identify the patients most at risk for SBP and to evaluate the associated short- and long-term survival.

Materials and methods

All consecutive ascitic cultures performed in patients with cirrhosis between January 2003 and December 2005 (first cohort) and between January 2013 and December 2014 (second cohort) at Erasmus MC, University Medical Center, Rotterdam, were included. Demographic, clinical, biochemical and survival data from patient hospital records were retrospectively studied to evaluate the prevalence, risk factors,

microbiology, and mortality of SBP. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in an approval by the ethical review board of the Erasmus MC in February 2017.

The diagnosis of cirrhosis was based on liver histology or a combination of clinical, biochemical, and radiologic findings.¹⁸ SBP was defined as a PMN count equal to or greater than 250/ μ l in ascites without evidence of an intra-abdominal source of infection.^{10,11} All ascites samples obtained during the two study periods were studied, implying that, if applicable, multiple samples per patient were taken into account. However, only the first positive culture per SBP episode, and thus one culture per SBP episode, was included in the analysis. Multidrug-antibiotic-resistant organisms (MDR) were defined, according to international guidelines, as an acquired resistance to at least three antibiotic classes.¹⁹ Nosocomial acquisition was defined as SBP diagnosed at least 48 hours after hospital admission. During the study periods, the standard antibiotic prophylaxis for SBP was norfloxacin 400 mg daily.^{10,11,20} The primary choice for empirical antibiotic treatment for SBP was ceftriaxone 2000 mg daily for five to seven days and the secondary choice was amoxicillin/clavulanic acid 1000/200 mg every eight hours for five to seven days, according to international guidelines.^{10,11,20}

All ascitic fluid samples were routinely analyzed in the central clinical laboratory with automated determination of the white blood cell count with differential. In addition, at least 10 ml of ascitic fluid was inoculated at bedside under sterile conditions in aerobic and anaerobic blood culture bottles (Bactec[®]) for culture in the central medical microbiology laboratory. For identification of positive cultures, the ascitic fluid was plated on agar and current identification methods were used. Susceptibility was determined with the VITEK[®] 2 system (VITEK AMS; bioMerieux Vitek Systems Inc, Hazelwood, MO, USA). Cultures collected until 2013 were called resistant using Clinical and Laboratory Standards Institute criteria; later cultures were called resistant using European Committee on Antimicrobial Susceptibility Testing criteria 2013.

Statistical analysis

A mean and standard deviation (SD) was computed for approximately normally distributed variables and compared using the Student's *t* test. Non-normally distributed continuous variables were summarized by their median and interquartile range (IQR), and compared using the Mann-Whitney ranks sum test. Categorical variables were expressed with percentages and compared using the Chi-square test. A two-sided *p* value <0.05 was considered significant. Patients were

followed up to a maximum of one year. This time frame was chosen based on the severity of decompensated advanced chronic liver disease. Survival was analyzed using the Kaplan-Meier method. The actual 30-day mortality and one-year mortality was calculated after the first ascitic fluid analysis; both liver transplantation and death were considered as events. The survival rates were compared using log-rank test. Univariable and multivariable Cox's proportional hazard analyses were carried out to identify independent predictors for 30-day mortality and one-year mortality after SBP. The variables selected for univariable analysis were based on previous studies: gender, age, etiology of liver disease, community- or nosocomial-acquired SBP, positive microbial ascites culture, causative micro-organism, antibiotic susceptibility, use of antibiotic prophylaxis, use of immunosuppressant drug, hepatocellular carcinoma (HCC), model for end-stage liver disease (MELD) score, albumin in serum, platelets in serum, and protein in ascites at time of ascites analyses. Variables with a p value of <0.10 in univariate analysis were included in the multivariate Cox's proportional hazards model.

Results

In the first (2003–2005) cohort of 125 patients, 343 ascitic fluid samples were obtained for analysis. In the second (2013–2014) cohort of 187 patients, 497 samples were obtained. The diagnosis of SBP was established in 132 of the total 840 (16%) ascitic fluid samples in 95 patients (Figure 1).

The total study population included 197 men and 115 women with a mean age of 56 years (± 12) and a mean MELD score of 19 (± 8). Norfloxacin was used by 12% and 10% of patients at the time of paracentesis in the first and second cohort, respectively ($p = 0.638$).

The clinical characteristics of patients with and without SBP are shown in Table 1. Patients with SBP had more frequent liver disease of autoimmune origin, more frequently used immunosuppressive drugs ($p = 0.012$) and had higher baseline MELD scores ($p = 0.020$).

Microbiology

In the two cohorts a culture-positive SBP was found in 23/56 (41%) and 33/76 (43%) episodes with SBP, respectively (Figure 1). The microbiological culture




Figure 1. Flowchart of study population. PMN: polymorphonuclear neutrophil.

Table 1. Clinical characteristics of the study population with and without SBP.

	SBP negative (n = 217)	SBP positive (n = 95)	p value
Male (%)	138 (64%)	59 (62%)	0.802
Age in years	57 (±12)	54 (±13)	0.036
Etiology of cirrhosis (%)			
Alcohol	92 (42%)	31 (33%)	0.045
Viral	51 (24%)	23 (24%)	
Autoimmune	28 (13%)	24 (25%)	
Other	46 (21%)	17 (18%)	
Child-Pugh class (%)			
A	25 (12%)	5 (5%)	0.215
B	74 (34%)	33 (35%)	
C	118 (54%)	57 (60%)	
MELD score	19 (±8)	21 (±8)	0.020
Creatinine (µmol/l)	96.8 (±1.7)	110.1 (±1.8)	0.052
Albumin (g/l)	29 (±6)	29 (±6)	0.912
INR	1.5 (±1.0)	2.0 (±1.5)	0.006
Bilirubin (µmol/l)	62 (±3)	76 (±4)	0.260
Thrombocytes (10 ⁹ /l)	107 (±2)	97 (±2)	0.304
Ascites protein (g/l)	10.6 (±2.0)	16.3 (±2.6)	0.009
Hepatocellular carcinoma	30 (14%)	11 (12%)	0.589
Diabetes mellitus	37 (17%)	11 (12%)	0.218
Use of immunosuppressant drug	21 (10%)	19 (20%)	0.012
Use of norfloxacin	26 (12%)	8 (8%)	0.353

SBP: spontaneous bacterial peritonitis; MELD: model for end-stage liver disease; INR: international normalized ratio. Significant *p* values (<0.050) are highlighted in boldface.

results are shown in Table 2. In the first cohort, 61% of culture-positive SBP was due to Gram-negative bacteria vs. 51% in the second cohort. *Candida albicans* was isolated in four cultures. Although the percentage SBP with Gram-positive organisms increased over time, the differences were not statistically significant (*p* = 0.122) (Table 2).

Antibiotic susceptibility patterns

In the first cohort 5/20 (25%) of the isolated bacteria were MDR versus 12/38 (32%) in the second cohort (Table 3). There was no significant change in prevalence of MDR organisms over time (*p* = 0.350). In the MDR organisms, the most frequently detected resistance mechanism was due to extended-spectrum beta-lactamase (ESBL) production (*Escherichia coli* *n* = 9, *Klebsiella* *n* = 1). Furthermore, methicillin-resistant *Staphylococcus aureus* (MRSA) (*n* = 4), intrinsically cephalosporin-resistant *Enterobacter* (*n* = 2), and vancomycin-resistant Enterococci (VRE)

Table 2. Microbiological findings in two cohorts of patients; 62 organisms were identified in 56 episodes of culture-positive spontaneous bacterial peritonitis.

	Cohort 2003–2005 (n = 23)	Cohort 2013–2014 (n = 39) ^a
Gram-negative bacteria	14 (61%)	20 (51%)
<i>Escherichia coli</i>	9	13
<i>Enterobacter aerogenes</i>	–	2
<i>Enterobacter cloacae</i>	–	2
<i>Pseudomonas aeruginosa</i>	2	–
<i>Klebsiella pneumoniae</i>	1	3
<i>Morganella morganii</i>	1	–
Aeromonas spp.	1	–
Gram-positive bacteria	6 (26%)	18 (46%)
<i>Staphylococcus aureus</i>	1	3
<i>Staphylococcus haemolyticus</i>	–	1
Staphylococcus (coagulase negative)	3	3
<i>Enterococcus faecium</i>	–	5
<i>Streptococcus oralis</i>	1	3
<i>Streptococcus anginosus</i>	–	1
<i>Streptococcus salivarius</i>	–	1
<i>Streptococcus viridans</i>	–	1
<i>Streptococcus pneumoniae</i>	1	–
Yeast	3 (13%)	1 (3%)
<i>Candida albicans</i>	3 (13%)	1 (3%)

^aIncluding six cultures showing two microorganisms.

Table 3. Antimicrobial susceptibility patterns of bacteria from culture-positive spontaneous bacterial peritonitis (SBP).

	Bacterial isolates in SBP (n = 58)		
	Cohort 2003–2005 (n = 20)	Cohort 2013–2014 (n = 38)	p value
Multidrug resistant	5 (25%)	12 (32%)	0.350
Norfloxacin resistant ^a	3 (15%)	9 (24%)	0.274
Ceftriaxon resistant ^a	3 (15%)	5 (13%)	0.952
Amoxicillin/clavulanic acid resistant ^a	8 (40%)	8 (21%)	0.254

^aIntrinsically and/or acquired antimicrobial resistance.

(*n* = 1) were found. There was no evidence that the risk of SBP caused by MDR organisms was related to a Gram-negative or Gram-positive microbiologic isolate (*p* = 0.192), a nosocomial acquisition of the infection (*p* = 0.677), a previous history of SBP (*p* = 0.245), or the use of antibiotic prophylaxis (*p* = 0.316).

Three of the 20 (15%) and six of the 38 (16%) isolated organisms were norfloxacin resistant, in the first and second cohort, respectively ($p = 0.274$).

Analysis with respect to ceftriaxone showed that in the first cohort 3/20 (15%) of bacterial isolates were resistant to this agent as compared to 5/38 (13%) in the second cohort. Amoxicillin/clavulanic acid-resistance was slightly more prevalent. In the first cohort 6/20 (30%) organisms were found to be resistant versus 8/38 (21%) in the second cohort. The frequency of ceftriaxone and amoxicillin/clavulanic acid-resistant organisms did not differ significantly between the cohorts ($p = 0.952$ and $p = 0.254$, respectively) (Table 3).

Survival

At one year, 153 patients had died (49%) and 48 patients had received a liver transplant (15%). Furthermore, 83 patients (27%) were alive after one year, while 28 patients (9%) were lost to follow-up. The median follow-up time of the patients not reaching the endpoint of death or liver transplantation was 365 days (IQR 12 days). The survival of patients with SBP did not differ significantly between the cohorts (log-rank $p = 0.442$), nor did survival differ significantly between the cohorts for patients without SBP (log-rank $p = 0.216$).

The median survival after the first ascites analyses was 168 days for SBP-negative patients and 77 days for SBP-positive patients (log-rank $p = 0.001$) (Figure 2). The 30-day mortality rate was 33% (32/95) for patients with SBP compared to 25% (55/217) for patients without SBP (Figure 2). Univariable and multivariable Cox-regression analyses were carried out to identify risk factors for 30-day mortality and one-year mortality after SBP. MELD score was the only independent

predictive factor for 30-day mortality (hazard ratio (HR) 1.106 per point, 95% CI 1.061–1.154, $p < 0.001$) and one-year mortality (HR 1.060 per point, 95% CI 1.030–1.091, $p < 0.001$) (Table 4).

Discussion

In this single-center study in the Netherlands, the microbiological characteristics and antibiotic susceptibility patterns of organisms causing SBP in liver cirrhosis patients were compared between the periods 2003–2005 and 2013–2014. No significant increase in Gram-positive bacteria was observed and Gram-negative bacteria remained the primary cause. Bacteria resistant to empirical treatment with third-generation cephalosporin accounted for 13%–15% of all causative pathogens.

There was no evidence that mortality was influenced by causative microorganisms, antibiotic susceptibility, use of prophylactic antibiotics, intensive care admission or a nosocomial acquisition of SBP. The main predictors for mortality were age, MELD score, and platelet count.

Our most important finding is that in the era of quinolone prophylaxis for SBP, we cannot confirm observations made elsewhere regarding a significant increase in Gram-positive and MDR organisms causing SBP.^{3,8,9,12,13} Although in the Netherlands antibiotics are used prudently, a rise in quinolone use in the last decades has been described.^{21,22}

Third-generation cephalosporin may poorly cover the causative pathogens in SBP, with reported antibiotic resistance rates ranging from 57% to 69%.^{9,23,24} However, our results show a susceptibility rate of 85–87%. We hypothesize this difference can be most likely attributed to different national antibiotic policies. High consumption of antibiotics has been related to higher rates of antibiotic resistance.¹⁵ The Netherlands has always had a restrictive national policy regarding antibiotic prescription and a conservative approach toward the prescription of new broad-spectrum antibiotic agents.^{15,16,25}

The microbiology and susceptibility patterns' differences can be hospital and region dependent, implying difficulties with recommending antibiotic treatment and prophylaxis in international guidelines. The results of this study in the Netherlands do not support the need to revise guidelines as previously proposed.^{3,23} Empirical antibiotic treatment should be based on known regional and national differences of antibiotic resistance patterns.

A previous study from our institution on the microbiology of SBP in the period 1987–1991, before the implementation of long-term quinolone prophylaxis in the relevant patient population, reported that causative




Figure 2. One-year mortality after first ascites analysis. Spontaneous bacterial peritonitis (SBP)-negative patients (solid line) have a median survival of 168 days and SBP-positive patients (dotted line) of 77 days (log-rank $p = 0.001$).

Table 4. Demographic and clinical factors after SBP in 95 patients predicting one-year mortality using Cox-regression analysis.

	Univariate			Multivariate		
	HR	95% CI	<i>p</i> value	HR	95% CI	<i>p</i> value
Gender			<0.001			
Female (ref.)	1					
Male	0.424	0.263–0.682				
Age (per year)	0.999	0.981–1.017	0.911			
Etiology			0.206			
Alcohol (ref.)	1					
Viral	0.841	0.444–1.596				
Autoimmune	1.545	0.836–2.858				
Other	0.837	0.431–1.625				
Acquisition SBP			0.078			
Community (ref.)	1					
Nosocomial	1.761	0.938–3.303				
Positive microbial ascites culture	1.406	0.886–2.231	0.148			
Causative microorganism type			0.561			
Gram-negative bacteria (ref.)	1					
Gram-positive bacteria	0.764	0.359–1.625				
Yeast	1.511	0.444–5.146				
Multidrug-resistant microorganism	1.989	0.972–4.073	0.060			
Antibiotic prophylaxis	0.662	0.286–1.531	0.335			
Immunosuppressant use	0.508	0.260–0.992	0.047			
HCC	0.971	0.465–2.026	0.937			
MELD score (per point)	1.060	1.030–1.091	<0.001	1.060	1.030–1.091	<0.001
Albumin in serum (per point)	0.967	0.927–1.008	0.109			
Platelets in serum (<150 × 10 ⁹ /l)	2.179	1.226–3.870	0.008			
Low protein in ascites (<15 g/l)	1.287	0.530–3.124	0.578			

SBP: spontaneous bacterial peritonitis; MELD: model for end-stage liver disease; HCC: hepatocellular carcinoma; HR: hazard ratio; CI: confidence interval. Significant *p* values in univariable analysis (<0.100) and in multivariable analysis (<0.050) are highlighted in boldface.

pathogens were isolated in 25 of 31 SBP episodes.²⁶ Gram-negative bacteria were detected in 60% of the episodes and Gram-positive bacteria in 40% of the episodes. Despite the small sample size, the proportions of causative pathogens seem comparable to those identified in the cohorts reported here.

Although optimizations in bacterial culture techniques have been implemented, an organism was isolated in a minority of all SBP episodes (40%). This is a stable percentage over the last decade and similar proportions have been documented in other studies.^{3,8,9} It may be expected that, with technologies arising from bacterial DNA detection and microbiome studies, more causative pathogens can be identified rapidly in the future for targeted antibiotic therapy.

There are a few limitations of the study regarding methodology. This study was designed as a retrospective, double-cohort study, which implied some laboratory and clinical data were missing. For instance, we

have no information about short-term antibiotic use prescribed by general practitioners and clinicians outside the hospital, but it is not to be expected that this would differ significantly between the cohorts. Furthermore, multiple tests have been performed that may lead to an increased risk of finding spurious significant results and results should be interpreted while keeping this in mind. Prospective cohort studies comparing multiple regions during a large time frame could provide more insight as to the microbiology of SBP in diverse regions over time.

In this study we show that the microbiology of pathogens causing SBP did not change significantly in our center over the last decade. These findings suggest that guidelines with respect to antibiotic prophylaxis and treatment of SBP should carefully take into account potential national and regional differences in the microorganisms causing SBP and antibiotic resistance patterns.

Acknowledgements

We would like to thank Joyce Kuiper and Dirk Jan Bac for their previous work, in particular the acquisition of data relevant to the present study.

Author contributions include the following: R.C. Oey: study concept and design, acquisition of data, analysis and interpretation of data, critical revision of the manuscript for important intellectual content, and finalizing the article; R.A. de Man: study concept and design, analysis and interpretation of data, critical revision of the manuscript for important intellectual content, and finalizing the article; N.S. Erler: statistical analysis, critical revision of the manuscript for important intellectual content, and approval of the final article; A. Verbon: interpretation of data, critical revision of the manuscript for important intellectual content, and approval of the final article; H.R. van Buuren: study concept and design, analysis and interpretation of data, critical revision of the manuscript for important intellectual content, and finalizing the article.

Ethics approval

The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in an approval by the ethical review board of the Erasmus MC in February 2017.

Informed consent

The ethical review board determined that written informed consent was not necessary to obtain from each patient included in this study due to its retrospective design.

Declaration of conflicting interests

None declared.

Funding

This study was sponsored by the Foundation for Liver and Gastrointestinal Research Rotterdam (SLO).

References

- Rimola A, García-Tsao G, Navasa M, et al. Diagnosis, treatment and prophylaxis of spontaneous bacterial peritonitis: A consensus document. International Ascites Club. *J Hepatol* 2000; 32: 142–153.
- Caly WR and Strauss E. A prospective study of bacterial infections in patients with cirrhosis. *J Hepatol* 1993; 18: 353–358.
- Fernández J, Navasa M, Gómez J, et al. Bacterial infections in cirrhosis: Epidemiological changes with invasive procedures and norfloxacin prophylaxis. *Hepatology* 2002; 35: 140–148.
- Tandon P and García-Tsao G. Renal dysfunction is the most important independent predictor of mortality in cirrhotic patients with spontaneous bacterial peritonitis. *Clin Gastroenterol Hepatol* 2011; 9: 260–265.
- Wiest R and García-Tsao G. Bacterial translocation (BT) in cirrhosis. *Hepatology* 2005; 41: 422–433.
- Wiest R, Lawson M and Geuking M. Pathological bacterial translocation in liver cirrhosis. *J Hepatol* 2014; 60: 197–209.
- Lutz P, Nischalke HD, Strassburg CP, et al. Spontaneous bacterial peritonitis: The clinical challenge of a leaky gut and a cirrhotic liver. *World J Hepatol* 2015; 7: 304–314.
- Cholongitas E, Papatheodoridis GV, Lahanas A, et al. Increasing frequency of Gram-positive bacteria in spontaneous bacterial peritonitis. *Liver Int* 2005; 25: 57–61.
- Friedrich K, Nüssle S, Rehlen T, et al. Microbiology and resistance in first episodes of spontaneous bacterial peritonitis: Implications for management and prognosis. *J Gastroenterol Hepatol* 2016; 31: 1191–1195.
- European Association for the Study of the Liver. EASL clinical practice guidelines on the management of ascites, spontaneous bacterial peritonitis, and hepatorenal syndrome in cirrhosis. *J Hepatol* 2010; 53: 397–417.
- Runyon BA. Practice guideline. Management of adult patients with ascites due to cirrhosis: Update 2012, http://www.aasld.org/sites/default/files/guideline_documents/adultascitesenhanced.pdf. Retrieval June 1st, 2017.
- Tandon P, Delisle A, Topal JE, et al. High prevalence of antibiotic-resistant bacterial infections among patients with cirrhosis at a US liver center. *Clin Gastroenterol Hepatol* 2012; 10: 1291–1298.
- Fernández J, Acevedo J, Castro M, et al. Prevalence and risk factors of infections by multiresistant bacteria in cirrhosis: A prospective study. *Hepatology* 2012; 55: 1551–1561.
- Alexopoulou A, Papadopoulos N, Eliopoulos DG, et al. Increasing frequency of gram-positive cocci and gram-negative multidrug-resistant bacteria in spontaneous bacterial peritonitis. *Liver Int* 2013; 33: 975–981.
- Goossens H, Ferech M, Vander Stichele R, et al. Outpatient antibiotic use in Europe and association with resistance: A cross-national database study. *Lancet* 2005; 365: 579–587.
- Cars O, Mölsted S and Melander A. Variation in antibiotic use in the European Union. *Lancet* 2001; 357: 1851–1853.
- Organisation for Economic Co-operation and Development (OECD) iLibrary. Health at a glance 2015. Prescribing in primary care, http://dx.doi.org/10.1787/health_glance-2015-46-en. Retrieval June 1st, 2017.
- Schuppan D and Afdhal NH. Liver cirrhosis. *Lancet* 2008; 371: 838–851.
- Magiorakos AP, Srinivasan A, Carey RB, et al. Multidrug-resistant, extensively drug-resistant and pan-drug-resistant bacteria: An international expert proposal for interim standard definitions for acquired resistance. *Clin Microbiol Infect* 2012; 18: 268–281.
- Stichting Werkgroep Antibioticabeleid. Peritonitis-primair (SBP) SWAB, <http://swabid.nl/node/7362>. Retrieval June 1st, 2017.
- Haeseker MB, Dukers-Muijers NH, Hoebe CJ, et al. Trends in antibiotic prescribing in adults in Dutch general practice. *PLoS One* 2012; 7: e51860.
- Tyrstrup M, van der Velden A, Engstrom S, et al. Antibiotic prescribing in relation to diagnoses and consultation rates in Belgium, the Netherlands and Sweden:

- Use of European quality indicators. *Scand J Prim Health Care* 2017; 35: 10–18.
23. Novovic S, Semb S, Olsen H, et al. First-line treatment with cephalosporins in spontaneous bacterial peritonitis provides poor antibiotic coverage. *Scand J Gastroenterol* 2012; 47: 212–216.
 24. Lutz P, Nischalke HD, Kramer B, et al. Antibiotic resistance in healthcare-related and nosocomial spontaneous bacterial peritonitis. *Eur J Clin Invest* 2017; 47: 44–52.
 25. Janknegt R, Monkelbaan JF, Stobberingh E, et al. Antibiotic policies in Dutch hospitals for the treatment of patients with serious infection. *J Antimicrob Chemother* 1994; 34: 1059–1069.
 26. Bac DJ, Siersema PD, Mulder PGH, et al. Spontaneous bacterial peritonitis: Outcome and predictive factors. *Eur J Gastroenterol Hepatol* 1993; 5: 635–640.