



Photos: HarvestPlus

SCIENCE BRIEF: BIOFORTIFICATION NO. 2 | February 2018

Biofortification addresses mineral and vitamin deficiencies

Malnutrition is estimated to contribute to 45% of all child deaths in developing countries. More than two billion people do not get enough essential vitamins and minerals because their diets are not properly balanced. Even when consumption of food staples is high, so that the poor do not go hungry, more nutritious foods are not affordable and getting more so.

The deficiencies of Vitamin A, iron and zinc are the most widespread and serious. These result in higher mortality and morbidity, reduced cognitive abilities and lower work performance.

HarvestPlus and its partners seek to reduce such “hidden hunger” by breeding staple foods that are high yielding and dense in minerals and vitamins (Bouis and Saltzman 2017a), a process known as biofortification. Biofortified varieties, developed through research at a central location, can be made available to multiple countries, and, once released, are available in national food systems year after year at no additional cost to farmers and consumers.

For more information on the 12 biofortified crops being developed at CGIAR centers, see Bouis and Saltzman (2017b). For information about progress on biofortification in general, see Bouis and Saltzman (2017a).

Vitamin A-Biofortified Maize: Exploiting Native Genetic Variation for Nutrient Enrichment

Vitamin A is important in immune function. Approximately one third of children under the age of five are at risk for Vitamin A deficiency (WHO 2009). This is the leading cause of preventable childhood blindness.

Nutrition trials in countries administering Vitamin A capsules resulted on average in a 24% reduction in child mortality. By breeding staple crops with higher amounts of Vitamin A, the supply of Vitamin A in our food sources can be sustainably increased.

Maize provides approximately 30% of the total calories of more than 4.5 billion people in developing countries, and is the most widely eaten food staple in Africa. Poor quality diets, dominated by food staples, are often deficient in minerals and vitamins, but maize can provide sufficient quantities of Provitamin A (proVA), which the body converts to Vitamin A.

Maize can grow in diverse environments and has a high potential to be bred to provide productive cultivars that are attractive to farmers and consumers, including for their nutritional properties.

HarvestPlus has collaborated with CIMMYT, IITA, EMBRAPA and other partners to develop maize varieties that are high yielding, profitable, acceptable to consumers, and also rich in Provitamin A carotenoids. Beta-carotene is the most common proVA carotenoid in plants. The breeding target for maize was set at 15 µg/g of beta-carotene, to provide an additional 50% of the estimated average requirement for Vitamin A in maize-eating regions. Maize with these levels of beta-carotene would be an effective contributor to reducing Vitamin A deficiency.

ProVA maize breeding is an excellent example of the effective use of both adapted and exotic genetic resources in crop improvement. Breeders found a wide range of levels of proVA carotenoids in temperate, tropical and subtropical germplasm during the initial screening studies (Ortiz-Monasterio et al. 2007; Menkir et al. 2008).

BOX 1

CIMMYT and IITA manage diverse germplasm collections of maize

Maize holdings at CIMMYT and IITA comprise 29,900 seed samples. These include the world's largest collection of maize landraces (varieties developed and nurtured by farmers), along with populations of maize's wild relatives, teosinte and *Tripsacum*, and many improved varieties.

The CIMMYT's Genetic Resources Program (www.cimmyt.org/germplasm-bank/) and IITA's Genetic Resources Center (grc-iita.org/) manage the maize germplasm collections, with coordination from the CGIAR Genebank Platform, under the policy umbrella provided by the International Treaty on PGRFA. The programs conserve maize genetic resources, and, together with the Global Maize Program, study, document and facilitate their use.

Most yellow maize grown and consumed throughout the world has on average only 2 µg/g proVA carotenoids. Tropical maize contains less beta-carotene than temperate maize, so the initial breeding sources of high proVA carotenoids were selected from temperate regions (Pixley et al. 2013; Menkir et al. 2017).

ProVA donors were crossed with elite germplasm of white maize that had high yield potential and good agronomic traits, such as disease resistance and drought tolerance (Box 2). For example, IITA breeders used the KU1414 and KU1409 lines from

Thailand, which had been derived from the downy mildew resistant Suwan 1 variety from the Philippines and a Thai Composite. The Thai Composite was developed by inter-crossing 36 maize varieties and cultivars introduced from the Caribbean, Mexico, Central America, South America and other areas.

Backcross breeding was then used to fix the desired orange color. Marker-assisted selection for the desired alleles of key proVA genes accelerated genetic gain and allowed to double, sometimes triple, the total concentration of proVA (Harjes et al. 2008; Yan et al. 2010).

Maize inbred lines with high levels of proVA and desirable agronomic and adaptive traits developed at CIMMYT (for tropical mid-altitudes), IITA (for tropical lowlands), and EMBRAPA in Brazil were then used to develop hybrids and open-pollinated varieties (OPVs) that have high levels of proVA content and are agronomically competitive in target countries. Extensive evaluations were conducted in collaboration with public and private sector partners in the Democratic Republic of Congo, Ghana, Mali, Malawi, Nigeria, Tanzania, Zambia and Zimbabwe. These evaluations led to the selection of promising hybrids and

BOX 2 The path to Vitamin A-enriched maize



Breeding crosses for the first wave (generation) of high proVA maize involved the elite white-grained inbred lines and temperate yellow lines with Provitamin A concentrations ranging from 5 to 13 µg/g.



Each F1 cross was back-crossed to its respective elite line. Selection during the first two cycles of inbreeding was primarily for deep yellow or orange color, as well as for resistances to lodging, and plant and ear diseases.



The best 10–20% of these new lines were selected to move to the next steps of hybrid formation.



The best lines were grown at multiple locations in target countries to ensure suitability and good performance across many environments and under different growing conditions. The breeding program followed a cyclic process to develop varieties that are competitive for crucial traits, including productivity of the seed, grain yield, disease resistance, drought tolerance, food processing quality, taste, and various characteristics that determine acceptability to farmers and consumers.



A second "dose" of high proVA was introduced, by crossing with novel higher provitamin proVA sources, followed by further inbreeding and selection.

Photos: CIMMYT



Vitamin A-enriched maize being harvested in Zambia
Photo: HarvestPlus

OPVs for further testing in national performance and farmer participatory on-farm trials. The testing schemes facilitated the release of more than 40 hybrids and OPVs with up to 90% of the proVA breeding target in eight countries in sub-Saharan Africa and Brazil (Andersson et al. 2017).

Breeder seed of the released proVA biofortified maize OPVs and hybrid parental lines has been multiplied and made available to national programs and private seed companies, who are tasked with producing and marketing these materials. Seed has also been supplied to community-based and other seed producers as well as startup national seed companies to help accelerate multiplication and commercialization of OPV products to farming communities in rural areas with limited market access.

Nutritional studies have shown that proVA biofortified maize is an efficacious source of Vitamin A. Consumption of orange maize has been demonstrated to improve total body Vitamin A stores as effectively as supplementation (Gannon et al. 2014), and significantly improve visual function in marginally Vitamin A deficient children (Palmer et al. 2016).

Conclusion

Crop diversity contributes to a stable, sustainable and diverse food production system and plays an important

BOX 3

Securing Crop Diversity Forever: The Global Genebank Partnership

The CGIAR Genebank Platform aims to conserve the diversity of plant genetic resources in the 11 crop genebanks managed by the CGIAR Centers, in partnership with the Crop Trust, under the International Treaty on Plant Genetic Resources for Food and Agriculture. The Genebank Platform supports the core activities of the CGIAR genebanks: conserving and making available crop and tree diversity. It ensures that the genebanks meet international standards, improve efficiency and facilitate more effective use within an enabling policy environment. For more information, visit <https://www.genebanks.org>.



Genebank Platform

role in improving nutritional outcomes. CGIAR, in partnership with the Crop Trust, is working towards ensuring that such crop diversity is conserved and available forever (Box 3).

There is a long and often complex history behind every modern crop cultivar, but their economic and other benefits far exceed the costs of the collection and conservation of germplasm (Robinson 2000), as well as those of the breeding programs that produced them. This progress described in improving the Provitamin A content of maize varieties and hybrids is just one example of what is made possible by the genetic variation preserved in the genebanks managed by CGIAR and national programs around the world.

Acknowledgments

The authors would like to thank the following for their input in this brief: Thanda Dhliwayo, Natalia Palacios-Rojas, Thokozile Ndhlela and Kevin Pixley of CIMMYT; Abebe Menkir, Bussie Maziya-Dixon, Wende Mengesha and Oladeji Alamu of IITA; Torbert Rocheford of Purdue University and; Maria Cristina Dias Paes and Paulo Evaristo Oliveira Guimarães of EMBRAPA.



References

- Andersson, M.S., Saltzman, A., Virk, P.S. and Pfeiffer, W.H. (2017) Progress update: crop development of biofortified staple food crops under HarvestPlus. *African Journal of Food, Agriculture, Nutrition and Development* 17(2): 11905–11935. Available online at <https://www.afjand.net/Volume17/No2/>.
- Bouis, H.E. and Saltzman, A. (2017a) Improving nutrition through biofortification: A review of evidence from HarvestPlus, 2003 through 2016. *Global Food Security* 12(2017): 49–58. Available online at <https://doi.org/10.1016/j.gfs.2017.01.009>.
- Bouis, H.E. and Saltzman, A. (2017b) Special issue devoted to Biofortification. *African Journal of Food, Agriculture, Nutrition and Development* 17(2). Available online at <https://www.afjand.net/Volume17/No2/>.
- Gannon, B., Kaliwile, C., Arscott, S., Schmaelzle, S., Chileshe, J., Kalungwana, N., Mosonda, M., Pixley, K., Masi, C. and Tanumihardjo, S.A. (2014) Biofortified orange maize is as efficacious as a vitamin A supplement in Zambian children even in the presence of high liver reserves of vitamin A: a community-based, randomized placebo-controlled trial. *American Journal of Clinical Nutrition* 100(6): 1541–50.
- Harjes, C.E., Rocheford, T.R., Bai, L., Brutnell, T.P., Vallabhaneni, R., Williams, M., Wurtzel, E.T., Kandianis, C.B., Sowinski, S.G., Stapleton, A.E., Yan, J. and Buckler, E.S. (2008) Natural genetic variation in lycopene epsilon cyclase tapped for maize biofortification. *Science* 319: 330–333.
- Menkir, A., Liu, W., White, W., Maziya-Dixon, B., and Rocheford, T. (2008) Carotenoid diversity in tropical-adapted yellow maize inbred lines. *Food Chemistry* 109(3): 521–529.
- Menkir, A., Maziya-Dixon, B., Mengesha, W., Rocheford, T. and Oladeji Alamu, E. (2017) Accruing genetic gain in pro-vitamin A enrichment from harnessing diverse maize germplasm. *Euphytica* 213(5): 105.
- Ortiz-Monasterio, I., Palacios-Rojas, N., Meng, E., Pixley, K., Trethowan, R., and Pena, R.J. (2007) Enhancing the mineral and vitamin content of wheat and maize through plant breeding. *Journal of Cereal Science* 46: 293–307.
- Palmer, A., Chileshe J., Hall, A., Barffour, M., Molobeka, N., West Jr., K.P. and Haskell, M.J. (2016) Short-term daily consumption of provitamin A carotenoid biofortified maize has limited impact on breast milk retinol concentrations in Zambia women enrolled in a randomized controlled feeding trial. *Journal of Nutrition* 146: 1783–92.
- Pixley, K., Palacios, N., Babu, R., Mutale, R., Surles, R., Simpungwe, E. (2013) Biofortification of maize with provitamin a carotenoids. In: *Carotenoids and Human Health*. Tanumihardjo, Sherry A. (Ed.) Humana Press. Chapter 17. p 271–292.
- Robinson, J. (2000) Genetic resources impact tracing study. Report prepared for the System-wide Genetic Resources Program (SGRP), Rome, Italy.
- WHO (2009) Global prevalence of vitamin A deficiency in populations at risk 1995–2005. WHO Global Database on Vitamin A Deficiency. Geneva, World Health Organization. http://www.who.int/nutrition/publications/micronutrients/vitamin_a_deficiency/9789241598019/en/.
- Yan, J., Kandianis, C.B., Harjes, C.E., Bai, L., Kim, E-H., Yang, X., Skinner, D.J., Fu, Z., Palacios, N., Li, J., DellaPenna, D., Brutnell, T., Buckler, E.S., Warburton, M.L., and Rocheford, T. (2010) Rare genetic variation at *Zea mays* crtRB1 increases β -carotene in maize grain. *Nature Genetics* 42: 322–329.

Recommended Citation

Menkir, Abebe; Palacios-Rojas, Natalia; Alamu, Oladeji; Dias Paes, Maria Cristina; Dhliwayo, Thanda; Maziya-Dixon, Bussie; Mengesha, Wende; Ndhlela, Thokozile; Oliveira Guimarães, Paulo Evaristo; Pixley, Kevin; Rocheford, Torbert. (2018) Vitamin A-Biofortified Maize: Exploiting Native Genetic Variation for Nutrient Enrichment. *Science Brief: Biofortification* No. 2 (February 2018). CIMMYT, IITA, EMBRAPA, HarvestPlus, and Crop Trust. Bonn, Germany.



CONTACT

Meike S. Andersson (HarvestPlus), Howarth Bouis (HarvestPlus), Nelissa Jamora (Crop Trust) and Michael Major (Crop Trust) are the editors of the *Science Brief: Biofortification* series. For more information, e-mail: science.briefs@croptrust.org or visit: croptrust.org/our-work/science-briefs/ • harvestplus.org • croptrust.org