

Range additivity, shorted operator and the Sherman-Morrison-Woodbury formula

M. Laura Arias^{a,1}, Gustavo Corach^{a,b,1,*}, Alejandra Maestri^{a,b,1}

^a*Instituto Argentino de Matemática “Alberto P. Calderón”
Saavedra 15, Piso 3 (1083), Buenos Aires, Argentina.*

^b*Dpto. de Matemática, Facultad de Ingeniería, Universidad de Buenos Aires.*

Abstract

We say that two operators A, B have the range additivity property if $R(A + B) = R(A) + R(B)$. In this article we study the relationship between range additivity, shorted operator and certain Hilbert space decomposition known as compatibility. As an application, we extend to infinite dimensional Hilbert space operators a formula of Fill and Fishkind related to the well-known Sherman-Morrison-Woodbury formula.

Keywords: operator ranges, shorted operator, Sherman-Morrison-Woodbury formula
2010 MSC: 46C07, 47A64

1. Introduction

In this paper we explore some results implied by range additivity of operators in a Hilbert space \mathcal{H} . Let $L(\mathcal{H})$ be the algebra of bounded linear operators on \mathcal{H} and $L(\mathcal{H})^+$ the cone of positive operators on \mathcal{H} . Consider the set

$$\mathcal{R} := \{(A, B) : A, B \in L(\mathcal{H}) \text{ and } R(A + B) = R(A) + R(B)\},$$

where $R(T)$ denotes the range of T . If $(A, B) \in \mathcal{R}$ we say that A, B satisfy the *range additivity property*. On the other side, we say that a positive operator $A \in L(\mathcal{H})^+$ and a closed subspace $\mathcal{S} \subseteq \mathcal{H}$ are *compatible* if $\mathcal{S} + (A\mathcal{S})^\perp = \mathcal{H}$; in [11] it is shown that A and \mathcal{S} are compatible if and only if there exists an idempotent operator $E \in L(\mathcal{H})$ such that $R(E) = \mathcal{S}$ and E is A -selfadjoint, in the sense that $\langle Ex, y \rangle_A = \langle x, Ey \rangle_A$ for $x, y \in \mathcal{H}$, where $\langle x, y \rangle_A = \langle Ax, y \rangle$. Notice that $\|x\|_A = \langle x, x \rangle_A^{1/2}$ is a seminorm, and that E behaves, with respect to this seminorm, as an orthogonal projection. So, A and \mathcal{S} are compatible if there is an A -orthogonal projection onto \mathcal{S} . One of the main results of the paper is that A and \mathcal{S} are compatible if and only if $(A, I - P_{\mathcal{S}}) \in \mathcal{R}$, where $P_{\mathcal{S}}$ denotes the classical orthogonal projection onto \mathcal{S} . In order to prove this assertion, and some other general facts on range additivity and compatibility, we explore some features of the shorted operator $[S]A$. This

*Corresponding author. Address: IAM-CONICET, Saavedra 15, Piso 3 (1083), Buenos Aires, Argentina.

Email addresses: lauraarias@conicet.gov.ar (M. Laura Arias), gcorach@fi.uba.ar (Gustavo Corach), amaestri@fi.uba.ar (Alejandra Maestri)

¹Partially supported by UBACYT 20020100100250; 2011-2014

operator has been defined by M. G. Krein [25] as

$$[\mathcal{S}]A := \max\{X \in L(\mathcal{H})^+ : X \leq A \text{ and } R(X) \subseteq \mathcal{S}\}.$$

He proved that the maximum for the Löwner ordering (i.e., $C \leq D$ if $\langle C\xi, \xi \rangle \leq \langle D\xi, \xi \rangle$ for every $\xi \in \mathcal{H}$) exists and then applied this construction for a parametrization of the selfadjoint extensions of semi-bounded operators. W. N. Anderson and G. E. Trapp [1] redefined and studied this operator, which can be used in the mathematical study of electrical networks. Here, we use the properties of the shorted operator in order to prove that, for $A, B \in L(\mathcal{H})^+$ such that $R(B)$ is closed, it holds that $(A, B) \in \mathcal{R}$ if and only if A and $N(B)$ are compatible, where $N(B)$ denotes the nullspace of B . In particular, for $B = I - P_{\mathcal{S}}$ we get the assertion above. However, this is not the first manifestation of a relationship between compatibility of A and \mathcal{S} and properties of $[\mathcal{S}]A$. In fact, Anderson and Trapp [1] proved that $[\mathcal{S}]A$ is the infimum, for the Löwner ordering, of the set $\{EAE^* : E \in L(\mathcal{H}), E^2 = E, N(E) = \mathcal{S}^\perp\}$. In [11, Prop. 4.2], [12, Prop. 3.4] it is proven that the infimum is attained if and only if A and \mathcal{S} are compatible. Moreover, it is shown that if $E \in L(\mathcal{H})$ is an idempotent operator such that $AE = E^*A$ and $R(E) = \mathcal{S}$, then $[\mathcal{S}^\perp]A = A(I - E)$. Here, we explore more carefully the properties of $[\mathcal{S}]A$ which are relevant for the compatibility of A and \mathcal{S} . Another result which may be relevant for updating theory is the extension of the well-known theorem of J. A. Fill and D. E. Fishkind [20] which says that, for $n \times n$ complex matrices A, B such that $rk(A + B) = rk(A) + rk(B)$ it holds that

$$(A + B)^\dagger = (I - S)A^\dagger(I - T) + SB^\dagger T,$$

where \dagger denotes the Moore-Penrose inverse, $S = (P_{N(B)^\perp}P_{N(A)})^\dagger$ and $T = (P_{N(A^*)}P_{N(B^*)^\perp})^\dagger$. Here, $rk(X)$ denotes the rank of the matrix X . This is a generalization of the famous Sherman-Morrison-Woodbury formula:

$$(A + YGZ^*)^{-1} = A^{-1} - A^{-1}Y(G^{-1} + Z^*A^{-1}Y)^{-1}Z^*A^{-1}, \quad (1)$$

provided that A and $A + YGZ^*$ are invertible; here A, Y, G, Z are matrices of suitable sizes. Around 1950. J. Sherman and W. J. Morrison [32], M. A. Woodbury [33] and M. S. Barlett [10], published different versions of formula (1); however it was first discovered by W. J. Duncan [18] and L. Guttman [23] but their finding did not attract much attention because it was not explicitly connected with applications. Indeed, the mentioned authors were interested in finding a formula for $(A + uv^*)^{-1}$ provided that A^{-1} is known. For an excellent survey on formula (1) and its applications, see [24]. Of course, for Hilbert space operators the rank hypothesis must be replaced by a different one. Since it is well-known that $rk(A + B) = rk(A) + rk(B)$ if and only if $R(A) \cap R(B) = \{0\}$ and $R(A^*) \cap R(B^*) = \{0\}$, we prove that Fill-Fishkind formula holds for $A, B \in L(\mathcal{H})$ such that $R(A)$ and $R(B)$ are closed, $R(A) \cap R(B) = R(A^*) \cap R(B^*) = \{0\}$ and $(A, B), (A^*, B^*) \in \mathcal{R}$.

We end this section introducing some notation. The direct sum of two closed subspaces \mathcal{S} and \mathcal{T} will be denoted by $\mathcal{S} \dot{+} \mathcal{T}$. If $\mathcal{H} = \mathcal{S} \dot{+} \mathcal{T}$ then $Q_{\mathcal{S} // \mathcal{T}}$ denotes the oblique projection with range \mathcal{S} and nullspace \mathcal{T} .

Acknowledgements: We wish to thank the referee whose remarks helped us to improve the readability of the paper.

2. Range additivity

Let \mathcal{H}, \mathcal{K} be Hilbert spaces. We say that $A, B \in L(\mathcal{H}, \mathcal{K})$ have the *range additivity property* if $R(A + B) = R(A) + R(B)$. We denote by \mathcal{R} the set of all these pairs (A, B) , i.e.,

$$\mathcal{R} := \{(A, B) : A, B \in L(\mathcal{H}, \mathcal{K}) \text{ and } R(A + B) = R(A) + R(B)\}.$$

We collect first some trivial or well-known facts about \mathcal{R} .

Proposition 2.1. *Let $A, B \in L(\mathcal{H}, \mathcal{K})$. Then*

1. $(A, B) \in \mathcal{R}$ if and only if $(B, A) \in \mathcal{R}$.
2. If $R(A) = \mathcal{K}$ and $A = C + D$ for some $C, D \in L(\mathcal{H}, \mathcal{K})$ then $(C, D) \in \mathcal{R}$.
3. If $\mathcal{H} = \mathcal{K}$ is finite dimensional and $A, B \in L(\mathcal{H})^+$ then $(A, B) \in \mathcal{R}$.
4. If $\mathcal{H} = \mathcal{K}$, $A, B \in L(\mathcal{H})^+$ and $R(A + B)$ or $R(A) + R(B)$ is closed, then $(A, B) \in \mathcal{R}$; in particular, if $A, B \in L(\mathcal{H})^+$, $R(A)$ is closed and $\dim R(B) < \infty$ then $(A, B) \in \mathcal{R}$.

Proof. Items 1 and 2 are trivial. Item 4 has been proven by P. A. Fillmore and J. P. Williams [21, Corollary 3] under the additional hypothesis that $R(A)$ and $R(B)$ are closed. In [6, Theorem 3.3] there is a proof without this hypothesis. Item 3 follows from item 4. \square

Proposition 2.2. *For $A, B \in L(\mathcal{H}, \mathcal{K})$ consider the following conditions:*

1. $\overline{R(A^*)} + \overline{R(B^*)}$ is closed.
2. $N(A) + N(B) = \mathcal{H}$
3. $(A, B) \in \mathcal{R}$.

The next implications hold: $1 \Leftrightarrow 2 \Rightarrow 3$. The converse $3 \Rightarrow 2$ holds if $R(A) \cap R(B) = \{0\}$.

Proof. See [7, Prop. 5.8]. For more a general result see Theorem 2.10. \square

Examples 2.3. 1. Consider $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix}$. Clearly, $(A, B) \in \mathcal{R}$ but $(A^*, B^*) \notin \mathcal{R}$.

2. There exist $C, D \in L(\mathcal{H})^+$ such that $R(C), R(D)$ are dense and $(C, D) \notin \mathcal{R}$. For this, consider $C, D \in L(\mathcal{H})^+$ with dense ranges such that $R(C) \cap R(D) = \{0\}$ (see [21]). Hence, as $N(C) + N(D) = \{0\} \neq \mathcal{H}$ then, by Proposition 2.2, $(C, D) \notin \mathcal{R}$.

Below we collect some useful characterizations of \mathcal{R} . Notice that the proof holds also for vector spaces and modules over a ring.

Proposition 2.4. *Given $A, B \in L(\mathcal{H})$, the following conditions are equivalent:*

1. $(A, B) \in \mathcal{R}$,
2. $R(A) \subseteq R(A + B)$,
3. $R(A - B) \subseteq R(A + B)$.

Proof. $1 \Rightarrow 2$. If $R(A + B) = R(A) + R(B)$, then, a fortiori, $R(A) \subseteq R(A + B)$.

$2 \Rightarrow 3$. For every $x \in \mathcal{H}$, $(A - B)x = -(A + B)x + 2Ax \in R(A + B)$.

$3 \Rightarrow 1$. For every $x \in \mathcal{H}$, $2Ax = (A - B)x + (A + B)x \in R(A + B)$ and $2Bx = -(A - B)x + (A + B)x \in R(A + B)$, and we get $R(A) + R(B) \subseteq R(A + B)$, i.e. $(A, B) \in \mathcal{R}$ \square

The next result of R.G. Douglas [17] will be frequently used in the paper.

Theorem 2.5. *Let $A \in \mathcal{L}(\mathcal{H}, \mathcal{K})$ and $B \in L(\mathcal{F}, \mathcal{K})$. The following conditions are equivalent:*

1. $R(B) \subseteq R(A)$.
2. *There is a positive number λ such that $BB^* \leq \lambda AA^*$.*
3. *There exists $C \in L(\mathcal{F}, \mathcal{H})$ such that $AC = B$.*

If one of these conditions holds then there is a unique operator $D \in L(\mathcal{F}, \mathcal{H})$ such that $AD = B$ and $R(D) \subseteq N(A)^\perp$.

This unique operator D is called the **reduced solution** of $AX = B$.

Corollary 2.6. *For $A, B \in L(\mathcal{H})$ the following conditions are equivalent:*

1. *the equation $AX = B$ has a solution in $L(\mathcal{H})$.*
2. $(A - B, B) \in \mathcal{R}$.

Corollary 2.7. *For $A, B \in L(\mathcal{H})^+$ it holds:*

1. $(A + B)^{1/2}X = A^{1/2}$ *has a solution.*
2. $(A^{1/2}, (A + B)^{1/2} - A^{1/2}) \in \mathcal{R}$.

Proof. In fact, it holds $A + B \geq A$ and Douglas' theorem applies. □

The next corollary complements Proposition 2.2. For a proof see [5, Prop. 4.13].

Corollary 2.8. *For $A, B \in L(\mathcal{H}, \mathcal{K})$ the following conditions are equivalent:*

1. $\overline{R(A^*)} + \overline{R(B^*)}$ *is closed;*
2. *equation $(A + B)X = A$ admits a solution which is an oblique (i.e., not necessarily orthogonal) projection in $L(\mathcal{H})$.*

Recall that $A, B \in L(\mathcal{H})^+$ are said to be *Thompson equivalent* (in symbols, $A \sim_T B$) if there exist positive numbers r, s such that $rA \leq B \leq sA$ (where $C \leq D$ means that $\langle Cx, x \rangle \leq \langle Dx, x \rangle$ for all $x \in \mathcal{H}$). By Douglas' theorem, $A \sim_T B$ if and only if $R(A^{1/2}) = R(B^{1/2})$. For a fixed $A \in L(\mathcal{H})^+$ the *Thompson component* of A is the convex cone $\{B \in L(\mathcal{H})^+ : A \sim_T B\}$. The following identity is due to Crimmins (see [21] for a proof): if $A, B \in L(\mathcal{H}, \mathcal{K})$ then $R(A) + R(B) = R((AA^* + BB^*)^{1/2})$. Using Crimmins' identity the following result is clear:

Proposition 2.9. *If $A, B \in L(\mathcal{H})^+$ then $(A, B) \in \mathcal{R}$ if and only if $(A + B)^2 \sim_T A^2 + B^2$.*

The next characterization of \mathcal{R} is less elementary than that of Proposition 2.2. Notice, however, that its proof is algebraic, so it also holds in the context of vector spaces, modules over a ring, and so on.

Theorem 2.10. *Let $A, B \in L(\mathcal{H})$. Then $R(A + B) = R(A) + R(B)$ if and only if $R(A) \cap R(B) \subseteq R(A + B)$ and $\mathcal{H} = A^{-1}(R(B)) + B^{-1}(R(A))$. In particular, if $R(A) \cap R(B) = \{0\}$ then $(A, B) \in \mathcal{R}$ if and only if $N(A) + N(B) = \mathcal{H}$.*

Proof. Let $T = A + B$, $\mathcal{W} = R(A) \cap R(B)$ and suppose that $R(T) = R(A) + R(B)$. Then $R(A) \subseteq R(T)$ and $R(B) \subseteq R(T)$ so that $\mathcal{W} \subseteq R(T)$. On the other hand, using again that $R(A)$ and $R(B)$ are subsets of $R(T)$ it holds $\mathcal{H} = T^{-1}(R(T)) = T^{-1}(R(A) + R(B)) = T^{-1}(R(A)) + T^{-1}(R(B))$. But it is easy to see that $T^{-1}(R(A)) = B^{-1}(R(A))$. Hence, $\mathcal{H} = T^{-1}(R(A)) + T^{-1}(R(B)) = A^{-1}(R(B)) + B^{-1}(R(A))$.

Conversely, suppose that $\mathcal{W} \subseteq R(T)$ and $\mathcal{H} = B^{-1}(R(A)) + A^{-1}(R(B))$. We shall prove that $R(B) = T(A^{-1}(R(B)))$. In fact, since $B^{-1}(R(A)) = B^{-1}(\mathcal{W})$ and $A^{-1}(R(B)) = A^{-1}(\mathcal{W})$ then

$$R(B) = B(\mathcal{H}) = B(B^{-1}(\mathcal{W}) + A^{-1}(\mathcal{W})) = \mathcal{W} + B(A^{-1}(\mathcal{W})),$$

because $\mathcal{W} \subseteq R(B)$. Moreover, $R(B) = \mathcal{W} + B(A^{-1}(\mathcal{W})) = \mathcal{W} + T(A^{-1}(\mathcal{W})) = T(A^{-1}(\mathcal{W}))$. In fact, for the second equality consider $y \in \mathcal{W} + B(A^{-1}(\mathcal{W}))$ then $y = w + Bx$ where $w \in \mathcal{W}$ and $x \in A^{-1}(\mathcal{W})$, so that $y = w - Ax + Tx$ where $w - Ax \in \mathcal{W}$ and $Tx \in T(A^{-1}(\mathcal{W}))$; the other inclusion is clear. Then the second equality holds.

To see that $\mathcal{W} + T(A^{-1}(\mathcal{W})) = T(A^{-1}(\mathcal{W}))$ it is sufficient to note that $\mathcal{W} \subseteq T(A^{-1}(\mathcal{W}))$. In fact, $T^{-1}(\mathcal{W}) = A^{-1}(\mathcal{W}) \cap B^{-1}(\mathcal{W}) \subseteq A^{-1}(\mathcal{W})$ then applying T to both sides of the inclusion $\mathcal{W} = TT^{-1}(\mathcal{W}) \subseteq T(A^{-1}(\mathcal{W}))$ because $\mathcal{W} \subseteq R(T)$.

Hence, $R(B) = T(A^{-1}(\mathcal{W})) = T(A^{-1}(R(B))) \subseteq R(T)$. Applying Proposition 2.4, $(A, B) \in \mathcal{R}$. □

One of the obstructions for range additivity for operators in Hilbert spaces is that $R(A)$ is, in general, non closed. Therefore, the identity $R(A + B) = R(A) + R(B)$ is not equivalent to $N(A^* + B^*) = N(A^*) \cap N(B^*)$, which is easier to check. On these matters, see the papers by P. Šemrl [31, §2] and G. Lešnjak and P. Šemrl [26], where they discuss different kinds of topological range additivity properties. See also the paper by J. Baksalary, P. Šemrl and G. P. H. Styan [9].

3. Shorted operators and range additivity

In his paper on selfadjoint extensions of certain unbounded operators [25], M. G. Krein defined what is nowadays known as a shorted operator. More precisely, if $A \in L(\mathcal{H})^+$ and \mathcal{S} is a closed subspace of \mathcal{H} , Krein proved that the set

$$\{C \in L(\mathcal{H})^+ : C \leq A \text{ and } R(C) \subseteq \mathcal{S}\}$$

admits a maximal element $[\mathcal{S}]A$. Moreover, Krein proved that

$$[\mathcal{S}]A = A^{1/2}P_{\mathcal{M}}A^{1/2},$$

if $\mathcal{M} = (A^{1/2})^{-1}(\mathcal{S})$. Krein constructed the shorted operators to find selfadjoint positive extensions of certain unbounded operators. For a modern exposition of Krein's ideas on these matters, see [8].

Later, W. N. Anderson and G. E. Trapp [1] rediscovered the operator $[\mathcal{S}]A$, proved many useful properties and showed its relevance in the theory of impedance matrices of networks. The papers by E. L. Pekarev [27], E. L. Pekarev and J. L. Smul'jan [28], T. Ando [3] and S. L. Eriksson and H. Leutwiler [19] contain many useful theorems about shorted operators.

A nice exposition of results about shorted operators in finite dimensional spaces is that of T. Ando [3]. It is worth mentioning the parallel sum, a binary operation defined on positive operators, which is also relevant in electrical network theory and which is related to shorted operators. If A, B are the impedance matrices of two n -port resistive networks then $A : B := A(A + B)^\dagger B$ is the impedance matrix of their parallel connection. For positive operators A, B on a Hilbert space \mathcal{H} , Fillmore and Williams [21] defined

$$A : B = A^{1/2} C^* D B^{1/2},$$

if C (resp. D) is the reduced solution of $(A + B)^{1/2} X = A^{1/2}$ (resp. $(A + B)^{1/2} X = B^{1/2}$).

Anderson and Trapp [1] proved that $A : B$ is the $(1, 1)$ entry of $[\mathcal{S}] \begin{pmatrix} A & A \\ A & A + B \end{pmatrix}$, where $\mathcal{S} = \mathcal{H} \oplus \{0\}$ and the matrix $\begin{pmatrix} A & A \\ A & A + B \end{pmatrix}$ is considered as an element of $L(\mathcal{H} \oplus \mathcal{H})^+$.

Thus, the parallel addition is a particular form of the shorted operation. Any extension to non necessarily positive operators of the parallel sum operation requires that (A, B) and (A^*, B^*) belong to \mathcal{R} , at least if one wants to keep the desirable commutativity $A : B = B : A$ [30, 10.1.6]. Indeed, Rao and Mitra say that A, B are *parallel summable* if $A(A + B)^- B$ is invariant for any generalized inverse of $A + B$. It turns out that this happens if and only if $(A, B) \in \mathcal{R}$ and $(A^*, B^*) \in \mathcal{R}$. This means that there is a strong relationship among Krein shorted operators, Douglas range inclusion and range additivity.

A warning about notation. The original notation by Krein is $A_{\mathcal{S}}$. Anderson and Trapp [1] used $\mathcal{S}(A)$. Ando [4] proposed $[\mathcal{S}]A$. This is coherent with a relevant construction $[B]A$ for $A, B \in L(\mathcal{H})^+$ that he defined and studied in [2], by generalizing a theorem of Anderson and Trapp that $([\mathcal{S}]A)x = \lim_{n \rightarrow \infty} (A : nP_{\mathcal{S}})x$ for every $x \in \mathcal{H}$. Ando defined the existence of $([B]A)x = \lim_{n \rightarrow \infty} (A : nB)x$ for every $x \in \mathcal{H}$ and proved many relevant results on this construction. In particular, it holds that $[\mathcal{S}]A = [B]A$ if $\mathcal{S} = R(B)$. Erikson and Leutwiler [19] used $Q_B A$ for Ando's $[B]A$. In [3], Ando has used $A_{/\mathcal{S}}$ for the shorted operator and $A_{\mathcal{S}} = A - A_{/\mathcal{S}}$. Corach, Maestripieri and Stojanoff used $\sum(P_{\mathcal{S}}, A)$ in [11] and $A_{/\mathcal{S}}$ in [12] to denote what we are denoting now $[\mathcal{S}^\perp]A$.

In the proposition below we collected some facts on the Krein shorted operators, mainly extracted from the paper [1] of Anderson and Trapp.

Proposition 3.1. *Given $A, B \in L(\mathcal{H})^+$ and closed subspaces \mathcal{S}, \mathcal{T} of \mathcal{H} the following properties hold:*

1. $R(A) \cap \mathcal{S} \subseteq R([\mathcal{S}]A) \subseteq R(([\mathcal{S}]A)^{1/2}) = R(A^{1/2}) \cap \mathcal{S}$; in particular, $R([\mathcal{S}]A)$ is closed if $R(A)$ is closed or, more generally, if $R(A) \cap \mathcal{S} = R(A^{1/2}) \cap \mathcal{S}$.
2. $N([\mathcal{S}]A) = N(P_{(A^{1/2})^{-1}(\mathcal{S})} A^{1/2}) = (A^{1/2})^{-1}(\overline{A^{1/2}(\mathcal{S}^\perp)}) (\supseteq N(A) + \mathcal{S}^\perp)$; equality holds if and only if $\overline{A^{1/2}(\mathcal{S}^\perp)} \cap R(A^{1/2}) = A^{1/2}(\mathcal{S}^\perp)$.
3. $[\mathcal{S}](A + B) \geq [\mathcal{S}]A + [\mathcal{S}]B$; equality holds if and only if $R((A - [\mathcal{S}]A + B - [\mathcal{S}]B)^{1/2}) \cap \mathcal{S} = \{0\}$.
4. $R((A - [\mathcal{S}]A)^{1/2}) \cap \mathcal{S} = \{0\}$. In particular, $\overline{R([\mathcal{S}]A)} \cap R(A - [\mathcal{S}]A) = \{0\}$.

Proof. 1. See [1, Corollary 4 of Theorem 1 and Corollary of Theorem 3]

2. See [12, Corollary 2.3]

3. See [1, Theorem 4].
4. See [1, Theorem 2].

□

Corollary 3.2. *Let $A, B \in L(\mathcal{H})^+$. Then:*

1. If $\mathcal{S} = \overline{R(B)}$ then $[\mathcal{S}]B = B$ and $[\mathcal{S}](A + B) = [\mathcal{S}]A + B$.
2. If $\mathcal{S} = R(B)$ is closed then $R([\mathcal{S}](A + B)) = \mathcal{S}$ and $N([\mathcal{S}](A + B)) = \mathcal{S}^\perp$.

Proof. 1. The identity $[\mathcal{S}]B = B$ can be checked through the definition of $[\mathcal{S}]B$; the identity $[\mathcal{S}](A + B) = [\mathcal{S}]A + B$ follows from items 3 and 4 in Proposition 3.1.

2. For every $C \in L(\mathcal{H})^+$ it holds $R([\mathcal{S}]C) \subseteq \mathcal{S}$, therefore $\mathcal{S} \supseteq R([\mathcal{S}](A + B)) = R([\mathcal{S}]A + [\mathcal{S}]B) = R([\mathcal{S}]A) + \mathcal{S} \supseteq \mathcal{S}$, where the second equality holds by Crimmins' identity. The kernel condition follows by taking orthogonal complement.

□

Proposition 3.3. *Let $A \in L(\mathcal{H})^+$ and let \mathcal{S} be a closed subspace of \mathcal{H} . The following conditions are equivalent:*

1. $([\mathcal{S}]A, A - [\mathcal{S}]A) \in \mathcal{R}$;
2. $R(A) = R(A - [\mathcal{S}]A) + R([\mathcal{S}]A)$;
3. $R([\mathcal{S}]A) \subseteq R(A)$;
4. $R(A^{1/2}) = \mathcal{M} \cap R(A^{1/2}) \oplus \mathcal{M}^\perp \cap R(A^{1/2})$, if $\mathcal{M} = (A^{1/2})^{-1}(\mathcal{S})$.

Proof. Notice that $N([\mathcal{S}]A) = (A^{1/2})^{-1}(\overline{A^{1/2}(\mathcal{S}^\perp)})$ and $N(A - [\mathcal{S}]A) = A^{-1}(\mathcal{S})$.

1 \Leftrightarrow 2 \Leftrightarrow 3. It follows from Proposition 2.4 and Proposition 3.1.

3 \Leftrightarrow 4. Assume that $R([\mathcal{S}]A) \subseteq R(A)$ and let $y = A^{1/2}x \in R(A^{1/2})$. Hence, $A^{1/2}x = P_{\mathcal{M}}A^{1/2}x + (I - P_{\mathcal{M}})A^{1/2}x$. Applying $A^{1/2}$ at both sides, we get that $Ax = A^{1/2}(I - P_{\mathcal{M}})A^{1/2}x + [\mathcal{S}]Ax$. Thus, since $R([\mathcal{S}]A) \subseteq R(A)$ we obtain that $A^{1/2}(I - P_{\mathcal{M}})A^{1/2}x \in R(A)$. Therefore, $A^{1/2}(I - P_{\mathcal{M}})A^{1/2}x = Az$ for some $z \in \mathcal{H}$. From this, $(I - P_{\mathcal{M}})A^{1/2}x - A^{1/2}z \in N(A) \cap \overline{R(A)} = \{0\}$, i.e., $(I - P_{\mathcal{M}})A^{1/2}x = A^{1/2}z \in R(A^{1/2}) \cap \mathcal{M}^\perp$. Therefore, $A^{1/2}x = P_{\mathcal{M}}A^{1/2}x + (I - P_{\mathcal{M}})A^{1/2}x \in \mathcal{M} \cap R(A^{1/2}) \oplus \mathcal{M}^\perp \cap R(A^{1/2})$ and implication 3 \Rightarrow 4 is proved.

Conversely, assume that $R(A^{1/2}) = \mathcal{M} \cap R(A^{1/2}) \oplus \mathcal{M}^\perp \cap R(A^{1/2})$. Hence, $R([\mathcal{S}]A) = R(A^{1/2}P_{\mathcal{M}}A^{1/2}) \subseteq A^{1/2}(\mathcal{M} \cap R(A^{1/2})) \subseteq R(A)$.

□

4. Compatibility and range additivity

Definition 4.1. *Given $A \in L(\mathcal{H})^+$ and \mathcal{S} a closed subspace of \mathcal{H} , we say that A and \mathcal{S} are compatible if $\mathcal{H} = \mathcal{S} + (AS)^\perp$.*

As shown in [11] the compatibility of A and \mathcal{S} means that there exists a (bounded linear) projection with image \mathcal{S} which is Hermitian with respect to the semi-inner product $\langle \cdot, \cdot \rangle_A$ defined by $\langle \xi, \eta \rangle_A = \langle A\xi, \eta \rangle$. It is worth mentioning that compatibility gives a kind of weak version of invariant subspaces. In fact, if A is a selfadjoint operator on \mathcal{H} and \mathcal{S} is a closed subspace, then \mathcal{S} is an invariant subspace for A if $AS \subseteq \mathcal{S}$, which means that $P_{\mathcal{S}}AP_{\mathcal{S}} = P_{\mathcal{S}}A$. On the other side, A and \mathcal{S} are compatible if and only if $R(P_{\mathcal{S}}AP_{\mathcal{S}}) = R(P_{\mathcal{S}}A)$; for a proof

of this fact see [11, Proposition 3.3]. In the recent paper [7, Proposition 2.9] it is proven that A and \mathcal{S} are compatible if and only if $(P_{\mathcal{S}}A, I - P_{\mathcal{S}}) \in \mathcal{R}$. In this section we shall complete this result by proving that A and \mathcal{S} are compatible if and only if $(A, I - P_{\mathcal{S}}) \in \mathcal{R}$, i.e., $R(A + P_{\mathcal{S}^\perp}) = R(A) + R(P_{\mathcal{S}^\perp}) = R(A) + \mathcal{S}^\perp$.

Proposition 4.2. [12, Theorem 3.8] *Let $A \in L(\mathcal{H})^+$ and \mathcal{S} a closed subspace of \mathcal{H} . The following conditions are equivalent:*

1. A and \mathcal{S} are compatible.
2. $R([\mathcal{S}^\perp]A) \subseteq R(A)$ and $N([\mathcal{S}^\perp]A) = N(A) + \mathcal{S}$.

Proposition 4.3. *Let $A, B \in L(\mathcal{H})^+$ with closed ranges. The next conditions are equivalent:*

1. A and $N(B)$ are compatible.
2. $N(A) + N(B)$ is closed.
3. B and $N(A)$ are compatible.
4. $R(A) + R(B)$ is closed.
5. $(A, B) \in \mathcal{R}$.

Proof. $1 \Leftrightarrow 2$. [11, Theorem 6.2].

$2 \Leftrightarrow 3$. Idem.

$2 \Leftrightarrow 4$. It follows from the general fact that, for closed subspaces \mathcal{S}, \mathcal{T} then $\mathcal{S} + \mathcal{T}$ is closed if and only if $\mathcal{S}^\perp + \mathcal{T}^\perp$ is closed. See [16, Theorem 13].

$4 \Rightarrow 5$. See [21, Corollary 3].

$5 \Rightarrow 4$. $R(A + B) = R(A) + R(B) = R(A^{1/2}) + R(B^{1/2}) = R((A + B)^{1/2})$ by Crimmins' identity. Then $R(A + B)$ is closed and so $R(A) + R(B)$ is closed. \square

Theorem 4.4. *Let $A, B \in L(\mathcal{H})^+$ and suppose that B has a closed range. The following conditions are equivalent:*

1. A and $N(B)$ are compatible.
2. $(A, B) \in \mathcal{R}$.
3. $R(B) \dot{+} \overline{AN(B)}$ is closed.

Proof. $1 \Leftrightarrow 2$. Let $\mathcal{S} = N(B)$. First observe that A and \mathcal{S} are compatible if and only if $A + B$ and \mathcal{S} are compatible. Indeed, $\mathcal{S} + ((A + B)\mathcal{S})^\perp = \mathcal{S} + (A\mathcal{S})^\perp$. Hence, by Proposition 4.2, A and \mathcal{S} are compatible if and only if $R([\mathcal{S}^\perp](A + B)) \subseteq R(A + B)$ and $N([\mathcal{S}^\perp](A + B)) = \mathcal{S} + N(A + B)$ or, equivalently, by Corollary 3.2, $\mathcal{S}^\perp \subseteq R(A + B)$ (notice that $N(A + B) = N(A) \cap N(B) \subseteq \mathcal{S}$). Summarizing, A and \mathcal{S} are compatible if and only if $R(B) = \mathcal{S}^\perp \subseteq R(A + B)$, i.e., $R(A + B) = R(A) + R(B)$.

$1 \Leftrightarrow 3$. It follows applying [16, Theorem 13]. \square

Corollary 4.5. *Let $A \in L(\mathcal{H})^+$ and \mathcal{S} a closed subspace of \mathcal{H} . The next conditions are equivalent:*

1. A and \mathcal{S} are compatible;
2. $(A, I - P_{\mathcal{S}}) \in \mathcal{R}$.
3. $\mathcal{S}^\perp \dot{+} \overline{A\mathcal{S}}$ is closed.

Proposition 4.6. *Let $A, B \in L(\mathcal{H})^+$ such that $R(A) \cap \overline{R(B)} = \{0\}$. Then, $(A, B) \in \mathcal{R}$ if and only if A and $N(B)$ are compatible.*

Proof. Since, $R(A) \cap R(B) = \{0\}$ then $R(A+B) = R(A) + R(B)$ if and only if $\mathcal{H} = N(A) + N(B)$. Now, $N(A) + N(B) = A^{-1}(\{0\}) + N(B) = A^{-1}(\overline{R(B)}) + N(B) = A^{-1}(N(B)^\perp) + N(B)$. Therefore, $R(A+B) = R(A) + R(B)$ if and only if $\mathcal{H} = A^{-1}(N(B)^\perp) + N(B)$, i.e., if and only if A and $N(B)$ are compatible. \square

The next example shows that the compatibility of A and $N(B)$ does not imply, in general, that $(A, B) \in \mathcal{R}$.

Example 4.7. Considering C and D as in Example 2.3.2, we define $A = \begin{pmatrix} 0 & 0 \\ 0 & C \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 0 \\ 0 & D \end{pmatrix}$. Clearly, $(A, B) \notin \mathcal{R}$. However, A and $N(B)$ are compatible.

Corollary 4.8. *Let $A \in L(\mathcal{H})^+$ and \mathcal{S} a closed subspace of \mathcal{H} . The following conditions are equivalent:*

1. $R(A) = R(A - [\mathcal{S}]A) + R([\mathcal{S}]A)$;
2. $A - [\mathcal{S}]A$ and $N([\mathcal{S}]A)$ are compatible;
3. A and $N([\mathcal{S}]A)$ are compatible.

Proof. 1 \Leftrightarrow 2. It follows from Proposition 3.1 and Proposition 4.6.

2 \Leftrightarrow 3. It follows from the fact that $A = [\mathcal{S}]A + A - [\mathcal{S}]A$. \square

5. The Fill-Fishkind formula

This last section is devoted to the Fill-Fishkind formula. In order to identify certain Moore-Penrose inverses of products of orthogonal projections, the next theorem (due to Penrose and Greville) will be helpful.

Theorem 5.1. *If $Q \in L(\mathcal{H})$ is an oblique projection then $Q^\dagger = P_{N(Q)^\perp} P_{R(Q)}$. Conversely, if M and N are closed subspaces of \mathcal{H} such that $P_M P_N$ has closed range, then $(P_M P_N)^\dagger$ is the unique oblique projection with range $R(P_N P_M)$ and nullspace $N(P_N P_M)$.*

Proof. For matrices, the proof appears in the paper by Penrose [29, Lemma 2.3] and Greville [22, Theorem 1]. For general Hilbert spaces, see [13, Theorem 4.1]. \square

We prove now the extension of the theorem of Fill and Fishkind [20, Theorem 3] mentioned in the introduction.

Theorem 5.2. *Let $A, B \in L(\mathcal{H}, \mathcal{K})$ such that $R(A), R(B)$ are closed, $R(A) \cap R(B) = R(A^*) \cap R(B^*) = \{0\}$ and $(A, B) \in \mathcal{R}$ and $(A^*, B^*) \in \mathcal{R}$. Hence,*

$$(A + B)^\dagger = (I - S)A^\dagger(I - T) + SB^\dagger T, \quad (2)$$

where

$$S := (P_{N(B)^\perp} P_{N(A)})^\dagger$$

and

$$T := (P_{N(A^*)} P_{N(B^*)^\perp})^\dagger.$$

Proof. We start with a series of claims which will be used in the proof.

Claim 1: All Moore-Penrose inverses in (2) are bounded, i.e., $R(A)$, $R(B)$, $R(A+B)$, $R((P_{N(B)^\perp}P_{N(A)})$ and $R(P_{N(A^*)}P_{N(B^*)^\perp})$ are closed. The first two by hypotheses, the third one by Proposition 2.2, and $R((P_{N(B)^\perp}P_{N(A)})$ and $R(P_{N(A^*)}P_{N(B^*)^\perp})$ are closed by Theorem 22 of F. Deutsch's paper [16] and since $N(A) + N(B) = \mathcal{H}$, by Proposition 2.2 and $R(A) + R(B) = R(A + B)$ is closed.

Claim 2: $S = Q_{P_{N(A)}(N(B)^\perp)//N(B)}$ and $T = Q_{R(B)//R(A)+R(A)^\perp \cap R(B)^\perp}$.

By Theorem 5.1, $S = Q_{P_{N(A)}(N(B)^\perp)//N(P_{N(A)}P_{N(B)^\perp})}$ and $T = Q_{P_{N(B^*)^\perp}P_{N(A^*)//N(P_{N(B^*)^\perp}P_{N(A^*)})}$.

We claim that $N(S) = N(B)$, $R(T) = R(B)$ and $N(T) = R(A) + R(A)^\perp \cap R(B)^\perp$. In fact, it holds $N(S) = N(P_{N(A)}P_{N(B)^\perp}) = N(B)$, because $N(A)^\perp \cap N(B)^\perp = R(A^*) \cap R(B^*) = \{0\}$; for T , notice that $R(P_{N(B^*)^\perp}P_{N(A^*)}) = N(B^*)^\perp = R(B)$ because $N(A^*) + N(B^*) = \mathcal{H}$ (it is dense because $R(A) \cap R(B) = \{0\}$ and it is closed because $R(A) + R(B)$ is closed); finally, $N(P_{N(B^*)^\perp}P_{N(A^*)}) = N(A^*)^\perp + N(B^*) \cap N(A^*) = R(A) + R(A)^\perp \cap R(B)^\perp$.

Claim 3: $\mathcal{H} = [R(A) \dot{+} R(B)] \oplus R(A)^\perp \cap R(B)^\perp$. It follows from the form of T in Claim 2.

Claim 4: $AS = 0$ and $TA = 0$ because $R(S) \subseteq N(A)$ and $R(A) \subseteq N(T)$, respectively. Moreover, $TB = B$ since $R(B) \subseteq R(T)$.

Claim 5: $BS = B$, or, which is the same, $S^*B^* = B^*$ because, by Claim 2, S^* is an oblique projection with range $R(B^*) = N(B)^\perp$. Moreover, $SB^\dagger B = S$.

Claim 6: $Q_1 := P_{R(A)}(I - T)$ is the oblique projection onto $R(A)$ with nullspace $R(B) + R(A)^\perp \cap R(B)^\perp$. In order to prove the claim first observe that $P_{R(A)}(I - T)$ is a projection because, by Claim 2, it holds $R(A) \subseteq N(T)$; on the other hand, $R(P_{R(A)}(I - T)) \subseteq R(A)$ and $N(P_{R(A)}(I - T)) \subseteq R(B) + R(A)^\perp \cap R(B)^\perp$. By Claim 3 the projection Q_1 with range $R(A)$ and nullspace $R(B) + R(A)^\perp \cap R(B)^\perp$ is well defined. Therefore, $Q_1 = P_{R(A)}(I - T)$, because if E_1, E_2 are oblique projections such that $R(E_1) \subseteq R(E_2)$ and $N(E_1) \subseteq N(E_2)$, it follows that $E_1 = E_2$.

We use now these technical results to prove formula (2). Recall that, from Penrose's characterization of the Moore-Penrose inverse of C it holds that $X = C^\dagger$ if and only if $CX = P_{R(C)}$, $XC = P_{R(C^*)}$ and $XCX = X$. Hence, we must prove:

- (2i) $(A + B)((I - S)A^\dagger(I - T) + SB^\dagger T) = P_{R(A+B)}$
- (2ii) $((I - S)A^\dagger(I - T) + SB^\dagger T)(A + B) = P_{R(A^*+B^*)}$.
- (2iii) $((I - S)A^\dagger(I - T) + SB^\dagger T)(A + B)((I - S)A^\dagger(I - T) + SB^\dagger T) = (I - S)A^\dagger(I - T) + SB^\dagger T$.

By Claims 4, 5 and 6, it holds:

$$\begin{aligned} (A + B)((I - S)A^\dagger(I - T) + SB^\dagger T) &= AA^\dagger(I - T) + BB^\dagger T \\ &= P_{R(A)}(I - T) + P_{R(B)}T = Q_1 + T, \end{aligned}$$

We must prove now that $Q_1 + T = P_{R(A+B)}$:

- (a) Since Q_1, T are projections and $Q_1 T = T Q_1 = 0$ it follows that $Q_1 + T$ is a projection.
- (b) Clearly, $R(Q_1 + T) \subseteq R(A + B)$. On the other side, as $(Q_1 + T)(A + B) = A + B$ we get the reverse inclusion, and so $R(Q_1 + T) = R(A + B)$.
- (c) $R(A + B)^\perp = R(A)^\perp \cap R(B)^\perp \subseteq N(Q_1 + T)$.

Hence, from (a), (b) and (c) we obtain that $Q_1 + T = P_{R(A+B)}$ which proves (2i).

For the proof of (2ii), by Claims 4 and 5, we get

$$\begin{aligned} ((I - S)A^\dagger(I - T) + SB^\dagger T)(A + B) &= (I - S)A^\dagger A + SB^\dagger B \\ &= (I - S)(I - P_{N(A)}) + S \\ &= I - (I - S)P_{N(A)}. \end{aligned}$$

Therefore,

- (a) As $P_{N(A)}S = S$ then it is straightforward that $(I - (I - S)P_{N(A)})^2 = I - (I - S)P_{N(A)}$.
- (b) Clearly, $N(A) \cap N(B) \subseteq N(A + B) \subseteq N(I - (I - S)P_{N(A)})$. If $x \in N(I - (I - S)P_{N(A)})$, since $(I - S)P_{N(A)} = P_{N(A)}(I - S)P_{N(A)}$, we have that $x = (I - S)P_{N(A)}x \in N(B)$ and $x = P_{N(A)}(I - S)P_{N(A)}x \in N(A)$, i.e., $x \in N(A) \cap N(B)$. Therefore, $N(I - (I - S)P_{N(A)}) = N(A) \cap N(B) = (R(A^*) + R(B^*))^\perp = R(A^* + B^*)^\perp$.
- (c) By Claim 2, $(I - (I - S)P_{N(A)})(A^* + B^*) = A^* + B^*$, i.e., $R(A^* + B^*) \subseteq R(I - (I - S)P_{N(A)})$.

Therefore, from (a), (b) and (c) we conclude that $((I - S)A^\dagger(I - T) + SB^\dagger T)(A + B) = I - (I - S)P_{N(A)} = P_{R(A^* + B^*)}$ which is (2ii).

Finally, in order to prove (2iii), observe that, from (2ii), it holds that $((I - S)A^\dagger(I - T) + SB^\dagger T)(A + B) = I - (I - S)P_{N(A)}$. It suffices to prove that $(I - S)P_{N(A)}((I - S)A^\dagger(I - T) + SB^\dagger T) = 0$. But $(I - S)A^\dagger(I - T) + SB^\dagger T = A^\dagger(I - T) + S(B^\dagger T - A^\dagger(I - T))$; obviously $P_{N(A)}A^\dagger = 0$ and $(I - S)P_{N(A)}S = 0$ because $R(S) \subseteq N(A)$ by Claim 4. This ends the proof. \square

Remark 5.3. Fill and Fishkind proved their formula under the hypothesis $rk(A + B) = rk(A) + rk(B)$ where A, B are $n \times n$ complex matrices and rk denotes the rank. It is well known that this rank additivity property is equivalent to $R(A) \cap R(B) = R(A^*) \cap R(B^*) = \{0\}$. On the other side, by Proposition 2.2, the hypotheses of the theorem of Fill and Fishkind are equivalent (for matrices) to $(A, B), (A^*, B^*) \in \mathcal{R}$. Thus, the hypotheses of our infinite-dimensional version are not stronger than those of the finite-dimensional one. For a quite different set of hypotheses for Fill-Fishkind formula in Hilbert spaces, see the paper by Deng [14].

References

- [1] W. N. Anderson and G. E. Trapp, Shorted operators II, SIAM J. Appl. Math. 28 (1975), 60-71.
- [2] T. Ando, Lebesgue-type decomposition of positive operators. Acta Sci. Math. (Szeged) 38 (1976), 253-260.
- [3] T. Ando, Generalized Schur complements, Linear Algebra Appl. 27 (1979), 173-186.
- [4] T. Ando, Schur complements and matrix inequalities: operator theoretic approach, in "The Schur complement and its Applications" (ed. F. Zhang), 137-162, Springer, New York, 2005.

- [5] J. Antezana, G. Corach, and D. Stojanoff, Bilateral shorted operators and parallel sums, *Linear Algebra Appl.* 414 (2006), 570-588.
- [6] M. L. Arias, G. Corach, M.C. Gonzalez, Additivity properties of operator ranges, *Linear Algebra Appl.* 439 (2013) 3581-3590.
- [7] M. L. Arias, G. Corach, M.C. Gonzalez, Saddle point problems, Bott-Duffin inverses, abstract splines and oblique projections, *Linear Algebra Appl.* 457 (2014), 61-75.
- [8] Y. Arlinskii, E. M. Tsekanovskiĭ, M. Krein's research on semi-bounded operators, its contemporary developments, and applications. *Modern analysis and applications. The Mark Krein Centenary Conference. Vol. 1: Operator theory and related topics*, 65-112, *Oper. Theory Adv. Appl.*, 190, Birkhauser Verlag, Basel, 2009.
- [9] J. Baksalary, P. Šemrl, G. P. H. Styan, A note on rank additivity and range additivity, *Linear Algebra Appl.* 237/238 (1996), 489-498.
- [10] M. S. Bartlett, An inverse matrix adjustment arising in discriminant analysis, *Ann. Math. Statistics* 22 (1951), 107-111.
- [11] G. Corach, A. Maestripieri, D. Stojanoff, Oblique projections and Schur complements, *Acta Sci. Math. (Szeged)* 67 (2001), 337-356.
- [12] G. Corach, A. Maestripieri and D. Stojanoff, Generalized Schur complements and oblique projections, *Linear Algebra Appl.* 341 (2002), 259-272.
- [13] G. Corach, A. Maestripieri, Polar decompositions of oblique projections, *Linear Algebra Appl.* 433 (2010), 511-519.
- [14] C. Y. Deng, A generalization of the Sherman-Morrison-Woodbury formula, *Appl. Math. Lett.* 24 (2011), 1561-1564.
- [15] C. A. Desoer, B. H. Whalen, A note on pseudoinverses. *J. Soc. Indust. Appl. Math.* 11 (1963), 442-447.
- [16] F. Deutsch, *The angle between subspaces of a Hilbert space: "Approximation theory, wavelets and applications"*, (Maratea 1994), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci. 454 (1995), 107-130.
- [17] R.G. Douglas, On majorization, factorization and range inclusion of operators in Hilbert spaces, *Proc. Amer. Math. Soc.* 17 (1966), 413-416.
- [18] W. J. Duncan, Some devices for the solution of large sets of simultaneous linear equations. *Philos. Mag. (7)* 35 (1944). 660-670.
- [19] S. L. Eriksson, H. Leutwiler, A potential theoretic approach to parallel addition, *Math. Ann.* 274 (1986), 301-317.
- [20] J. A. Fill, D. E. Fishkind, The Moore-Penrose generalized inverse for sums of matrices, *SIAM J. Matrix Anal. Appl.* 21 (1999), 629-635.

- [21] P. A. Fillmore, J. P. Williams, On operator ranges, *Advances in Math.* 7 (1971), 254-281.
- [22] T. N. E. Greville, Solutions of the matrix equation $XAX = X$, and relations between oblique and orthogonal projectors, *SIAM J. Appl. Math.* 26 (1974) 828-832.
- [23] L. Guttman, Enlargement methods for computing the inverse matrix. *Ann. Math. Statistics* 17 (1946), 336-343.
- [24] W. W. Hager, Updating the inverse of a matrix. *SIAM Rev.* 31 (1989), 221-239.
- [25] M. G. Krein, The theory of self-adjoint extensions of semibounded Hermitian operators and its applications, *Mat. Sb. (N.S.)* 20(62) (1947), 431-495.
- [26] G. Lešnjak, P. Šemrl, Quasidirect addition of operators, *Linear and Multilinear Algebra* 40 (1996), 373-377.
- [27] E. L. Pekarev, Shorts of operators and some extremal problems. *Acta Sci. Math. (Szeged)* 56 (1992), 147-163.
- [28] E.L. Pekarev, J.L. Smul'jan, Parallel addition and parallel subtraction of operators, *Math. USSR Izvestija* 10 (1976), 351-370.
- [29] R. Penrose, A generalized inverse for matrices, *Proc. Cambridge Philos. Soc.* 51 (1955), 406-413.
- [30] C. R. Rao, S. K. Mitra, *Generalized inverse of matrices and its applications.* John Wiley, New York-London-Sydney, 1971.
- [31] P. Šemrl, Automorphism of $B(H)$ with respect to minus partial order, *J. Math. Anal. Appl.* 369 (2010), 205-213.
- [32] J. Sherman, W.J. Morrison, Adjustment of an inverse matrix corresponding to a change in one element of a given matrix, *Ann. Math. Statistics*, 21 (1950), 124-127.
- [33] M. A. Woodbury, Inverting modified matrices, *Statistical Research Group, Memo. Rep. no. 42*, Princeton University, Princeton, N. J., 1950.