

Accepted Manuscript

Ca_v2.1 voltage activated calcium channels and synaptic transmission in familial hemiplegic migraine pathogenesis

Osvaldo D. Uchitel, Carlota González Inchauspe, Francisco J. Urbano, Mariano N. Di Guilmi

PII: S0928-4257(11)00042-8

DOI: [10.1016/j.jphysparis.2011.10.004](https://doi.org/10.1016/j.jphysparis.2011.10.004)

Reference: PHYSIO 494

To appear in: *Journal of Physiology - Paris*



Please cite this article as: Uchitel, O.D., Inchauspe, C.G., Urbano, F.J., Di Guilmi, M.N., Ca_v2.1 voltage activated calcium channels and synaptic transmission in familial hemiplegic migraine pathogenesis, *Journal of Physiology - Paris* (2011), doi: [10.1016/j.jphysparis.2011.10.004](https://doi.org/10.1016/j.jphysparis.2011.10.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Cav2.1 voltage activated calcium channels and synaptic transmission in familial hemiplegic migraine pathogenesis

Oswaldo D. Uchitel, Carlota González Inchauspe, Francisco J. Urbano and Mariano N. Di Guilmi

Instituto de Fisiología, Biología Molecular y Neurociencias (CONICET). Departamento de Fisiología, Biología Molecular y Celular, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina.

Corresponding author: Dr. O.D. Uchitel,

E-mail: odu@fbmc.fcen.uba.ar

Facultad de Ciencias Exactas y Naturales. Pabellón 2, piso 2. Ciudad Universitaria. Buenos Aires 1428. Argentina.

FAX: (+54-11) 4576 3321

TEL: (+54-11) 4576 3368

Abstract

Studies on the genetic forms of epilepsy, chronic pain, and migraine caused by mutations in ion channels have given crucial insights into the molecular mechanisms, pathogenesis, and therapeutic approaches to complex neurological disorders. In this review we focus on the role of mutated Cav2.1 (i.e., P/Q-type) voltage-activated Ca²⁺ channels, and on the ultimate consequences that mutations causing familial hemiplegic migraine type-1 (FHM1) have in neurotransmitter release. Transgenic mice harbouring the human pathogenic FHM1 mutation R192Q or S218L (KI) have been used as models to study neurotransmission at several central and peripheral synapses. FHM1 KI mice are a powerful tool to explore presynaptic regulation associated with expression of Cav2.1 channels. Mutated Cav2.1 channels activate at more hyperpolarizing potentials and lead to a gain-of-function in synaptic transmission. This gain-of-function might underlie alterations in the excitatory/ inhibitory balance of synaptic transmission, favoring a persistent state of hyperexcitability in cortical neurons that would increase the susceptibility for cortical spreading depression (CSD), a mechanism believed to initiate the attacks of migraine with aura.

Keywords: Cav2.1 (P/Q-type) Ca²⁺ channels, familial hemiplegic migraine, cortical spreading depression, R192Q and S218L knock in mice, synaptic transmission.

*Index****Migraine and Familial Hemiplegic Migraine (FHM)***

- a. FHM type 2: Glial cells have a central role in migraine***
- b. FHM type 3: Sodium channels and migraine***
- c. FHM type 1: A central role of voltage activated calcium channels in migraine***
 - c.1. Cav2.1 (P/Q-type) calcium channels***
 - c.2. Familial Hemiplegic Migraine type 1.***
 - c.3. Functional consequences of FHM1 gene mutations: Studies in heterologous systems***
 - c.4. Functional consequences of FHM1 gene mutations: Studies in mice models including FHM1 CACNA1A Knock-in mutations***
 - c.5. Effects of FHM1 mutation on synaptic plasticity***
- d. Conclusions***
- e. Acknowledgements***
- f. Bibliography cited***

Migraine and Familial Hemiplegic Migraine (FHM)

Migraine is a common, chronic neurovascular disorder, typically characterized by recurrent attacks (1-3 days) of disabling headaches with associated autonomic symptoms. Twelve percent of the general population has on average one to two migraine attacks per month and treatments are frequently unsatisfactory. The etiology of migraine is multifactorial (for reviews, see Marmura and Silberstein 2011, Goadsby et al. 2002 and Pietrobon and Striessnig 2003). The migraine pain is likely to be triggered by activation of the trigeminovascular system, which primarily modulate sensory signal transmission through the activation of trigeminal afferents to meningeal blood vessels, the trigeminal nerve, and brainstem nuclei.

In 20% of cases, the migraine headache is preceded by a visual hallucination/illusion known as *aura*. There is growing evidence from animal models suggesting that CSD is the electrophysiological event underlying migraine aura (Haerter et al. 2005, Lauritzen 1994). In experimental animals, CSD is an intense and steady depolarization of neuronal and glial cell membranes that last for less than one minute, which can spread to contiguous cortical areas of the brain at a rate of 3 to 5 mm/min, regardless of functional cortical divisions or arterial territories. Evoked when local extracellular K^+ concentrations exceed a critical threshold, CSD is associated with disruption of membrane ionic gradients, massive influxes of Ca^{2+} and Na^+ , and massive K^+ efflux with concomitant glutamate release. All such ionic unbalance is believed to depolarize adjacent neurons and glia, thereby facilitating its spread. An advancing wave of brief excitation would then be followed by a longer-lasting inhibition of spontaneous and evoked neuronal activity that traverses the cortex (Somjen 2001, Goadsby 2007, Pietrobon 2005, Lauritzen 1994, Cutrer et al. 1998, Hadjikhani et al. 2001 and Bowyer et al. 2001). In humans, evidence for a causal role of CSD in the aura comes from functional magnetic resonance imaging (fMRI) performed during migraine attacks with aura (Hadjikhani et al 2001)

The nature and mechanisms of the primary brain dysfunction leading to the activation of the meningeal trigeminal nociceptors remain incompletely understood (for reviews see Charles 2009, Goadsby et al 2009a, Levy et al 2009) and it is controversial whether CSD can initiate the migraine headache cascade by itself. Animal studies support the idea that CSD may also initiate the headache mechanisms, but the connection between CSD and headache in patients (particularly those with migraine without aura) remains an open question (Ayata 2009, Pietrobon and Striessnig 2003). On the other hand, there is experimental animal evidence that CSD might activate the trigeminal sensory system, presumably by depolarizing perivascular trigeminal terminals at meningeal and dural blood vessels (Haerter et al. 2005, Ayata et al. 2006, Bolay et al. 2002). Zhang et al (2010, 2011) demonstrated, for the first time, that the induction of CSD by focal stimulation *in vivo* of the rat visual

cortex can lead to long-lasting activation of the nociceptors innervating the meninges in the trigeminal ganglion.

Current views of migraine describe it primarily as a multi-factorial brain disorder with an estimated heritability as high as 50% (Mulder *et al.* 2003). Genetic research in the field of migraines has mainly focused on the identification of genes involved in familial hemiplegic migraine (FHM), a rare monogenic subtype of migraines with aura. Genes for three monogenic subtypes of migraine have been identified so far (van den Maagdenberg *et al.* 2007): *CACNA1A* (FHM1, encoding the pore-forming α_1 subunit of $Ca_v2.1$ calcium channels; Ophoff *et al.* 1996), *ATP1A2* (FHM2, encoding the α_2 -subunit of sodium-potassium (Na^+ , K^+) pumps present in glial cell; De Fusco *et al.* 2003), and *SCN1A* (FHM3, encoding the pore-forming α_1 -subunit of neuronal $Na_v1.1$ voltage gated sodium channels, Dichgans *et al.* 2005).

More recently, in some patients suffering from migraine an homozygous mutation in the SLC4A4 gene (NBCe1, encoding the electrogenic Na^+ - HCO_3^- co-transporter was found; Suzuki *et al.* 2010). The immunohistological and functional analyses of these mutants demonstrate that the nearly total loss of NBCe1 activity in astrocytes can cause migraine potentially through dysregulation of synaptic pH.

A link between a common form of migraine and a genetic mutation of the TWIK-related spinal cord (TRESK) two-pore domain potassium channel (K2P), encoded by *KCNK18*, was recently reported (Lafreniere *et al.* 2010). By screening the *KCNK18* gene in a large multigenerational family with typical migraine with aura, inherited in a dominant fashion, Lafreniere *et al.* (2010) identified a mutation in TRESK (F139WfsX24). They also identified prominent TRESK expression in migraine salient areas such as the trigeminal ganglion. The F139WfsX24 mutation produces a frame shift that prematurely truncates the channel in the second transmembrane domain and causes a complete loss of TRESK function. In addition, the mutant subunit suppresses wild-type channel function through a dominant negative effect, thus, explaining the dominant penetrance of this allele. It is expected that the TRESK mutation would make the trigeminal neurons more easily excitable. Increasing TRESK activity might help to diminish the excitability of these neurons, and may lessen migraine severity or incidence, whereas decreasing activity (through *KCNK18* mutations) may increase the risk of migraine.

A recent genome-wide association study in a large clinic-based sample of European individuals with migraine identified a genetic variant on chromosome 8q22.1 associated with migraine (Anttila *et al.* 2011). Furthermore, in a new study including 5,122 migraineurs and 18,108 non-migraineurs, several single nucleotide polymorphism associations (SNPs) specific for migraine were identified (Chasman *et al.* 2011) establishing a link of migraine with LRP1 (modulator of glutamate signaling) and TRPM8 receptors (related to neuropathic pain models).

a. FHM type 2: Glial cells have a central role in migraine

Mutations in the *ATP1A2* gene are responsible for at least 20 percent of FHM cases. The FHM2 *ATP1A2* gene encodes the α_2 subunit of a Na^+ , K^+ pump ATPase (De Fusco *et al.* 2003, Marconi *et al.* 2003). This catalytic subunit binds Na^+ , K^+ , and ATP, and utilizes ATP hydrolysis to extrude Na^+ ions out of the cell while moving K^+ ions in. Na^+ pumping provides the steep Na^+ gradient essential for the transport of glutamate and Ca^{2+} . The gene is predominantly expressed in neurons at the neonatal stage

and in glial cells at the adult (De Fusco *et al.* 2003, Vanmolkot *et al.* 2003). An important function of this specific ATPase in adults is to modulate the key reuptake of potassium and glutamate from the synaptic cleft into the glia (Moskowitz *et al.* 2004). When this mechanism fails, it leads to elevated extracellular levels of glutamate and potassium, and accordingly, to an increased susceptibility to CSD (Pietrobon 2007, Koenderink *et al.* 2005). All FHM2 mutations studied in heterologous expression systems result in a “loss-of-function” or a kinetically altered Na⁺, K⁺ pump (Pietrobon 2007, Tavraz *et al.* 2008, Vanmolkot *et al.* 2006, De Fusco *et al.* 2003, Segall *et al.* 2005).

Leo *et al.* (2011) have reported the generation of the first mouse model of FHM2, a knock-in mutant harboring the human W887R-ATP1A2 mutation which decreases induction threshold and increases the velocity of propagation of CSD. The authors suggested a relatively minor role of the glial Na⁺,K⁺ pump in K⁺ clearance meaning that the reduced CSD threshold in FHM2 knockin mice would not be primarily due to impaired K⁺ reuptake. Instead their evidence point to an inefficient astrocyte clearance of glutamate and a consequent increased cortical excitatory neurotransmission.

b. FHM type 3: Sodium channels and migraine

FHM type 3 is caused by missense mutations in SCN1A, the genes encoding the pore-forming subunits of the neuronal voltage-gated Na⁺ channel Nav1.1 (Dichgans *et al.* 2005).

Kahlig *et al.* (2008) investigated the functional consequences of two mutations linked to FHM3:

Q1489K and L263V. Q1489K mutation, when compared to WT-Nav1.1, causes a predominantly loss-of-function phenotype, as deduced from a more rapid onset of slow inactivation, delayed recovery from fast and slow inactivation and a greater loss of channel availability during repetitive stimulation.

However, opposite effects of the same mutation on tsA-201 cells and rat cultured neurons transfected with human Nav1.1 were observed by Cestele *et al.* (2008). L263V exhibited biophysical abnormalities compatible with a gain-of-function mutation: increased persistent (not inactivating) current, delayed entry into fast and slow inactivation, depolarizing shifts in the steady-state voltage dependence of both fast and slow inactivation, accelerated recovery from fast inactivation, and greater channel availability during repetitive stimulation Kahlig *et al.* (2008). Notably, greater channel availability during repetitive stimulation, as seen for L263V mutation, has also been observed for various epilepsy-associated SCN1A mutations (Rhodes *et al.* 2005) and might correlate with the unusually high prevalence of epilepsy observed in L263V mutation carriers. The coincidence of migraine with epilepsy suggests that these two neurological disorders may share common mechanisms in L263V mutation carriers.

Nav1.1 is the major target for epileptogenic mutations and different mutations in this gene are known to be associated with epilepsy and febrile seizures (Meisler and Kearney 2005, Avanzini *et al.* 2007).

However, interestingly, just few cases of seizures have been reported in FHM patients (Dichgans *et al.* 2005, Vanmolkot *et al.* 2007). Thus, FHM mutations could be modifying Nav1.1 differently compared to purely epileptogenic mutations.

Nav1.1 is the major target for epileptogenic mutations and different mutations in this gene are known to be associated with epilepsy and febrile seizures (Meisler and Kearney 2005, Avanzini *et al.* 2007).

However, interestingly, just few cases of seizures have been reported in FHM patients (Dichgans *et al.* 2005, Vanmolkot *et al.* 2007). Thus, FHM mutations could be modifying Nav1.1 differently compared to purely epileptogenic mutations.

Nav1.1 is the major target for epileptogenic mutations and different mutations in this gene are known to be associated with epilepsy and febrile seizures (Meisler and Kearney 2005, Avanzini *et al.* 2007).

However, interestingly, just few cases of seizures have been reported in FHM patients (Dichgans *et al.* 2005, Vanmolkot *et al.* 2007). Thus, FHM mutations could be modifying Nav1.1 differently compared to purely epileptogenic mutations.

Studies using heterologous expression and functional analysis of recombinant Nav1.1 channels suggest that epilepsy mutations in Nav1.1 may cause either gain-of-function or loss-of-function effects that are consistent with either increased or decreased neuronal excitability (Ragsdale 2008, Cestele et al 2008). However, most Nav1.1 mutations have their ultimate epileptogenic effects by reducing Nav1.1-mediated whole cell sodium currents in GABAergic neurons, resulting in widespread loss of brain inhibition, an ideal background for the genesis of epileptic seizures (for review: Catterall et al 2010, Ragsdale 2008).

GABAergic interneurons are particularly sensitive to the loss of Nav1.1 in gene-targeted mice (Yu et al. 2006, Ogiwara et al. 2007), probably because Nav1.1 is expressed at higher levels in GABAergic interneurons than in glutamatergic pyramidal neurons. Hence, epileptogenic mutations may cause seizures because of decreased inhibition in neuronal circuits.

Because Nav1.1 channels are crucial for the generation and propagation of action potentials, the overall effects of FHM3 mutations are most likely an increased frequency of neuronal firing plus an enhanced neuronal excitability and neurotransmitter release.

c. FHM type 1: A central role of voltage activated calcium channels in migraine

Voltage-gated calcium channels can be classified based on their biophysical characteristics, into high-voltage-activated (HVA) and low-voltage-activated (LVA) channels. LVA Ca^{2+} channels encompass the family of T-type channels with three members (Cav3.1 through Cav3.3). They activate at a relatively negative threshold of around -60 mV, have a small conductance, and inactivate rapidly. They are inhibited by drugs like mibefradil (Ro 40-5967) or 2-octanol (Llinas 1988; Perez-Reyes 2003; Urbano et al., 2009; Bisagno et al., 2010). HVA Ca^{2+} channels include seven members (Cav1.1 to 1.4, Cav2.1 to 2.3) encoded by distinct genes each with multiple splice variants (Catterall 2000). HVA channels typically have an activation threshold of around -30 mV, with the exception of Cav1.3, which activates at around -50 mV (Xu and Lipscombe 2001). Cav1.1 to 1.4 channels, also referred to as L-type channels, are inhibited by three classes of drugs, dihydropyridines, phenylalkylamines, and benzothiazepines (Striessnig et al. 1998). Selective inhibitors for specific channels have also been described: Cav2.1 (P/Q-type) channels are selectively blocked by ω -agatoxin IVA (Mintz et al. 1992), while Cav2.2 (N-type) channels are inhibited by ω -conotoxin GVIA (Plummer et al. 1989, Aosaki and Kasai 1989) and Cav2.3 (R-type) channels are blocked by SNX-482 (Newcomb et al. 1998, Tottene et al. 2000).

HVA Ca^{2+} channels are heteromultimers of a pore forming $\alpha 1$ subunit that co-assembles with ancillary β , $\alpha 2\delta$ (and in some cases a γ subunit) into a functional channel complex (Baumgart and Perez-Reyes 2009, Catterall et al 2005). In contrast, LVA Ca^{2+} channels are $\alpha 1$ subunit monomers. The $\alpha 1$ subunit defines the channel subtype, whereas the ancillary subunits modulate $\alpha 1$ subunit function and surface expression. Vertebrates express four different types of calcium channel β subunits ($\beta 1$ through $\beta 4$),

four different types of $\alpha 2\delta$ subunits ($\alpha 2\delta 1$ through $\alpha 2\delta 4$), and as many as eight different γ subunits (Baumgart and Perez-Reyes 2009, Black 2003).

P/Q-type, N-type and, to some extent, R-type channels are highly expressed at presynaptic nerve terminals where their activities evoke neurotransmitter release (Katz et al. 1996, Iwasaki et al. 2000, Nudler et al. 2003, Trimmer and Rhodes 2004).

c.1. Cav2.1 (P/Q-type) calcium channels

Cav2.1 channel contains a pore-forming α_{1A} subunit and several regulatory subunits, including intracellular β subunits (Cav $\beta 1-4$) that bind to the intracellular loop between transmembrane domains I and II of $\alpha 1A$. The effect of the regulatory subunits is essential for increasing the expression levels and modulating the voltage-dependent activation and inactivation of Cav2.1 channels (Baumgart and Perez-Reyes 2009, Birnbaumer et al 1998, Catterall 2000, Di Guilmi et al., 2011, Jones 2002, Stotz et al 2004, Uchitel et al., 2010) as well as targeting the channels to the plasma membrane. Cav2.1 channels are located throughout the mammalian brain and spinal cord (Westenbroek et al. 1995) at presynaptic terminals (Catterall 1998, Wu et al. 1999, Mintz et al. 1995) and at somatodendritic membranes. The CACNA1A gene encoding Cav2.1 channels undergoes alternative splicing at multiple loci in an age-, gender-, and species-dependent manner (Bourinet et al 1999, Chang et al 2007, Chaudhuri et al 2005, Kanumilli et al 2006, Soong et al 2002). This mechanism results in multiple Cav2.1 splice variants with different outcomes in neuronal distribution and subcellular localization, biophysical properties, and sensitivity to ω -AgaIVA. Further functional diversity of Cav2.1 channels is generated by the combination of Cav2.1 $\alpha 1$ subunits with various auxiliary β and $\alpha 2\delta$ subunits (Luvisetto et al 2004). Indeed, evidence exists for large functional and pharmacological diversity of native Cav2.1 channels (Mermelstein et al 1999, Mintz et al 1992, Randall and Tsien 1995, Tottene et al 1996, Tringham et al 2007, Usowicz et al 1992). They play a prominent role in initiating action-potential-evoked neurotransmitter release both at peripheral neuromuscular junctions and central synapses, mainly within the cerebellum, brainstem, and cerebral cortex (Katz et al. 1996, Catterall 1998, Iwasaki et al. 2000). Even when pharmacological and electrophysiological studies in brain slices from rodents have revealed that P/Q-, N- and R-type channels cooperate in controlling release at many central excitatory synapses, P/Q-type channels have a dominant role, partly because of their more efficient coupling to the exocytotic machinery (González Inchauspe et al. 2004, Li et al 2007, Matsushita et al. 2002, Qian and Noebels 2001, 2000, Wu et al 1999, Mintz et al. 1995).

At many central synapses, including the inhibitory synapses between Purkinje cells and deep cerebellar nuclei neurons, there is a developmental change in the Ca^{2+} channel types mediating synaptic transmission, whereby the relative contribution of Cav2.1 channels increases with post-natal age, until it becomes exclusively dependent on P/Q-type channels (Iwasaki et al 2000). At certain synapses, N- or R-type Ca^{2+} channels fail to fully compensate the lack of Cav2.1 channels in Cav2.1^{-/-} and other

loss-of-function Cav2.1 mouse mutants (González Inchauspe et al 2004, 2007, Giugovaz et al 2011, Pietrobon 2005). Among the presynaptic Ca^{2+} channels, Cav2.1 channels are also unique in their capacity for interacting and being modulated in a complex manner by a number of Ca^{2+} -binding proteins (Catterall and Few 2008). As a result, Cav2.1 channels may exhibit either Ca^{2+} -dependent inactivation (CDI) or Ca^{2+} -dependent facilitation (CDF). Moreover, Ca^{2+} -dependent regulation of presynaptic Cav2.1 channels may play a crucial role in short-term synaptic plasticity during trains of action potentials (Adams et al 2010, González Inchauspe et al 2004, Mochida et al 2008, Muller *et al.* 2008, Takago *et al.* 2005, Xu and Wu 2005, González Inchauspe et al 2004, Tsujimoto et al 2002, Cuttle et al 1998, Takahashi *et al.* 1996).

Cav2.1 channels are expressed in those brain structures implicated in the pathogenesis of migraine, including the cerebral cortex, the trigeminal ganglia, and brainstem nuclei involved in the central control of nociception, their expression being particularly high in the cerebellum (Pietrobon and Striessnig 2003). In the cerebral cortex, excitatory synaptic transmission at pyramidal cell synapses in different cortical areas depends predominantly on Cav2.1 channels (Ali and Nelson 2006, Iwasaki et al 2000, Koester and Sakmann 2000, Rozov et al 2001, Tottene et al 2009, Zaitsev et al 2007). They also mediate about 40% of the AP-evoked Ca^{2+} influx in dendritic spines and shaft of layer 2/3 cortical pyramidal neurons (Koester and Sakmann 2000) and contribute to the regulation of the intrinsic firing of the same neurons via activation of different Ca^{2+} -dependent K^+ (K_{Ca}) channels (Pineda et al 1998). However, it is still a central problem in Ca^{2+} channel neurobiology studies to be able to translate their genetic complexity to their functional role in synaptic transmission. The availability of several mouse models with mutations in Ca^{2+} channel genes have provided a valuable tool to define Ca^{2+} channel functions. It would be expected that changes in the functional properties of Cav2.1 channels alter the behavior of neuronal networks.

c.2. Familial Hemiplegic Migraine type 1.

FHM1 is an autosomal-dominant subtype of migraine with aura, caused by a spontaneous missense mutation in the *CACNA1A* gene encoding the ion-conducting, pore-forming α_{1A} subunit of Cav2.1 channels (Pietrobon 2005, Ophoff et al. 1996). FHM1 missense mutations reported so far produce substitutions of conserved amino acids in important functional regions of the Cav2.1 channel including the pore lining and the voltage sensors (Cuenca-Leon et al 2008, Pietrobon 2007, Thomsen et al 2007) and have been associated with a wide range of clinical phenotypes (Haan et al. 2005, Kors et al. 2004, Alonso et al. 2004). Aside from the characteristic transient hemiparesis, typical attacks of FHM1 are identical to those of the common forms of migraine with aura (Pietrobon and Striessnig 2003, Pietrobon 2005, Thomsen et al. 2002). Different clinical phenotypes include pure types of FHM1 (Ophoff *et al.* 1996), combinations of FHM1 with various degrees of cerebellar ataxia (Ducros et al. 2001, Ophoff et al. 1996), or fatal coma due to excessive cerebral edema (Kors et al. 2001, 2002).

Disorders not associated with FHM are episodic ataxia type 2 (Zafeiriou et al 2009, Cuenca-Leon et al 2009, Cricchi et al 2007, Jen et al. 2004, 2007, 2008, Ophoff et al. 1996), progressive ataxia (Yue et al. 1997), spinocerebellar ataxia type 6 (Tsou et al 2011, Restituito et al 2000, Ishikawa et al 1999, Zhuchenko et al. 1997) and absence (Imbrici et al. 2004) and generalized epilepsy (Haan et al. 2005, Jouvenceau et al. 2001). In addition FHM1 mutations were also found in family members who had only “normal” no-paretic migraine but no FHM. This suggests that gene mutations for FHM may also be responsible for the common forms of migraine, probably due to different genetic and no-genetic modulating factors. Hence, FHM is considered a valid monogenic model for studying the pathogenetic mechanisms involved in the genetically more complex common forms of migraine (Ferrari *et. al.* 2008, Ducros et al 2001).

c.3. Functional consequences of FHM1 gene mutations: Studies in heterologous systems

Biophysical analysis of FHM1 Ca²⁺ channel dysfunction in heterologous systems show that the FHM1 mutations alter many biophysical properties of human Cav2.1 channels in a complex way, which is rather debateable since both loss-of-function and gain-of-function phenotypes have been reported (Cao et al. 2004, Barrett et al. 2005, Cao and Tsien 2005, Tottene et al. 2002, Kraus et al. 2000, 1998, Hans et al. 1999). The most consistent effect revealed by the analysis of single-channel properties of human Cav2.1 channels carrying FHM1 mutations was an increase in channel open probability and in single channel Ca²⁺ influx over a broad voltage range, mainly due to a shift of channel activation to more negative voltages (Tottene et al. 2002, 2005, Hans et al. 1999). Shifts to lower voltages of activation of the mutant channels were also revealed by measuring whole-cell currents in heterologous expression systems and transfected neurons (Adams et al 2009, Melliti et al 2003, Mullner et al 2004, Serra et al 2009, Tottene et al 2002, 2005, Weiss et al 2008). As a result, channels open with smaller depolarizations. However, the overall change in calcium influx at presynaptic terminals is difficult to predict since there have been contradictory results. For instance, Cao and colleagues (2004) found evidence for reduced calcium influx at the whole cell level in transfections of cultured mouse hippocampal neurons. What happens in the mutant brain will therefore be determined by the delicate interplay between the functional effects of a particular mutation, the different channel properties and density, the type of channel subunits, and the direct and indirect cellular environment. Since the biophysical effects of FHM1 mutations can be affected by a number of factors, including Cav2.1 splice-variation (Adams et al 2009), β subunit coexpression (Mullner et al. 2004), and the nature of the expression system (Kraus et al 1998), it seems pertinent to study the functional consequences of gene mutations on P/Q-type Ca²⁺ and synaptic transmission in their native neuronal environment and at their endogenous level of expression in knock-in mouse models carrying human pathogenic mutations.

c.4. Functional consequences of FHM1 gene mutations: Studies in mice models including FHM1 CACNA1A Knock-in mutations

The generation of two knock-in FHM1 mouse models carrying either the human pathogenic R192Q or S218L missense mutations allowed the first analysis of the functional consequences of FHM1 mutations on Cav2.1 channels and synaptic transmission in neurons expressing the channels at the endogenous physiological level (Maagdenberg et al 2010, 2004, Kaja et al. 2010, Tottene et al 2009, González Inchauspe et al 2010, Adams et al 2010).

The R192Q KI mice exhibit no overt clinical phenotype or structural abnormalities. This situation is very similar in humans, where the R192Q mutation causes only mild FHM attacks, very similar to the common forms of migraine, albeit with hemiparesis, but without any other neurological symptoms. The S218L mutation causes a severe migraine phenotype combined with slowly progressive cerebellar ataxia and atrophy, epileptic seizures, coma or profound stupor, and severe, sometimes fatal, cerebral edema which can be triggered by a trivial head trauma (Chan et al 2008, Kors et al 2001, van den Maagdenberg et al 2010).

As noted before, CSD is implicated in the pathophysiology of FHM1. In both R192Q and S218L KI mice, a lower threshold for CSD induction and an increased velocity of CSD propagation was reported (van denMaagdenberg *et al.* 2004, 2010, Tottene *et al.* 2009, Eikermann-Haerter et al. 2009 *a*). In agreement with the higher incidence of migraine in females, the velocity of propagation and the frequency of CSDs were larger in females than in males of both mutant strains (Eikermann-Haerter *et al.* 2009 *a,b*). The sex difference was abrogated by ovariectomy and enhanced by orchietomy, suggesting that female and male gonadal hormones exert reciprocal effects on CSD susceptibility. A female predominance is also described for familial (5:2 ratio) and sporadic (4.25:1 ratio) hemiplegic migraine (Eriksen et al 2006, Thomsen et al 2007, 2002). All these findings underscore the complex synergistic interactions between genetic and hormonal factors determining migraine susceptibility.

Accordingly to the larger gain-of-function observed *in vitro*, the S218L KI mice shows higher CSD susceptibility, *in vivo*, compared to the R192Q mice. Experimentally induced CSD induces pure hemiplegia in R192Q mutant mice, whereas S218L mutants additionally develop coma and often fatal seizures (van den Maagdenberg et al. 2010, Eikermann-Haerter et al. 2009). Recently, Eikermann-Haerter et al (2011) have provided evidence for an enhanced subcortical SD susceptibility in both FHM1 mutant mice compared to WT. Whereas the facilitated subcortical spread appeared limited to the striatum in R192Q mutants, hippocampal and thalamic spread was detected in the S218L mutants with an allele-dosage effect. Their findings suggest a role for subcortical SD as a potential mechanism to explain hemiplegia, seizures, and coma in FHM1.

Other studies in knock in animals revealed multiple gain-of-function effects. In cerebellar granule cells and cortical pyramidal cells of R192Q and S218L KI mice, the Cav2.1 Ca²⁺ current density was larger

than in WT neurons in a wide range of relatively mild depolarizations, reflecting activation of mutant mouse Cav2.1 channels at 8–9 mV more negative voltages than the corresponding WT channels. P/Q current densities were similar in KI and WT neurons at higher voltages (that elicit maximal Cav2.1 channel open probability), indicating similar densities of functional Cav2.1 channels (van den Maagdenberg et al 2004, 2010, Tottene et al 2009). Thus, in two different FHM1 mouse models and two different types of neurons the functional consequences of the FHM1 mutations on native neuronal mouse Cav2.1 channels were quite similar to those on single recombinant human Cav2.1 channels (Tottene et al 2002, 2005). In agreement with the lower threshold of activation of human S218L Cav2.1 channels compared to that of human R192Q CaV2.1 channels (Tottene et al 2005), the gain of function of the P/Q current at low voltages was larger in S218L than R192Q KI mice (van den Maagdenberg et al 2004, 2010). Similarly, a negative shift in the activation voltage of presynaptic Cav2.1 mutated channels was confirmed at the calyx of Held synapse of KI S218L mice (Di Guilmi et al, unpublished results).

The first indication that the gain of function of Cav2.1 channels produced by FHM1 mutations could lead to enhanced evoked neurotransmitter release was obtained at the neuromuscular junction (NMJ), where evoked neurotransmission was unaltered at physiological Ca^{2+} ion concentrations but increased at 0.2 mM Ca^{2+} in KI mice (Kaja et al 2005, van den Maagdenberg et al 2004, 2010). Recently, cortical excitatory neurotransmission was investigated in neuronal microcultures and in brain slices from homozygous R192Q KI mice (Tottene et al 2009). The results show increased synaptic strength at physiological Ca^{2+} concentration, presumably, due to enhanced action potential-evoked Ca^{2+} influx through mutant Cav2.1 channels and enhanced probability of glutamate release at cortical pyramidal cell (PC) synapses. Short-term synaptic depression during trains of action potentials was enhanced. Neither amplitude nor frequency of miniature postsynaptic currents were altered, indicating the absence of homeostatic compensatory mechanisms at excitatory synapses onto pyramidal cells (Tottene et al 2009).

Characteristically, CSD arises spontaneously in response to specific triggers that somehow create in the cortex of migraineurs the conditions for initiating the positive feedback cycle that overwhelms the regulatory mechanisms controlling cortical $[\text{K}^+]_o$ and ignites CSD. Insights into how this might occur have been provided by the interesting finding that, in contrast with the enhanced glutamatergic excitatory neurotransmission at PC synapses, inhibitory GABAergic neurotransmission at fast spiking (FS) interneurons was not altered in R192Q KI mice, despite being mediated by Cav2.1 channels (Tottene et al 2009). Given the evidence that the magnitude (or even the presence) of the negative shift in activation of human Cav2.1 channels produced by FHM1 mutations may depend on the particular recombinant Cav2.1 $\alpha 1$ splice variant and/or Cav2.1 β subunit (Mullner *et al.* 2004, Adams *et al.* 2009), a possible explanation for the unaltered inhibitory transmission at the FS interneurons may be

the presence of a Cav2.1 isoform that is little affected by the mutation or a near saturation of the presynaptic Ca^{2+} sensors. Evoked transmitter release was also found to be normal at the neuromuscular junction and at the calyx of Held synapse of R192Q KI mice when studied in normal Ca/Mg concentration in the external solution (González Inchauspe et al 2010). This finding was unexpected since mutated R192Q Cav2.1 channels at the calyx of Held nerve terminals are active at lower membrane potentials as predicted. Interestingly, transmitter release at the FS interneurons, at the neuromuscular junction and at the calyx of Held is triggered by a relatively brief action potential. In each of these R192Q KI preparations transmitter release is not affected. Thus, one factor that may be controlling the expression of a synaptic gain of function in the FHM1 KI mice is the duration of the action potential (AP) triggering transmitter release. To address this question we used the calyx of Held, a giant glutamatergic synapse in the mammalian auditory brainstem where, due to its size and accessibility, it is possible to make direct patch-clamp recordings from the presynaptic nerve terminal and its postsynaptic target, the principal neurons of the medial nucleus of the trapezoid body (MNTB). The large size of the calyx of Held allows it to harbor hundreds of active zones and thus a single presynaptic AP releases hundreds of quanta, generating a large excitatory postsynaptic current (EPSC) that rapidly depolarizes the MNTB neuron to threshold. A single MNTB principal neuron receives input from only one calyx of Held presynaptic terminal, and glutamatergic synaptic transmission mediated by AMPA receptors is triggered almost exclusively by Cav2.1 channels in mature mice (for a review, Schneggenburger and Forsythe 2006). The presynaptic Ca^{2+} currents (I_{pCa}) from the calyx of Held terminals recorded in brainstem slices also activate at more hyperpolarizing potentials in R192Q KI than in WT mice (González Inchauspe et al 2010). Assuming presynaptic Ca^{2+} currents through Cav2.1 channels can be modeled by Hodgkin–Huxley equations, a shift to more negative activation voltages by the FHM1 mutation theoretically should generate larger Ca^{2+} currents during APs (Borst and Sakmann 1999). However, the short duration of the calyx of Held APs (half-width: 0.45 ± 0.02 ms) elicited I_{pCa} with similar amplitudes in both WT and R192Q KI mice. Only when the AP duration was prolonged to 2 ms (typical mean half-width of pyramidal neuron APs), the shift in the activation by the R192Q mutation was sufficient to cause greater presynaptic Ca^{2+} influx (González Inchauspe et al 2010). According to the similarity in I_{pCa} amplitudes evoked by physiological calyx of Held APs, there were no differences in mean amplitudes of EPSCs between WT and R192Q KI mice. However, when synaptic transmission was triggered by long duration APs (obtained by inhibiting K^+ channels with 4-Aminopyridine and tetraethylammonium chloride), FHM1 R192Q KI mice did show a significant increase in both amplitude and area of EPSCs compared to WT mice (González Inchauspe et al., unpublished results). Thus, the expression of FHM1 mutations might vary according to the shape of the APs in charge of triggering synaptic transmission, adding to the complexity of the pathophysiology of migraine.

c.5. Effects of FHM1 mutation on synaptic plasticity

The dynamics and strength of neural circuits are essential for encoding and processing information in the CNS and rely on short and long forms of synaptic plasticity. In several synapses, repetitive stimulation causes short term depression (STD). Short-term depression of synaptic release has been traditionally accredited to depletion of the release-ready vesicle pool (von Gersdorff and Borst 2002, Wong et al. 2003, Zucker and Regehr 2002, Wang and Kaczmarek 1998), although other mechanisms like receptor desensitization (Wong et al. 2003) may be involved. At the large calyx of Held synapse, it has been shown that short-term synaptic plasticity is also achieved by a mechanism involving the regulation of presynaptic Cav2.1 Ca^{2+} currents (Cuttle et al 1998, Tsujimoto et al 2002), like calcium channel inactivation (Xu and Wu 2005, Muller *et al* 2008, Forsythe et al. 1998, Di Guilmi et al. 2011) and calcium channel inhibition by presynaptic metabotropic glutamate receptor (von Gersdorff et al. 1997, Takahashi *et al.* 1996) or AMPA receptors (Takago *et al.* 2005). Hennig et al. (2008) have developed a model of synaptic depression at the calyx of Held synaptic terminal that combines many of the mechanisms involved in short term depression and plasticity, including vesicle recycling, facilitation, activity-dependent vesicle retrieval and multiple mechanisms affecting calcium channel activity and release probability. The efficacy of synaptic transmission during repetitive stimulation is also determined by the rate of recovery from STD, owed to the replenishment of the readily releasable pool of synaptic vesicles, which is dynamically regulated by Ca^{2+} influx through voltage-gated Ca^{2+} channels in an activity dependent manner (Zucker and Regehr 2002, Wang and Kaczmarek 1998). High-frequency stimulation of presynaptic terminals significantly enhances the rate of replenishment. Therefore, mutations or neuromodulators capable of altering the activation/inactivation properties of calcium channels (Di Guilmi et al. 2011; Uchitel et al. 2010) may also alter STD and/or vesicle replenishment. Recordings in connected MNTB neurons at the calyx of Held synapses show that EPSC amplitudes depress to the same extent and with similar kinetics in WT and R192Q KI mice, but, interestingly, the rate of recovery of EPSCs amplitudes was significantly faster in R192Q KI compared to WT synapses (González Inchauspe et al., unpublished results). Ca^{2+} -dependent facilitation (CDF) of Cav2.1 channels is an important mechanism required for normal synaptic plasticity at fast synapses in the mammalian CNS. As an example, at the calyx of Held synapse of Cav2.1 KO mice, lack of pair pulse facilitation of presynaptic Ca^{2+} channels was in correlation with absence of facilitation of synaptic transmission (González Inchauspe et al 2004, 2007). In the same animal model, pair pulse facilitation of end plate potential was also impaired in the neuromuscular junction (Urbano et al 2003). Short-term facilitation of synaptic release has been attributed to enhanced vesicle release resulting from the accumulation of intracellular Ca^{2+} in presynaptic terminals during repetitive action potentials (APs), where the buildup of residual Ca^{2+} enhances binding to those sensor proteins directly mediating vesicle fusion and transmitter release

(Atluri et al 1996, Zucker and Regehr 2002). There may exist at least another mechanism of facilitation, driven by a Ca^{2+} sensor with high Ca^{2+} affinity that can detect modest, transient levels of Ca^{2+} , most likely near the pore of presynaptic Ca^{2+} channels (Atluri et al 1996). A recent study using recombinant Cav2.1 channels expressed in cultured superior cervical ganglion neurons demonstrated that the Ca^{2+} -dependent facilitation (CDF) and Ca^{2+} -dependent inactivation (CDI) of Cav2.1 channels are mediated through neuronal Ca^{2+} sensor proteins (CaSs) like calmodulin (CaM), which binds the Cav2.1 subunit carboxyl terminus (Mochida et al 2008). CaM-mediated CDF and CDI are robust forms of Cav2.1 channel modulation in which CaM interacts with the Cav2.1 carboxyl terminus in a bipartite regulatory processes where CDF is mediated by a local increase in Ca^{2+} and CDI through a global increase in Ca^{2+} (Tsujiimoto et al 2002, Chaudhuri et al 2007). Adams et al (2010) provide supporting evidence for the hypothesis that Ca^{2+} influx at Purkinje fibers boutons induces calcium-dependent facilitation of Cav2.1 channels as a means to enhance Ca^{2+} influx during subsequent APs and achieve synaptic facilitation at the cerebellar parallel fiber (PF)-to-Purkinje cell (PC) central synapse. Interestingly, they demonstrated that mutations associated with human diseases affect this process. They observed that FHM1 gain-of-function missense mutations (R192Q and S218L) significantly occlude CDF of Cav2.1 currents in recombinant and native systems and that this alteration correlates with reduced short term synaptic facilitation at cerebellar PF-to-PC synapses. These findings suggest that CDF of Cav2.1 is an important mechanism required for normal synaptic plasticity at a fast synapse in the mammalian CNS. Additional evidence provided by two-photon imaging supported the fact that Cav2.1 FHM1 mutant presynaptic channels are in a constitutively facilitated state that basally increases presynaptic Ca^{2+} influx and transmitter release. Thus, observed changes in synaptic plasticity at the PF-to-PC synapse may result from a larger initial Ca^{2+} influx through basally facilitated mutant channels relative to unfacilitated WT channels in Purkinje fibers boutons. Overall, these results provide evidence that FHM1 mutations directly affect both Cav2.1 channel CDF and synaptic plasticity and that together they likely contribute toward the pathophysiology underlying FHM1. The findings also suggest that Cav2.1 channel CDF is an important mechanism required for normal synaptic plasticity at a fast synapse in the mammalian CNS.

d. Conclusions

Calcium channels containing the I_A subunit (Cav2.1 or P/Q-type) are expressed throughout the human and mammalian brain with a higher concentration in the cerebellum and are localized in most presynaptic terminals (Catterall 1998, Wu et al. 1999, Mintz et al. 1995) as well as in the cell body and dendrites of many neurons (Volsen et al. 1995, Westenbroek et al. 1995). In many central synapses, Cav2.1 channels are preferentially located at the release sites and are more effectively coupled to neurotransmitter release than other Ca^{2+} channel types (Li et al 2007, Matsushita et al. 2002, Qian and

Noebels 2001, 2000, Dunlap et al. 1995, Mintz et al. 1995, Wu et al. 1999, Iwasaki et al. 2000). At these synapses, the action potential-evoked Ca^{2+} influx and the local Ca^{2+} increase that triggers neurotransmitter release are mainly determined by the kinetics of opening and closing, the open probability and the unitary conductance of $\text{Ca}_v2.1$ channels (Borst and Sakmann 1998, Sabatini and Regehr 1999, Meinrenken et al. 2002, 2003).

In addition to the types of voltage-dependent Ca^{2+} channels mediating Ca^{2+} entry at the terminal, the relationship between presynaptic Ca^{2+} influx and neurotransmitter release is a critical aspect characterizing the Ca^{2+} induced-release of neurotransmitter. Neurotransmitter release is a power function of presynaptic Ca^{2+} current, with a power number between 3 and 4 (Dodge and Rahamimoff 1967, Bollmann et al. 2000, Schneggenburger and Neher 2000, Mintz et al. 1995). So, even small changes in presynaptic Ca^{2+} influx have a large effect on the number of vesicles released by an action potential (Sabatini and Regehr 1999, Meinrenken et al. 2002). The impact of a relatively small change in the kinetics of activation of presynaptic Ca^{2+} channels on the Ca^{2+} current and neurotransmitter release evoked by an action potential at a central synapse has been shown by Borst and Sakmann (1998). The importance of the kinetics of activation of presynaptic Ca^{2+} channels in determining the fraction of channels that open during a short action potential and consequently, in Ca^{2+} current waveform and neurotransmitter release has been also described by Sabatini and Regehr (1999) and our group (González Inchauspe et al. 2010). Liu et al. (2003) have shown that different kinetics of inactivation and especially voltage-dependence of steady-state inactivation of Ca^{2+} channels have a striking impact on the amount and temporal pattern of Ca^{2+} influx in response to repetitive firing waveforms.

The specifically dominant and efficient roles of $\text{Ca}_v2.1$ channels in controlling fast neurotransmitter release from central excitatory synapses suggest that the human and mouse $\text{Ca}_v2.1$ channelopathies and their episodic neurological symptoms, ranging from migraine to absence epilepsy and ataxia, might be primarily synaptic diseases. These different disorders probably arise from a disruption of the finely tuned balance between excitation and inhibition in neuronal circuits of specific brain regions: the cortex in the case of migraine, the thalamus in the case of absence epilepsy and the cerebellum in the case of ataxia. Synapses are affected differently depending on intrinsic differences in the relative reliance on P/Q channels and shape of the action potential.

The mutations in the $\text{Ca}_v2.1$ channels linked to FHM1 affect the biophysical properties and the density of presynaptic calcium currents as well as several other properties of synaptic transmission, like short term plasticity. A gain of function was observed in R192Q and S218L KI mice, that includes an increased Ca^{2+} influx through $\text{Ca}_v2.1$ channels as a consequence of mutant channels that open at lower voltages than WT channels (González Inchauspe et al. 2010, Tottene et al. 2002, 2005, 2009, van den Maagdenberg et al. 2004, 2010). Given the three-fourth power dependence of neurotransmitter release

on intracellular Ca^{2+} concentration, small changes in amplitude or time course of Ca^{2+} influx at the release sites are expected to be very effective in modulating transmitter output at those synapses where the Ca^{2+} sensors are not saturated during an action potential (Schneggenburger and Neher 2000). This predicts that, at such synapses, $\text{Ca}_v2.1$ channels that open more readily and at lower voltages because of the R192Q mutation will lead to an increased action potential-evoked Ca^{2+} influx and a consequent increase in neurotransmitter release. At the calyx of Held in P11 and older mice, transmitter release is triggered exclusively by P/Q-type Ca^{2+} channels. The hyperpolarizing shift in both the activation and inactivation of $\text{Ca}_v2.1$ channels had little effect on presynaptic Ca^{2+} currents evoked by the short duration APs of the same calyces. In consequence, similar EPSC amplitudes, release probability and pair pulse facilitation were observed in R192Q KI mice compared to WT ones. These results contrast with the increased release probability of the glutamatergic pyramidal cell synapses recently reported by Tottene et al. (2009). Nevertheless, they agree with the normal transmitter release observed at the fast spiking interneuron inhibitory synapses and at the neuromuscular junction studied in the same animal model (Kaja et al 2005, Tottene et al 2009).

Interestingly, R192Q KI mice did show a significant enhancement in both I_{pCa} and EPSCs at the calyx of Held synapse when currents were elicited by long duration APs. These findings strongly suggest that synapses driven by larger amplitude and short duration APs (e.g., calyx of Held and interneurons APs) are less affected by the mutation-induced hyperpolarizing shift in voltage-dependence of Ca^{2+} channel activation, than those driven by longer duration APs (e.g., pyramidal neurons APs). The differences in AP durations that trigger cortical excitatory and inhibitory synapses might explain the unaltered inhibitory neurotransmission observed by Tottene et al. (2009) at the fast spiking (FS) interneuron-pyramidal cell (PC) synapses. It may also explain the gain of function observed at the PC-FS interneuron excitatory synapses, since several types of interneurons and other neurons that display fast spiking behavior have APs with short half-widths durations (Ali et al. 2007), while PCs depicts long APs. Additional factors like a faster recovery of vesicle recycling during high frequency

transmission could also contribute to the increased excitability in FHM1 mutant mice. Although an established model explaining migraine attacks is still lacking, a favored hypothesis considers that the abnormal balance of cortical excitation-inhibition and the resulting persistent state of hyperexcitability of neurons in the cerebral cortex may be associated with the increased susceptibility for cortical spreading depression (CSD), which is believed to initiate the episodes of migraine with aura (van den Maagdenberg et al 2010, Tottene et al 2009, Haerter et al. 2005, Pietrobon 2005, Lauritzen 1994, Welch 1998, Flippen and Welch 1997). Several results also indicate that the underlying mechanism changing synaptic strength by the R192Q mutation might be related to the variability in calcium currents according to the shape of the action potentials.

Genetic migraine models are useful in unraveling the triggering mechanisms for migraine attacks and in identifying novel migraine preventive targets and therapies. The FHM data support a key role of CSD in the pathogenesis of migraine aura and possibly migraine headache. It also points to cortical hyperexcitability as the basis for vulnerability to CSD. Cortical spreading depression seems to be a common mechanism underlying different forms of FHM (Moskowitz et al. 2004, Goadsby 2007). Mutations in the FHM1 CACNA1 gene cause increased neuronal release of neurotransmitters, more specifically glutamate in the cortex (Pietrobon et al. 2010) that can induce, maintain, and propagate CSD. On the other hand mutations in the FHM2 sodium-potassium pump gene cause reduced reuptake of K^+ and glutamate from the synaptic cleft into the glial cell (Moskowitz et al 2004), whereas mutations in the FHM3 sodium channel gene result in hyperexcitability, excessive neuronal firing and most likely increased release of neurotransmitters in the synaptic cleft. The overall result is an increased level of glutamate and K^+ in the synaptic cleft resulting in an elevated propensity for CSD (van den Maagdenberg et al. 2004), which supports the concept that the migrainous brain is hyperexcitable (Aurora and Wilkinson 2007). This fact could easily explain the aura of FHM attacks. More controversial, however, is whether the enhanced tendency for CSD might also be responsible for triggering the headache phase. Lastly, the most important question that remains to be answered is whether the same mechanisms also are involved in the common forms of migraine with and without aura.

More than 15 years after the introduction of sumatriptan, triptans continue to be a mainstay for acute migraine therapy. Drugs selectively targeting the circuits now implicated in migraine, such as glutamate, nitric oxide, triptans, 5-hydroxytryptamine (5-Ht) 1B/1D receptor agonists (Ferrari et al 2001) and calcitonin gene-related peptide (CGRP) receptor antagonists have demonstrated efficacy in both preventing (topiramate) and aborting (triptans, CGRP receptor antagonists) acute migraine attacks. Among these, CGRP receptor antagonists, such as olcegepant (BiBn4096Bs, Olesen et al 2004) and telcagepant (Ho et al 2008, 2009, Williams et al 2006) may be the closest to availability for widespread clinical use (Ho et al 2010). Novel therapeutic strategies that consider CSD and cortical hyperexcitability as key targets of preventive migraine treatment promise to improve the lives of the large number of migraine sufferers. Notably, although drugs with migraine preventive activity belong to a wide range of pharmacological classes (e.g., anti-epileptics and blockers of calcium channels and serotonergic, beta-adrenergic, and histaminergic receptors), they all seem to share anti-CSD activity (Akerman & Goadsby 2005, Ayata et al. 2006). A particularly interesting compound is Tonabersat (SB-220453), which inhibits CSD, CSD-induced nitric oxide (NO) release and cerebral vasodilation (Smith et al. 2000). If proven effective, it would be the first migraine prophylactic agent developed on the basis of the CSD hypothesis (Goadsby et al 2009b). Other examples of potential compounds effective as preventive therapy for chronic migraine are Memantine, an activity-dependent *N*-methyl-D-

aspartate receptor blocker, and Topiramate (Diener et al 2007, Silberstein et al 2007, Bussone et al. 2005, Storer and Goadsby 2004, 2005), both inhibiting CSD (Goadsby et al 2009b, Akerman and Goadsby 2004). In an attempt to correlate the preventive action of a range of medications with an effect on CSD, Ayata et al. (2006) have studied topiramate, valproate, amitriptyline, propranolol and methysergide. Topiramate, valproate and amitriptyline inhibited CSD induction at doses that are typically used in rodent studies, none of which was similar to the dose that is used in migraine prevention. The dose of propranolol used was four times that reported in previous rat studies (Smits et al. 1980), whereas methysergide had no effect on the speed of propagation (Ayata et al. 2006). CSD inhibition might thus be a good model for the study of migraine to help in developing preventive drugs.

e. Acknowledgements

This work was supported by the following grants: UBACYT X-223; PICT BID 1728 OC.AR.PICT 2005 N 32113; FONCYT (ANPCyT) PICT BID 1728 OC.AR.PICT 2006 N 199 (Argentina) and Wellcome Trust #084636 (UK), (to Dr. Uchitel); FONCYT (ANPCyT) PICT 2007–1009, PICT 2008–2019 and PIDRI-PRH 2007 (to Dr. Urbano). In addition, Dr. F.J. Urbano is a 2011 fellow of the John Simon Guggenheim Memorial Foundation.

f. Bibliography cited

Adams PJ, Garcia E, David LS, Mulatz KJ, Spacey SD, Snutch TP (2009) Ca(V)2.1 P/Q-type calcium channel alternative splicing affects the functional impact of familial hemiplegic migraine mutations: implications for calcium channelopathies. *Channels (Austin)* 3:110-121

Adams PJ, Rungta RL, Garcia E, van den Maagdenberg AM, MacVicar BA and Snutch TP (2010) Contribution of calcium-dependent facilitation to synaptic plasticity revealed by migraine mutations in the P/Q-type calcium channel. *PNAS* 107: 18694 - 18699

Akerman S and Goadsby PJ (2004) Topiramate inhibits cortical spreading depression in rat and cat: a possible contribution to its preventive effect in migraine. *Cephalalgia* 24, 783-784.

Akerman S. and Goadsby PJ (2005) Topiramate inhibits cortical spreading depression in rat and cat: impact in migraine aura. *NeuroReport* 16, 1383-1387.

Ali AB and Nelson C (2006) Distinct Ca²⁺ channels mediate transmitter release at excitatory synapses displaying different dynamic properties in rat neocortex. *Cereb Cortex* 16: 386-393.

- Ali AB, Bannister AP and Thomson AM (2007) Robust correlations between action potential duration and the properties of synaptic connections in layer 4 interneurons in neocortical slices from juvenile rats and adult rat and cat. *J. Physiol.* 580: 149-169.
- Alonso I, Barros J, Tuna A, Seixas A, Coutinho P, Sequeiros J and Silveira I (2004) *Clin. Genet.* 65: 70-72.
- Anttila V, Wessman M, Kallela M and Palotie A (2011) Towards an understanding of genetic predisposition to migraine. *Genome Med.* 3(3): 17.
- Aosaki T, Kasai H (1989) Characterization of two kinds of high-voltage-activated Ca-channel currents in chick sensory neurons. Differential sensitivity to dihydropyridines and omega-conotoxin GVIA. *Pflügers Arch* 414: 150-156.
- Atluri PP, Regehr WG (1996) Determinants of the time course of facilitation at the granule cell to Purkinje cell synapse. *J Neurosci* 16:5661-5671.
- Aurora SK and Wilkinson F (2007) The brain is hyperexcitable in migraine. *Cephalalgia* 27:1442-1453.
- Avanzini G, Franceschetti S and Mantegazza M (2007) Epileptogenic channelopathies: experimental models of human pathologies. *Epilepsia* 48 Suppl 2: 51-64.
- Ayata C, Jin H, Kudo C, Dalkara T and Moskowitz MA (2006) Suppression of cortical spreading depression in migraine prophylaxis. *Ann Neurol* 59: 652-661.
- Ayata C (2009) Spreading depression: from serendipity to targeted therapy in migraine prophylaxis. *Cephalalgia* 29:1095-1114
- Barrett CF, Cao YQ and Tsien RW (2005) Gating deficiency in a familial hemiplegic migraine type 1 mutant P/Q-type calcium channel. *J. Biol. Chem.* 280: 24064-24071.
- Baumgart JP and Perez-Reyes E (2009) Voltage-gated Ca²⁺ channels. In: Kew JN, Davies C (eds) *Ion channels: from structure to function*. Oxford University Press, London, pp. 104-130
- Birnbaumer L et al. (1998) Structures and functions of calcium channel β subunits. *J Bioenerg Biomembr* 30:357-375.
- Black JL 3rd (2003) The voltage-gated calcium channel γ subunits: a review of the literature. *J Bioenerg Biomembr* 35:649-660.
- Bisagno V, Raineri M, Peskin V, Wikinski SI, Uchitel OD, Llinás RR, Urbano FJ (2010) Effects of T-type calcium channel blockers on cocaine-induced hyperlocomotion and thalamocortical GABAergic abnormalities in mice. *Psychopharmacology (Berl)*. 212(2): 205-214.
- Bolay H, Reuter U, Dunn AK, Huang Z, Boas DA, Moskowitz MA (2002) Intrinsic brain activity triggers trigeminal meningeal afferents in a migraine model. *Nat Med* 8: 136-142.
- Bollmann JH, Sakmann B and Borst JGG (2000) Calcium Sensitivity of Glutamate Release in a Calyx-Type Terminal. *Science* 289: 953-957.

- Borst JGG and Sakmann B (1998) Calcium current during a single action potential in a large presynaptic terminal of the rat brainstem. *J. Physiol.* 506: 143-157.
- Borst JG and Sakmann B (1999) Effect of changes in action potential shape on calcium currents and transmitter release in a calyx-type synapse of the rat auditory brainstem. *Philos Trans R Soc Lond B Biol Sci* 354: 347-355.
- Bourinet E, Soong TW, Sutton K, Slaymaker S, Mathews E, Monteil A, Zamponi GW, Nargeot J, Snutch TP (1999) Splicing of alpha 1A subunit gene generates phenotypic variants of P- and Q-type calcium channels. *Nat Neurosci* 2:407-415
- Bowyer SM, Aurora KS, Moran JE, Tepley N and Welch KM. (2001) Magnetoencephalographic fields from patients with spontaneous and induced migraine aura. *Ann. Neurol.* 50, 582-587.
- Bussone G. et al. (2005) Topiramate 100 mg/day in migraine prevention: a pooled analysis of double-blinded randomised controlled trials. *Int. J. Clin. Pract.* 59, 961-968
- Cao YQ, Piedras-Renteria ES, Smith GB, Chen G, Harata NC, Tsien RW (2004) Presynaptic Ca^{2+} channels compete for channel type-preferring slots in altered neurotransmission arising from Ca^{2+} channelopathy. *Neuron* 43: 387-400.
- Cao Y-Q and Tsien RW (2005) Effects of familial hemiplegic migraine type 1 mutations on neuronal P/Q-type Ca^{2+} channel activity and inhibitory synaptic transmission. *PNAS* 102: 2590-2595
- Catterall WA (1998) Structure and function of neuronal Ca^{2+} channels and their role in neurotransmitter release. *Cell Calcium* 24: 307-323.
- Catterall WA (2000) Structure and regulation of voltage-gated Ca^{2+} channels. *Annu Rev Cell Dev Biol* 16:521-555.
- Catterall WA, Perez-Reyes E, Snutch TP et al (2005) International Union of Pharmacology. XLVIII. Nomenclature and structure–function relationships of voltage-gated calcium channels. *Pharmacol Rev* 57:411-425
- Catterall WA and Few AP (2008) Calcium channel regulation and presynaptic plasticity. *Neuron* 59:882-901
- Catterall WA, Kalume F and Oakley JC (2010) $Na_v1.1$ channels and epilepsy. *J. Physiol.* 588: 1849-1859.
- Cestèle S, Scalmani P, Rusconi R, Terragni B, Franceschetti S and Mantegazza M (2008) Self-Limited Hyperexcitability: Functional Effect of a Familial Hemiplegic Migraine Mutation of the $Na_v1.1$ (SCN1A) Na^+ Channel. *J. Neurosci.* 28: 7273 - 7283.
- Chan YC, et al. (2008) Electroencephalographic changes and seizures in familial hemiplegic migraine patients with the CACNA1A gene S218L mutation. *J Clin Neurosci* 15:891-894.
- Chang SY, Yong TF, Yu CY, Liang MC, Pletnikova O, Troncoso J, Burgunder JM, Soong TW (2007) Age and gender-dependent alternative splicing of P/Q-type calcium channel EF-hand. *Neuroscience* 145:1026-1036
- Charles A (2009) Advances in the basic and clinical science of migraine. *Ann Neurol* 65:491-498

Chasman DI, Schurks M, Anttila V, de Vries B, Schminke U, Launer LJ, Terwindt GM, van den Maagdenberg AM, Fendrich K, Volzke H, Ernst F, Griffiths LR, Buring JE, Kallela M, Freilinger T, Kubisch C, Ridker PM, Palotie A, Ferrari MD, Hoffmann W, Zee RY and Kurth T (2011) Genome-wide association study reveals three susceptibility loci for common migraine in the general population. *Nat Genet.* 43(7): 695-698.

Chaudhuri D, Alseikhan BA, Chang SY, Soong TW, Yue DT (2005) Developmental activation of calmodulin-dependent facilitation of cerebellar P-type Ca²⁺ current. *J Neurosci* 25: 8282-8294.

Chaudhuri D, Issa JB, Yue DT (2007) Elementary mechanisms producing facilitation of Cav2.1 (P/Q-type) channels. *J Gen Physiol* 129:385-401.

Cricchi F, Di Lorenzo C, Grieco GS, Rengo C, Cardinale A, Racaniello M, Santorelli FM, Nappi G, Pierelli F, Casali C (2007) Early-onset progressive ataxia associated with the first CACNA1A mutation identified within the I-II loop. *J Neurol Sci* 254:69-71.

Cuenca-Leon E, Corominas R, Fernandez-Castillo N, Volpini V, Del Toro M, Roig M, Macaya A, Cormand B (2008) Genetic analysis of 27 Spanish patients with hemiplegic migraine, basilar-type migraine and childhood periodic syndromes. *Cephalalgia* 28:1039-1047

Cuenca-Leon E, Banchs I, Serra SA, Latorre P, Fernandez-Castillo N, Corominas R, Valverde MA, Volpini V, Fernandez-Fernandez JM, Macaya A, Cormand B (2009) Late-onset episodic ataxia type 2 associated with a novel loss-of-function mutation in the CACNA1A gene. *J Neurol Sci* 280:10-14.

Cutrer FM, Sorensen AG, Weisskoff RM, Ostergaard L, Sanchez del Rio M, Lee EJ, Rosen BR and Moskowitz MA (1998) Perfusion-weighted imaging defects during spontaneous migraineous aura. *Ann. Neurol.* 43, 25-31.

Cuttle MF, Tsujimoto T, Forsythe ID, Takahashi T (1998) Facilitation of the presynaptic calcium current at an auditory synapse in rat brainstem. *J Physiol* 512:723-729.

De Fusco M, Marconi R, Silvestri L, et al. (2003) Haploinsufficiency of ATP1A2 encoding the Na⁺/K⁺ pump α_2 subunit associated with familial hemiplegic migraine type 2. *Nat Genet.* 33:192-196.

Dichgans M, Freilinger T, Eckstein G, et al. (2005) Mutation in the neuronal voltage-gated sodium channel *SCN1A* in familial hemiplegic migraine. *Lancet.* 366: 371-377.

Diener HC, Bussone G, Van Oene JC, et al. (2007) Topiramate reduces headache days in chronic migraine: a randomized, double-blind, placebo-controlled study. *Cephalalgia* 27: 814-823.

Di Guilmi MN, Urbano FJ, González Inchauspe C and Uchitel OD (2011) Pregabalin modulation of neurotransmitter release is mediated by change in intrinsic activation/inactivation properties of Ca_v2.1 calcium channels. *J. Pharmacology and Experimental Therapy* 336(3): 973-982.

Dodge FA and Rahamimoff R (1967) Co-operative action of calcium ions in transmitter release at the neuromuscular junction. *J. Physiol.* 193: 419-432.

Ducros A, Denier C, Joutel A, Cecillon M, Lescoat C, Vahedi K, Darcel F, Vicaut E, Bousser MG and Tournier-Lasserre E. (2001) The clinical spectrum of familial hemiplegic migraine associated with mutations in a neuronal calcium channel. *N. Engl. J. Med.* 345: 17-24.

Dunlap K, Luebke JI and Turner TJ (1995) Exocytotic Ca^{2+} channels in mammalian central neurons. *Trends Neurosci.* 18: 89-98

Eikermann-Haerter K, Baum MJ, Ferrari MD, van den Maagdenberg AM, Moskowitz MA & Ayata C (2009a). Androgenic suppression of spreading depression in familial hemiplegic migraine type 1 mutant mice. *Ann Neurol* 66, 564-568.

Eikermann-Haerter K, Dilekoz E, Kudo C, Savitz SI, Waeber C, Baum MJ, Ferrari MD, van den Maagdenberg AMJM, Moskowitz MA & Ayata C (2009b). Genetic and hormonal factors modulate spreading depression and transient hemiparesis in mouse models of familial hemiplegic migraine type 1. *J Clin Invest* 119, 99-109.

Eikermann-Haerter K, Yuzawa I, Qin T, Wang Y, Baek K, Kim YR, Hoffmann U, Dilekoz E, Waeber C, Ferrari MD, van den Maagdenberg AM, Moskowitz MA and Ayata C (2011) Enhanced subcortical spreading depression in Familial Hemiplegic Migraine Type 1 mutant mice. *The Journal of Neuroscience* 31: 5755-5763.

Eriksen MK, Thomsen LL and Olesen J (2006). Implications of clinical subtypes of migraine with aura. *Headache.* 46:286-297.

Ferrari MD, van den Maagdenberg AMJM, Frants RR and Goadsby PJ (2008) *Molecular Neurology. Migraine as a cerebral ionopathy with impaired central sensory processing.* Waxman, S.G. (editor): 439-461.

Ferrari MD, Roon K I, Lipton R B & Goadsby P J (2001) Oral triptans (serotonin, 5-HT_{1B/1D} agonists) in acute migraine treatment: a metaanalysis of 53 trials. *Lancet* 358, 1668-1675

Flippen C and Welch KM (1997) Imaging the brain of migraine sufferers. *Curr Opin Neurol* 10(3): 226-30.

Forsythe ID, Tsujimoto T, Barnes-Davies M, Cuttle MF, and Takahashi T (1998) Inactivation of presynaptic calcium current contributes to synaptic depression at a fast central synapse. *Neuron* 20(4): 797-807.

Giugovaz-Tropper B, Gonzalez-Inchauspe C, Di Guilmi MN, Urbano FJ, Forsythe ID and Uchitel OD (2011) P/Q-type calcium channel ablation in a mice glycinergic synapse mediated by multiple types of Ca^{2+} channels alters transmitter release and short term plasticity. *Neuroscience*

Goadsby PJ, Lipton RB and Ferrari MD (2002) Migraine- current understanding and treatment. *N. Engl. J. Med.* 346, 257-270.

Goadsby PJ (2007) Recent advances in understanding migraine mechanisms, molecules and therapeutics. *Trends Mol Med* 13: 39- 44.

Goadsby PJ, Charbit AR, Andreou AP, Akerman S, Holland PR (2009a) Neurobiology of migraine. *Neuroscience* 161:327-341.

Goadsby PJ, Ferrari MD, Csanyi A et al (2009b) Randomized, double-blind, placebo-controlled, proof-of-concept study of the cortical spreading depression inhibiting agent tonabersat in migraine prophylaxis. *Cephalalgia.* 29: 742-750.

González Inchauspe C, Martini FJ, Forsythe ID, Uchitel OD (2004) Functional compensation of P/Q by N-type channels blocks short term plasticity at the calyx of held presynaptic terminal. *J Neurosci* 24:10379-10383.

González Inchauspe C, Forsythe ID and Uchitel OD (2007). Changes in synaptic transmission properties due to the expression of N-type calcium channels at the calyx of Held synapse of mice lacking P/Q-type calcium channels. *Journal of Physiology* 584, 835 - 851

González Inchauspe C, Urbano FJ, Di Guilmi MN, Forsythe ID, Ferrari MD, van den Maagdenberg AM, Uchitel OD (2010) Gain of function in FHM-1 Ca(V)2.1 knock-in mice is related to the shape of the action potential. *J Neurophysiol.* 104(1): 291-299.

Haan J, Kors EE, Vanmolkot KR, van den Maagdenberg AM, Frants RR. and Ferrari MD (2005). Migraine genetics: an update. *Curr Pain Headache Rep* 9, 213-220.

Hadjikhani N, Sanchez Del Rio M, Wu O, Schwartz D, Bakker D, Fischl B, Kwong KK, Cutrer FM, Rosen BR, Tootell RB et al. (2001). Mechanisms of migraine aura revealed by functional MRI in human visual cortex. *Proc. Natl. Acad. Sci. USA* 98, 4687- 4692.

Haerter K, Ayata C and Moskowitz MA (2005) Cortical spreading depression: A model for understanding migraine biology and future drug targets. *Headache Currents* 2, 97-103.

Hans M, Luvisetto S, Williams ME, Spagnolo M, Urrutia A, Tottene A, Brust PF, Johnson EC, Harpold MM, Stauderman KA et al. (1999) Functional consequences of mutations in the human α_1A calcium channel subunit linked to familial hemiplegic migraine. *J Neurosci* 19: 1610-1619.

Hennig MH, Postlethwaite M, Forsythe ID and Graham BP (2008) Interactions between multiple sources of short-term plasticity during evoked and spontaneous activity at the rat calyx of Held. *J. Physiol.* 586: 3129 - 3146.

Ho TW et al.(2008) Randomized controlled trial of an oral CGRP antagonist, MK-0974, in acute treatment of migraine. *Neurology* 70: 1004-1012.

Ho TW, Ferrari MD, Dodick DW, et al.(2009) Efficacy and tolerability of MK-0974 (telcagepant), a new oral antagonist of calcitonin gene-related peptide receptor, compared with zolmitriptan for acute migraine: a randomised, placebo-controlled, parallel-treatment trial. *Lancet.* 372: 2115-2123.

Ho TW, Edvinsson L and Goadsby PJ (2010) CGRP and its receptors provide new insights into migraine pathophysiology. *Nat Rev Neurol.* 6(10): 573-82.

Imbrici P, Jaffe SL, Eunson LH, Davies NP, Herd C, Robertson R, Kullmann DM, Hanna MG (2004) Dysfunction of the brain calcium channel $Ca_v2.1$ in absence epilepsy and episodic ataxia. *Brain* 127: 2682-2692.

Ishikawa K, Fujigasaki H, Saegusa H, Ohwada K, et al (1999) Abundant expression and cytoplasmic aggregations of α_1A voltage-dependent calcium channel protein associated with neurodegeneration in spinocerebellar ataxia type 6. *Hum Mol Genet* 8:1185-1193

Iwasaki S, Momiyama A, Uchitel OD and Takahashi T (2000) Developmental changes in calcium channel types mediating central synaptic transmission. *J Neurosci* 20:59-65

Jen J, Kim GW and Baloh RW (2004). Clinical spectrum of episodic ataxia type 2. *Neurology* 62: 17-22.

Jen JC, Graves TD, Hess EJ, Hanna MG, Griggs RC & Baloh RW (2007) Primary episodic ataxias: diagnosis, pathogenesis and treatment. *Brain* 130: 2484-2493.

Jeng CJ, Sun MC, Chen YW & Tang CY (2008) Dominant-negative effects of episodic ataxia type 2 mutations involve disruption of membrane trafficking of human P/Q-type Ca²⁺ channels. *J Cell Physiol* 214: 422-433.

Jones SW (2002) Calcium channels: When is a subunit not a subunit? *J Physiol* 545:334.

Jouveneau A., Eunson LH, Spauschus A, Ramesh V, Zuberi SM, Kullmann DM and Hanna MG (2001) Human epilepsy associated with dysfunction of the brain P/Q-type calcium channel. *Lancet* 358: 801-807.

Kahlig KM, Rhodes TH, Pusch M, Freilinger T, Pereira-Monteiro JM, Ferrari MD, van den Maagdenberg AMJM, Dichgans M and George ALG Jr (2008) Divergent sodium channel defects in familial hemiplegic migraine. *PNAS* 105: 9799 - 9804

Kaja S, van de Ven RC, Broos LA, Veldman H, van Dijk JG, Verschuuren JJ, Frants RR, Ferrari MD, van den Maagdenberg AM and Plomp JJ (2005) Gene dosage-dependent transmitter release changes at neuromuscular synapses of *CACNA1A* R192Q knockin mice are non-progressive and do not lead to morphological changes or muscle weakness. *Neuroscience* 135: 81-95.

Kaja S, Van de Ven RCG, Broos LAM, Frants RR, Ferrari MD, Van den Maagdenberg AMJM and Plomp JJ (2010) Severe and Progressive Neurotransmitter Release Aberrations in Familial Hemiplegic Migraine Type 1 *Cacna1a* S218L Knock-in Mice. *J Neurophysiol* 104: 1445-1455.

Kanumilli S, Tringham EW, Payne CE, Dupere JR, Venkateswarlu K, Usowicz MM (2006) Alternative splicing generates a smaller assortment of CaV2.1 transcripts in cerebellar Purkinje cells than in the cerebellum. *Physiol Genomics* 24:86-96.

Katz E, Ferro PA, Weisz G and Uchitel OD (1996) Calcium channels involved in synaptic transmission at the mature and regenerating mouse neuromuscular junction. *J. Physiol.* 497: 687 - 697.

Koenderink JB, Zifarelli G, Qiu LY, Schwarz W, De Pont JJ, et al. (2005) Na,KATPase mutations in familial hemiplegic migraine lead to functional inactivation. *Biochim Biophys Acta* 1669: 61-68.

Koester HJ, Sakmann B (2000) Calcium dynamics associated with action potentials in single nerve terminals of pyramidal cells in layer 2/3 of the young rat neocortex. *J Physiol* 529(Pt 3):625-646

Kors EE, Terwindt GM, Vermeulen FL, Fitzsimons RB, Jardine PE, Heywood P, Love S, van den Maagdenberg AM, Haan J, Frants RR, Ferrari MD (2001) Delayed cerebral edema and fatal coma after minor head trauma: Role of *CACNA1A* calcium channel subunit gene and relationship with familial hemiplegic migraine. *Annals of Neurology* 49: 753-760.

Kors EE, van den Maagdenberg AM, Plomp JJ, Frants RR and Ferrari MD (2002). *Curr. Opin. Neurol.* 15: 311-316.

Kors EE, Melberg A, Vanmolkot KR, Kumlien E, Haan J, Raininko R, Flink R, Ginjaar HB, Frants RR, Ferrari MD and van den Maagdenberg AM (2004) *Neurology* 63: 1136-1137.

Kraus RL, Sinnegger MJ, Glossmann H, Hering S and Striessnig J (1998) Familial hemiplegic migraine mutations change $\alpha 1A$ Ca^{2+} channel kinetics. *J Biol Chem* 273: 5586-5590.

Kraus RL, Sinnegger M, Koschak A, Glossmann H, Stenirri S, Carrera P and Striessnig J (2000) *J. Biol. Chem.* 275, 9239-9243

Lafreniere RG, Cader MZ, Poulin JF, Andres-Enguix I, Simoneau M, Gupta N, Boisvert K, Lafreniere F, McLaughlan S, Dube MP, Marcinkiewicz MM, Ramagopalan S, Ansorge O, Brais B, Sequeiros J, Pereira-Monteiro JM, Griffiths LR, Tucker SJ, Ebers G and Rouleau GA (2010) A dominant-negative mutation in the TRESK potassium channel is linked to familial migraine with aura. *Nat Med* 16(10): 1157-60.

Lauritzen M (1994) Pathophysiology of the migraine aura. The spreading depression theory. *Brain* 117, 199-210.

Leo L, Gherardini L, Barone V, De Fusco M, Pietrobon D, Pizzorusso T and Casari G (2011) Increased susceptibility to cortical spreading depression in the mouse model of familial hemiplegic migraine type 2. *PLoS Genet.* 7(6): e1002129

Levy D, Strassman AM, Burstein R (2009) A critical view on the role of migraine triggers in the genesis of migraine pain. *Headache* 49:953-957

Li L, Bischofberger J, Jonas P (2007) Differential gating and recruitment of P/Q-, N-, and R-type Ca^{2+} channels in hippocampal mossy fiber boutons. *J Neurosci* 27:13420-13429.

Liu Z, Ren J and Murphy TH (2003) Decoding of synaptic voltage waveforms by specific classes of recombinant high-threshold Ca^{2+} channels. *J. Physiol.* 553: 473-488.

Llinás R (1988) The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science* 242: 1654-1664.

Luisetto S, Fellin T, Spagnolo M, Hivert B, Brust PF, Harpold MM, Stauderman KA, Williams ME, Pietrobon D (2004) Modal gating of human $CaV2.1$ (P/Q-type) calcium channels: I. The slow and the fast gating modes and their modulation by beta subunits. *J Gen Physiol* 124:445-461

Marconi R, De Fusco M, Aridon P, Plewnia K, Rossi M, Carapelli S et al. (2003) Familial hemiplegic migraine type 2 is linked to 0.9Mb region on chromosome 1q23. *Ann Neurol* 53: 376-381.

Marmura MJ and Silberstein SD (2011) Current Understanding and Treatment of Headache Disorders: Five New Things. *Neurology* 76: S31 - S36.

Matsushita K, Wakamori M, Rhyu IJ, Arii T, Oda S, Mori Y, Imoto K (2002) Bidirectional alterations in cerebellar synaptic transmission of tottering and rolling Ca^{2+} channel mutant mice. *J Neurosci* 22: 4388-4398.

Meinrenken CJ, Borst JGG and Sakmann B (2002) Calcium Secretion Coupling at Calyx of Held Governed by Nonuniform Channel-Vesicle Topography. *J. Neurosci.* 22: 1648-1667.

Meinrenken CJ, Borst JG, Sakmann B (2003) Local routes revisited: the space and time dependence of the Ca^{2+} signal for phasic transmitter release at the rat calyx of Held. *J Physiol* 547: 665-689.

- Meisler MH and Kearney JA. (2005). Sodium channel mutations in epilepsy and other neurological disorders. *J Clin Invest* 115: 2010-2017.
- Melliti K, Grabner M, Seabrook GR (2003) The familial hemiplegic migraine mutation R192Q reduces G-protein-mediated inhibition of P/Q-type (Ca_v2.1) calcium channels expressed in human embryonic kidney cells. *J Physiol* 546:337-347
- Mermelstein PG, Foehring RC, Tkatch T, Song WJ, Baranauskas G, Surmeier DJ (1999) Properties of Q-type calcium channels in neostriatal and cortical neurons are correlated with beta subunit expression. *J Neurosci* 19:7268-7277
- Mintz IM, Adams ME and Bean BP (1992) P-type calcium channels in rat central and peripheral neurons. *Neuron* 9: 85-95
- Mintz IM, Venema VJ, Swiderek KM, Lee TD, Bean BP, Adams ME (1992) P-type calcium channels blocked by the spider toxin omega-Aga-IVA. *Nature* 355: 827-829
- Mintz IM, Sabatini BL, Regehr WG (1995) Calcium control of transmitter release at a cerebellar synapse. *Neuron* 15: 675-688.
- Mochida S, Few AP, Scheuer T, Catterall WA (2008) Regulation of presynaptic Ca_v2.1 channels by Ca²⁺ sensor proteins mediates short-term synaptic plasticity. *Neuron* 57: 210-216.
- Moskowitz MA, Bolay H, Dalkara (2004) T. Deciphering migraine mechanisms: clues from familial hemiplegic migraine genotypes. *Ann Neurol*. 55: 276-280.
- Mulder EJ, Van Baal C, Gaist D, Kallela M, Kaprio J, Svensson DA, Nyholt DR, Martin NG, MacGregor AJ, Cherkas LF, Boomsma DI and Palotie A (2003). Genetic and environmental influences on migraine: a twin study across six countries. *Twin Res*. 6(5): 422-31.
- Mullner C, Broos LA, van den Maagdenberg AM, Striessnig J (2004) Familial hemiplegic migraine type 1 mutations K1336E, W1684R, and V1696I alter Ca_v2.1 Ca²⁺ channel gating: evidence for beta-subunit isoform-specific effects. *J Biol Chem* 279: 51844-51850.
- Muller M, Felmy F and Schneggenburger R (2008) A limited contribution of Ca²⁺ current facilitation to paired-pulse facilitation of transmitter release at the rat calyx of Held. *J Physiol (Lond)* 586: 5503-5520
- Newcomb R, Szoke B, Palma A, Wang G, Chen X, Hopkins W, Cong R, Miller J, Urge L, Tarczy-Hornoch K, Loo JA, Dooley DJ, Nadasdi L, Tsien RW, Lemos J, Miljanich G (1998) Selective peptide antagonist of the class E calcium channel from the venom of the tarantula *Hysterocrates gigas*. *Biochemistry* 37: 15353-15362.
- Nudler S, Piriz J, Urbano FJ, Rosato-Siri MD, Renteria ES and Uchitel OD (2003) Ca²⁺ channels and synaptic transmission at the adult, neonatal, and P/Q-type deficient neuromuscular junction. *Ann N Y Acad Sci*. 998: 11-17.
- Ogiwara I, Miyamoto H, Morita N, Atapour N, Mazaki E, Inoue I, Takeuchi T, Itohara S, Yanagawa Y, Obata K, Furuichi T, Hensch TK, Yamakawa K (2007) Na(v) 1.1 localizes to axons of parvalbumin-positive inhibitory interneurons: a circuit basis for epileptic seizures in mice carrying an Scn1a gene mutation. *J Neurosci* 27: 5903-5914.

Olesen J. *et al* (2004). Calcitonin gene-related peptide receptor antagonist BIBN 4096 BS for the acute treatment of migraine. *N. Engl. J. Med.* 350: 1104-1110.

Ophoff RA, Terwindt GM, Vergouwe MN, van Eijk R, Oefner PJ, Hoffman SMG, Lamerdin JE, Mohrenweiser HW, Bulman DE, Ferrari M *et al.* (1996) Familial hemiplegic migraine and episodic ataxia type-2 are caused by mutations in the Ca²⁺ channel gene CACNL1A4. *Cell* 87: 543-552.

Perez-Reyes E (2003) Molecular physiology of low-voltage-activated t-type calcium channels. *Physiol Rev* 83(1): 117-161.

Pietrobon D and Striessnig J (2003) Neurobiology of migraine. *Nat Rev Neurosci* 4: 386-398.

Pietrobon D (2005) Function and dysfunction of synaptic calcium channels: insights from mouse models. *Curr Opin Neurobiol* 15:257-265

Pietrobon D (2005). Migraine: new molecular mechanisms. *Neuroscientist* 11, 373-386.

Pietrobon D (2007) Familial hemiplegic migraine. *Neurotherapeutics* 4:274-284.

Pietrobon D (2010) Insights into migraine mechanisms and Cav2.1 calcium channel function from mouse models of familial hemiplegic migraine. *J Physiol* 588: pp 1871-1878.

Pineda JC, Waters RS and Foehring RC (1998) Specificity in the interaction of HVA Ca²⁺ channel types with Ca²⁺ dependent AHPs and firing behavior in neocortical pyramidal neurons. *J. Neurophysiol.* 79, 2522-2534.

Plummer MR, Logothetis DE and Hess P (1989) Elementary properties and pharmacological sensitivities of calcium channels in mammalian peripheral neurons. *Neuron* 2: 1453-1463.

Qian J, Noebels JL (2001) Presynaptic Ca²⁺ channels and neurotransmitter release at the terminal of a mouse cortical neuron. *J Neurosci* 21:3721-3728

Ragsdale DS (2008) How do mutant Nav1.1 sodium channels cause epilepsy? *Brain Res Rev.* 58(1): 149-59.

Randall A, Tsien RW (1995) Pharmacological dissection of multiple types of Ca²⁺ channel currents in rat cerebellar granule neurons. *J Neurosci* 15:2995-3012

Restituito S, Thompson RM, Eliet J, Raïke RS, Riedl M, Charnet P, Gomez CM (2000) The polyglutamine expansion in spinocerebellar ataxia type 6 causes a β subunit-specific enhanced activation of P/Q-type calcium channels in *Xenopus* oocytes. *J Neurosci* 20:6394-6403

Rhodes TH, *et al.* (2005) Sodium channel dysfunction in intractable childhood epilepsy with generalized tonic-clonic seizures. *J Physiol* 569:433-445.

Rozov A, Burnashev N, Sakmann B, Neher E (2001) Transmitter release modulation by intracellular Ca²⁺ buffers in facilitating and depressing nerve terminals of pyramidal cells in layer 2/3 of the rat neocortex indicates a target cell-specific difference in presynaptic calcium dynamics. *J Physiol* 531:807-826

Sabatini BL and Regehr WG (1999) Timing of synaptic transmission. *Annu Rev Physiol.* 61: 521-42.

Schneggenburger R and Forsythe ID (2006). The calyx of Held. *Cell Tissue Res.* 326: 311-337.

Schneggenburger R and Neher E (2000) Intracellular calcium dependence of transmitter release rates at a fast central synapse. *Nature* 406: 889-893.

Segall L, Mezzetti A, Scanzano R, Gargus JJ, Purisima E and Blostein R (2005) Alterations in the 2 isoform of Na,K-ATPase associated with familial hemiplegic migraine type 2. *Proc Natl Acad Sci USA.* 102: 11106-11111.

Serra SA, Fernandez-Castillo N, Macaya A, Cormand B, Valverde MA, Fernandez-Fernandez JM (2009) The hemiplegic migraine-associated Y1245C mutation in CACNA1A results in a gain of channel function due to its effect on the voltage sensor and Gprotein-mediated inhibition. *Pflugers Arch* 458:489-502.

Silberstein SD, Lipton RB, Dodick DW, et al. (2007) Efficacy and safety of topiramate for the treatment of chronic migraine: a randomized, double-blind, placebo-controlled trial. *Headache* 47: 170-180.

Smith M.I. et al. (2000) Repetitive cortical spreading depression in a gyrencephalic feline brain: inhibition by the novel benzoylaminobenzopyran SB-220453. *Cephalalgia* 20, 546-553

Smits JF et al. (1980) Is the antihypertensive effect of propranolol caused by an action within the central nervous system? *J. Pharmacol. Exp. Ther.* 215, 221-225

Somjen GG (2001). Mechanisms of spreading depression and hypoxic spreading depression-like depolarization. *Physiol Rev* 81, 1065-1096.

Soong TW, DeMaria CD, Alvania RS, Zweifel LS, Liang MC, Mittman S, Agnew WS, Yue DT (2002) Systematic identification of splice variants in human P/Q-type channel $\alpha 1(2.1)$ subunits: implications for current density and Ca^{2+} -dependent inactivation. *J Neurosci* 22:10142-10152

Storer RJ and Goadsby PJ (2004) Topiramate inhibits trigeminovascular neurons in the cat. *Cephalalgia* 24, 1049-1056.

Storer RJ and Goadsby PJ (2005) Topiramate has a locus of action outside of the trigeminocervical complex. *Neurology* 64 (Suppl 1), A150-A151.

Stotz SC, Jarvis SE, Zamponi GW (2004) Functional roles of cytoplasmic loops and pore lining transmembrane helices in the voltage-dependent inactivation of HVA calcium channels. *J Physiol* 554:263-273.

Striessnig J, Grabner M, Mitterdorfer J, Hering S, Sinnegger MJ, Glossmann H (1998) Structural basis of drug binding to L Ca^{2+} channels. *Trends Pharmacol Sci* 19: 108-115.

Suzuki M, Van Paesschen W, Stalmans I, Horita S, et al. (2010) Defective membrane expression of the Na^{+} - HCO_3^{-} cotransporter NBCe1 is associated with familial migraine. *Proc Natl Acad Sci U S A.* 107(36): 15963-8.

Takago H, Nakamura Y & Takahashi T (2005). G protein-dependent presynaptic inhibition mediated by AMPA receptors at the calyx of Held. *Proc Natl Acad Sci U S A* 102, 7368-7373.

Takahashi T, Forsythe I, Tsujimoto T, Barnes-Davies M and Onodera K (1996). Presynaptic calcium current modulation by a metabotropic glutamate receptor. *Science* 274, 594-597.

Tavraz NN, Friedrich T, Durr KL, Koenderink JB, Bamberg E, et al. (2008) Diverse functional consequences of mutations in the Na⁺/K⁺-ATPase alpha2-subunit causing familial hemiplegic migraine type 2. *J Biol Chem* 283: 31097-31106.

Thomsen LL, Eriksen MK, Roemer SF, Andersen I, Olesen J and Russell MB (2002). A population-based study of familial hemiplegic migraine suggests revised diagnostic criteria. *Brain* 125: 1379-1391.

Thomsen LL, Kirchmann M, Bjornsson A, Stefansson H, Jensen RM, Fasquel AC, Petursson H, Stefansson M, Frigge ML, Kong A, Gulcher J, Stefansson K, Olesen J (2007) The genetic spectrum of a population-based sample of familial hemiplegic migraine. *Brain* 130:346-356

Tottene A, Moretti A, Pietrobon D (1996) Functional diversity of P-type and R-type calcium channels in rat cerebellar neurons. *J Neurosci* 16:6353-6363

Tottene A, Volsen S, Pietrobon D (2000) Alpha(1E) subunits form the pore of three cerebellar R-type calcium channels with different pharmacological and permeation properties. *J Neurosci* 20: 171-178.

Tottene A, Fellin T, Pagnutti S, Luvisetto S, Striessnig J, Fletcher C, Pietrobon D (2002) Familial hemiplegic migraine mutations increase Ca²⁺ influx through single human CaV2.1 channels and decrease maximal Cav2.1 current density in neurons. *Proc Natl Acad Sci USA* 99: 13284-13289.

Tottene A, Pivotto F, Fellin T, Cesetti T, van den Maagdenberg AMJM, Pietrobon D (2005) Specific kinetic alterations of human CaV2.1 calcium channels produced by mutation S218L causing familial hemiplegic migraine and delayed cerebral edema and coma after minor head trauma. *J Biol Chem* 280: 17678-17686.

Tottene A, Conti R, Fabbro A, Vecchia D, Shapovalova M, Santello M, van den Maagdenberg AM, Ferrari MD and Pietrobon D (2009) Enhanced excitatory transmission at cortical synapses as the basis for facilitated spreading depression in Ca_v2.1 knock in migraine mice. *Neuron* 61: 762-773.

Trimmer JS, Rhodes KJ (2004) Localization of voltage-gated ion channels in mammalian brain. *Annu Rev Physiol* 66:477-519

Tringham EW, Payne CE, Dupere JR, Usowicz MM (2007) Maturation of rat cerebellar Purkinje cells reveals an atypical Ca²⁺ channel current that is inhibited by omega-agatoxin IVA and the dihydropyridine (-)-(S)-Bay K8644. *J Physiol* 578:693-714

Tsou WL, Soong BW, Paulson HL and Rodriguez-Lebron E (2011) Splice isoform-specific suppression of the Ca(V)2.1 variant underlying spinocerebellar ataxia type 6. *Neurobiol Dis.* 43: 533-542,

Tsujimoto T, Jeromin A, Saitoh N, Roder JC, Takahashi T (2002) Neuronal calcium sensor 1 and activity-dependent facilitation of P/Q-type calcium currents at presynaptic nerve terminals. *Science* 295:2276-2279.

Uchitel OD, Di Guilmi MN, Urbano FJ, Gonzalez-Inchauspe C (2010) Acute modulation of calcium currents and synaptic transmission by gabapentinoids. *Channels (Austin)* 4(6): 490-496.

Urbano FJ, Piedras-Rentería ES, Jun K, Shin HS, Uchitel OD and Tsien RW (2003). Altered properties of quantal neurotransmitter release at endplates of mice lacking P/Q-type Ca^{2+} channels. *Proc. Natl. Acad. Sci.* 100, 3491-3496.

Urbano FJ, Bisagno V, Wikinski SI, Uchitel OD, Llinás RR (2009) Cocaine Acute “Binge” Administration Results in Altered Thalamocortical Interactions in Mice. *Biol Psychiatry* 66: 769-776.

Usowicz MM, Sugimori M, Cherskey B, Llinas R (1992) P-type calcium channels in the somata and dendrites of adult cerebellar Purkinje cells. *Neuron* 9:1185-1199

van den Maagdenberg AM, Pietrobon D, Pizzorusso T, Kaja S, Broos LA, Cesetti T, van de Ven RC, Tottene A, van der Kaa J, Plomp JJ, Frants RR and Ferrari MD (2004) A *Ca_v1a* knock-in migraine mouse model with increased susceptibility to cortical spreading depression. *Neuron* 41: 701-710.

van den Maagdenberg AM, Pizzorusso T, Kaja S, Terpolilli N, Shapovalova M, Hoebeek FE, Barrett CF, Gherardini L, van de Ven RC, Todorov B, Broos LA, Tottene A, Gao Z, Fodor M, De Zeeuw CI, Frants RR, Plesnila N, Plomp JJ, Pietrobon D & Ferrari MD (2010). High cortical spreading depression susceptibility and migraine-associated symptoms in *Cav2.1 S218L* mice. *Ann Neurol* 67, 85-98.

van den Maagdenberg AM, Haan J, Terwindt GM and Ferrari MD (2007) Migraine: gene mutations and functional consequences. *Curr. Opin. Neurol.* 20:299-305

Vanmolkot KRJ, Kors EE, Hottenga JJ, Terwindt GM, Haan J, Hoefnagels WAJ et al. (2003). Novel mutations in the Na^{+},K^{+} -ATPase pump gene *ATP1A2* associated with Familial Hemiplegic Migraine and Benign Familial Infantile Convulsions. *Annals of Neurology* 54: 360-366.

Vanmolkot KR, Kors EE, Turk U, et al (2006) Two de novo mutations in the Na,K -ATPase gene *ATP1A2* associated with pure familial hemiplegic migraine. *Eur J Hum Genet* 14: 555-560.

Vanmolkot KR, et al. (2007) The novel p.L1649Q mutation in the *SCN1A* epilepsy gene is associated with familial hemiplegic migraine: Genetic and functional studies. *Hum Mut* 28:522.

Volsen SG, Day NC, McCormack AL, Smith W, Craig PJ, Beattie R, Ince PG, Shaw PJ, Ellis SB and Gillespie A (1995) The expression of neuronal voltage-dependent calcium channels in human cerebellum. *Brain Res Mol Brain Res* 34: 271-82.

Von Gersdorff H and Borst JG (2002). Short-term plasticity at the calyx of Held. *Neuroscience* 3, 53-64.

Von Gersdorff H, Schneggenburger R, Weis S and Neher E (1997). Presynaptic Depression at a Calyx Synapse: The Small Contribution of Metabotropic Glutamate Receptors. *The Journal of Neuroscience* 17 (21): 8137-8146.

Wang L-Y and Kaczmarek LK (1998). High-frequency firing helps replenish the readily releasable pool of synaptic vesicles. *Nature* 394, 384-388

Weiss N, Sandoval A, Felix R, Van den Maagdenberg A, De Waard M (2008) The S218L familial hemiplegic migraine mutation promotes de/inhibition of $Ca(v)2.1$ calcium channels during direct G-protein regulation. *Pflugers Arch* 457:315-326.

Welch M (1998) Current opinions in headache pathogenesis: introduction and synthesis. *Curr Opin Neurol* 11: 193-197.

Welch M (1998) The occipital cortex as a generator of migraine aura. *Cephalalgia Suppl* 22: 15-21.

Westenbroek RE, Sakurai T, Elliott EM, Hell JW, Starr TVB, Snutch TP, Catterall WA (1995) Immunochemical identification and subcellular distribution of the $\alpha 1A$ subunits of brain calcium channels. *J Neurosci* 15: 6403-6418.

Williams TM *et al.* (2006) Non-peptide calcitonin related peptide receptor antagonists from a benzodiazepinone lead. *Bioorg. Med. Chem. Lett.* 16: 2595-2598.

Wong AYC, Graham BP, Billups B and Forsythe ID (2003) Distinguishing between presynaptic and postsynaptic mechanisms of short-term depression during action potential trains. *J. Neurosci.* 23, 4868-4877.

Wu LG, Westenbroek RE, Borst JGG, Catterall WA, Sakmann B (1999) Calcium channel types with distinct presynaptic localization couple differentially to transmitter release in single calyx-type synapses. *J Neurosci* 19: 726-736.

Xu W and Lipscombe D (2001) Neuronal CaV1 D.3. $\alpha(1)$ L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci* 21: 5944-5951.

Xu J and Wu LG (2005) The decrease in the presynaptic calcium current is a major cause of short-term depression at a calyx-type synapse. *Neuron* 46:633-645.

Yu FH, Mantegazza M, Westenbroek RE, Robbins CA, Kalume F, Burton KA, Spain WJ, McKnight GS, Scheuer T, Catterall WA (2006) Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic epilepsy in infancy. *Nat Neurosci* 9:1142-1149.

Yue Q, Jen JC, Nelson SF and Baloh RW (1997) Progressive ataxia due to a missense mutation in a calcium-channel gene. *Am J Hum Genet* 61: 1078-1087.

Zafeiriou DI, Lehmann-Horn F, Vargiami E, Teflioudi E, Ververi A, Jurkat-Rott K (2009) Episodic ataxia type 2 showing ictal hyperhidrosis with hypothermia and interictal chronic diarrhea due to a novel CACNA1A mutation. *Eur J Paediatr Neurol* 13:191-193.

Zaitsev AV, Povysheva NV, Lewis DA, Krimer LS (2007) P/Qtype, but not N-type, calcium channels mediate GABA release from fast-spiking interneurons to pyramidal cells in rat prefrontal cortex. *J Neurophysiol* 97:3567-3573

Zhang XC, Levy D, Nosedá R, Kainz V, Jakubowski M and Burstein R (2010) Activation of Meningeal Nociceptors by Cortical Spreading Depression: Implications for Migraine with Aura. *The Journal of Neuroscience* 30(26): 8807- 8814.

Zhang X, Levy D, Kainz V, Nosedá R, Jakubowski M & Burstein R (2011). Activation of central trigeminovascular neurons by cortical spreading depression. *Ann Neurol* 69: 855-865.

Zhuchenko O, Bailey J, Bonnen P, Ashizawa T, Stockton DW, Amos C, Dobyns WB, Subramony SH, Zoghbi HY, Lee C (1997) Autosomal dominant cerebellar ataxia (SCA6) associated with small polyglutamine expansions in the α_1A -voltage-dependent calcium channel. *Nat Genet* 15:62-69.

Zucker RS and Regehr WG (2002) Short-term synaptic plasticity. *Ann. Rev. Physiol.* 64, 355-405.

ACCEPTED MANUSCRIPT

Highlights

Mutations in Cav2.1 Ca^{2+} channels cause familial hemiplegic migraine type 1 (FHM1).

$\text{Ca}_v2.1$ channels with the human FHM1 mutations activate at more hyperpolarizing potentials.

FHM1 generates a gain-of-function in synaptic transmission, affecting also synaptic plasticity.

FHM1 increases the susceptibility for cortical spreading depression (CSD).

ACCEPTED MANUSCRIPT