

Report on LAA-D-12-00856:  
Further refinements of the Heinz inequality  
by M. S. Moslehian, R. Kaur, M. Singh and C. Conde

**Overview** In this paper, the authors give several refinement inequalities of the Heinz inequality:

$$2\|A^{1/2}XB^{1/2}\| \leq \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \leq \|AX + XB\|.$$

In section 2, they recall the Hermite-Hadamard inequality:

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}$$

for a convex function  $f$ , and its refinements in Theorems 2.1, 2.3, 2.4. In section 3, they apply the above results to the convex function

$$F(\nu) := \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\|$$

for  $\nu \in [0, 1]$  and have improvement of Kittaneh's inequalities in Theorem 3.2 and other refinements. In section 4, they obtain refinements of the Heinz inequality for matrices. In Theorem 4.1, inequality (4.2) with two parameters is proved by the standard argument: checking the positive semidefiniteness of the relevant matrices  $Y$  and  $Z$ . By Theorem 4.1 they give Corollaries 4.2, 4.3 as refinements of the Heinz inequality. They also give a new estimation (4.4) in Theorem 4.4 which is of interest and implies Corollaries 4.5 and 4.8. In 4.5, 4.6, 4.8, used is the observation that  $t\alpha + s\beta \leq (t-1)\alpha + (s+1)\beta$  when  $\alpha \leq \beta$ , which is not interesting to the referee.


**Conclusion** I think that all argument are clear and that the proof of (4.4) is interesting so that I would like to recommend its publication in LAA.

**Comments**

Page 9, line 4: remove 'the matrix'.

Page 9, line-9: remove 'matrix'; assume that  $A$  is . . . .

**AUTHOR QUERY FORM**

	<b>Journal: LAA</b>  <b>Article Number: 12062</b>	<b>Please e-mail or fax your responses and any corrections to:</b> <b>E-mail: <a href="mailto:corrections.esch@elsevier.sps.co.in">corrections.esch@elsevier.sps.co.in</a></b> <b>Fax: +31 2048 52799</b>
---	---	---

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

<b>Location in article</b>	<b>Query / Remark: click on the Q link to go</b> <b>Please insert your reply or correction at the corresponding line in the proof</b>
<p><a href="#">Q1</a></p> <p><a href="#">Q2</a></p> <p><a href="#">Q3</a></p> <p><a href="#">Q4</a></p>	<p>Please check the e-mail address for the corresponding author that has been added here, and correct if necessary.</p> <p>Please check whether the designated corresponding author and tel./fax numbers are correct, and amend if necessary.</p> <p>Please note that Refs. [5,22] are listed but not cited in the text. Kindly check.</p> <p>Please update Ref. [20], if available.</p> <div data-bbox="257 1488 824 1590" style="border: 1px solid black; padding: 10px; margin-top: 20px;"> <p style="color: red;">Please check this box if you have no corrections to make to the PDF file</p> <input data-bbox="710 1506 782 1567" type="checkbox"/> </div>

Thank you for your assistance.



ELSEVIER

Contents lists available at SciVerse ScienceDirect

# Linear Algebra and its Applications

journal homepage: [www.elsevier.com/locate/laa](http://www.elsevier.com/locate/laa)



## Further refinements of the Heinz inequality

Rupinderjit Kaur<sup>a</sup>, Mohammad Sal Moslehian<sup>b,\*</sup>, Mandeep Singh<sup>a</sup>,  
Cristian Conde<sup>c</sup>

<sup>a</sup> Department of Mathematics, Sant Longowal Institute of Engineering and Technology, Longowal 148106, Punjab, India

<sup>b</sup> Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures (CEAAS), Ferdowsi University of Mashhad, P.O. Box 1159, Mashhad 91775, Iran

<sup>c</sup> Instituto de Ciencias, Universidad Nacional de Gral. Sarmiento, J. M. Gutierrez 1150, (B1613GSX) Los Polvorines and Instituto Argentino de Matemática “Alberto P. Calderón”, Saavedra 15 3 piso, (C1083ACA) Buenos Aires, Argentina

### ARTICLE INFO

#### Article history:

Received 3 September 2012

Accepted 8 January 2013

Available online xxxx

Q1 Submitted by Hans Schneider

Dedicated in the memory of Professors  
Michael Neumann and Uriel Rothblum.

#### AMS classification:

15A60

47A30

47A64

47B15

#### Keywords:

Heinz inequality

Convex function

Hermite–Hadamard inequality

Positive definite matrix

Unitarily invariant norm

### ABSTRACT

The celebrated Heinz inequality asserts that  $2\| |A|^{1/2}XB^{1/2} \| \leq \| |A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu \|$  for  $X \in \mathbb{B}(\mathcal{H})$ ,  $A, B \in \mathbb{B}(\mathcal{H})_+$ , every unitarily invariant norm  $\| \cdot \|$  and  $\nu \in [0, 1]$ . In this paper, we present several improvement of the Heinz inequality by using the convexity of the function  $F(\nu) = \| |A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu \|$ , some integration techniques and various refinements of the Hermite–Hadamard inequality. In the setting of matrices we prove that

$$\begin{aligned} & \left\| \left| A^{\frac{\alpha+\beta}{2}} XB^{1-\frac{\alpha+\beta}{2}} + A^{1-\frac{\alpha+\beta}{2}} XB^{\frac{\alpha+\beta}{2}} \right\| \right\| \\ & \leq \frac{1}{|\beta - \alpha|} \left\| \int_{\alpha}^{\beta} (A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu) d\nu \right\| \\ & \leq \frac{1}{2} \left\| \left| A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha + A^\beta XB^{1-\beta} + A^{1-\beta}XB^\beta \right\| \right\|, \end{aligned}$$

for real numbers  $\alpha, \beta$ .

© 2013 Elsevier Inc. All rights reserved.

## 30 1. Introduction

31 Let  $\mathbb{B}(\mathcal{H})$  denote the  $C^*$ -algebra of all bounded linear operators acting on a complex separable  
32 Hilbert space  $(\mathcal{H}, \langle \cdot, \cdot \rangle)$ . In the case when  $\dim \mathcal{H} = n$ , we identify  $\mathbb{B}(\mathcal{H})$  with the full matrix algebra

Q2 \* Corresponding author. Tel.: +98 9151140894; fax: +98 511 8644594.

E-mail addresses: [rupinder\\_grewal\\_86@yahoo.co.in](mailto:rupinder_grewal_86@yahoo.co.in) (R. Kaur), [moslehian@um.ac.ir](mailto:moslehian@um.ac.ir), [moslehian@ferdowsi.um.ac.ir](mailto:moslehian@ferdowsi.um.ac.ir), [moslehian@member.ams.org](mailto:moslehian@member.ams.org), [moslehian@yahoo.com](mailto:moslehian@yahoo.com) (M.S. Moslehian), [msrawla@yahoo.com](mailto:msrawla@yahoo.com) (M. Singh), [cconde@ungs.edu.ar](mailto:cconde@ungs.edu.ar) (C. Conde).

0024-3795/\$ - see front matter © 2013 Elsevier Inc. All rights reserved.

<http://dx.doi.org/10.1016/j.laa.2013.01.012>

33  $\mathcal{M}_n$  of all  $n \times n$  matrices with entries in the complex field. The cone of positive operators is denoted  
 34 by  $\mathbb{B}(\mathcal{H})_+$ . A unitarily invariant norm  $\|\cdot\|$  is defined on a norm ideal  $\mathfrak{J}_{\|\cdot\|}$  of  $\mathbb{B}(\mathcal{H})$  associated with  
 35 it and has the property  $\|UXV\| = \|X\|$ , where  $U$  and  $V$  are unitaries and  $X \in \mathfrak{J}_{\|\cdot\|}$ . Whenever we  
 36 write  $\|X\|$ , we mean that  $X \in \mathfrak{J}_{\|\cdot\|}$ . The operator norm on  $\mathbb{B}(\mathcal{H})$  is denoted by  $\|\cdot\|$ .

37 The arithmetic–geometric mean inequality for two positive real numbers  $a, b$  is  $\sqrt{ab} \leq (a+b)/2$ ,  
 38 which has been generalized in the context of bounded linear operators as follows. For  $A, B \in \mathbb{B}(\mathcal{H})_+$   
 39 and an unitarily invariant norm  $\|\cdot\|$  it holds that

$$2\|A^{1/2}XB^{1/2}\| \leq \|AX + XB\|.$$

40 For  $0 \leq \nu \leq 1$  and two nonnegative real numbers  $a$  and  $b$ , the *Heinz mean* is defined as

$$H_\nu(a, b) = \frac{a^\nu b^{1-\nu} + a^{1-\nu} b^\nu}{2}.$$

41 The function  $H_\nu$  is symmetric about the point  $\nu = \frac{1}{2}$ . Note that  $H_0(a, b) = H_1(a, b) = \frac{a+b}{2}$ ,

42  $H_{1/2}(a, b) = \sqrt{ab}$  and

$$H_{1/2}(a, b) \leq H_\nu(a, b) \leq H_0(a, b) \quad (1.1)$$

43 for  $0 \leq \nu \leq 1$ , i.e., the Heinz means interpolates between the geometric mean and the arithmetic  
 44 mean. The generalization of (1.1) in  $\mathbb{B}(\mathcal{H})$  asserts that for operators  $A, B, X$  such that  $A, B \in \mathbb{B}(\mathcal{H})_+$ ,  
 45 every unitarily invariant norm  $\|\cdot\|$  and  $\nu \in [0, 1]$  the following double inequality due to Bhatia and  
 46 Davis [3] holds

$$2\|A^{1/2}XB^{1/2}\| \leq \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \leq \|AX + XB\|. \quad (1.2)$$

47 Indeed, it has been proved that  $F(\nu) = \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\|$  is a convex function of  $\nu$  on  $[0, 1]$   
 48 with symmetry about  $\nu = 1/2$ , which attains its minimum there at and its maximum at  $\nu = 0$  and  
 49  $\nu = 1$ .

50 The second part of the previous inequality is one of the most essential inequalities in the operator  
 51 theory, which is called *the Heinz inequality*; see [11]. The proof given by Heinz [12] is based on the  
 52 complex analysis and is somewhat complicated. In [19], McIntosh showed that the Heinz inequality is  
 53 a consequence of the following inequality

$$\|A^*AX + XBB^*\| \geq 2\|AXB\|,$$

54 where  $A, B, X \in \mathbb{B}(\mathcal{H})$ . In the literature, the above inequality is called the *arithmetic–geometric mean*  
 55 *inequality*. Fujii et al. [10] proved that the Heinz inequality is equivalent to several other norm inequal-  
 56 ities such as the *Corach–Porta–Recht inequality*  $\|AXA^{-1} + A^{-1}XA\| \geq 2\|X\|$ , where  $A$  is a selfadjoint  
 57 invertible operator and  $X$  is a selfadjoint operator; see also [7]. Audenaert [2] gave a singular value  
 58 inequality for Heinz means by showing that if  $A, B \in \mathcal{M}_n$  are positive semidefinite and  $0 \leq \nu \leq 1$ ,  
 59 then  $s_j(A^\nu B^{1-\nu} + A^{1-\nu}B^\nu) \leq s_j(A+B)$  for  $j = 1, \dots, n$ , where  $s_j$  denotes the  $j$ th singular value. Also,  
 60 Yamazaki [25] used the classical Heinz inequality  $\|AXB\|^r \|X\|^{1-r} \geq \|A^\nu XB^{1-\nu}\|$  ( $A, B, X \in \mathbb{B}(\mathcal{H})$ ,  $A \geq$   
 61  $0, B \geq 0, r \in [0, 1]$ ) to characterize the chaotic order relation and to study isometric Aluthge trans-  
 62 formations.

63 For a detailed study of these and associated norm inequalities along with their history of origin,  
 64 refinements and applications, one may refer to [3,4,6,13–16].

65 It should be noticed that  $F(1/2) \leq F(\nu) \leq \frac{F(0)+F(1)}{2}$  provides a refinement to the Jensen inequality  
 66  $F(1/2) \leq \frac{F(0)+F(1)}{2}$  for the function  $F$ . Therefore it seems quite reasonable to obtain a new refinement  
 67 of (1.2) by utilizing a refinement of Jensen's inequality. This idea was recently applied by Kittaneh [18]  
 68 in virtue of the Hermite–Hadamard inequality (2.1).

69 One of the purposes of the present article is to obtain some new refinements of (1.2), from different  
 70 refinements of inequality (2.1). We also aim to give a unified study and further refinements to the  
 71 recent works for matrices.

72 **2. The Hermite–Hadamard inequality and its refinements**73 For a convex function  $f$ , the double inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2} \quad (2.1)$$

74 is known as the *Hermite–Hadamard* (H-H) inequality. This inequality was first published by Hermite  
75 in 1883 in an elementary journal and independently proved in 1893 by Hadamard. It gives us an  
76 estimation of the mean value of the convex function  $f$ ; see [17,20].77 There is an extensive amount of literature devoted to this simple and nice result, which has many  
78 applications in the theory of special means from which we would like to refer the reader to [21].  
79 Interestingly, each of two sides of the H-H inequality characterizes convex functions. More precisely, if  
80  $J$  is an interval and  $f : J \rightarrow \mathbb{R}$  is a continuous function, whose restriction to every compact subinterval  
81  $[a, b]$  verifies the first inequality of (2.1) then  $f$  is convex. The same works when the first inequality is  
82 replaced by the second one.83 Applying the H-H inequality, one can obtain the well-known geometric–logarithmic–arithmetic  
84 inequality

$$H_{1/2}(a, b) \leq L(a, b) \leq H_0(a, b),$$

85 where  $L(a, b) = \int_0^1 a^t b^{1-t} dt$ . An operator version of this has been proved by Hiai and Kosaki [14],  
86 which says that for  $A, B \in \mathbb{B}(\mathcal{H})_+$ ,

$$\| \|A^{1/2}XB^{1/2}\| \| \leq \| \| \int_0^1 A^\nu XB^{1-\nu} d\nu \| \| \leq \frac{1}{2} \| \|AX + XB\| \|,$$

87 which is another refinement of the arithmetic–geometric operator inequality.

88 Throughout this paper we will use the following notation: For  $a, b \in \mathbb{R}$  and  $t \in [0, 1]$ , let

$$m_f(a, b) = \frac{1}{b-a} \int_a^b f(x) dx,$$

89 and

$$[a, b]_t = (1-t)a + tb.$$

90 If  $f$  is an integrable function on  $[a, b]$  then

$$\frac{1}{b-a} \int_a^b f(x) dx = \int_0^1 f(ta + (1-t)b) dt = \int_0^1 f(tb + (1-t)a) dt,$$

91 and if  $f$  is convex on  $[a, b]$  we get

$$\frac{1}{b-a} \int_a^b f(x) dx = \int_0^1 F_{(a,b)}(t) dt,$$

92 where  $F_{(a,b)}(t) = \frac{1}{2} \left( f\left(a + \frac{t(b-a)}{2}\right) + f\left(b - \frac{t(b-a)}{2}\right) \right)$ ; see [1, Theorem 1.2].

93 In this section we collect various refinements of the H-H inequality for convex functions.

94 **Theorem 2.1** [8,23]. If  $f : [a, b] \rightarrow \mathbb{R}$  is a convex function and  $H_t, G_t$  are defined on  $[0, 1]$  by

$$H_t(a, b) = \frac{1}{b-a} \int_a^b f\left(\left[\frac{a+b}{2}, x\right]_t\right) dx,$$

95 and

$$G_t(a, b) = \frac{1}{2(b-a)} \int_a^b [f([x, a]_t) + f([x, b]_t)] dx,$$

96 then  $H_t$  and  $G_t$  are convex, increasing and

$$f\left(\frac{a+b}{2}\right) = H_0(a, b) \leq H_t(a, b) \leq H_1(a, b) = m_f(a, b), \quad (2.2)$$

$$m_f(a, b) = G_0(a, b) \leq G_t(a, b) \leq G_1(a, b) = \frac{f(a) + f(b)}{2} \quad (2.3)$$

97 for all  $t \in [0, 1]$ . Furthermore,

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{2}{b-a} \int_{\frac{3a+b}{4}}^{\frac{a+3b}{4}} f(x) dx \leq \int_0^1 H_t(a, b) dt \\ &\leq \frac{1}{2} \left( f\left(\frac{a+b}{2}\right) + m_f(a, b) \right) \leq m_f(a, b) \end{aligned}$$

98 and

$$\begin{aligned} \frac{2}{b-a} \int_{\frac{3a+b}{4}}^{\frac{a+3b}{4}} f(x) dx &\leq \frac{1}{2} \left( f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right) \leq \int_0^1 G_t(a, b) dt \\ &\leq \frac{1}{2} \left( f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right) \leq \frac{f(a) + f(b)}{2}. \end{aligned} \quad (2.4)$$

99 **Remark 2.2.** (1) From (2.4) we get that

$$m_f(a, b) \leq \frac{1}{2} \left( f\left(\frac{a+b}{2}\right) + \frac{f(a) + f(b)}{2} \right) \leq \frac{f(a) + f(b)}{2},$$

100 which is the well-known Bullen's inequality; see [21, p. 140]. As an immediate consequence,  
101 from the previous inequality, we note that the first inequality is stronger than the second one  
102 in (2.1), i.e.

$$m_f(a, b) - f\left(\frac{a+b}{2}\right) \leq \frac{f(a) + f(b)}{2} - m_f(a, b).$$

103 (2) We note some properties of  $H_t$  and  $G_t$  useful in the next sections. For  $\mu \in [0, 1]$  we get

$$104 \quad (a) \quad H_t(\mu, 1 - \mu) = \frac{1}{1-2\mu} \int_{\mu}^{1-\mu} f\left(\left[\frac{1}{2}, x\right]_t\right) dx = \frac{1}{2\mu-1} \int_{1-\mu}^{\mu} f\left(\left[\frac{1}{2}, x\right]_t\right) dx = H_t(1 - \mu, \mu).$$

$$105 \quad (b) \quad G_t(\mu, 1 - \mu) = \frac{1}{2(1-2\mu)} \int_{\mu}^{1-\mu} [f([x, \mu]_t) + f([x, 1 - \mu]_t)] dx = G_t(1 - \mu, \mu).$$

106 Recently, the following result was proved:

107 **Theorem 2.3** [24]. If  $f$  is a convex function defined on an interval  $J$ ,  $a, b \in J^\circ$  with  $a < b$  and the mapping  
108  $T_t$  is defined by

$$T_t(a, b) = \frac{1}{2} \left( f\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) + f\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) \right),$$

109 then  $T_t$  is convex and increasing on  $[0, 1]$  and

$$f\left(\frac{a+b}{2}\right) \leq T_\eta(a, b) \leq T_\xi(a, b) \leq T_\lambda(a, b) \leq \frac{f(a) + f(b)}{2},$$

110 for all  $\eta \in (0, \xi)$ ,  $\lambda \in (\xi, 1)$ , where  $T_\xi(a, b) = m_f(a, b)$ .

111 In [9], the author asked whether for a convex function  $f$  on an interval  $J$  there exist real numbers  $l$ ,  
112  $L$  such that

$$f\left(\frac{a+b}{2}\right) \leq l \leq \frac{1}{b-a} \int_a^b f(x)dx \leq L \leq \frac{f(a)+f(b)}{2}.$$

113 An affirmative answer to this question is given as follows.

114 **Theorem 2.4** [9]. Assume that  $f : [a, b] \rightarrow \mathbb{R}$  is a convex function. Then

$$f\left(\frac{a+b}{2}\right) \leq l(\lambda) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq L(\lambda) \leq \frac{f(a)+f(b)}{2} \tag{2.5}$$

115 for all  $\lambda \in [0, 1]$ , where

$$l(\lambda) = \lambda f\left(\frac{\lambda b + (2-\lambda)a}{2}\right) + (1-\lambda)f\left(\frac{(1+\lambda)b + (1-\lambda)a}{2}\right)$$

116 and

$$L(\lambda) = \frac{1}{2}(f(\lambda b + (1-\lambda)a) + \lambda f(a) + (1-\lambda)f(b)).$$

117 **Remark 2.5.** Applying inequality (2.5) for  $\lambda = \frac{1}{2}$  we get

$$\begin{aligned} f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2} \left( f\left(\frac{3a+b}{4}\right) + f\left(\frac{a+3b}{4}\right) \right) \leq m_f(a, b) \\ &\leq \frac{1}{2} \left( f\left(\frac{a+b}{2}\right) + \frac{f(a)+f(b)}{2} \right) \leq \frac{f(a)+f(b)}{2}. \end{aligned}$$

118 This result has been obtained by Akkouchi in [1].

119 **3. Refinements of the Heinz inequality for operators**

120 In this section we use the convexity of  $F(\nu) = |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu|||$ ;  $\nu \in [0, 1]$  and the  
121 different refinements of inequality (2.1) described in the previous section.

122 **Theorem 3.1.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$ . Then for any  $t, \mu \in [0, 1]$  and any  
123 unitary invariant norm  $|||\cdot|||$ ,

$$\begin{aligned} 2|||A^{1/2}XB^{1/2}||| &\leq \frac{1}{1-2\mu} \int_\mu^{1-\mu} F([1/2, x]_t) dx \\ &\leq \frac{1}{1-2\mu} \int_\mu^{1-\mu} |||A^x XB^{1-x} + A^{1-x}XB^x||| dx \\ &\leq \frac{1}{2(1-2\mu)} \int_\mu^{1-\mu} [F([x, \mu]_t) + F([x, 1-\mu]_t)] dx \\ &\leq |||A^\mu XB^{1-\mu} + A^{1-\mu}XB^\mu||| \end{aligned}$$

124 **Proof.** For  $\mu \neq \frac{1}{2}$  the inequalities follows by applying inequalities (2.2) and (2.3) on the interval  
125  $[\mu, 1-\mu]$  if  $0 \leq \mu < \frac{1}{2}$  or  $[1-\mu, \mu]$  if  $\frac{1}{2} < \mu \leq 1$ . Finally

$$\lim_{\mu \rightarrow \frac{1}{2}} \frac{1}{2(1-2\mu)} \int_\mu^{1-\mu} (F([x, \mu]_t) + F([x, 1-\mu]_t)) dx = 2|||A^{1/2}XB^{1/2}|||$$

126 completes the proof.  $\square$

127 Applying Theorem 2.1 to the function  $F$  on the interval  $[\mu, \frac{1}{2}]$  or  $[\frac{1}{2}, \mu]$  for  $\mu \in [0, 1]$  we obtain  
128 the following refinement of [18, Theorem 2 and Corollary 1].

129 **Theorem 3.2.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$ . Then for every  $\mu \in [0, 1]$  and every  
130 unitarily invariant norm  $||| \cdot |||$ ,

$$\begin{aligned} 2|||A^{1/2}XB^{1/2}||| &\leq |||A^{\frac{2\mu+1}{4}}XB^{\frac{3-2\mu}{4}} + A^{\frac{3-2\mu}{4}}XB^{\frac{2\mu+1}{4}}||| \\ &\leq \frac{4}{1-2\mu} \int_{\frac{(2\mu+3)}{8}}^{\frac{(2\mu+3)}{8}} |||A^xXB^{1-x} + A^{1-x}XB^x|||dx \leq \int_0^1 H_t(1/2, \mu)dt \\ &\leq \frac{1}{2}|||A^{\frac{2\mu+1}{4}}XB^{\frac{3-2\mu}{4}} + A^{\frac{3-2\mu}{4}}XB^{\frac{2\mu+1}{4}}||| + \frac{1}{1-2\mu} \int_{\mu}^{1/2} F(x)dx \\ &\leq \frac{2}{1-2\mu} \int_{\mu}^{1/2} |||A^xXB^{1-x} + A^{1-x}XB^x|||dx = G_0(1/2, \mu) \leq \int_0^1 G_t(1/2, \mu)dt \\ &\leq \frac{1}{2} \left( |||A^{\frac{2\mu+1}{4}}XB^{\frac{3-2\mu}{4}} + A^{\frac{3-2\mu}{4}}XB^{\frac{2\mu+1}{4}}||| + |||A^{\mu}XB^{1-\mu} + A^{1-\mu}XB^{\mu}||| + F(1/2) \right) \\ &\leq \frac{1}{2} |||A^{\mu}XB^{1-\mu} + A^{1-\mu}XB^{\mu}||| + |||A^{1/2}XB^{1/2}||| \\ &\leq |||A^{\mu}XB^{1-\mu} + A^{1-\mu}XB^{\mu}|||. \end{aligned}$$

131 Now, we have the following refinement of the first part of the the Heinz inequality via certain  
132 sequences.

133 **Theorem 3.3.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$  and for  $n \in \mathbb{N}_0$ ,

$$\begin{aligned} x_n(F, a, b) &= \frac{1}{2^n} \sum_{i=1}^{2^n} F \left( a + \left( i - \frac{1}{2} \right) \frac{b-a}{2^n} \right), \\ y_n(F, a, b) &= \frac{1}{2^n} \left( \frac{F(a) + F(b)}{2} + \sum_{i=1}^{2^n-1} F \left( [a, b]_{\frac{i}{2^n}} \right) \right). \end{aligned}$$

134 Then

135 (1) For  $\mu \in [0, 1/2]$  and for every unitarily invariant norm  $||| \cdot |||$ ,

$$\begin{aligned} 2|||A^{1/2}XB^{1/2}||| &= x_0(F, \mu, 1-\mu) \leq \dots \leq x_n(F, \mu, 1-\mu) \\ &\leq \frac{1}{1-2\mu} \int_{\mu}^{1-\mu} |||A^xXB^{1-x} + A^{1-x}XB^x|||dx \\ &\leq y_n(F, \mu, 1-\mu) \leq \dots \leq y_0(F, \mu, 1-\mu) = F(\mu) \end{aligned}$$

136 (2) For  $\mu \in [1/2, 1]$  and for every unitarily invariant norm  $||| \cdot |||$ ,

$$\begin{aligned} 2|||A^{1/2}XB^{1/2}||| &= x_0(F, 1-\mu, \mu) \leq \dots \leq x_n(F, 1-\mu, \mu) \\ &\leq \frac{1}{2\mu-1} \int_{1-\mu}^{\mu} |||A^xXB^{1-x} + A^{1-x}XB^x|||dx \\ &\leq y_n(F, 1-\mu, \mu) \leq \dots \leq y_0(F, 1-\mu, \mu) = F(\mu) \end{aligned}$$

137 Applying the Theorem 2.4, we obtain the following refinement.



138 **Theorem 3.4.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$  and  $\alpha, \beta \in [0, 1]$  and  $\|\cdot\|$  be a  
139 unitarily invariant norm. Then

$$F\left(\frac{\alpha + \beta}{2}\right) \leq I(\lambda) \leq \frac{1}{b-a} \int_a^b F(x) dx \leq L(\lambda) \leq \frac{F(\alpha) + F(\beta)}{2}$$

140 for all  $\lambda \in [0, 1]$ , where

$$I(\lambda) = \lambda F\left(\frac{\lambda\beta + (2-\lambda)\alpha}{2}\right) + (1-\lambda)F\left(\frac{(1+\lambda)\beta + (1-\lambda)\alpha}{2}\right)$$

141 and

$$L(\lambda) = \frac{1}{2}(F(\lambda\beta + (1-\lambda)\alpha) + \lambda F(\alpha) + (1-\lambda)F(\beta)).$$

142 Finally, using the refinement presented in Theorem 2.3 we get the following statement.

143 **Theorem 3.5.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$ . For  $a, b \in (0, 1)$  with  $a < b$  let  $T_t$  be  
144 the mapping defined in [0, 1] by

$$T_t(a, b) = \frac{1}{2} \left( F\left(\frac{1+t}{2}a + \frac{1-t}{2}b\right) + F\left(\frac{1-t}{2}a + \frac{1+t}{2}b\right) \right).$$

145 Then, there exists  $\xi \in (0, 1)$  such that for any  $\mu \in (0, 1)$  and any unitary invariant norm  $\|\cdot\|$ ,

$$\begin{aligned} 2\| |A|^{1/2}XB^{1/2} \| \leq T_\eta(\mu, 1-\mu) &\leq T_\xi(\mu, 1-\mu) = \frac{1}{1-2\mu} \int_\mu^{1-\mu} F(x) dx \\ &\leq T_\lambda(\mu, 1-\mu) \leq \| |A^\mu XB^{1-\mu} + A^{1-\mu}XB^\mu \|, \end{aligned}$$

146 where  $\eta \in [0, \xi]$  and  $\lambda \in [\xi, 1]$ .

147 From the generalization of the H-H inequality due to Vasić and Lacković, we get

148 **Theorem 3.6.** Let  $A, B, X$  be operators such that  $A, B \in \mathbb{B}(\mathcal{H})_+$  and let  $p, q$  be positive numbers and  
149  $0 \leq \alpha < \beta \leq 1$ . Then the double inequality

$$F\left(\frac{p\alpha + q\beta}{p+q}\right) \leq \frac{1}{2y} \int_{c-y}^{c+y} F(t) dt \leq \frac{pF(\alpha) + qF(\beta)}{p+q}$$

150 holds for  $c = \frac{p\alpha + q\beta}{p+q}$ ,  $y > 0$  if and only if  $y \leq \frac{\beta-\alpha}{p+q} \min\{p, q\}$ .

#### 151 4. Refinement of the Heinz inequality for matrices

152 In what follows, the capital letters  $A, B, X, \dots$  denote arbitrary elements of  $\mathcal{M}_n$ . By  $\mathbb{P}_n$  we denote  
153 the set of positive definite matrices. The Schur product of two matrices  $A = [a_{ij}]$  and  $B = [b_{ij}]$  in  $\mathcal{M}_n$   
154 is the entrywise product and denoted by  $A \circ B$ . We shall state the following preliminary result, which  
155 is needed to prove our main results.

156 If  $X = [x_{ij}]$  is positive semidefinite, then for any matrix  $Y$ , we have

$$\| |X \circ Y| \| \leq \| |Y| \| \max_i x_{ii} \tag{4.1}$$

157 for every unitarily invariant norm  $\|\cdot\|$ . For a proof of this, the reader may be referred to [12].

158 **Theorem 4.1.** Let  $A, B \in \mathbb{P}_n$  and  $X \in M_n$ . Then for any real numbers  $\alpha, \beta$  and any unitarily invariant  
159 norm  $\|\cdot\|$ ,

$$\begin{aligned} & \left\| \left\| A^{\frac{\alpha+\beta}{2}} X B^{1-\frac{\alpha+\beta}{2}} + A^{1-\frac{\alpha+\beta}{2}} X B^{\frac{\alpha+\beta}{2}} \right\| \right\| \leq \frac{1}{|\beta-\alpha|} \left\| \int_{\alpha}^{\beta} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \\ & \leq \frac{1}{2} \left\| \left\| A^{\alpha} X B^{1-\alpha} + A^{1-\alpha} X B^{\alpha} + A^{\beta} X B^{1-\beta} + A^{1-\beta} X B^{\beta} \right\| \right\|. \end{aligned} \tag{4.2}$$

160 **Proof.** Without loss of generality assume that  $\alpha < \beta$ . We shall first prove the result for the case  
161  $A = B$ . Since the norms considered here are unitarily invariant, so we can assume that  $A$  is diagonal,  
162 i.e.  $A = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_n)$ .

163 Note that

$$A^{\frac{\alpha+\beta}{2}} X A^{1-\frac{\alpha+\beta}{2}} + A^{1-\frac{\alpha+\beta}{2}} X A^{\frac{\alpha+\beta}{2}} = Y \circ \left( \int_{\alpha}^{\beta} (A^{\nu} X A^{1-\nu} + A^{1-\nu} X A^{\nu}) d\nu \right),$$

164 where  $Y$  is a Hermitian matrix. If  $X = [x_{ij}]$  and  $Y = [y_{ij}]$ , then

$$\left[ \lambda_i^{\frac{\alpha+\beta}{2}} x_{ij} \lambda_j^{1-\frac{\alpha+\beta}{2}} + \lambda_i^{1-\frac{\alpha+\beta}{2}} x_{ij} \lambda_j^{\frac{\alpha+\beta}{2}} \right] = \left[ y_{ij} \int_{\alpha}^{\beta} (\lambda_i^{\nu} x_{ij} \lambda_j^{1-\nu} + \lambda_i^{1-\nu} x_{ij} \lambda_j^{\nu}) d\nu \right],$$

165 whence

$$\begin{aligned} y_{ij} &= \frac{\lambda_i^{\frac{\alpha+\beta}{2}} \lambda_j^{1-\frac{\alpha+\beta}{2}} + \lambda_i^{1-\frac{\alpha+\beta}{2}} \lambda_j^{\frac{\alpha+\beta}{2}}}{\int_{\alpha}^{\beta} (\exp(\log(\lambda_i)\nu + \log(\lambda_j)(1-\nu)) + \exp(\log(\lambda_i)(1-\nu) + \log(\lambda_j)\nu)) d\nu} \\ &= \frac{\lambda_i^{\frac{\beta-\alpha}{2}} (\lambda_i^{\alpha} \lambda_j^{1-\beta} + \lambda_i^{1-\beta} \lambda_j^{\alpha}) \lambda_j^{\frac{\beta-\alpha}{2}} (\log \lambda_i - \log \lambda_j)}{\lambda_i^{\beta} \lambda_j^{1-\beta} - \lambda_i^{1-\beta} \lambda_j^{\beta} - \lambda_i^{\alpha} \lambda_j^{1-\alpha} + \lambda_i^{1-\alpha} \lambda_j^{\alpha}} \\ &= \frac{\lambda_i^{\frac{\beta-\alpha}{2}} (\log \lambda_i - \log \lambda_j) \lambda_j^{\frac{\beta-\alpha}{2}}}{\lambda_i^{\beta-\alpha} - \lambda_j^{\beta-\alpha}}, \quad \text{for } i \neq j \end{aligned}$$

166 and  $y_{ii} = \frac{1}{\beta-\alpha} > 0$ . By (4.1), it is enough to show that the matrix  $Y$  is positive semidefinite, or  
167 equivalently the matrix

$$y'_{ij} = \begin{cases} \frac{\log \lambda_i - \log \lambda_j}{\lambda_i^{\beta-\alpha} - \lambda_j^{\beta-\alpha}} & \text{if } i \neq j \\ \frac{1}{(\beta-\alpha)\lambda_i^{\beta-\alpha}} & \text{if } i = j \end{cases}$$

168 is positive semidefinite. On taking  $\lambda_i^{\beta-\alpha} = s_i$ , we get

$$(\beta-\alpha)y'_{ij} = \begin{cases} \frac{\log s_i - \log s_j}{s_i - s_j} & \text{if } i \neq j \\ \frac{1}{s_i} & \text{if } i = j, \end{cases}$$

169 which is a positive semidefinite matrix, since the matrix on the right hand side is the Löwner matrix  
170 corresponding to the matrix monotone function  $\log x$ ; see [4, Theorem 5.3.3]. This proves the first  
171 inequality in (4.2) for the case  $A = B$ .

172 The second inequality will follow on the same lines. We indeed have

$$\int_{\alpha}^{\beta} (A^{\nu} X A^{1-\nu} + A^{1-\nu} X A^{\nu}) d\nu = Z \circ (A^{\alpha} X B^{1-\alpha} + A^{1-\alpha} X B^{\alpha} + A^{\beta} X B^{1-\beta} + A^{1-\beta} X B^{\beta}),$$

173 where  $Z$  is the Hermitian matrix with entries

$$z_{ij} = \begin{cases} \frac{\lambda_i^{\beta-\alpha} - \lambda_j^{\beta-\alpha}}{(\log \lambda_i - \log \lambda_j)(\lambda_i^{\beta-\alpha} + \lambda_j^{\beta-\alpha})} & \text{if } i \neq j \\ \frac{(\beta-\alpha)}{2} & \text{if } i = j. \end{cases}$$

174 On taking  $\lambda_i^{\beta-\alpha} = e^{t_i}$  we conclude that  $Z$  is positive semidefinite if and only if so is the following  
175 matrix

$$\frac{2}{\beta - \alpha} z'_{ij} = \begin{cases} \frac{\tanh((t_i - t_j)/2)}{(t_i - t_j)/2} & \text{if } i \neq j \\ 1 & \text{if } i = j. \end{cases}$$

176 The right hand side matrix is positive semidefinite since the function  $f(x) = \frac{\tanh x}{x}$  is positive definite;  
177 see [4, Example 5.2.11]. This proves the second inequality in (4.2) for the case  $A = B$ .

178 The general case follows on replacing  $A$  by  $\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$  and  $X$  by  $\begin{bmatrix} 0 & X \\ 0 & 0 \end{bmatrix}$ .  $\square$

179 The first corollary provides some variants of [18, Theorems 2 and 3]. It should be noticed that

$$\lim_{\mu \rightarrow 1/2} \left( \frac{2}{|1 - 2\mu|} \left\| \int_{\mu}^{1/2} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \right) = 2 \left\| A^{1/2} X B^{1/2} \right\|$$

180 and

$$\lim_{\mu \rightarrow 0} \left( \frac{1}{|\mu|} \left\| \int_0^{\mu} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \right) = \|AX + XB\|.$$

181 **Corollary 4.2.** Let  $A, B \in \mathbb{P}_n, X \in M_n, \mu$  be a real number and  $\|\cdot\|$  be any unitarily invariant norm.  
182 Then

$$\begin{aligned} \left\| A^{\frac{2\mu+1}{4}} X B^{\frac{3-2\mu}{4}} + A^{\frac{3-2\mu}{4}} X B^{\frac{2\mu+1}{4}} \right\| &\leq \frac{2}{|1 - 2\mu|} \left\| \int_{\mu}^{1/2} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \\ &\leq \frac{1}{2} \left\| A^{\mu} X B^{1-\mu} + A^{1-\mu} X B^{\mu} + 2A^{1/2} X B^{1/2} \right\|, \end{aligned}$$

$$\begin{aligned} \left\| A^{\frac{\mu}{2}} X B^{1-\frac{\mu}{2}} + A^{1-\frac{\mu}{2}} X B^{\frac{\mu}{2}} \right\| &\leq \frac{1}{|\mu|} \left\| \int_0^{\mu} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \\ &\leq \frac{1}{2} \left\| AX + XB + A^{\mu} X B^{1-\mu} + A^{1-\mu} X B^{\mu} \right\|. \end{aligned}$$

183 The following consequence provides a matrix analogue of (1.1).

184 **Corollary 4.3.** Let  $A, B \in \mathbb{P}_n$  and  $X \in M_n$ . Then for any  $0 \leq \alpha < \beta \leq 1$  with  $\alpha + \beta \leq 2$  and any  
185 unitarily invariant norm  $\|\cdot\|$ ,

$$\begin{aligned} 2 \left\| A^{1/2} X B^{1/2} \right\| &\leq \left\| A^{\frac{\alpha+\beta}{2}} X B^{1-\frac{\alpha+\beta}{2}} + A^{1-\frac{\alpha+\beta}{2}} X B^{\frac{\alpha+\beta}{2}} \right\| \\ &\leq \frac{1}{|\beta - \alpha|} \left\| \int_{\alpha}^{\beta} (A^{\nu} X B^{1-\nu} + A^{1-\nu} X B^{\nu}) d\nu \right\| \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2} \left\| \|A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha + A^\beta XB^{1-\beta} + A^{1-\beta}XB^\beta\| \right\| \\ &\leq \frac{1}{2} \left\| \|A^\alpha XB^{1-\alpha} + A^{1-\alpha}XB^\alpha\| \right\| + \frac{1}{2} \left\| \|A^\beta XB^{1-\beta} + A^{1-\beta}XB^\beta\| \right\| \\ &\leq \|AX + XB\|. \end{aligned}$$

186 **Proof.** Applying the triangle inequality, the properties of the function  $f(\nu) = \| \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \|$   
187 and Theorem 4.1 we get the required inequalities.  $\square$

188 It is shown in [18, Corollary 3] that

$$\| \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \| \leq 4r_0 \| \|A^{1/2}XB^{1/2}\| \| + (1 - 2r_0) \| \|AX + XB\| \|. \tag{4.3}$$

189 A natural generalization of (4.3) would be

$$\| \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \| \leq \| \|4r_0A^{1/2}XB^{1/2} + (1 - 2r_0)(AX + XB)\| \|$$

190 for  $0 \leq \nu \leq 1$  and  $r_0 = \min\{\nu, 1 - \nu\}$  with  $A, B \in \mathbb{P}_n$  and  $X \in M_n$ , which in fact is not true, in  
191 general. The following counterexample justifies this:

192 Take  $X = \begin{bmatrix} 52.39 & 38.71 & 12.36 \\ 32.86 & 35.38 & 64.82 \\ 91.79 & 99.45 & 66.10 \end{bmatrix}$ ,  $A = \begin{bmatrix} 92.315 & 87.791 & 71.090 \\ 87.791 & 120.130 & 83.340 \\ 71.090 & 83.340 & 103.610 \end{bmatrix}$ ,

193  $B = \begin{bmatrix} 118.482 & 23.249 & 112.676 \\ 23.249 & 10.343 & 38.224 \\ 112.676 & 38.224 & 156.551 \end{bmatrix}$  and  $\nu = 0.4680$ . Then  $\text{tr}|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu| = 78135.5$ , while

194  $\text{tr}|4r_0A^{1/2}XB^{1/2} + (1 - 2r_0)(AX + XB)| = 78125.4$ .

195 We shall, however, present another result, which is a possible generalization of (4.3).  
196

197 **Theorem 4.4.** Let  $A, B \in \mathbb{P}_n$  and  $X \in M_n$ . Then for  $\nu \in [0, 1]$  and for every unitarily invariant norm  
198  $\| \cdot \|$ ,

$$\| \|A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu\| \| \leq \| \|4r_1(\nu)A^{1/2}XB^{1/2} + (1 - 2r_1(\nu))(AX + XB)\| \|, \tag{4.4}$$

199 where  $r_1(\nu) = \min\{\nu, \left|\frac{1}{2} - \nu\right|, 1 - \nu\}$ .

200 **Proof.** First, we consider the case  $\nu \in [0, 1/2]$ . Notice that by some simple algebraic or geometrical  
201 arguments, we may conclude that  $0 \leq r_1 \leq 1/4$ . Again, by following a similar way as in Theorem 4.1,  
202 we can write the matrix

$$A^\nu XA^{1-\nu} + A^{1-\nu}XA^\nu = W \circ (4r_1A^{1/2}XA^{1/2} + (1 - 2r_1)(AX + XA)),$$

203 where  $W$  is a Hermitian matrix with entries

$$w_{ij} = \begin{cases} \frac{\lambda_i^\nu (\lambda_i^{1-2\nu} + \lambda_j^{1-2\nu}) \lambda_j^\nu}{4r_1 \lambda_i^{1/2} \lambda_j^{1/2} + (1-2r_1)(\lambda_i + \lambda_j)} & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

204 Now, observe that  $0 \leq \frac{4r_1}{1-2r_1} \leq 2$  and  $0 \leq 1 - 2\nu \leq 1$ , so the matrix  $W$  is positive semidefinite; see  
205 [6, Theorem 5.2, p. 225]. On repeating the same argument as in Theorem 4.1, the required inequality  
206 (4.4) follows.

207 Finally, if  $\nu \in [\frac{1}{2}, 1]$  let  $\mu = 1 - \nu \in [0, \frac{1}{2}]$ , then by the previous case we have

$$\begin{aligned} |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| &= |||A^{1-\mu}XB^\mu + A^\mu XB^{1-\mu}||| \\ &\leq |||4r_1(\mu)A^{\frac{1}{2}}XB^{\frac{1}{2}} + (1 - 2r_1(\mu))(AX + XB)|||, \end{aligned}$$

208 where  $r_1(\mu) = \min \left\{ \mu, \left| \frac{1}{2} - \mu \right|, 1 - \mu \right\} = r_1(\nu)$ .  $\square$

209 From the previous theorem, we deduce a new refinement of the Heinz inequality for matrices.

210 **Corollary 4.5.** Let  $A, B \in \mathbb{P}_n$  and  $X \in M_n$ . Then for  $\nu \in [0, 1]$  and for every unitarily invariant norm  
211  $||| \cdot |||$ ,

$$\begin{aligned} |||A^\nu XB^{1-\nu} + A^{1-\nu}XB^\nu||| &\leq |||4r_1(\nu)A^{1/2}XB^{1/2} + (1 - 2r_1(\nu))(AX + XB)||| \\ &\leq 4r_1(\nu)|||A^{1/2}XB^{1/2}||| + (1 - 2r_1(\nu))|||AX + XB||| \\ &\leq 2(2r_1(\nu) - 1)|||A^{1/2}XB^{1/2}||| + 2(1 - r_1(\nu))|||AX + XB||| \\ &\leq |||AX + XB|||, \end{aligned}$$

212 where  $r_1(\nu) = \min \left\{ \nu, \left| \frac{1}{2} - \nu \right|, 1 - \nu \right\}$ .

213 As a direct consequence of Theorem 4.4, we obtain the following refinement of an inequality  
214 (see [7]).

215 **Corollary 4.6.** Let  $A, B \in \mathbb{P}_n$ ,  $X \in M_n$ ,  $r \in [\frac{1}{2}, \frac{3}{2}]$  and  $t \in (-2, 2]$ . Then for every unitarily invariant  
216 norm  $||| \cdot |||$ ,

$$\begin{aligned} |||A^r XB^{2-r} + A^{2-r}XB^r||| &\leq |||4sAXB + (1 - 2s)(A^{3/2}XB^{1/2} + A^{1/2}XB^{3/2})||| \\ &\leq 4s|||AXB||| + (1 - 2s)|||A^{3/2}XB^{1/2} + A^{1/2}XB^{3/2}||| \\ &\leq 4s|||AXB||| + (1 - 2s)\frac{2}{t+2}|||A^2X + tAXB + XB^2||| \\ &\leq 2(2s - 1)|||AXB||| + \frac{4(1 - s)}{t+2}|||A^2X + tAXB + XB^2||| \\ &\leq \frac{2}{t+2}|||A^2X + tAXB + XB^2||| \end{aligned}$$

217 in which  $s = \min \left\{ r - \frac{1}{2}, |1 - r|, \frac{3}{2} - r \right\}$ .

218 **Proof.** Let  $Y = A^{1/2}XB^{1/2} \in M_n$  and  $\nu = r - \frac{1}{2} \in [0, 1]$ . It follows from Theorem 4.4 that

$$\begin{aligned} |||A^r XB^{2-r} + A^{2-r}XB^r||| &= |||A^r A^{-1/2}YB^{-1/2}B^{2-r} + A^{2-r}A^{-1/2}YB^{-1/2}B^r||| \\ &= |||A^\nu YB^{1-\nu} + A^{1-\nu}YB^{1-\nu}||| \\ &\leq |||4r_1(\nu)A^{1/2}YB^{1/2} + (1 - 2r_1(\nu))(AY + YB)||| \\ &= |||4r_1(\nu)AXB + (1 - 2r_1(\nu))(A^{3/2}XB^{1/2} + A^{1/2}XB^{3/2})|||, \end{aligned}$$

219 where  $r_1(\nu) = \min \left\{ \nu, \left| \frac{1}{2} - \nu \right|, 1 - \nu \right\}$ . Let  $s = r_1 \left( r - \frac{1}{2} \right)$ . Applying the triangle inequality and  
220 Zhan's inequality, we obtain

$$\begin{aligned} |||A^r XB^{2-r} + A^{2-r}XB^r||| &\leq |||4sAXB + (1 - 2s)(A^{3/2}XB^{1/2} + A^{1/2}XB^{3/2})||| \\ &\leq 4s|||AXB||| + (1 - 2s)|||A^{3/2}XB^{1/2} + A^{1/2}XB^{3/2}||| \end{aligned}$$

$$\begin{aligned} &\leq 4s\| |AXB| \| + \frac{2(1-2s)}{t+2} \| |A^2X + tAXB + XB^2| \| \\ &\leq 2(2s-1)\| |AXB| \| + \frac{4(1-s)}{t+2} \| |A^2X + tAXB + XB^2| \| \\ &\leq \frac{2}{t+2} \| |A^2X + tAXB + XB^2| \|. \quad \square \end{aligned}$$

## 221 Acknowledgments

222 The first author would like to sincerely thank the “Council of Scientific & Industrial Research  
223 Human Resource Development Group, India” for grant of Senior Research Fellowship vide  
224 09/797(0009)/2012-EMR-I. The second author was supported by a grant from Ferdowsi University  
225 of Mashhad (No. MP91290MOS).

## 226 References

- 227 [1] M. Akkouchi, Some remarks on the Jensen–Hadamard inequalities and applications, *Ann. Univ. Ferrara Sez. VII (N.S.)* 49 (2003)  
228 1–10.
- 229 [2] K.M.R. Audenaert, A singular value inequality for Heinz means, *Linear Algebra Appl.* 422 (2007) 279–283.
- 230 [3] R. Bhatia, C. Davis, More matrix forms of the arithmetic–geometric mean inequality, *SIAM J. Matrix Anal. Appl.* 14 (1993) 132–136.
- 231 [4] R. Bhatia, *Positive Definite Matrices*, Princeton University Press, New Jersey, 2007.
- 232 [5] R. Bhatia, F. Kittaneh, Clarkson inequalities with several operators, *Bull. London Math. Soc.* 36 (6) (2004) 820–832.
- 233 [6] R. Bhatia, K.R. Parthasarathy, Positive definite functions and operator inequalities, *Bull. London Math. Soc.* 32 (2000) 214–228.
- 234 [7] C. Conde, A. Seddik, M.S. Moslehian, Operator inequalities related to the Corach–Porta–Recht inequality, *Linear Algebra Appl.*  
235 436 (9) (2012) 3008–3017.
- 236 [8] S.S. Dragomir, Two mappings in connection to Hadamard’s inequalities, *J. Math. Anal. Appl.* 167 (1) (1992) 49–56.
- 237 [9] A. El Farissi, Simple proof and refinement of Hermite–Hadamard inequality, *J. Math. Inequal.* 4 (3) (2010) 365–369.
- 238 [10] J.I. Fujii, M. Fujii, T. Furuta, M. Nakamoto, Norm inequalities equivalent to Heinz inequality, *Proc. Amer. Math. Soc.* 118 (1993)  
239 827–830.
- 240 [11] E. Heinz, Beiträge zur Störungstheorie der Spektralzerlegung, *Math. Ann.* 123 (1951) 415–438 (in German).
- 241 [12] R.A. Horn, C.R. Johnson, *Topics in Matrix Analysis*, Cambridge University Press, 1990.
- 242 [13] F. Hiai, H. Kosaki, Means for matrices and comparison of their norms, *Indiana Univ. Math. J.* 48 (1999) 899–936.
- 243 [14] F. Hiai, H. Kosaki, Comparison of various means for operators, *J. Funct. Anal.* 163 (1999) 300–323.
- 244 [15] F. Hiai, H. Kosaki, *Means of Hilbert Space Operators*, Lecture Notes in Mathematics, vol. 1820, Springer, New York, 2003.
- 245 [16] H. Kosaki, Arithmetic–geometric mean and related inequalities for operators, *J. Funct. Anal.* 156 (1998) 429–451.
- 246 [17] E. Kikianty, Hermite–Hadamard inequality in the geometry of Banach spaces, Ph.D. thesis, Victoria University, 2010. Available  
247 from: <<http://eprints.vu.edu.au/15793/1/EderKikiantyThesis.pdf>>.
- 248 [18] F. Kittaneh, On convexity of the Heinz means, *Integral Equations Operator Theory* 68 (2010) 519–527.
- 249 [19] A. McIntosh, Heinz inequalities and perturbation of spectral families, *Macquarie Mathematical Reports*, Macquarie University,  
250 1979.
- 251 [20] M.S. Moslehian, Matrix Hermite–Hadamard type inequalities, *Houston J. Math.*, in press, [arXiv:1203.5300](https://arxiv.org/abs/1203.5300).
- 252 [21] J. Pečarić, F. Proschan, Y.L. Tong, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press Inc., 1992.
- 253 [22] W.T. Sulaiman, Some mappings in connection to Hadamard’s inequality, *Int. J. Math. Anal. (Ruse)* 4 (37–40) (2010) 1829–1836.
- 254 [23] G.-S. Yang, M.-C. Hong, A note on Hadamard’s inequality, *Tamkang J. Math.* 28 (1) (1997) 33–37.
- 255 [24] F. Wang, A family associated with Hadamard’s inequality on a hypercube, *ISRN Math. Anal.* (2011) Article ID 594758.
- 256 [25] T. Yamazaki, Characterizations of  $\log A \geq \log B$  and normaloid operators via Heinz inequality, *Integral Equations Operator*  
257 *Theory* 43 (2) (2002) 237–247.