



<b>Title</b>	The association of night-time systolic blood pressure with ultrasound markers of subclinical cardiac and vascular disease
<b>Author(s)</b>	O'Flynn, Anne Marie; Ho, Emily; Dolan, Eamon; Curtin, Ronan J.; Kearney, Patricia M.
<b>Publication date</b>	2017
<b>Original citation</b>	O'Flynn, A. M., Ho, E., Dolan, E., Curtin, R. J. and Kearney, P. M. (2017) 'The association of night-time systolic blood pressure with ultrasound markers of subclinical cardiac and vascular disease', Blood Pressure Monitoring, 22(1), pp. 18-26. doi: 10.1097/MBP.0000000000000223
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="https://journals.lww.com/bpmonitoring/Fulltext/2017/02000/The_association_of_night_time_systolic_blood.4.aspx">https://journals.lww.com/bpmonitoring/Fulltext/2017/02000/The_association_of_night_time_systolic_blood.4.aspx</a> <a href="http://dx.doi.org/10.1097/MBP.0000000000000223">http://dx.doi.org/10.1097/MBP.0000000000000223</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2017, the Authors. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal. <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a> <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/6342">http://hdl.handle.net/10468/6342</a>

Downloaded on 2019-01-07T05:57:23Z



# UCC

University College Cork, Ireland  
Coláiste na hOllscoile Corcaigh

# The association of night-time systolic blood pressure with ultrasound markers of subclinical cardiac and vascular disease

Anne Marie O'Flynn<sup>a</sup>, Emily Ho<sup>d</sup>, Eamon Dolan<sup>c</sup>, Ronan J. Curtin<sup>b</sup> and Patricia M. Kearney<sup>a</sup>

**Introduction** The aim of this study was to examine the association of night-time systolic blood pressure (BP) with subclinical cardiac dysfunction measured by global longitudinal strain (GLS) and subclinical vascular damage measured by carotid intima-media thickness (CIMT) and carotid plaques.

**Methods** GLS was measured by speckle-tracking analysis of echocardiogram images. CIMT was measured at the distal 1 cm of the common carotid artery. The presence of carotid plaques was recorded. Philips QLAB cardiac and vascular ultrasound quantification software was used for analysis. The association of night-time systolic BP with GLS, CIMT and carotid plaques was assessed using linear and logistic regression.

**Results** Fifty (response rate 63%) individuals took part in this study. In univariable models, night-time systolic BP was significantly associated with GLS [ $\beta$  coefficient 0.85 for every 10 mmHg increase, 95% confidence interval (CI): 0.3–1.4] and carotid plaques (odds ratio 1.9 for every 10 mmHg increase, 95% CI: 1.1–3.2). Univariable analysis of daytime systolic BP did not show any statistically significant associations. In age-adjusted and sex-adjusted models, the association for night-time systolic BP and GLS remained significant ( $\beta$  coefficient 0.68 for every 10 mmHg increase, 95% CI: 0.1–1.3). The association for carotid plaques was no

longer statistically significant. In multivariable models, findings were diminished.

**Discussion** Our results suggest a trend towards an association between night-time systolic BP and subclinical cardiac and vascular disease. When assessing ambulatory blood pressure monitoring results, the absolute night-time systolic BP seems to be a better prognostic parameter than daytime systolic BP, but ultimately a large randomised controlled trial involving chronotherapy is necessary to fully address this. *Blood Press Monit* 22:18–26 Copyright © 2017 Wolters Kluwer Health, Inc. All rights reserved.

*Blood Pressure Monitoring* 2017, 22:18–26

**Keywords:** ambulatory blood pressure monitoring, carotid intima-media thickness, global longitudinal strain, hypertension, ultrasonography

<sup>a</sup>Department of Epidemiology and Public Health, University College Cork, <sup>b</sup>Department of Cardiology, Cork University Hospital, Cork, <sup>c</sup>Department of Geriatrics, Connolly Hospital, Blanchardstown, Ireland and <sup>d</sup>Department of Cardiology, Sandwell and West Birmingham Hospitals NHS Trust, Birmingham, UK

Correspondence to Anne Marie O'Flynn, MB, BAO, BCh, Department of Epidemiology and Public Health, 4th Floor, Western Gateway Building, University College Cork, Western Road, Cork, Ireland  
Tel: +353 86 079 3133; fax: +353 21 420 5469; e-mail: am.oflynn@ucc.ie

Received 22 June 2016 Revised 22 September 2016

Accepted 23 September 2016

## Introduction

Cardiovascular disease remains the leading cause of mortality worldwide [1] and hypertension is the risk factor with the greatest population-attributable risk [2,3]. Prevalence rates of hypertension are high, whereas control rates are low [4]. Ambulatory blood pressure monitoring (ABPM) measures blood pressure (BP) over 24–48 h and has been shown to be superior to office BP for the prediction of clinical events [5,6]. Night-time

systolic BP is a stronger predictor of events than daytime systolic BP [7].

Although it is unclear whether night-time BP should be a specific therapeutic target, chronotherapy has shown some promising results. In one study, patients were randomized to either take all of their antihypertensive medications in the morning or to take at least one of them at night. The decrease in nocturnal BP was associated with a reduced risk of total cardiovascular events. Similarly, patients with chronic kidney disease who took at least one antihypertensive at night had a lower hazard ratio of total cardiovascular events than those taking all of their medications in the morning [8,9].

The association of night-time BP with subclinical target organ damage has been investigated [10,11]. This would seem intuitive, given the greater association of night-time

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website ([www.bpmonitoring.com](http://www.bpmonitoring.com)).

This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

BP with clinical events, the continuum of cardiovascular disease [12] and subclinical disease being a prognostic marker for future cardiovascular events [13]. Many studies have focused on dipping status rather than the absolute BP level [14–16]. We have previously shown that absolute night-time systolic BP is better associated than dipping status with subclinical cardiac and vascular damage documented by electrocardiogram left ventricular hypertrophy (LVH) voltage criteria and microalbuminuria in the Mitchelstown Cohort Study [17].

Echocardiography can measure the subclinical cardiac consequences of hypertension such as increased left atrial (LA) size and LVH. Speckle-tracking echocardiography has enabled the quantification of strain, which is a dimensionless measure of myocardial deformation. Global longitudinal strain (GLS) is a measure of the myocardial systolic deformation over the longitudinal axis [18]. There is emerging evidence for the prognostic importance of this measure [19,20]. It offers incremental prognostic information in the assessment of left ventricular (LV) function, particularly when the ejection fraction is near normal [21]. Few studies have examined the association of night-time BP and GLS measured by speckle-tracking analysis [22,23].

Carotid intima–media thickness (CIMT) measured by ultrasound is a marker of subclinical vascular damage and is recognized to be associated with cardiovascular risk factors and with the incidence of myocardial infarction and stroke [24,25]. There is evidence for the validity of CIMT as a suitable surrogate measure of atherosclerotic disease [26,27]. The addition of carotid plaques to risk prediction models including CIMT improves performance [28,29].

The present study aims to build on previous work by examining the association of night-time systolic BP with ultrasound markers of subclinical cardiovascular disease including abnormal GLS, CIMT and carotid plaques in a sample from the Mitchelstown Cohort Study.

## Methods

In 2010, the Mitchelstown Cohort Study recruited 2047 participants from a single large primary care centre, the Livinghealth Clinic in Mitchelstown, a town in the south of Ireland [30]. Of these, 1207 (response rate 59%) also underwent 24-h ABPM. These individuals provide the sample for the present study. On the basis of the initial ABPM results, the sample was divided into four groups: normotension, isolated nocturnal hypertension, isolated daytime hypertension and day–night hypertension. Twenty participants were selected randomly from each group and invited to attend for echocardiogram and carotid ultrasound in 2014. This study therefore includes analysis of baseline ABPM data and follow-up imaging data.

## Height and weight measurements

A trained researcher carried out the physical measures. Height and weight were measured without footwear using a Seca (Hamburg, Germany) measuring and weighing station. BMI and body surface area (BSA) were calculated.

## Study blood pressure

At the baseline visit after the participant had been in a relaxed seated position for at least 5 min, three BP readings were taken on the right arm, 1 min apart, using an OMRON M7 digital automatic BP monitor (OMRON Healthcare Europe B.V., Hoofddorp, the Netherlands). The average of the second and third BP reading was defined as the study BP.

## Ambulatory blood pressure monitoring

ABPM measurements were performed at baseline using the MEDITECH (Budapest, Hungary) ABPM-05 and data were stored using the dabl ABPM system (dabl Ltd, Dublin, Ireland). The monitors were programmed to record the BP every 30 min throughout the 24-h period. Participants kept a diary of the times they went to bed and got up. Diary times were used to calculate the mean daytime and night-time BP. The mean 24-h BP was calculated as the mean of all the readings throughout the 24-h period.

## Echocardiography

A Philips iE33 ultrasound machine (Philips Healthcare, Guildford, UK) and a S5-1 phased array transducer were used for image acquisition. All scans were carried out by the same operator (I.B.). I.B. is accredited in transthoracic echocardiography by the British Society of Echocardiography. A standard echocardiogram protocol was used. Parasternal long and short axis, apical four-chamber, two-chamber and three-chamber views were obtained. Optimization of frame rate was carried out by reducing the sector depth and width.

LV wall thickness and diameters were measured from the parasternal long axis view. LV mass was calculated using the Devereux formula [31]. LV mass was indexed for BSA. LVH was defined as LV mass more than 115 g/m<sup>2</sup> in men and more than 95 g/m<sup>2</sup> in women [32]. LV volumes and ejection fraction were calculated from the apical four-chamber view using the single-plane method of discs [33]. LV volumes were indexed for BSA. LA volume was calculated from the apical four-chamber view using the single-plane method of discs. LA volume was indexed for BSA [32].

Diastolic function parameters were measured in the apical four-chamber view. Mitral inflow early (*E*) and late (*A*) velocities and *E* wave deceleration time were obtained by pulse-wave Doppler with the sample volume at the mitral valve tips. Peak diastolic mitral annular (*e'*) velocity was measured from the septal and lateral mitral annulus and averaged. The *E/e'* ratio was then calculated. Diastolic dysfunction was defined as  $E/A \leq 0.7$  or

deceleration time  $> 260$  ms; or  $E/A > 0.7$  and  $\leq 1.5$  and  $e'$  velocity  $< 7$  cm/s; or  $E/A > 1.5$  and  $e'$  velocity  $< 7$  cm/s or deceleration time  $< 140$  ms [34].

The acquired images were saved using digital media and speckle-tracking GLS analysis was carried out offline by a single reader (A.M.O.F.) from apical four-chamber, two-chamber and three-chamber views using Philips QLAB cardiac and vascular ultrasound quantification software (version 9.0; Philips). The region of interest was identified by the selection of three points, one on either side of the mitral valve annulus and one at the apex for each view. Adequate tracking was confirmed visually and, if deemed inadequate, the region of interest was edited. If inadequate tracking persisted, problematic segments were excluded. If more than two segments in a single view had to be excluded, the entire study was excluded from speckle-tracking analysis [32]. GLS average was obtained from 17 ventricular segments represented on a bulls-eye plot from these views. Normal cut-off was taken as  $-19.7\%$  [35].

### Carotid ultrasound

A Philips Cx50 portable ultrasound machine and an L12-3 linear array transducer were used for image acquisition. All scans were carried out by the same operator (A.M.O.F.). A.M.O.F. received formal training in CIMT image acquisition. Patients were examined in the supine position with their head tilted to the opposite side. A Meijer arc was used to ensure optimal positioning. A thorough transverse and longitudinal scan of the extracranial carotid arteries was carried out to evaluate for the presence of atherosclerotic plaques. Ultrasound images of the distal portion of the far wall of both common carotid arteries were obtained for assessment of CIMT. Far-wall still frames were taken from anterior, lateral and posterior angles. Three still frame images were taken from each angle [36].

The acquired images were saved using digital media and measurement was carried out offline by a single reader (A.M.O.F.) using Philips QLAB cardiac and vascular ultrasound quantification software (version 9.0; Philips). Measurements were made on the three still frame images from each angle over a length of 1 cm at the distal common carotid artery (CCA). The reference point for the commencement of the measurement was where the CCA began to dilate before the bifurcation. A mean measurement from each angle was obtained. The mean of these means was obtained to yield the measurement for each side. The mean of the left and right was then taken as the CIMT. Normal cut-off was taken as the 75th percentile [37]. The presence of plaques in the extracranial carotid arteries was recorded as a binary variable. A plaque was defined according to the Mannheim Consensus as a focal protrusion into the blood vessel of at least 50% of the thickness of the adjacent IMT or focal IMT more than 1.5 mm [38].

### Reproducibility

Intraobserver and interobserver reproducibility was assessed before carrying out the study. Ten patients undergoing routinely indicated echocardiography were asked to consent to have their images analysed by speckle-tracking analysis. All echocardiograms were carried out by the same operator (I.B.). Speckle-tracking analysis was carried out offline by A.M.O.F. and repeated for intraobserver reproducibility. An independent observer (E.H.) also analysed the images for interobserver reproducibility. Ten healthy volunteers underwent a carotid ultrasound twice within the same week carried out by the same investigator (A.M.O.F.), who also analysed the images for intraobserver reproducibility. An independent observer (E.H.) analysed the second set of images for interobserver reproducibility.

The intraobserver intraclass coefficient (ICC) for GLS was 0.93 [95% confidence interval (CI): 0.70–0.98]. The interobserver ICC was 0.86 (95% CI: 0.51–0.96). The intraobserver ICC for CIMT was 0.91 (95% CI: 0.69–0.98). The interobserver ICC was 0.97 (95% CI: 0.88–0.99).

### Statistical analysis

Statistical analysis was carried out using Stata 12 (StataCorp LP, College Station, Texas, USA). Continuous variables are described by mean  $\pm$  SD. Categorical variables are described using proportions. Univariable linear and logistic regression analysis was used to compare echocardiographic and carotid ultrasound findings between groups.

The association of baseline night-time, daytime and study systolic BP with GLS and CIMT was assessed using univariable and multivariable linear regression. The association of baseline night-time, daytime and study systolic BP with carotid plaques was assessed using logistic regression. Regression models were initially adjusted for sex and age. Multivariable regression analysis was also carried out with adjustments applied for sex, age, BMI, smoking status, diabetes mellitus and total cholesterol. Interaction terms including antihypertensive medications with night-time, daytime and study systolic BPs, respectively, were also included in the models.

### Ethical considerations

The study was approved by the Clinical Research Ethics Committee of the Cork Teaching Hospitals and was carried out in accordance with the Declaration of Helsinki. All participants provided informed consent.

### Results

Fifty (overall response rate 63%) individuals took part in this study. The mean period of follow-up was 3.9 years. The mean age of the participants was 60 years and 26 (52%) were men. Baseline characteristics are presented in Table 1, together with the baseline characteristics of the full cohort and those who underwent ABPM.

Table 1 Baseline characteristics

	Total cohort (n = 2047)	Sample with satisfactory ABPM (n = 1072)	Sample (n = 50)
Age	59 ± 6	60 ± 6	60 ± 5
Male	1008 (49)	567 (47)	26 (52)
Education category			
Primary	537 (28)	335 (30)	7 (15.5)
Secondary	936 (49)	547 (48)	22 (49)
Tertiary	435 (23)	253 (22)	16 (35.5)
IPAQ category			
Low	932 (49)	571 (50)	31 (62)
Moderate	566 (30)	318 (28)	19 (38)
High	420 (22)	248 (22)	0 (0)
Smoking status			
Nonsmoker	1002 (51)	615 (53)	27 (54)
Former smoker	671 (34)	378 (32)	18 (36)
Current smoker	292 (15)	175 (15)	5 (10)
Medical history			
Hypertension	567 (29)	407 (34)	19 (40)
Myocardial infarction	49 (2)	33 (3)	2 (4)
Stroke	22 (1)	13 (1)	0 (0)
Heart failure	8 (0.4)	6 (0.5)	1 (2)
Diabetes	174 (9)	114 (10)	5 (10)
Medication			
Antihypertensive	584 (29)	405 (35)	21 (42)
Cholesterol lowering	711 (36)	457 (39)	21 (42)
BMI (kg/m <sup>2</sup> )	29 ± 5	30 ± 5	29 ± 5
Waist circumference	97 ± 13	97 ± 14	100 ± 13
LDL (mmol/l)	3.2 ± 0.9	3.2 ± 0.9	3.2 ± 0.8
Creatinine (µmol/l)	71 ± 16	72 ± 16	74 ± 17
ACR (mg/mmol)	0.7 ± 2.1	0.8 ± 2.1	0.6 ± 0.7
eGFR (mls/min)	90 ± 13	89 ± 13	87 ± 13
Cystatin C	0.83 ± 0.18	0.83 ± 0.18	0.87 ± 0.14
Study systolic	130 ± 17	134 ± 18	134 ± 16
Study diastolic	80 ± 10	83 ± 10	82 ± 10
Daytime systolic	–	131 ± 14	132 ± 12
Daytime diastolic	–	77 ± 9	78 ± 9
Night-time systolic	–	112 ± 14	117 ± 14
Night-time diastolic	–	63 ± 8	66 ± 9
Twenty-four hour systolic	–	124 ± 13	126 ± 11
Twenty-four hour diastolic	–	72 ± 8	74 ± 8

Note some missing data.

Values are given as mean ± SD or n (%).

ABPM, ambulatory blood pressure monitoring; ACR, albumin : creatinine ratio; eGFR, estimated glomerular filtration rate; IPAQ, International Physical Activity Questionnaire; LDL, low-density lipoprotein.

The echocardiogram and carotid ultrasound findings by BP strata are presented in Table 2. Speckle-tracking echocardiography analysis was not possible in one study because of poor image quality. In univariable models, night-time systolic BP was significantly associated with GLS ( $\beta$  coefficient 0.85 for every 10 mmHg increase, 95% CI: 0.3–1.4) and carotid plaques (odds ratio 1.9 for every 10 mmHg increase, 95% CI: 1.1–3.2). Univariable analysis of daytime systolic BP did not show any statistically significant associations (Table 3). In age-adjusted and sex-adjusted models, the association for night-time systolic BP and GLS remained significant ( $\beta$  coefficient 0.68 for every 10 mmHg increase, 95% CI: 0.11–1.25). The association for carotid plaques was no longer statistically significant (Table 4). In multivariable models, findings were attenuated (Table 5). Interaction terms including antihypertensive medications with night-time, daytime and study systolic BPs, respectively, were included in the

models, but were not statistically significant (data not shown).

## Discussion

We have shown an association of night-time systolic BP with two measures of subclinical cardiovascular disease, GLS and carotid plaques, in a community-based middle-aged population. These associations were attenuated in multivariable models. No such associations were found for daytime or study systolic BPs and subclinical cardiovascular disease.

Cumulative BP exposure over a 25-year period is associated with subclinical systolic and diastolic dysfunction assessed by speckle-tracking echocardiography in middle age [39]. Uncontrolled 24-h BP has also been shown to be associated with abnormal GLS, whereas uncontrolled office BP was not, in treated hypertensive patients [40]. Few studies have specifically addressed the association of night-time BP and GLS. Kalaycioglu *et al.* [22] found a significant reduction in GLS in nondippers compared with dippers in a sample of 86 treated hypertensive diabetic patients with a mean age of 57.8 years. They also reported night-time systolic BP to be associated independently with GLS and GLS rate in linear regression models adjusted for age, sex and LV mass index [22]. In untreated hypertensive patients, Tadic *et al.* [23] found reduced two-dimensional and three-dimensional LV GLS and reduced atrial longitudinal strain in nondippers compared with dippers. Acar *et al.* [41] also examined LA strain in dippers and nondippers and found reduced atrial function in nondippers. Others have examined right heart mechanics and found reduced function in nondippers [42]. Our results suggest an association between increased night-time systolic BP and subclinical LV systolic dysfunction. We also found night-time systolic BP to have a stronger association than daytime systolic BP and study BP with other echocardiographic markers of subclinical cardiac damage such as LA volume and LV mass (Supplementary Table 1, Supplemental digital content 1, <http://links.lww.com/BPMJ/A27>).

CIMT does improve cardiovascular risk prediction models, but the overall impact is small [43]. There are concerns regarding measurement methods [44]. Moreover, increasing IMT is recognized as a normal ageing phenomenon that further complicates interpretation of CIMT measurements [45–47]. The use of CIMT is therefore not without controversy and therefore European and American guidelines no longer recommend routine measurement of CIMT in clinical practice for the assessment of cardiovascular risk [48,49]. The addition of plaques has been shown to improve the predictive performance of CIMT [28,50]. Findings on the association of night-time BP and CIMT are conflicting. Cuspidi *et al.* [10] reported no difference in CIMT or plaques between those with nocturnal normotension and nocturnal hypertension. However, Wang *et al.* [51] reported an association between

**Table 2 Echocardiogram and carotid ultrasound characteristics**

	Total sample (n = 50)	Normotension (n = 11)	Isolated nocturnal hypertension (n = 14)	Isolated daytime hypertension (n = 14)	Sustained day–night hypertension (n = 11)
<b>Echocardiogram</b>					
LA volume index (ml/m <sup>2</sup> )	32.0 ± 9.4	27.7 ± 7.3	32.6 ± 9.7	31.3 ± 7.7	36.2 ± 12.0*
LV mass index (g/m <sup>2</sup> )	99.3 ± 24.2	85.5 ± 19.7	101.5 ± 24.7	97.8 ± 17.5	112.4 ± 29.9*
Left ventricular hypertrophy	16 (32)	2 (18)	5 (36)	4 (29)	5 (45)
LV end diastolic volume (ml/m <sup>2</sup> )	48.3 ± 8.4	45.8 ± 7.6	48.6 ± 10.3	48.2 ± 7.0	50.5 ± 8.7
LV end systolic volume (ml/m <sup>2</sup> )	17.2 ± 5.2	17.7 ± 6.8	17.5 ± 7.0	16.3 ± 2.9	17.6 ± 3.6
LV ejection fraction (%)	65.1 ± 6.2	64.5 ± 7.2	64.9 ± 7.5	66.0 ± 5.0	65.0 ± 5.3
Global longitudinal strain (%)	−21.2 ± 3.0	−22.6 ± 2.9	−21.0 ± 2.9	−21.9 ± 2.8	−19.2 ± 2.6*
Abnormal GLS	16 (33)	1 (10)	5 (36)	3 (21)	7 (64)
E/A	0.98 ± 0.3	1.04 ± 0.2	0.95 ± 0.4	0.89 ± 0.2	1.05 ± 0.3
E wave deceleration time (ms)	253.9 ± 74.2	224 ± 54	282 ± 90.1	255.4 ± 60.0	246 ± 80.3
E' (cm/s)	7.7 ± 1.8	8.2 ± 1.0	7.3 ± 2.0	7.8 ± 2.2	7.9 ± 1.6
E/e'	9.5 ± 1.9	9.4 ± 0.9	9.9 ± 1.9	8.6 ± 2.1	10.2 ± 2.3
Diastolic dysfunction	30 (60)	4 (36)	10 (71)	10 (71)	6 (55)
<b>Carotid ultrasound</b>					
Common carotid IMT (mm)	0.72 ± 0.13	0.64 ± 0.10	0.76 ± 0.14*	0.71 ± 0.14	0.76 ± 0.12*
Common carotid IMT > 75th percentile	12 (24)	1 (9)	4 (29)	3 (21)	4 (36)
Plaques	33 (66)	5 (45)	12 (86)*	8 (57)	8 (73)

GLS analysis not possible in one study because of poor image quality.

Values are given as mean ± SD or n (%).

GLS, global longitudinal strain; IMT, intima–media thickness; LA, left atrium; LV, left ventricle.

P values represent comparison with the normotensive group.

\*P < 0.05.

**Table 3 Univariable linear and logistic regression results**

	GLS [ $\beta$ coefficients (95% CI)]	P-value	CIMT [ $\beta$ coefficient (95% CI)]	P-value	Plaques [OR (95% CI)]	P-value
Sex (male vs. female)	2.70 (1.15–4.24)	0.001	0.09 (0.02–0.16)	0.02	6.5 (1.7–24.7)	0.006
Age	0.05 (−0.11–0.21)	0.5	0.009 (0.002–0.015)	0.009	1.2 (1.0–1.3)	0.03
Antihypertensives (yes vs. no)	−0.91 (−2.65–0.84)	0.3	−0.03 (−0.11–0.05)	0.4	0.3 (0.1–1.1)	0.06
BMI	0.20 (0.03–0.37)	0.02	−0.004 (−0.01–0.004)	0.3	1.1 (0.96–1.2)	0.2
Smoking status (current vs. non/ex)	−0.18 (−3.05–2.69)	0.9	0.18 (0.06–0.29)	0.004	2.2 (0.2–21.4)	0.5
Diabetes mellitus (yes vs. no)	3.16 (0.43–5.88)	0.02	0.02 (−0.11–0.15)	0.7	2.1 (0.2–20.9)	0.5
Total cholesterol	0.05 (−0.89–0.99)	0.9	−0.02 (−0.06–0.03)	0.4	0.67 (0.34–1.3)	0.2
Night-time SBP	0.85 (0.3–1.4)	0.003	0.02 (−0.002–0.05)	0.08	1.9 (1.1–3.2)	0.03
Daytime SBP	0.47 (−0.24–1.17)	0.2	0.02 (−0.01–0.05)	0.3	1.2 (0.7–2.0)	0.4
Study SBP	0.53 (0.01–1.06)	0.05	0.01 (−0.01–0.03)	0.4	1.16 (0.8–1.7)	0.4

$\beta$  Coefficients and odds ratios for night-time SBP, daytime SBP and study SBP represent per 10 mmHg increase.

CI, confidence interval; CIMT, carotid intima–media thickness; GLS, global longitudinal strain; OR, odds ratio; SBP, systolic blood pressure.

nocturnal hypertension and CIMT in patients with chronic kidney disease. Cuspidi *et al.* [52] have recently reported the results of a meta-analysis that examined the association of nondipping with carotid atherosclerosis and found higher CIMT and greater prevalence of plaques in nondippers. Although we found no association between night-time systolic BP and CIMT, we did find an association with carotid plaques in univariable analysis. We measured CIMT at the distal 1 cm of the common carotid artery as the guidelines recommend, while we assessed all of the extracranial carotid vessels for plaques [36]. This may have contributed towards the differential findings for CIMT and plaques as they likely reflect different stages of atherosclerosis and plaques are more likely to develop in areas of turbulent flow such as the bifurcation [53]. In addition, we may not have had sufficient power to detect an association between night-time BP and CIMT.

The evidence for the prognostic importance of night-time BP is compelling [7]. However, the potential underlying mechanisms are unclear and include altered sympathetic nervous system activity, disturbed baroreflex sensitivity, increased sodium sensitivity and obstructive sleep apnoea [7]. It may be that night-time BP is subject to less variability and more accurately represents true BP [54]. Reverse causality is also possible and elevated night-time BP may merely be a marker of more severe end organ damage. Cuspidi and colleagues have recently carried out a systematic review and meta-analysis on the association of nocturnal hypertension with subclinical cardiac and carotid disease documented by ultrasound and found increased LV mass index and CIMT in those with nocturnal hypertension compared with those with nocturnal normotension [55]. They acknowledge the cross-sectional nature of existing data

Table 4 Sex-adjusted and age-adjusted linear and logistic regression results

	GLS model 1		GLS model 2		GLS model 3	
	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value
Sex (male vs. female)	2.21 (0.64–3.77)	0.007	2.61 (0.89–4.33)	0.004	2.44 (0.64–4.23)	0.009
Age	−0.05 (−0.20–0.10)	0.5	0.01 (−0.14–0.16)	0.9	0.002 (−0.15–0.15)	1.0
Night-time SBP	0.68 (0.11–1.25)	0.02	–	–	–	–
Daytime SBP	–	–	0.08 (−0.61–0.78)	0.8	–	–
Study SBP	–	–	–	–	0.17 (−0.40–0.74)	0.6
	CIMT model 1		CIMT model 2		CIMT model 3	
	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value	[ $\beta$ coefficient (95% CI)]	<i>P</i> -value
Sex (male vs. female)	0.07 (−0.01–0.14)	0.07	0.07 (−0.01–0.14)	0.09	0.08 (−0.0003–0.16)	0.05
Age	0.01 (0.0001–0.01)	0.05	0.01 (0.001–0.01)	0.02	0.01 (0.001–0.01)	0.02
Night-time SBP	0.01 (−0.02–0.03)	0.6	–	–	–	–
Daytime SBP	–	–	0.01 (−0.02–0.04)	0.6	–	–
Study SBP	–	–	–	–	−0.005 (−0.03–0.02)	0.7
	Plaques model 1 [OR (95% CI)]		Plaques model 2 [OR (95% CI)]		Plaques model 3 [OR (95% CI)]	
	<i>P</i> -value	<i>P</i> -value	<i>P</i> -value	<i>P</i> -value	<i>P</i> -value	<i>P</i> -value
Sex (male vs. female)	4.8 (1.16–19.86)	0.03	6.11 (1.39–26.75)	0.02	8.4 (1.64–43.01)	0.01
Age	1.11 (0.97–1.28)	0.1	1.14 (0.99–1.31)	0.06	1.15 (1.00–1.32)	0.05
Night-time SBP	1.47 (0.80–2.69)	0.2	–	–	–	–
Daytime SBP	–	–	0.95 (0.55–1.65)	0.9	–	–
Study SBP	–	–	–	–	0.80 (0.49–1.30)	0.4

$\beta$  Coefficients and odds ratios for night-time SBP, daytime SBP and study SBP represent per 10 mmHg increase.

Model 1: night-time SBP adjusted for sex and age.

Model 2: daytime SBP adjusted for sex and age.

Model 3: study SBP adjusted for sex and age.

CI, confidence interval; CIMT, carotid intima–media thickness; GLS, global longitudinal strain; OR, odds ratio; SBP, systolic blood pressure.

and that the causal relationship between nocturnal hypertension and subclinical cardiovascular disease remains unproven. Our study provides some prospective data on subclinical target organ damage and our results suggest that elevated night-time BP may contribute more than daytime or office BP towards cardiac and vascular end organ damage. However, large prospective randomized trials with interventions aimed at normalizing night-time BP are required to resolve the questions that remain on the importance of night-time BP as a therapeutic target. The methods of a large prospective open-label blinded randomized-controlled trial have recently been published, which may help to answer this question. The treatment in morning versus evening trial aims to randomize 10 269 hypertensive patients to either morning or evening dosing of antihypertensive medications. The primary end-point is vascular death or hospitalization for the composite of nonfatal myocardial infarction or nonfatal stroke [56].

### Limitations

This is a small study and the sample size may have provided insufficient power to detect true associations between night-time systolic BP and target organ damage in multivariable models and for CIMT in particular. The ultrasound machines used were in fulltime clinical use and could only be used for research scans at the end of the working day; therefore, collection of imaging data was limited by access. This resulted in a prolonged data collection period. Given these feasibility issues, a decision was taken after 50 participants had been recruited to

proceed with data analysis. On the basis of our results, we had 86% power to detect a true association between night-time systolic BP and GLS. However, for CIMT, we had just 33% power and would have required a sample size of 162 to avoid a type II error [57].

Selection bias is another limitation as those who took part in this study were more likely to self-report a history of hypertension; therefore, the sample is not representative of the full Mitchelstown Cohort (40 vs. 29% previous doctor diagnosis of hypertension). Similarly, they were more likely to report being on antihypertensive medication at baseline (42 vs. 29%). Also those who took part in the study were better educated (35% had tertiary education compared with 23% of the full cohort) and less physically active (all of those taking part had low or moderate physical activity levels and nobody had high physical activity levels). In addition, the use of antihypertensive medications in the sample increased between 2010 and 2014 from 42% ( $n=21$ ) to 61% ( $n=30$ ), which may have impacted the results, although one would expect that increased use of antihypertensive medications would be more likely to influence results towards the null hypothesis. Although interaction terms did not suggest effect modification by antihypertensive therapy, we repeated the univariable regression analysis between night-time systolic BP and ultrasound markers of subclinical cardiovascular disease by treatment status. We found that the association between night-time systolic BP and subclinical cardiovascular disease was greater for untreated individuals (data not shown). This needs to

Table 5 Multivariable adjusted linear and logistic regression results

	GLS model 1		GLS model 2		GLS model 3	
	[ $\beta$ coefficient (95% CI)]	P-value	[ $\beta$ coefficient (95% CI)]	P-value	[ $\beta$ coefficient (95% CI)]	P-value
Sex (male vs. female)	1.79 (-0.03-3.6)	0.05	1.97 (0.03-3.9)	0.05	2.02 (0.03-4.01)	0.05
Age	0.002 (-0.17-0.17)	1.0	0.06 (-0.10-0.21)	0.5	0.06 (-0.10-0.22)	0.5
BMI	0.06 (-0.13-0.26)	0.5	0.11 (-0.08-0.30)	0.2	0.11 (-0.08-0.30)	0.2
Smoking (current vs. non/ex)	-0.01 (-2.82-2.80)	1.0	0.27 (-2.61-3.15)	0.9	0.28 (-2.60-3.16)	0.8
Diabetes mellitus (yes vs. no)	1.91 (-0.75-4.56)	0.2	2.16 (-0.57-4.89)	0.1	2.16 (-0.61-4.93)	0.1
Total cholesterol	0.28 (-0.62-1.17)	0.5	0.38 (-0.54-1.29)	0.4	0.38 (-0.54-1.31)	0.4
Night-time SBP	0.52 (-0.13-1.16)	0.1	-	-	-	-
Daytime SBP	-	-	0.04 (-0.66-0.75)	0.9	-	-
Study SBP	-	-	-	-	-0.01 (-0.61-0.58)	1.0
	CIMT model 1		CIMT model 2		CIMT model 3	
	[ $\beta$ coefficient (95% CI)]	P-value	[ $\beta$ coefficient (95% CI)]	P-value	[ $\beta$ coefficient (95% CI)]	P-value
Sex (male vs. female)	0.07 (-0.01-0.15)	0.07	0.07 (-0.01-0.15)	0.1	0.07 (-0.01-0.16)	0.08
Age	0.01 (-0.001-0.01)	0.08	0.01 (0.004-0.01)	0.04	0.01 (0.0003-0.1)	0.04
BMI	-0.003 (-0.01-0.004)	0.4	-0.003 (-0.01-0.004)	0.4	-0.0003 (-0.01-0.004)	0.5
Smoking (current vs. non/ex)	0.15 (0.03-0.27)	0.02	0.16 (0.04-0.28)	0.01	0.16 (0.04-0.28)	0.01
Diabetes mellitus (yes vs. no)	0.02 (-0.10-0.13)	0.8	0.02 (-0.09-0.14)	0.7	0.02 (-0.10-0.14)	0.7
Total cholesterol	-0.01 (-0.05-0.03)	0.5	-0.01 (-0.05-0.03)	0.6	-0.01 (-0.05-0.03)	0.6
Night-time SBP	0.01 (-0.02-0.03)	0.5	-	-	-	-
Daytime SBP	-	-	0.01 (-0.02-0.04)	0.5	-	-
Study SBP	-	-	-	-	0.001 (-0.02-0.95)	0.2
	Plaques model 1 [OR (95% CI)]		Plaques model 2 [OR (95% CI)]		Plaques model 3 [OR (95% CI)]	
	P-value	P-value	P-value	P-value	P-value	P-value
Sex (male vs. female)	4.66 (0.78-27.80)	0.09	5.70 (0.84-38.50)	0.07	6.57 (0.93-46.49)	0.06
Age	1.17 (0.99-1.38)	0.07	1.19 (1.01-1.40)	0.03	1.21 (1.02-1.42)	0.03
BMI	1.10 (0.93-1.30)	0.3	1.13 (0.95-1.33)	0.2	1.13 (0.96-1.34)	0.1
Smoking (current vs. non/ex)	6.81 (0.30-157.32)	0.2	10.28 (0.38-280.48)	0.2	9.16 (0.39-217.16)	0.2
Diabetes mellitus (yes vs. no)	0.61 (0.04-10.57)	0.7	0.74 (0.04-12.38)	0.8	0.93 (0.05-16.23)	1.0
Total cholesterol	0.63 (0.25-1.56)	0.3	0.69 (0.29-1.64)	0.4	0.70 (0.29-1.70)	0.4
Night-time SBP	1.31 (0.65-2.66)	0.5	-	-	-	-
Daytime SBP	-	-	0.86 (0.46-1.62)	0.6	-	-
Study SBP	-	-	-	-	0.78 (0.45-1.37)	0.4

$\beta$  Coefficients and odds ratios for night-time SBP, daytime SBP and study SBP represent per 10 mmHg increase.

Model 1: night-time SBP adjusted for sex, age, BMI, smoking, diabetes mellitus and total cholesterol.

Model 2: daytime SBP adjusted for sex, age, BMI, smoking, diabetes mellitus and total cholesterol.

Model 3: study SBP adjusted for sex, age, BMI, smoking, diabetes mellitus and total cholesterol.

CI, confidence interval; CIMT, carotid intima-media thickness; GLS, global longitudinal strain; OR, odds ratio; SBP, systolic blood pressure.

be interpreted with caution, given the small numbers included in the analysis. However, future studies should ideally recruit those who are not on antihypertensive therapy. Although our study provides some prospective imaging data in a community-based sample, the lack of ultrasound data at baseline recruitment means that it is not possible to draw inference on the temporal relationship for the observed associations of night-time systolic BP with GLS and carotid plaques.

In addition, night-time BP profiles are not fully reproducible [58,59]. We could assess the reproducibility of BP profiles in 47 of those who took part in this study. There were no significant differences in the mean BP levels between 2010 and 2014. However, the reproducibility of dipping status was poor, with just 24% maintaining the same profile after 4 years. The reproducibility of BP profiles defined by the absolute BP level was better, but still just 40% (Supplementary Tables 2-4, Supplemental digital content 2, <http://links.lww.com/BPMJ/A28>). These findings highlight the limitations of applying thresholds

and categories to continuously distributed risk factors but also the need to be cautious when interpreting results of studies examining nocturnal BP profiles on the basis of a single ABPM recording.

The strength of this study lies in the novel question addressed as limited studies have assessed the association of night-time BP and GLS [22,23]. In addition, these participants will continue to be followed up for the Mitchelstown Cohort Study; thus, further prospective data will be available over time.

## Conclusion

This study suggests a trend towards an association between night-time systolic BP with markers of sub-clinical cardiac and vascular disease, although the sample size limits interpretation of the multivariable analysis. We did not find any significant association between these markers and daytime systolic BP. When assessing ABPM results, the absolute night-time BP seems to be a better prognostic parameter than daytime systolic BP. However,



whether normalizing night-time BP improves prognosis remains unanswered and only a large randomized-controlled trial involving chronotherapy can address this.

## Acknowledgements

The authors wish to thank the study participants without whom this work would not be possible. They also thank Mr Ivan Bhatti, who carried out the echocardiograms.

## Conflicts of interest

Dr Anne Marie O'Flynn is currently receiving funding from a Health Research Board Ireland research training fellowship for healthcare professionals (reference HPF/2012/14) and has also received the John Feely research bursary from the Irish Heart Foundation to support this work. She has also received payment unrelated to the submitted work through her institution for the development of the European Society of Cardiology e-learning platform. Dr Ronan Curtin has received funding and subsistence support for lectures and activities outside of the submitted work from A. Menarini, Daichi Sankyo, Astra Zeneca, Bayer, Bristol Myers Squibb, Pfizer and Servier pharmaceutical companies. Professor Patricia Kearney has received grants from the Health Research Board Ireland and the European Union FP7 for work outside of the submitted work. Drs Emily Ho and Eamon Dolan have no conflicts of interest.

## References

- GBD 2013 Mortality and Causes of Death Collaborators. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *The Lancet* 2015; **385**:117–171.
- Roth GA, Nguyen G, Forouzanfar MH, Mokdad AH, Naghavi M, CJL Murray. Estimates of global and regional premature cardiovascular mortality in 2025. *Circulation* 2015; **132**:1270–1282.
- Poulter NR, Prabhakaran D, Caulfield M. Hypertension. *The Lancet* 2015; **386**:801–812.
- Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, *et al.* PURE (Prospective Urban Rural Epidemiology) Study investigators. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *JAMA* 2013; **310**:959–968.
- Dolan E, Stanton A, Thijs L, Hinedi K, Atkins N, McClory S, *et al.* Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension* 2005; **46**:156–161.
- Clement DL, de Buyzere ML, de Bacquer DA, de Leeuw PW, Duprez DA, Fagard RH, *et al.* Prognostic value of ambulatory blood-pressure recordings in patients with treated hypertension. *N Engl J Med* 2003; **348**:2407–2415.
- Hansen TW, Li Y, Boggia J, Thijs L, Richart T, Staessen JA. Predictive role of the night-time blood pressure. *Hypertension* 2011; **57**:3–10.
- Hermida RC, Ayala DE, Mojón A, Fernández JR. Decreasing sleep-time blood pressure determined by ambulatory monitoring reduces cardiovascular risk. *J Am Coll Cardiol* 2011; **58**:1165–1173.
- Hermida RC, Ayala DE, Mojón A, Fernández JR. Bedtime dosing of antihypertensive medications reduces cardiovascular risk in CKD. *J Am Soc Nephrol* 2011; **22**:2313–2321.
- Cuspidi C, Sala C, Valerio C, Negri F, Mancia G. Nocturnal blood pressure in untreated essential hypertensives. *Blood Press* 2011; **20**:335–341.
- Perez-Lloret S, Toblli JE, Cardinali DP, Malateste JC, Milei J. Nocturnal hypertension defined by fixed cut-off limits is a better predictor of left ventricular hypertrophy than non-dipping. *Int J Cardiol* 2008; **127**:387–389.
- Dzau VJ, Antman EM, Black HR, Hayes DL, Manson JE, Plutzky J, *et al.* The cardiovascular disease continuum validated: clinical evidence of improved patient outcomes: part I: pathophysiology and clinical trial evidence (risk factors through stable coronary artery disease). *Circulation* 2006; **114**:2850–2870.
- Mancia G, Fagard R, Narkiewicz K, Redon J, Zanchetti A, Bohm M, *et al.* 2013 ESH/ESC guidelines for the management of arterial hypertension: the Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 2013; **34**:2159–2219.
- Cuspidi C, Giudici V, Negri F, Sala C. Nocturnal nondipping and left ventricular hypertrophy in hypertension: an updated review. *Expert Rev Cardiovasc Ther* 2010; **8**:781–792.
- Pierdomenico SD, Lapenna D, Guglielmi MD, Costantini F, Romano F, Schiavone C, *et al.* Arterial disease in dipper and nondipper hypertensive patients. *Am J Hypertens* 1997; **10** (Pt 1):511–518.
- Roman MJ, Pickering TG, Schwartz JE, Cavallini MC, Pini R, Devereux RB. Is the absence of a normal nocturnal fall in blood pressure (nondipping) associated with cardiovascular target organ damage? *J Hypertens* 1997; **15**:969–978.
- O'Flynn AM, Dolan E, Curtin RJ, O'Brien E, Perry JJ, Kearney PM. Night-time blood pressure and target organ damage: a comparative analysis of absolute blood pressure and dipping status. *J Hypertens* 2015; **33**:2257–2264.
- Blessberger H, Binder T. Non-invasive imaging: two dimensional speckle tracking echocardiography: basic principles. *Heart* 2010; **96**:716–722.
- Russo C, Jin Z, Elkind MS, Rundek T, Homma S, Sacco RL, *et al.* Prevalence and prognostic value of subclinical left ventricular systolic dysfunction by global longitudinal strain in a community-based cohort. *Eur J Heart Fail* 2014; **16**:1301–1309.
- Holland DJ, Marwick TH, Haluska BA, Leano R, Hordern MD, Hare JL, *et al.* Subclinical LV dysfunction and 10-year outcomes in type 2 diabetes mellitus. *Heart* 2015; **101**:1061–1066.
- Kalam K, Otahal P, Marwick TH. Prognostic implications of global LV dysfunction: a systematic review and meta-analysis of global longitudinal strain and ejection fraction. *Heart* 2014; **100**:1673–1680.
- Kalaycioglu E, Gokdeniz T, Aykan AC, Gul I, Ugur M, Gursoy OM, *et al.* The influence of dipper/nondipper blood pressure patterns on global left ventricular systolic function in hypertensive diabetic patients: a speckle tracking study. *Blood Press Monit* 2014; **19**:263–270.
- Tadic M, Cuspidi C, Majstorovic A, Slijivic A, Pencic B, Ivanovic B, *et al.* Does a nondipping pattern influence left ventricular and left atrial mechanics in hypertensive patients? *J Hypertens* 2013; **31**:2438–2446.
- O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson SK Jr. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. Cardiovascular Health Study Collaborative Research Group. *N Engl J Med* 1999; **340**:14–22.
- Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, *et al.* Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: The Atherosclerosis Risk in Communities (ARIC) Study, 1987–1993. *Am J Epidemiol* 1997; **146**:483–494.
- Peters SA, Grobbee DE, Bots ML. Carotid intima-media thickness: a suitable alternative for cardiovascular risk as outcome? *Eur J Cardiovasc Prev Rehabil* 2011; **18**:167–174.
- Iwakiri T, Yano Y, Sato Y, Hatakeyama K, Marutsuka K, Fujimoto S, *et al.* Usefulness of carotid intima-media thickness measurement as an indicator of atherosclerosis: findings from autopsy analysis. *Atherosclerosis* 2012; **225**:359–362.
- Nambi V, Chambless L, Folsom AR, He M, Hu Y, Mosley T, *et al.* Carotid intima-media thickness and presence or absence of plaque improves prediction of coronary heart disease risk: the ARIC (Atherosclerosis Risk in Communities) study. *J Am College of Cardiol* 2010; **55**:1600–1607.
- Polak JF, Szklo M, Kronmal RA, Burke GL, Shea S, Zavodni AEH, *et al.* The value of carotid artery plaque and intima-media thickness for incident cardiovascular disease: the multi-ethnic study of atherosclerosis. *J Am Heart Assoc* 2013; **2**:e000087.
- Kearney PM, Harrington JM, Mc Carthy VJ, Fitzgerald AP, Perry JJ. Cohort Profile: The Cork and Kerry Diabetes and Heart Disease Study. *Int J Epidemiol* 2013; **42**:1253–1262.
- Devereux RB, Reichek N. Echocardiographic determination of left ventricular mass in man. Anatomic validation of the method. *Circulation* 1977; **55**:613–618.
- Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, *et al.* Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2015; **16**:233–270.

- 33 Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, *et al.* Recommendations for chamber quantification. *Eur J Echocardiogr* 2006; **7**:79–108.
- 34 Russo C, Jin Z, Sera F, Lee ES, Homma S, Rundek T, *et al.* Left ventricular systolic dysfunction by longitudinal strain is an independent predictor of incident atrial fibrillation: a community-based cohort study. *Circ Cardiovasc Imaging* 2015; **8**:e003520.
- 35 Yingchoncharoen T, Agarwal S, Popovic ZB, Marwick TH. Normal ranges of left ventricular strain: a meta-analysis. *J Am Soc Echocardiogr* 2013; **26**:185–191.
- 36 Stein JH, Korcarz CE, Hurst RT, Lonn E, Kendall CB, Mohler ER, *et al.* Use of carotid ultrasound to identify subclinical vascular disease and evaluate cardiovascular disease risk: a consensus statement from the American Society of Echocardiography carotid intima–media thickness task force. Endorsed by the society for vascular medicine. *J Am Soc Echocardiogr* 2008; **21**:93–111.
- 37 Aminbakhsh A, Mancini GB. Carotid intima–media thickness measurements: what defines an abnormality? a systematic review. *Clin Invest Med* 1999; **22**:149–157.
- 38 Touboul PJ, Hennerici MG, Meairs S, Adams H, Amarenco P, Bornstein N, *et al.* Mannheim carotid intima–media thickness and plaque consensus (2004–2006–2011). An update on behalf of the advisory board of the 3rd, 4th and 5th Watching the Risk symposia, at the 13th, 15th and 20th European Stroke Conferences, Mannheim, Germany, 2004, Brussels, Belgium, 2006, and Hamburg, Germany, 2011. *Cerebrovasc Dis* 2012; **34**:290–296.
- 39 Kishi S, Teixido-Tura G, Ning H, Venkatesh BA, Wu C, Almeida A, *et al.* Cumulative blood pressure in early adulthood and cardiac dysfunction in middle age: The CARDIA Study. *J Am Coll Cardiol* 2015; **65**:2679–2687.
- 40 Sera F, Jin Z, Russo C, Lee ES, Schwartz JE, Rundek T, *et al.* Ambulatory blood pressure control and subclinical left ventricular dysfunction in treated hypertensive subjects. *J Am Coll Cardiol* 2015; **66**:1408–1409.
- 41 Acar G, Bulut M, Arslan K, Alizade E, Ozkan B, Alici G, *et al.* Comparison of left atrial mechanical function in nondipper versus dipper hypertensive patients: a speckle tracking study. *Echocardiography* 2013; **30**:164–170.
- 42 Tadic M, Cuspidi C, Pencic B, Ivanovic B, Scepanovic R, Marjanovic T, *et al.* Circadian blood pressure pattern and right ventricular and right atrial mechanics: a two- and three-dimensional echocardiographic study. *J Am Society of Hypertens* 2014; **8**:45–53.
- 43 Den Ruijter HM, Peters SA, Anderson TJ, Britton AR, Dekker JM, Eijkemans MJ, *et al.* Common carotid intima–media thickness measurements in cardiovascular risk prediction: a meta-analysis. *JAMA* 2012; **308**:796–803.
- 44 Peters SA, Bots ML. Carotid intima–media thickness studies: study design and data analysis. *J Stroke* 2013; **15**:38–48.
- 45 Mancini GBJ, Dahlöf B, Diez J. Surrogate markers for cardiovascular disease: structural markers. *Circulation* 2004; **109** (Suppl 1):IV22–IV30.
- 46 Lakatta EG, Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: part I: aging arteries: a 'set up' for vascular disease. *Circulation* 2003; **107**:139–146.
- 47 Najjar SS, Scuteri A, Lakatta EG. Arterial aging: is it an immutable cardiovascular risk factor? *Hypertension* 2005; **46**:454–462.
- 48 Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, *et al.* 2016 European Guidelines on cardiovascular disease prevention in clinical practice: The Sixth Joint Task Force of the European Society of Cardiology and Other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of 10 societies and by invited experts). Developed with the special contribution of the European Association for Cardiovascular Prevention & Rehabilitation (EACPR). *Eur Heart J* 2016; **37**:2315–2381.
- 49 Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB Sr, Gibbons R, *et al.* 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014; **63**:2935–2959.
- 50 Inaba Y, Chen JA, Bergmann SR. Carotid plaque, compared with carotid intima–media thickness, more accurately predicts coronary artery disease events: a meta-analysis. *Atherosclerosis* 2012; **220**:128–133.
- 51 Wang C, Deng WJ, Gong WY, Zhang J, Zhang QZ, Ye ZC, Lou T. Nocturnal hypertension correlates better with target organ damage in patients with chronic kidney disease than a nondipping pattern. *J Clin Hypertens (Greenwich)* 2015; **17**:792–801.
- 52 Cuspidi C, Sala C, Tadic M, Gherbesi E, Grassi G, Mancia G. Nondipping pattern and carotid atherosclerosis: a systematic review and meta-analysis. *J Hypertens* 2016; **34**:385–392.
- 53 Finn AV, Kolodgie FD, Virmani R. Correlation between carotid intimal/medial thickness and atherosclerosis: a point of view from pathology. *Arterioscler Thromb Vasc Biol* 2010; **30**:177–181.
- 54 Pickering TG. What is the true blood pressure? Smirk revisited. *J Clin Hypertens* 2005; **7**:421–424.
- 55 Cuspidi C, Sala C, Tadic M, Gherbesi E, Grassi G, Mancia G. Nocturnal hypertension and subclinical cardiac and carotid damage: an updated review and meta-analysis of echocardiographic studies. *J Clin Hypertens (Greenwich)* 2016; **18**:913–920.
- 56 Rorie DA, Rogers A, Mackenzie IS, Ford I, Webb DJ, Williams B, *et al.* Methods of a large prospective, randomised, open-label, blinded end-point study comparing morning versus evening dosing in hypertensive patients: the treatment in morning versus evening (TIME) study. *BMJ Open* 2016; **6**: e010313.
- 57 Dupont WD, Plummer WD Jr. Power and sample size calculations for studies involving linear regression. *Control Clin Trials* 1998; **19**:589–601.
- 58 Cuspidi C, Meani S, Salerno M, Valerio C, Fusi V, Severgnini B, *et al.* Reproducibility of nocturnal blood pressure fall in early phases of untreated essential hypertension: a prospective observational study. *J Human Hypertens* 2004; **18**:503–509.
- 59 White WB, Larocca GM. Improving the utility of the nocturnal hypertension definition by using absolute sleep blood pressure rather than the 'dipping' proportion. *Am J Cardiol* 2003; **92**:1439–1441.