

On some species of *Astyanax* reported erroneously from Argentina.

Juan Marcos Mirande¹ & Stefan Koerber²

¹ CONICET-Fundación Miguel Lillo, Miguel Lillo 251, 4000 Tucumán, Argentina | mcmirande@gmail.com

² Friesenstr. 11, 45476 Muelheim, Germany | pecescrillos@koerber-germany.de

Abstract

Several species of *Astyanax* have been reported from Argentina which currently are considered not to be distributed so far South or whose identity as *Astyanax* is doubtful. These records have been copied during decades based on historic accounts or erroneous determination. We conclude that *Astyanax bimaculatus*, *A. fasciatus*, *A. paranahybae*, *A. scabripinnis*, and *A. taeniatus* shall be excluded from the faunal lists of Argentina and, at least in the case of *A. fasciatus*, also of Uruguay.

Resumen

Varias especies de *Astyanax* han sido reportadas para Argentina, de las que actualmente no se considera que estén distribuidas tan al sur o cuya identidad en *Astyanax* es dudosa. Estos registros han sido repetidos durante décadas basándose en cuestiones históricas o identificaciones erróneas. Nosotros concluimos que *Astyanax bimaculatus*, *A. fasciatus*, *A. paranahybae*, *A. scabripinnis* y *A. taeniatus* deberían ser excluidas de las listas de fauna de Argentina y, al menos en el caso de *A. fasciatus*, también de Uruguay.

Introduction

Astyanax is the most diverse genus of freshwater fishes inhabiting Argentina. Most of its diversity in this country was described in the last 20 years, in which this genus was intensely studied across most basins and valuable information was produced on some problematic species complexes. Especially, advances on the resolution of the *A. eigenmanniorum* complex, with detailed comparisons with type material and topotypes, permitted the description of many species of the genus (e.g. *Astyanax chico*, *A. endy*, *A. hermosus*, *A. puka*, and *A. troya*). Also, some species not assignable to any complex were discovered and described recently (e.g. *A. latens*, *A. paris*).

However, this high and growing known diversity of the genus in Argentina is also positively biased by species that are cited but are not present in this country. Those wrong citations are either due to repeated mentions of originally misidentified specimens or to lack of nomenclatural updates after taxonomic changes proposed in literature.

The aim of this paper is to discuss the situation of species of *Astyanax* usually cited in Argentina but, according to our observations and/or recent literature, are not present in this country and should be removed from lists of species.

Astyanax bimaculatus (Linnaeus, 1758)

original	<i>Salmo bimaculatus</i>
type locality	America Meridionali
synonyms	<i>Tetragonopterus maculatus</i> Müller & Troschel, 1845 Brasilia, Guiana et Surinam ? <i>Tetragonopterus gronovii</i> Valenciennes, 1850 Surinam; rivières de La Guyane ? <i>Tetragonopterus linnaei</i> Valenciennes, 1850 Cayenne [French Guiana] ? <i>Tetragonopterus wappi</i> Valenciennes, 1850 [Essequibo River, Guyana] ? <i>Tetragonopterus vittatus</i> Castelnau, 1855 Bahia [Brazil] ? <i>Poecilurichthys brevoortii</i> Gill, 1858 [Island of] Trinidad <i>Tetragonopterus bartletti</i> Günther, 1866 Upper Amazon ? <i>Tetragonopterus orientalis</i> Cope, 1870 Pará [Brazil]

- ?*Tetragonopterus bahiensis* Steindachner, 1877 | Bahia, Brazil
Astyanax bimaculatus borealis Eigenmann, 1908 | Rio Magdalena and Cauca
Astyanax bimaculatus novae Eigenmann, 1911 | Rio Sapon, Prazeres, Bahia [Brazil]
Astyanax rupununi Fowler, 1914 | Rupununi River, British Guiana
Astyanax bimaculatus incaicus Tortonese, 1942 | Rio Zamora, Ecuador

Ringuelet et al. (1967) and López et al. (2003) mentioned as first record for Argentina 'Cuvier & Valenciennes (1848) sub *Tetragonopterus orbignyanus*, in part'. *Astyanax orbignyanus* (Valenciennes, 1850) was described from La Plata and other localities in Argentina and is currently considered to be a valid species by Buckup (2003). The type material of *A. orbignyanus* is still available in Paris (Bertin 1948) and since the times of Eigenmann probably nobody has examined these specimens.

Eigenmann (1921) treated *A. orbignyanus* as a synonym of two different congeners and assigned part of the type material to *Astyanax abramis* and part to *A. bimaculatus* [the nominal subspecies]. Pozzi (1945), Ringuelet & Arámburu (1962), Ringuelet et al. (1967), López et al. (1987) and López et al. (2003) followed Eigenmann (1921) or copied from each other and thus initiated the ongoing erroneous listings of *Astyanax bimaculatus* as a species from Argentina. In the same paper Eigenmann (1921) described the new subspecies *Astyanax bimaculatus paraguayensis*, raising automatically *A. bimaculatus* to the rank of the subspecies *A. b. bimaculatus*. This new subspecies was listed by Pozzi (1945), Ringuelet et al. (1967), and López et al. (1987). As *Astyanax bimaculatus paraguayensis* was preoccupied in *Astyanax* by *Bertoniolus paraguayensis* Fowler, 1918 a replacement name was proposed by Géry (1972): *Astyanax bimaculatus asuncionensis*.

López et al. (2003) did not mention *Astyanax bimaculatus paraguayensis*, *Astyanax bimaculatus asuncionensis* or *Astyanax asuncionensis*. Since then, *Astyanax asuncionensis* Géry, 1972 was treated as a valid species from Argentina by Azpelicueta et al. (2003), Casciotta et al. (2003), Mirande et al. (2004), Miquelarena & Menni (2005), Miquelarena et al. (2005), Mirande et al. (2006), Mirande et al. (2007), Almirón et al. (2010), Mirande (2010), and Soneira et al. (2010). Miquelarena & Menni (2005) did not include *A. bimaculatus* in their key to the *Astyanax* from Argentina. The first record for Uruguay was done by Messner (1962) sub *A. b. bimaculatus*. Nion et al. (2002) listed *A. bimaculatus*. Serra (2012) has excluded *A. bimaculatus* from the faunal list of Uruguay. Also, Buckup (2003) mentioned the distribution of *A. bimaculatus* as "Panama to Amazon basin". *Astyanax asuncionensis* is readily distinguished from *A. bimaculatus* by the absence of maxillary teeth (vs. presence). We assume that most of the Argentinean records of *Astyanax bimaculatus* can be assigned to *Astyanax asuncionensis* Géry, 1972.

***Astyanax fasciatus* (Cuvier, 1819)**

original	<i>Chalceus fasciatus</i>
type locality	Brazil
synonyms	<p><i>Tetragonopterus viejita</i> Valenciennes, 1850 Laguna de Maracaibo, Venezuela</p> <p><i>Salmo lambari</i> Natterer, 1859 Irisanga, Brazil</p> <p><i>Tetragonopterus aeneus</i> Günther, 1860 freshwaters of Oaxaca, Mexico</p> <p><i>Tetragonopterus humilis</i> Günther, 1864 Lake Amatitlan, Guatemala</p> <p>? <i>Tetragonopterus microphthalmus</i> Günther, 1864 Pacific coast of Guatemala</p> <p><i>Tetragonopterus panamensis</i> Günther, 1864 Pacific coast of Panama</p> <p><i>Tetragonopterus belizianus</i> Bocourt, 1868 environments de Belize</p> <p><i>Tetragonopterus cobanensis</i> Bocourt, 1868 </p> <p><i>Tetragonopterus finitimus</i> Bocourt, 1868 </p> <p><i>Tetragonopterus oaxacanensis</i> Bocourt, 1868 </p> <p><i>Astyanax carolinae</i> Gill, 1870 </p> <p><i>Tetragonopterus cuvieri</i> Lütken, 1875 </p> <p><i>Tetragonopterus oerstedii</i> Lütken, 1875 </p> <p><i>Tetragonopterus rutilus</i> var. <i>jequitinhonhae</i> Steind., 1877 </p> <p><i>Astyanax albeolus</i> Eigenmann, 1908 </p> <p>? <i>Tetragonopterus macrophthalmus</i> Regan, 1908 </p> <p><i>Astyanax regani</i> Meek, 1909 </p> <p><i>Astyanax grandis</i> Meek & Hildebrand, 1912 </p> <p><i>Astyanax heterurus</i> Eigenmann & Wilson, 1914 </p> <p><i>Astyanax aeneus</i> var. <i>costaricensis</i> Meek, 1914 </p>

? *Astyanax hanstroemi* Dahl, 1943 |
Astyanax fasciatus orteguasae Fowler, 1943 |

Astyanax fasciatus is actually a complex of species whose diversity is far to be completely discovered yet. Apparently this species has been cited from Argentina since Cuvier & Valenciennes (1849) have published their doubts about the identities of *Astyanax rutilus* and *A. fasciatus*.

The type locality of *A. rutilus* is in the Río Paraná, whereas *Astyanax fasciatus* was primarily described from the Rio São Francisco basin in Brazil, but subsequently cited for most basins from Mexico to central Argentina.

Melo & Buckup (2006) restricted *Astyanax fasciatus* to the Rio São Francisco basin, leaving most other populations as *A. aff. fasciatus*. They said that *A. rutilus* is probably a valid species and mentioned some differences in the number of vertebrae, supporting that idea. Also, Melo & Buckup (2006: fig. 5) showed a male specimen of *A. fasciatus* having an elongation of the dorsal fin that was neither observed by us nor mentioned in any population of *Astyanax* living in Argentina. Therefore, since that paper, specimens from Argentina once identified as *A. fasciatus* are named *A. rutilus* by most authors. Protogino et al. (2006) described *Astyanax aramburui*, which distinguishes from *A. rutilus* mainly by the presence of bony hooks in all fins of males. At least until a revision of the *A. fasciatus* complex might be conducted, we assume that the specimens from Argentina and Uruguay once identified as *A. fasciatus* are either *A. aramburui* or *A. rutilus*.

***Astyanax paranahybae* Eigenmann, 1911**

type locality	Rio Paranahyba, Brazil
distribution	Paraná River basin, Brazil

This species was cited by Ringuelet et al. (1967) based on a single examined specimen from Santa Fe city, with no collection number. Their citation was subsequently repeated by several other authors (e.g. Liotta, 2005) but no additional material of *A. paranahybae* was cited for Argentina. According to Ringuelet et al. (1967), meristic counts of the specimen they examined are similar to *A. eigenmanniorum*, but it has 3-4 maxillary teeth (instead of 1).

Astyanax paranahybae is known by a single specimen, which was examined by Garutti & Britski (2000). They mentioned that the posterior premaxillary row has 4 teeth in the left premaxilla and 5 in the right one and that it has only 7 branched dorsal-fin rays (vs. 9 in other species of *Astyanax*). Vari & Castro (2007) discussed the validity of *A. paranahybae* and suggested that it could be actually a member of *Bryconamericus* or *Piabina*, according to these data and the overall form of body.

There are just a few species of *Astyanax* in Argentina having 3 maxillary teeth, but no one has the combination of characters mentioned by Ringuelet et al. (1967). However, is not clear if the characters mentioned by Ringuelet et al. (1967) were observed by them or if they just copied the diagnosis by Eigenmann (1921). Ringuelet et al. (1967) did not mention how many premaxillary teeth and dorsal-fin rays had the specimen they examined, but the combination of characters provided by them is compatible with the discussion by Vari & Castro (2007) about this species. Therefore, and considering that no other specimen of this species was consequently cited, we consider *Astyanax paranahybae*, if valid, to be absent in Argentina. Even it is possible that the material examined by Ringuelet et al. (1967) was actually a specimen of *Bryconamericus*.

***Astyanax scabripinnis* (Jenyns, 1842)**

original	<i>Tetragonopterus scabripinnis</i>
type locality	Rio de Janeiro, Brazil

The subspecies *Astyanax scabripinnis paranae* was treated as an 'especie limítrofe', meaning 'from neighbouring countries', by Ringuelet et al. (1967) despite the fact that they cited Pozzi (1945) as the first record from Argentina. López et al. (2003) mention as first record Fernández Santos et al. (1987) who reported *Astyanax scabripinnis paranae* from the El Palmar National Park in Entre Ríos (fig. 1). Additionally they referred to more specimens reported by Miquelarena et al. (1997) from the Uruguaí stream in the Paraná basin of Misiones (fig. 2). Actually in both, the mentioned paper of Miquelarena

et al. (1997) and in Miquelarena et al. (2009) these specimens were addressed as the nominal species *Astyanax scabripinnis* and not as the subspecies *A. s. paranae* as indicated by López et al. (2003). *Astyanax paranae* Eigenmann, 1914 was treated as a valid species by Buckup (2003) and the record by Fernández Santos et al. (1987) would thus be the first one from Argentina for this species. These specimens should be existing yet at the MACN and deserve an examination. Anyhow, if treated as the subspecies as listed by López et al. (2003) and considered currently as a valid species this record cannot refer to *Astyanax scabripinnis*.

The only remaining record for *Astyanax scabripinnis* is based on the specimens reported from the Uruguaí subbasin in Misiones. The Uruguaí stream was separated from the Río Paraná by a waterfall until the construction of a hydroelectric dam and has developed its own ichthyofauna, different at least in part from the ones found in the ríos Paraná and Iguazú.

For the specimens from the Uruguaí we consider that these might have been confused with a later described species from the *Astyanax scabripinnis* complex sensu Ingenito & Duboc (2014). Considering that the arroyo Uruguaí shows an extremely diverse composition of species, possessing species endemic to this subbasin, sharing some with the río Paraná and others with the río Iguazú basin above the falls, the two species which might be candidates for such a confusion are *Astyanax leonidas* from the Uruguaí itself and *A. troya* from the río Paraná. Either would more probably be found in the Uruguaí than the real *A. scabripinnis* or *A. paranae*. Until an examination of the above mentioned specimens proves us wrong, *A. scabripinnis* should not be listed in faunal lists for Argentina.




fig. 1. *Astyanax scabripinnis paranae* from Fernández Santos et al. (1987)



fig. 2. *Astyanax scabripinnis* from Miquelarena et al. (1997)

Species of *Astyanax* reported from Argentina in the chronological order of their formal descriptions.

	Ringuelet et al. (1967)	López et al. (2003)	Miquelarena & Menni (2005)	valid for Argentina
<i>A. bimaculatus</i> (Linnaeus, 1758)	X	X	-	-
<i>A. fasciatus</i> (Cuvier, 1819)	X	X	X	-
<i>A. abramis</i> (Jenyns, 1842)	X	X	X	X
<i>A. rutilus</i> (Jenyns, 1842)	-	-	-	X
<i>A. scabripinnis</i> (Jenyns, 1842)	-	X	X	-
<i>A. taeniatus</i> (Jenyns, 1842)	-	X	-	-
<i>A. orbignyanus</i> Cuvier & Valenciennes, 1850	-	-	?	X
<i>A. cordovae</i> (Günther, 1880)	X	X	X	X
<i>A. correntinus</i> (Holmberg, 1891)	X	X	X	X
<i>A. lineatus</i> (Perugia, 1891)	X	X	X	X
<i>A. erythropterus</i> (Holmberg, 1891)	X	X	X	X
<i>A. eigenmanniorum</i> (Cope, 1894)	X	X	X	X
<i>A. jacuhiensis</i> (Cope, 1894)	-	-	?	?
<i>A. allenii</i> (Eigenmann & Mc Atee, 1907)	X	-	X	X
<i>A. pelegrini</i> Eigenmann, 1907	X	X	X	X
<i>A. gymnogenys</i> Eigenmann, 1911	X	X	X	X
<i>A. paranahybae</i> Eigenmann, 1911	X	X	-	-
<i>A. paranae</i> Eigenmann, 1914	X	-	-	-
<i>A. stenocephalus</i> Messner, 1962	-	-	-	X
<i>A. asuncionensis</i> Gery, 1972 (incl. <i>A. bimaculatus</i> parag.)	X	-	X	X
<i>A. ojara</i> Azpelicueta & García, 2000		X	X	X
<i>A. ita</i> Almirón, Azpelicueta & Casciotta, 2002		X	X	X
<i>A. leonidas</i> Azpelicueta, Casciotta & Almirón, 2002		X	X	X
<i>A. paris</i> Azpelicueta, Almirón & Casciotta, 2002		X	X	X
<i>A. troya</i> Azpelicueta, Casciotta & Almirón, 2002		X	X	X
<i>A. pynandi</i> Casciotta, Almirón et al. 2003			X	X
<i>A. saguazu</i> Casciotta, Almirón & Azpelicueta, 2003			X	X
<i>A. tupi</i> Azpelicueta et al., 2003			X	X
<i>A. latens</i> Mirande, Aguilera & Azpelicueta, 2004			X	X
<i>A. chico</i> Casciotta & Almirón, 2004			-	X
<i>A. hermosus</i> Miquelarena, Protogino & López, 2005			X	X
<i>A. tumbyaensis</i> Miquelarena & Menni, 2005			X	X
<i>A. pampa</i> Casciotta, Almirón & Azpelicueta, 2005				X
<i>A. endy</i> Mirande, Aguilera & Azpelicueta, 2006				X
<i>A. aramburui</i> Protogino, Miquelarena & López, 2006				X
<i>A. puka</i> Mirande, Aguilera & Azpelicueta, 2007				X

***Astyanax taeniatus* (Jenyns, 1842)**

original *Tetragonopterus taeniatus*
 type locality Socego, Rio de Janeiro state, Brazil

López et al. (2003) assigned the first record of this species to Géry (1978) from the South of Buenos Aires province. Géry actually only informed about the fact that Lueling had found *A. taeniatus* in a laguna near Guaminí. In 1981 Lueling himself published a report about the collection he had conducted in late 1975 in the lakes of Guaminí, known as the Lagunas Encadenas del Oeste.




fig. 3. *Astyanax taeniatus* from Lueling (1981)

He provided a photo (fig. 3) of a specimen he had determined as *A. taeniatus*, but no further details or data. The only members of the genus subsequently reported from localities South of the río Salado basin are *Astyanax eigenmanniorum* (e.g. Almirón et al. 1997, Casciotta et al. 1999) and *A. pampa* (Casciotta et al. 2005, Fulvio Pérez 2008). Miquelarena & López (1995) conducted another ichthyological inventory of the Lagunas Encadenadas del Oeste and recorded *A. cf. eigenmanniorum*, but explicitly stated that they could not confirm the presence of *A. taeniatus*. Buckup (2003) restricted the distribution of *A. taeniatus* to the “Paraíba do Sul River drainage and coastal rivers of Rio de Janeiro and Espírito Santo States” in Brazil.

Combining the above given information and despite the fact that so far we had no opportunity to examine Lueling's voucher material, if still available, we tentatively remove *Astyanax taeniatus* from the faunal list of freshwater fishes found in Argentina.

References

- Almirón, A., M. Azpelicueta, J. Casciotta & A. López Cazorla (1997): Ichthyogeographic boundary between the Brazilian and Austral subregions in South America. *Biogeographica* 73 (1): 23-30
- Almirón, A.E., J.R. Casciotta, M. Azpelicueta & M. Loureiro (2010): Redescription of *Astyanax stenohalinus* Messner, 1926 (Characiformes: Characidae), a poorly known species from Argentina and Uruguay. *Zootaxa* 2434: 60-68
- Azpelicueta, M., J.M. Mirande, A.E. Almirón & J.R. Casciotta (2003): A new species of *Astyanax* (Characiformes, Characidae) from Paraná river basin in Argentina. *Revista del Museo de La Plata* 15 (166): 1-12
- Bertin, L. (1948): Catalogue des types de poissons du Muséum national d'Histoire Naturelle. 3e partie. Ostariophysaires (Characiniiformes, Gymnotiformes). 3: 1-49
- Buckup, P.A. (2003): Genera Incertae Sedis in Characidae: *Astyanax*. 106-169. In: Reis, R.E., S.O. Kullander & C.J. Ferraris (eds.): CLOFFSCA - Check list of the freshwater fishes of South and Central America. Edipucrs, Porto Alegre. 729 pp.
- Casciotta, J.R., A.E. Almirón & M. Azpelicueta (2003): A new species of *Astyanax* from río Uruguay basin, Argentina (Characiformes: Characidae). *Ichthyological Exploration of Freshwaters* 14 (4): 329-334
- Casciotta, J.R., A.E. Almirón & M. Azpelicueta (2005): *Astyanax pampa* (Characiformes, Characidae), a new species from the southernmost boundary of the Brazilian subregion, Argentina. *Revue Suisse de Zoologie* 112 (2): 401-408
- Casciotta, J.R., A.E. Almirón & J.A. Bechara (2003): Los peces de la laguna Iberá. Colección Universitaria Ciencias Naturales. 1-203
- Casciotta, J.R., A.E. Almirón, J.A. Bechara, J.P. Roux & F.J. Ruiz Díaz (2003): *Astyanax pynandi* sp.n. (Characiformes, Characidae) from the Estreros del Iberá wetland, Argentina. *Revue Suisse de Zoologie* 110 (4): 807-816
- Casciotta, J., A. Almirón, A. Cione & M. Azpelicueta (1999): Brazilian freshwater fish assemblages from southern Pampean area, Argentina. *Biogeographica* 75 (2): 67-78
- Cuvier, G. & A. Valenciennes (1849): Histoire naturelle des poissons. Salmonoïdes. 22: 1-532
- Eigenmann, C. H. (1921): The American Characidae. *Memoirs of the Museum of Comparative Zoology* 43 (3): 209-310, 28 pls. (various numbers)

- Fernández Santos, J.O., I.R. Wais, A. Puig & R. Larrea (1982): Observaciones sobre la íctiofauna del Parque Nacional El Palmar. Anales de Parques Nacionales 15: 77-105
- Fulvio Pérez, C.H. (2008): Fish, southernmost record of *Astyanax pampa* (Ostariophysi, Characiformes, Characidae). CheckList 4 (4): 424-426
- Garutti, V. & H.A. Britski (2000): Descrição de uma espécie nova de *Astyanax* (Teleostei: Characidae) da bacia do Alto Rio Paraná e considerações sobre as demais espécies do gênero na bacia. Comunicações do Museu de Ciências e Tecnologia da PUCRS, Sér. Zoologia 13: 65-88
- Géry, J. (1972): Corrected and supplemented descriptions of certain characoid fishes described by Henry W. Fowler, with revisions of several of their genera. Studies on the Neotropical Fauna 7: 1-35
- Géry, J. (1978): Results of Dr. K.H. Lüling's research in Argentina in 1975: The relationships of the naked characid, *Gymnocharacinus bergii* (Pisces, Cypriniformes, Characoidei) from northern Patagonia. Zoologische Anzeiger 201 (5/6): 403-409
- Ingenito, L.F.S. & L.F. Duboc (2014): A new species of *Astyanax* (Ostariophysi: Characiformes: Characidae) from the upper rio Iguaçu basin, southern Brazil. Neotropical Ichthyology 12 (2): 281-290
- Liotta, J. (2005): Distribución geográfica de los peces de aguas continentales de la República Argentina. ProBiota Serie Documentos 3. 701 p.
- López, H.L., R.C. Menni & A.M. Miquelarena (1987): Lista de los peces de agua dulce de la Argentina. Biología Acuática 12: 1-50
- López, H.L., A.M. Miquelarena & R.C. Menni (2003): Lista comentada de los peces continentales de la Argentina. ProBiota - Serie Técnica y Didáctica 5: 1-85
- Lueling, K.H. (1981): Wissenschaftliche Ergebnisse des Forschungsaufenthaltes Dr. K.H. Lueling in Argentinien 1975/76. 1. Ichthyologische und gewässerkundliche Beobachtungen und Untersuchungen an der Seenplatte von Guanini (suedl. Prov. Buenos Aires, Argentinien). Zoologische Beiträge 27 (1): 1-24
- Melo, F.A.G. de & P.A. Buckup (2006): *Astyanax henseli*, a new name for *Tetragonopterus aeneus* Hensel, 1870 from southern Brazil (Teleostei: Characiformes). Neotropical Ichthyology 4 (1): 45-52
- Messner, E. (1962): Lista de los peces Tetragonopterinae (fam. Characidae) del Uruguay. Boletín de la Asociación Latino Americana de Ictiología y Herpetología 2 (1): 4-5
- Miquelarena, A.M. & H.L. López (1995): Fishes of the Lagunas Encadenadas (Province of Buenos Aires, Argentina), a wetland of international importance. Freshwater Forum 5(1): 48-53
- Miquelarena, A.M. & R.C. Menni (2005): *Astyanax tumbayaensis*, a new species from northwestern Argentina highlands (Characiformes: Characidae) with a key to the Argentinean species of the genus and comments on their distribution. Revue Suisse de Zoologie 112 (3): 661-676
- Miquelarena, A.M., L.C. Protogino & H.L. López (1997): Fishes from the Arroyo Uruguaí (upper Paraná basin, Misiones, Argentina) before impoundment of the dam. Revue française d'Aquariologie Herpétologie 24 (3-4): 65-72
- Miquelarena, A.M., L.C. Protogino & H.L. López (2005): *Astyanax hermosus*, a new species from the Primero River basin, Córdoba, Argentina (Characiformes, Characidae). Revue Suisse de Zoologie 112 (1): 13-20
- Miquelarena, A.M., L.C. Protogino, J. Ponte Gómez & H.L. López (2009): Catálogo de la Colección Ictiológica del Instituto de Limnología "Dr. Raúl A. Ringuelet" (Parte I). Biología Acuática 25: 1-126
- Mirande, J.M. (2010): Phylogeny of the family Characidae (Teleostei: Characiformes) from characters to taxonomy. Neotropical Ichthyology 9 (1): 385-568
- Mirande, J.M., G. Aguilera & M. Azpelicueta (2004): A new species of *Astyanax* (Characiformes, Characidae) from the upper río Bermejo basin, Salta, Argentina. Revue Suisse de Zoologie 111 (1): 213-223
- Mirande, J.M., G. Aguilera & M. Azpelicueta (2007): A new species of *Astyanax* (Characiformes: Characidae) from the endorheic Río Salí basin, Tucumán, northwestern Argentina. Zootaxa 1646: 31-39
- Mirande, J.M., M. Azpelicueta & G. Aguilera (2006): Redescription of *Astyanax correntinus* (Holmberg, 1891) (Teleostei: Characiformes: Characidae), more than one hundred years after original description. Zoologische Abhandlungen des Museums für Tierkunde Dresden 55: 9-15
- Nion, H., C. Rios & C. Menses (2002): Peces del Uruguay. Lista sistemática y nombres comunes. Dinara - Infopescas, Montevideo. 105 pp.
- Pozzi, A.J. (1945): Sistemática y distribución de los peces de agua dulce de la República Argentina. Gaea 7 (2): 239-292
- Protogino, L.C., A.M. Miquelarena & H.L. López (2006): A new species of *Astyanax* (Teleostei, Characiformes, Characidae), with breeding tubercles, from the Paraná and Uruguay river basins. Zootaxa 1297: 1-16
- Ringuelet, R.A. & R.H. Arámburu (1962): Peces argentinos de agua dulce. Claves de reconocimiento y caracterización de familias y subfamilias, con glosario explicativo. Agro 3 (7): 1-90
- Ringuelet, R.A., R.H. Arámburu & A. Alonso de Arámburu (1967): Los peces argentinos de agua dulce. Comisión de investigación científica. Provincia de Buenos Aires. La Plata. 602 p.
- Serra, W.S. (2012): Revisión sistemática del grupo de especies "*Astyanax bimaculatus*" (Ostariophysi: Characiformes) presentes en Uruguay. Thesis. Universidad de la República. Uruguay. 44 p.
- Soneira, P., J. Casciotta, A. Almirón, L. Ciotek & P. Giorgis (2010): Redescription of *Astyanax erythropterus* (Holmberg, 1891) (Teleostei: Characiformes: Characidae) from the La Plata basin in Argentina. Neotropical Ichthyology 8 (4): 779-785
- Vari, R.P. & R.M.C. Castro (2007): New species of *Astyanax* (Ostariophysi: Characiformes: Characidae) from the upper Rio Paraná system, Brazil. Copeia 2007 (1): 150-162

recommended form for reference:

Mirande, J.M. & S. Koerber (2015):

On some species of *Astyanax* reported erroneously from Argentina.

Ichthyological Contributions of PecesCriollos 35: 1-8

available as pdf-file at www.pecescriollos.de since 02.Mar.2015