

Genome Sequence of the Melanin-Producing Extremophile *Aeromonas salmonicida* subsp. *pectinolytica* Strain 34mel^T

M. Elisa Pavan,^a Esteban E. Pavan,^b Nancy I. López,^{a,c} Laura Levin,^d M. Julia Pettinari^{a,c}

Departamento de Química Biológica, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina^a; Biomedical Technologies Laboratory, Department of Electronics, Information and Bioengineering, Politecnico di Milano, Milan, Italy^b; IQUIBICEN-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina^c; Departamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Buenos Aires, Argentina^d

M.E.P. and E.E.P. contributed equally to this work.

The genome of *Aeromonas salmonicida* subsp. *pectinolytica* strain 34mel^T, isolated from a heavily polluted river, contains several genomic islands and putative virulence genes. The identification of genes involved in resistance to different kinds of stress sheds light on the mechanisms used by this strain to thrive in an extreme environment.

Received 29 July 2013 Accepted 19 August 2013 Published 12 September 2013

Citation Pavan ME, Pavan EE, López NI, Levin L, Pettinari MJ. 2013. Genome sequence of the melanin-producing extremophile *Aeromonas salmonicida* subsp. *pectinolytica* strain 34mel^T. *Genome Announc.* 1(5):e00675-13. doi:10.1128/genomeA.00675-13.

Copyright © 2013 Pavan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to M. Julia Pettinari, jul@qb.fcen.uba.ar.

Aeromonas salmonicida subsp. *pectinolytica* strain 34mel^T is an extremophile isolated from the water of a heavily polluted river that receives the effluents of hundreds of tanneries and other industries, as well as urban sewage and fuel hydrocarbons (1). This strain produces abundant melanin and is highly resistant to heavy metals and other pollutants.

Phenotypic characteristics, DNA-DNA hybridization, and 16S rRNA gene sequence analysis determined the classification of 34mel as the type strain of *A. salmonicida* subsp. *pectinolytica*, a new subspecies of *A. salmonicida* (2). Genetic information for this subspecies is scarce and mostly restricted to 16S rRNA genes and a few other sequences. Multilocus sequence typing showed that this subspecies is the most phylogenetically distant among the five subspecies of *A. salmonicida* (3).

The sequence was obtained using a whole-genome shotgun strategy with a Roche 454 GS FLX Titanium pyrosequencer at INDEAR, Argentina, achieving ~22-fold coverage. Assembly was done using Newbler version 2.6 and generated 309 contigs, the largest of which has 327,967 bases. The draft genome is 4,774,629 bases in length, with a mean G+C content of 58.47%.

Annotation was performed using the NCBI Prokaryotic Genomes Automatic Annotation Pipeline (PGAAP) and Rapid Annotations using Subsystems Technology (RAST) (4), revealing 4,245 predicted coding regions and 100 RNA genes, including one 16S rRNA gene. The annotation covered 525 RAST subsystems, including 54% of the coding sequences, and classified 1,085 open reading frames (ORFs) as hypothetical proteins.

Whole-genome analysis using IslandViewer (5) revealed the presence of 12 genomic islands (GI). The largest GI could be related to environmental adaptability, as it contains a mercury resistance operon. This operon has previously been found in plasmids and in GIs in both environmental and pathogenic bacteria (6, 7).

Apart from the mercury resistance genes, many others that could allow strain 34mel^T to cope with toxic compounds were

found, such as those involved in resistance to chromium, molybdenum, copper, cobalt, arsenic, zinc, and cadmium, including genes coding for several efflux pumps. Many genes involved in responses to other kinds of environmental stress were also detected, such as those related to oxidative stress: genes coding for catalases, superoxide dismutases, peroxidases, and alkylhydroperoxidases.

A relevant characteristic of *A. salmonicida* subsp. *pectinolytica* is its production of large amounts of melanin. Enzymes involved in the homogentisate pathway were found in the genome of 34mel^T along with other gene products that might be related to melanin biosynthesis (8, 9). Pectinases, amylases, proteases, and other enzymes that might be of interest for biotechnological applications were also identified.

Although many *A. salmonicida* strains are well-known fish pathogens, there are no reports describing pathogenicity for *A. salmonicida* subsp. *pectinolytica*. However, a search for virulence-related genes in 34mel^T yielded genes coding for hemolysins, toxins, chitinases, a collagenase, and adhesion proteins, among others.

The analysis of the genome of *A. salmonicida* subsp. *pectinolytica* will contribute to understanding the lifestyle of this bacterium that inhabits an extremely polluted environment.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [ARYZ00000000](http://www.ncbi.nlm.nih.gov/nuccore/ARYZ00000000). The version described in this paper is version ARYZ01000000.

ACKNOWLEDGMENTS

N.I.L., M.J.P., and L.L. are career investigators from CONICET.

This work was partially supported by the University of Buenos Aires.

REFERENCES

- Pavan ME, Carbonelli D, Zorzópulos J. 2002. Molecular approach to microbial diversity studies in the heavily polluted Matanza River. *Rev. Mus. Argent. Cienc. Nat.* 4:139–144.

2. Pavan ME, Abbott SL, Zorzópolos J, Janda JM. 2000. *Aeromonas salmonicida* subsp. *pectinolytica* subsp. nov., a new pectinase-positive subspecies isolated from a heavily polluted river. *Int. J. Syst. Evol. Microbiol.* 50(Pt 3):1119–1124.
3. Roger F, Marchandin H, Jumas-Bilak E, Kodjo A, colBVH Study Group, Lamy B. 2012. Multilocus genetics to reconstruct aeromonad evolution. *BMC Microbiol.* 12:62. doi:10.1186/1471-2180-12-62.
4. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. doi:10.1186/1471-2164-9-75.
5. Dhillon BK, Chiu TA, Laird MR, Langille MG, Brinkman FS. 2013. IslandViewer update: improved genomic island discovery and visualization. *Nucleic Acids Res.* 41:W129–W132. doi:10.1093/nar/gkt394.
6. Sen D, Van der Auwera GA, Rogers LM, Thomas CM, Brown CJ, Top EM. 2011. Broad-host-range plasmids from agricultural soils have IncP-1 backbones with diverse accessory genes. *Appl. Environ. Microbiol.* 77:7975–7983. doi:10.1128/AEM.05439-11.
7. Reva ON, Bezuidt O. 2012. Distribution of horizontally transferred heavy metal resistance operons in recent outbreak bacteria. *Mob. Genet. Elements* 2:96–100.
8. Hunter RC, Newman DK. 2010. A putative ABC transporter, *hatABCDE*, is among molecular determinants of pyomelanin production in *Pseudomonas aeruginosa*. *J. Bacteriol.* 192:5962–5971. doi:10.1128/JB.01021-10.
9. Chai B, Wang H, Chen X. 2012. Draft genome sequence of high-melanin-yielding *Aeromonas media* strain WS. *J. Bacteriol.* 194:6693–6694.