

Construction of reference chromosome-scale pseudomolecules for potato: Integrating the potato genome with genetic and physical maps

Sanjeev Kumar Sharma^{*1}, Daniel Bolser^{§1,2}, Jan de Boert[†], Mads Sønderkær[‡], Walter Amoros^{**}, Martin Federico Carboni^{§§}, Juan Martín D'Ambrosio^{§§}, German de la Cruz^{††}, Alex Di Genova^{‡‡}, David S. Douches^{***}, Maria Eguluz^{§§§}, Xiao Guo^{†††}, Frank Guzman^{§§§3}, Christine A. Hackett^{‡‡‡}, John P. Hamilton^{****}, Guangcun Li^{†††}, Ying Li^{§§§§}, Roberto Lozano^{§§§}, Alejandro Maass^{‡‡}, David Marshall^{††††}, Diana Martinez^{§§§}, Karen McLean^{*}, Nilo Mejía^{‡‡‡}, Linda Milne^{††††}, Susan Munive^{**}, Istvan Nagy^{****4}, Olga Ponce^{§§§}, Manuel Ramirez^{§§§}, Reinhard Simon^{**}, Susan J. Thomson^{§§§§§}, Yerisf Torres^{§§§}, Robbie Waugh[†], Zhonghua Zhang^{§§§§}, Sanwen Huang^{§§§§}, Richard G. F. Visser[†], Christian W. B. Bachem[†], Boris Sagredo^{††††}, Sergio E. Feingold^{§§}, Gisella Orjeda^{§§§}, Richard E. Veilleux^{‡‡‡‡}, Merideth Bonierbale^{**}, Jeanne M. E. Jacobs^{§§§§§}, Dan Milbourne^{****}, David Michael Alan Martin[§], Glenn J. Bryan^{*5}

^{*}Cell and Molecular Sciences, The James Hutton Institute, Dundee DD2 5DA, United Kingdom

[§]Division of Biological Chemistry and Drug Discovery, College of Life Sciences, University of Dundee, Dundee DD1 5EH, United Kingdom

[†]Laboratory of Plant Breeding, Department of Plant Sciences, Wageningen-UR, Wageningen, The Netherlands

[‡]Department of Biotechnology, Chemistry and Environmental Engineering, Aalborg University, Aalborg, Denmark

^{**}International Potato Center (CIP), Lima 12, Peru

^{§§}Laboratorio de Agrobiotecnología, Instituto Nacional de Tecnología Agropecuaria (INTA) cc276 (7620) Balcarce, Argentina

^{††}Laboratorio de Genética y Biotecnología Vegetal, Universidad Nacional San Cristobal de Huamanga, Ayacucho, Perú

^{‡‡}Mathomics, Centro de Regulación Genómica & Centro de Modelamiento Matemático, Universidad de Chile, Santiago, Chile

^{***}Department of Crop and Soil Sciences, Michigan State University, Michigan, United States of America

^{§§§}Genomics Research Unit, Facultad de Ciencias, Universidad Peruana Cayetano Heredia, Lima 31, Peru

^{†††}Institute of Vegetables, Shandong Academy of Agricultural Sciences, Jinan 250100, China

^{‡‡‡}Biomathematics and Statistics Scotland, Dundee DD2 5DA, United Kingdom

^{****}Department of Plant Biology, Michigan State University, Michigan, United States of America

^{§§§§}Institute of Vegetables and Flowers, Chinese Academy of Agricultural Sciences, Beijing 100081, China

^{††††}Information and Computational Sciences, The James Hutton Institute, Dundee DD2 5DA, United Kingdom

^{‡‡‡‡}INIA-La Platina, Santiago, Chile

^{****}Crops Environment and Land Use Programme, Teagasc, Carlow, Ireland

^{§§§§§}The New Zealand Institute for Plant & Food Research Ltd., Christchurch 8120, New Zealand

^{†††††}INIA-Rayentué, Rengo, Chile

^{‡‡‡‡‡}Department of Horticulture, Virginia Tech, Virginia, United States of America

Footnotes:

¹S. K. Sharma and D. Bolser contributed equally to this work

²Current Address: The EMBL-European Bioinformatics Institute, Wellcome Trust Genome Campus, Cambridge CB10 1SD, United Kingdom

³Current Address: Departamento de Genética, Universidade Federal do Rio Grande do Sul - UFRGS, Rio Grande do Sul, Brazil

⁴Current Address: Department of Molecular Biology and Genetics, Aarhus University, Slagelse DK-4200, Denmark

⁵Corresponding Author

Running Title: Reference potato pseudomolecules

Keywords: Solanaceae, genome anchoring, scaffold orientation, sequence tagged sites, pseudomolecules

Corresponding author:

Glenn J. Bryan
Cell and Molecular Sciences
The James Hutton Institute
Invergowrie
Dundee DD2 5DA
Scotland, United Kingdom

Tel: +44 (0)1382 568735

Email: glenn.bryan@hutton.ac.uk

ABSTRACT

The genome of potato, a major global food crop, was recently sequenced. The work presented here details the integration of the potato reference genome (DM) with a new STS marker based linkage map and other physical and genetic maps of potato and the closely related species tomato. Primary anchoring of the DM genome assembly was accomplished using a diploid segregating population, which was genotyped with several types of molecular genetic markers to construct a new ~936 cM linkage map comprising 2,469 marker loci. *In silico* anchoring approaches employed genetic and physical maps from the diploid potato genotype RH and tomato. This combined approach has allowed 951 superscaffolds to be ordered into pseudomolecules corresponding to the 12 potato chromosomes. These pseudomolecules represent 674 Mb (~93%) of the 723 Mb genome assembly and 37,482 (~96%) of the 39,031 predicted genes. The superscaffold order and orientation within the pseudomolecules is closely collinear with independently constructed high density linkage maps. Comparisons between marker distribution and physical location reveal regions of greater and lesser recombination, as well as regions exhibiting significant segregation distortion. The work presented here has led to a greatly improved ordering of the potato reference genome superscaffolds into chromosomal 'pseudomolecules'.

Abbreviations

AFLP: amplified fragment length polymorphism; AGI: The Arabidopsis Genome Initiative; AGP: Accessioned Golden Path; BAC: bacterial artificial chromosome; Chr: chromosome; D: Goniocalyx cultivar group accession (CIP 703825) used as pollen parent in F₁ and BC generations of the DMDD progeny; DArT: Diversity Arrays Technology; DM: doubled

monoploid potato clone (DM 1-3 516 R44); DMB: DM superscaffolds; GBrowse: Genome Browser; HSP: high-scoring segment pairs; IRGSP: International Rice Genome Sequencing Project; LG: linkage group; OPA: oligonucleotide pool assay; PEMP: Paired-end and Mate-pair; PGSC: Potato Genome Sequencing Consortium; PM: Pseudomolecule; POPA: potato OPA; QTL: Quantitative trait locus; RH: dihaploid potato clone RH89-039-16; SGN: SOL Genomics Network; SNP: single nucleotide polymorphism; SSR: simple sequence repeat; STS: sequence-tagged site; UHD: Ultra-high density; WGP: Whole Genome Profiling; WGS: whole genome shotgun

Keywords: Solanaceae, genome anchoring, scaffold orientation, sequence tagged sites, pseudomolecules

Introduction

Genome sequencing of crop plants has become increasingly routine, primarily due to the reduction in cost and increase in throughput brought about by continuing advances in sequencing technologies. First reports on the whole genome sequences of plants, such as *Arabidopsis thaliana* (The Arabidopsis Genome Initiative 2000) and rice (International Rice Genome Sequencing Project 2005), were mainly accomplished with the use of clone based (e.g., 'BAC by BAC') strategies. In this approach, a library of BAC clones is mapped onto chromosomes using molecular markers, the aim being to generate a clone-based physical map with a 'minimum tiling path'. This assures good genome coverage while minimizing the sequencing effort. More recently, plant genome sequencing has been based on whole genome shotgun (WGS) approaches involving conventional Sanger sequencing, next generation sequence technologies, or a combination of both (Hamilton and Buell 2012). The

WGS approach does not require a physical map and there is no pre-assumption of the position of the resulting sequence assemblies. Several research groups have developed 'scaffolding' algorithms to assemble these typically short sequence contigs into larger constructs (Miller *et al.* 2010). However, due to the genome size and complexity of most crop plants, scaffolds typically remain un-oriented and without chromosomal coordinates, despite being well annotated for gene content. A reference genome sequence requires that the products of the assembly process (contigs and scaffolds) be globally ordered and oriented to generate chromosomal 'pseudomolecules' (PMs). In the absence of a clone-based physical map or genetic map of the reference sequenced genotype, this represents a significant and challenging task. One widely adopted approach has been to link the sequence assembly to a genetic map using the presence of mapped sequence-tagged site (STS) genetic markers (Green and Green 1991) in the genome sequence. For example, a set of 409 molecular markers was used to order 69% of the assembled 487 Mb grapevine genome along the 19 grape linkage groups (The French-Italian Public Consortium for Grapevine Genome Characterization 2007). The link between the genome sequence and its genetic maps is critical in moving between trait loci and candidate genes underlying such loci. Successful genetic anchoring of a plant genome sequence assembly using maps developed in the reference sequenced genotype depends on marker density and distribution, as well as map accuracy and resolution. Other approaches can also be implemented to augment the anchoring process, including comparative analysis with physical and genetic maps of closely related species.

The Potato Genome Sequencing Consortium (Potato Genome Sequencing Consortium 2011) has published the genome of the doubled monoploid *Solanum tuberosum* group Phureja DM1-3 516 R44 (hereafter referred to as DM). At the time the genome

sequencing was initiated, DM did not have a physical map, nor was there any pre-existing genetic map for this genotype. Therefore a genome anchoring strategy was developed which included the generation of a segregating bi-parental mapping population involving DM as a parent, and generation of a dense STS-based genetic map. Other genetic mapping resources, such as the ultra-high density (UHD) map of diploid potato genotype RH 89-039-16 (Van Os *et al.* 2006), and tomato-EXPEN 2000 genetic reference map (Fulton *et al.* 2002) were also used.

We describe for the first time in detail the generation of an integrated *de novo* genetic/physical map of potato and significant refinements to the previously published assembly. Our combined map orders the genome sequence into 12 chromosomal pseudomolecules corresponding to each of the 12 potato chromosomes, and is linked to previously existing potato and Solanaceae mapping resources. The work represents the assimilation of various data types which required complex interpretation for correct ordering and orientation of superscaffolds. This process involved considerable manual curation, driven largely by a novel approach for visualization of mate-pair sequences from large genomic clones (BAC and fosmid) and long insert 454 reads (20 kb and 8 kb). This allowed us to assign robust orientations to many superscaffolds and also allowed the inclusion of many superscaffolds that remained unanchored when the reference genome sequence was published (Potato Genome Sequencing Consortium 2011). This resource will facilitate exploitation of the potato genome sequence for genetic analysis and crop improvement, and our approach can serve as a guide for others wishing to engage in genome sequencing of genotypes which lack physical or genetic maps.

MATERIALS AND METHODS

Genetic cross/population construction

A segregating diploid potato population (BC₁) derived from the reference sequence clone DM 1-3 516 R44 (DM) was developed. The homozygous DM clone ($2n=2x=24$) was generated by chromosome doubling of a monoploid ($2n=1x=12$) derived from a heterozygous accession of *S. tuberosum* Group Phureja (Paz and Veilleux 1999). A heterozygous diploid clonal accession (CIP 703825, referred to as D) belonging to the *Solanum tuberosum* diploid Andigenum Group Goniocalyx cultivar group (Spooner *et al.* 2007, Ovchinnikova *et al.* 2011) was crossed to DM. The direction of the cross (DM×D) was chosen because DM is male sterile. One of the resulting F₁ hybrids (DM/D, CIP 305156.17) was used as stylar parent in a backcross with D as pollen parent. The mapping population comprising 180 backcross progeny clones (hereafter referred to as DMDD) was raised in the greenhouse for DNA extraction and pathogen testing, and is also maintained pathogen-free *in vitro* (<https://research.cip.cgiar.org/confluence/display/dm/Home>) at the International Potato Center (CIP), Peru.

Plant material and genomic DNA extraction

Genomic DNA from 180 progeny clones of the mapping population and the pedigree parents was isolated using standard protocols (Herrera and Ghislain 2000). DNA concentration was estimated with a TBS-380 Fluorometer (Turner BioSystems, USA) with PicoGreen® reagent using salmon sperm DNA at 500 ng/ml as a reference. All DNA samples were normalized to a final concentration of 250 ng/μl and distributed among members of the PGSC mapping

group in order to perform multi-location genotyping by employing DArT, SSR, SNP and AFLP markers.

Marker identification, development and analysis

Simple sequence repeat (SSR) markers: SSR markers were designed from an early draft of the assembled potato genome superscaffolds (DM assembly version 1). Markers were selected from a masked copy of the genome to avoid placement in repetitive DNA. In addition to these SSR markers (labelled 'PM'), previously reported sets of SSRs from Stwax (potato *waxy* gene, Veilleux *et al.* 1995), STM (Milbourne *et al.* 1998), STI (Feingold *et al.* 2005), st_ (Tang *et al.* 2008a) and STG (Ghislain *et al.* 2009) were also used in linkage mapping. In total, 356 SSRs (Table S1A) were tested for polymorphism. In brief, 5-25 ng template DNA was added to PCR mix containing 1.5-2.5 mM MgCl₂, 0.16-0.25 mM dNTP, 0.25-1.0 U Taq polymerase, with the following primer combinations; for acrylamide gel analysis, 0.2-0.25 μM forward primer, 0.2-0.25 μM reverse primer, plus 0.2 mM cresol red and 6% sucrose; for ABI3130lx Genetic Analyser (Applied Biosystems), 0.2-0.25 μM reverse primer, 0.15-0.25 μM forward primer, 0.05-0.25 μM labelled (FAM, HEX, NED or PET) forward primer; for 4300 LI-COR DNA Analyser (LI-COR Biosciences), 0.2 μM or 22 pM forward primer, 0.2 μM or 15 pM reverse primers, 25 pM 700 or 800 IRDye labelled M13 forward primer. PCR reactions were conducted under optimised conditions: in brief, 4 min denature at 94°, 35 cycles of 30 sec at 94°, 30 sec at T_a (annealing temperature determined experimentally for each SSR primer combination), 30 sec at 72°, 1 cycle of 4 min at 72°; or 3 min denature at 94°, 36 cycles of 15 sec at 94°, 30 sec at 58-52° with touchdown of -0.5° for first 12 cycles, 30 sec at 72°, 1 cycle of 5 min at 72°; or 4 min denature at 94°, 30-33 cycles of

1 min at 94°, 1 min at T_a, 1 min at 72°, 1 cycle of 4 min at 72°. SSRs were resolved either by denaturing acrylamide gel electrophoresis and silver staining according to Creste *et al.* (2001), capillary electrophoresis following standard procedures for the ABI3130lx Genetic Analyser using Genscan 400 ROX or Genscan 500 LIZ size ladder, or by electrophoresis on the 4300 LI-COR DNA Analyser system (LI-COR Biosciences) using the LI-COR IRDye 50–350 bp size standard. Polymorphic markers were scored directly from silver stained gels; using GeneMarker 1.4 (SoftGenetics, State College, PA; www.softgenetics.com), GeneMapper 4.0 (Applied Biosystems) or Genographer (www.genographer.com) for ABI3130 lx; or the SAGA Generation 2 software (LI-COR, USA), and Cross Checker v.2.9.1 (Buntjer 1999) for LI-COR. SSRs were scored, where possible, as co-dominant markers, and if this was not possible, as dominant markers.

Single nucleotide polymorphism (SNP) markers: A custom filtering pipeline was developed to select 1,920 SNPs from a set of 69,011 high confidence SolCAP SNPs (Hamilton *et al.* 2011) that were incorporated into five 384-plex (5 x 384) Illumina GoldenGate™ oligonucleotide pool assays (OPAs; Fan *et al.* 2003), hereafter referred to as POPA (potato OPAs). Hamilton *et al.* (2011) identified these SNPs by comparing RNA-Seq and EST sequences from six potato cultivars (Atlantic, Premier, Snowden, Bintje, Kennebec and Shepody) to the draft DM potato reference genome. Our filtering pipeline involved finding non-repetitive positions on the DM assembly, avoiding overlapping SNPs that may have interfered with the Illumina SNP genotyping assay, and striving to cover the genome as fully as possible. Additionally, a POPA containing SNPs derived from pre-existing potato ESTs in the public databases was also designed and used. Table S1B shows details of 2,304 SNPs, derived from pre-existing potato ESTs (POPA1) and SolCAP markers (POPA2-6), employed in the study. Genotyping was performed using an Illumina BeadXpress platform following the

recommendations of the manufacturer (GoldenGate Genotyping Assay, Illumina VeraCode Manual, VC-901-1001). All reagents, unless stated otherwise in the standard protocol, were provided by Illumina. The data files were processed and genotypes called using Genome Studio software.

Amplified Fragment Length Polymorphism (AFLP) markers: AFLP analysis was carried out according to the procedures described by Vos *et al.* (1995) using the restriction enzyme combination *EcoRI* and *MseI*. AFLP fragments were separated on a LI-COR 4300 DNA Sequencer (LI-COR Biosciences) using 4.5% polyacrylamide denaturing gels (acrylamide:bisacrylamide, 19:1) as described in the user manual. LI-COR size standard ladder was loaded into each lane to facilitate the semi-automatic analysis of the gel and the sizing of the fragments. The names of the markers indicate the enzymes used, the selective nucleotides, and the size of the fragment; for instance, EACTMAAC_205.0 is an AFLP marker derived from a primer combination with the enzymes *EcoRI* and *MseI*, selective nucleotides ACT and AAC, and a mobility that corresponds to a fragment with an estimated size of 205 bp. Polymorphic bands were manually scored following the intensity degree and the parent backcross pattern. The details of the enzyme combinations, selective nucleotides and adapter sequences are provided in Table S1C.

Diversity Arrays Technology (DArT) markers: Representations from 180 DMDD progeny clones and the pedigree parents (DM, DM/D, D) were obtained by subjecting DNA from each clone to double restriction enzyme digestion (*PstI/TaqI*) and ligation to *PstI* adaptors for reducing genome complexity followed by PCR amplification for preparation of targets (Wenzl *et al.* 2004). Cy3 labelled representations (targets), mixed in an ExpressHyb buffer containing cy5-labelled polylinker fragment of the plasmid used for library preparation (as a reference), were denatured and hybridised to a high resolution potato genotyping array

containing 7,680 DArT probes (Sliwka *et al.* 2012). After overnight hybridization at 62°, arrays were washed and scanned with 20 µm resolution at 543 nm (cy3) and 488 nM (FAM) on a LS300 confocal laser scanner (Tecan, Grödig, Austria) to detect fluorescent signals emitted from the hybridised fragments. The data from the scanned images were extracted and analysed using the DArTsoft 7.4 software (Diversity Arrays Technology P/L, Canberra, Australia). The logarithm of the ratio between the two background-subtracted averages of feature pixels in the cy3 and cy5 channels ($\log_2[\text{cy3}/\text{cy5}]$) was used as a measure of the difference in abundance of the corresponding DNA fragment in the two representations hybridized to the array. The $\log_2[\text{cy3}/\text{FAM}]$ and $\log_2[\text{cy5}/\text{FAM}]$ values, approximate measures of the amount of hybridization signal per amount of DNA spotted on the array, were used for quality-control purposes. The unique signal pattern obtained by hybridizing each sample pair (individual clone and reference) to the genotyping array was recorded as '0' or '1'. All DArTs were sequenced and are available from [Spud DB site \(http://potato.plantbiology.msu.edu/\)](http://potato.plantbiology.msu.edu/); the detailed methodology is published on the Diversity Arrays Technology website (<http://www.diversityarrays.com>).

Linkage map construction

The SSR, SNP, AFLP and DArT genotyping data for 180 DMDD progeny clones was combined and screened for polymorphic markers. JoinMap[®]4 (Van Ooijen 2006) was used both to assign markers to linkage groups and to order markers within linkage groups. The backcross parents and offspring were coded according to the CP population type (outbreeder full-sib family after two independent meioses). A female-male combined DMDD map was generated including markers informative in one or both parents. Linkage groups were

formed using the Independence LOD parameter under 'population grouping' with a range from 2 to 15. Prior to grouping and ordering markers within linkage groups, loci or progeny clones with $\geq 20\%$ missing values were removed along with all identically segregating loci. The regression mapping algorithm with modified settings (recombination frequency threshold < 0.45 , LOD threshold > 0.05) was used to order loci within each linkage group. All linkage groups were subjected to three rounds of mapping. Recombination frequencies were converted into map distances using the 'Kosambi' mapping function.

Locating STS markers on the DM assembly

Sequence based markers (STS) were aligned to the reference genome assembly using SSAHA2 (Ning *et al.* 2001) or BLAST. The total set of alignments was processed as follows. **First**, alignments caused by short repetitive sequences were removed using a custom depth/coverage filter. In detail, any alignment covering a region of the query or target sequence that overlapped with five or more other competing alignments in that region was removed if this depth threshold was exceeded over 20% or more of the alignment length. In this way alignments spanning short repeats were not penalised, but alignments largely composed of likely repeats were removed. **Second**, short alignments were grouped by sequence into 'hits' that allowed for indels. **Third**, where applicable, the relative distance and orientation of the forward and reverse reads for the marker was taken into consideration. Pairs of forward and reverse reads with an incorrect orientation or implausible separation were removed. **Finally**, only markers with a unique, high scoring alignment position on the genome assembly were selected as anchor points in the physical

map. The final positions of all the STS markers (SSRs, SNPs and DArTs) are provided in Table S2.

Integration of additional sequence-based and physical resources

DM BAC- and Fosmid-end sequences, RH BAC-end sequences, and tomato BAC- and Fosmid-end sequences were aligned to the DM superscaffolds using SSAHA2 (Ning *et al.* 2001). The resulting alignments were filtered as described above. Roche 454 Paired-end (PE) reads from 14 and 20 kb insert-size libraries from DM, representing 0.7 and 1.0 Gb of raw data, respectively, were aligned to the superscaffold sequences using Newbler (Margulies *et al.* 2006) with all the default settings. Un-sequenced BAC clones from the RH physical map (de Boer *et al.* 2012) were positioned on the superscaffolds using BLAST alignment of their WGP sequence tags. For each BAC, the alignment hits of the individual 25 nt tags were processed to retain only unique hits. The aligned BAC clones that carried AFLP markers provided the link between the DM superscaffolds and the RH UHD genetic map (Van Os *et al.* 2006). In addition, sequenced RH BAC clones and RH BAC-end sequences were used for anchoring and scaffolding of the DM sequences. Finally, sequences from the available tomato pseudomolecules (v2.40, The Tomato Genome Sequencing Consortium 2012) were aligned using ATAC (Istrail *et al.* 2004).

Manual scaffolding using the 'link-peak' strategy

All Paired-End and Mate-Pair (PEMP) sequence data that could be reliably mapped to the DM superscaffolds was combined to compute a composite directional link-score across each superscaffold. In detail, the link-score combined PEMP that had unique, high-scoring

alignments for both ends of each mate pair sequence, but with the two end sequences aligning to different non-adjointing superscaffolds. A reciprocally high link-score between the ends of a pair of superscaffolds indicated a probable scaffolding link between them. The composite directional link-score is calculated in a sliding window along the length of a superscaffold (the source) as follows:

1. All mate pairs with one end aligning in that window and the other corresponding mate pair end reliably mapping to another superscaffold (the target) are selected. These are designated as unsatisfied mate pairs.
2. These mate pairs are grouped according to the target superscaffold.
3. For each target superscaffold group, a score is calculated by summing the value for each mate pair in that group (see below for details of how the value is determined).
4. The link-peak score is the highest score of all the target groups.

Different link-score values were empirically assigned to the different PEMP sequence libraries with higher scores assigned to DM based libraries over RH and tomato based libraries, and higher values given to longer sequences which have more accurate alignments. In addition to accumulating link-evidence from consistent unsatisfied PEMPs, a noise-score was calculated for unsatisfied PEMPs that suggested links to multiple different target superscaffolds. The noise score allowed spurious, high-scoring links caused by repeats to be identified. In this way the evidence for links between pairs of superscaffolds could be conveniently described as a continuous value in wiggle format (<https://www.genome.ucsc.edu/goldenPath/help/wiggle.html>) which allows for visualisation as tracks in GBrowse, alongside genetic and physical evidence from other sources.

Visualisation of integrated genetic and physical map

The integrated genetic and physical maps of the DM genome were visualised with the software 'DMAP' (DMA Martin, manuscript in preparation). The figures produced by the DMAP software take as input the AGP file describing the pseudomolecule architecture, a GFF file describing the sequence positions of the markers on the superscaffolds, and the JoinMap output file from linkage mapping for each linkage group. As there are many more markers than can be coherently visualised on a printed figure, DMAP employs a selection and layout algorithm where only a user determined maximum number of labels are displayed.

DM chromosome idiogram figures were reproduced from the potato reference genome publication (Potato Genome Sequencing Consortium 2011), and were aligned by orienting the short arms towards the start of the pseudomolecule sequence, except for chromosomes 5 and 11, where the pseudomolecule sequence begins in the long arm (Tang *et al.* 2009; Potato Genome Sequencing Consortium 2011).

Identification of centromere positions and pericentromeric regions

Centromere positions were determined with the sequence information provided by Gong *et al.* (2012). For chromosomes 4, 6, 9, 10, 11 and 12, the DM superscaffolds covering the centromere locations were identified from the major peaks in the CENH3 chromatin immunoprecipitation sequence read plots on the DM V2.1.10 pseudomolecule sequences. Satellite repeat analysis was performed by searching for the repeats in the DM sequence at <http://yh.genomics.org.cn/potato/search.jsp> and by evaluating the repeat coverage through dot plot alignment of candidate DM sequences with the repeat sequence. Additionally,

centromere positions were also indirectly inferred from the marker density in RH UHD genetic map (van Os *et al.* 2006).

The revised physical positions of all of the Illumina Potato 8303 Infinium array SNPs, reported by Felcher *et al.* (2012) using their customized version (2.1.11) of potato reference pseudomolecules (PMs), were obtained for the latest version (4.03) of PMs (Table S3). Graphs depicting the progression of genetic distance and recombination rate versus physical distance were calculated for all of the SNPs included in the current PMs and D84 and DRH genetic maps, using the MareyMap package (Rezvoy *et al.* 2007). The pericentromeric heterochromatin regions of the DM PMs were identified in these plots from the absence of genetic recombination between the SNP markers in such regions. In addition, AFLP markers from the marker-dense pericentromeric bins of the RH genetic maps were used to define heterochromatin boundaries in the PMs (Park *et al.* 2007), especially in cases where the genetic maps of Felcher *et al.* (2012) offered limited resolution.

BAC assembly and comparison with pseudomolecules

A total of 96 DM BACs spanning scaffolding gaps on chromosome 4 were selected (using DM BAC-end hits; Potato Genome Sequencing Consortium 2011). The BACs were picked from the library and end-sequenced to verify correct selection. Eighty-two verified BACs were further processed and grouped into six normalized pools as well as a composite master pool containing all 82 BACs. Each of the six BAC pools was subjected to Roche 454 single-end (SE) shotgun sequencing and the master pool to 3 kb paired-end (PE) sequencing. SE data for each pool were combined with the PE data and were assembled together using the Newbler

GSAssembler (Margulies *et al.* 2006). The sequences were deposited in the EBI Short Read Archive (accession number: ERP000934).

Candidate BAC scaffolds containing BAC-end sequences were identified with BLAST, filtering hits with a minimum match length of 400 bases and bit score exceeding 700 before manual curation. BAC scaffolds were matched to genomic superscaffolds with MUMmer (Kurtz *et al.* 2004). Matching regions were filtered to retain only matches longer than 1000bp with >97% identity. Data was expressed graphically with matches as edges and BAC end sequences, superscaffolds and BAC scaffolds as nodes using the graphical exchange format. Code was written in Python with the pygexf library and visualisation performed with Gephi (<http://www.gephi.org>). In addition, BAC ends were linked by a BAC label as a node. Assemblies which linked superscaffolds with sequence data could then be readily observed as cycles containing a BAC label in the graph. BAC-oriented GFF files were generated and visualised with R.

RESULTS AND DISCUSSION

DM genome assembly: a brief summary

The potato nuclear genome involved generation of ~96.6 Gb of raw sequence, which assembled into 66,254 'superscaffolds' comprising a net sequence assembly of 727 Mb, 117 Mb less than the estimated genome size of 844 Mb. Superscaffold length is inversely proportional to the numerical value in the name of each DM superscaffold (DMB), where the largest DMB (7.1 Mb) bears the ID 'PGSC0003DMB000000001' and the smallest (100 bp) 'PGSC0003DMB000066254'. About 94% of the assembled genome is non-gapped sequence and over 90% of the genome (N_{90}) is represented by 443 superscaffolds which are equal to

or larger than 349 kb. The anchoring strategy preferentially targeted the larger superscaffolds. At the time of publication 649 superscaffolds equalling 623 Mb (86%) of the assembled genome and 90% of the 39,031 estimated genes were anchored (Potato Genome Sequencing Consortium 2011). Draft pseudomolecules for the 12 chromosomes had been constructed but superscaffolds were mostly un-oriented. Since the original publication, continuous efforts have been made to perform further anchoring and orientation of the DM superscaffolds in order to generate the revised and improved genome pseudomolecules presented here (version 4.03).

Genetic analysis of the mapping population

The DMDD mapping population was genotyped for AFLP, SSR, SNP and DArT markers. Twenty two AFLP primer pairs (*EcoRI/MseI*) amplified 213 detectable fragments. A total of 356 SSR loci were assayed. Out of 2,304 POPA SNPs and 7,680 DArTs interrogated, 2,160 and 2,174 yielded genotype data, respectively. The compiled set of 4,903 markers was screened for presence of polymorphism, data integrity and concordance between parental and progeny genotypes, as well as meeting the missing data threshold (<20%) and other standard quality control checks. These data filtering and quality measures resulted in considerable reduction in the total number of markers used for linkage mapping to 2,597 which comprised 187 AFLPs, 234 SSRs, 367 SNPs and 1,809 DArTs. After excluding co-segregating markers, we used a subset of 1,864 uniquely segregating loci for linkage grouping; 1,751 unique loci were incorporated into a combined parental linkage map with the 12 expected linkage groups (LG), while the remaining 113 remained unmapped. The 12 chromosomal linkage groups span 936.2 cM with an average marker spacing of 0.54 cM per

interval. The individual linkage groups ranged in size from 62.9 cM (Chr11) to 101.8 cM (Chr03). A combination of the use of previously mapped SSR markers (Veilleux *et al.* 1995, Milbourne *et al.* 1998, Feingold *et al.* 2005, Tang *et al.* 2008a, Ghislain *et al.* 2009) and other available resources such as the RH genetic map (Van Os *et al.* 2006), whole genome profiling (WGP) (de Boer *et al.* 2012) and the tomato-EXPEN 2000 maps (Fulton *et al.* 2002), allowed orientation and assignment of all 12 linkage groups to their respective chromosomes. Table 1 shows the summary statistics of linkage mapping in the DMDD cross.

Departure from Mendelian segregation has been frequently observed in potato crosses. Markers showing segregation distortion were not excluded from the mapping process and most could be mapped to their appropriate linkage groups. The frequency of segregation distortion was highly variable among different chromosomes with the most significant distorted regions observed on chromosomes 1 and 4. Previous potato mapping studies have also shown varying levels of segregation distortion (Gebhardt *et al.* 1991, Felcher *et al.* 2012). Figure S1 shows genome-wide distribution of levels of segregation distortion for all STS markers employed in DMDD.

Linkage map based (direct) anchoring

The linkage map of DMDD is predominantly composed of STS markers. The primary map based anchoring strategy involved locating these sequence-based markers in the DM superscaffolds. SNPs and previously unpublished SSR markers (prefixed with 'PM') used in the DMDD linkage map were designed *a priori* against genome superscaffolds so their unique positions in the relevant superscaffolds were known. The positions of DArT and

previously reported SSRs were determined using the bioinformatics alignment and filtering pipeline illustrated in Figure 1.

Co-segregating markers removed during linkage map construction were included in the anchoring process as such genetically redundant markers represent distinct, but physically linked sites in the genome. The complete set of STS markers was filtered for unique and unambiguous marker-assembly sequence alignments as described. The combined sequence and genetic map coordinates for these unique STS markers were used to assign and order superscaffolds for constructing a framework physical map. The integrated genetic and physical anchoring strategy is shown in Figure 2. Using this strategy, we anchored 1,730 (1,305 DARts, 345 SNPs and 80 SSRs) of the 2,292 mapped, including co-segregating, STS markers to their unique positions on the DM superscaffolds. This approach anchored 54.2% (394 Mb) of the DM genome assembly arranged into 334 superscaffolds (Table 2). The proportion of genetic markers anchored on the genome sequence from each marker-category was 96% (SNPs), 28% (SSRs) and 76% (DARts). Mapped AFLP fragments were not used in the anchoring process, due to a lack of sequence information. Table S2 contains genomic positions for all the STS markers employed in the study. Genetic and physical coordinates for the DMDD mapped markers, including 718 co-segregating markers, are provided in Table S4.

***In silico* approach based (indirect) anchoring**

The DMDD-based framework physical map was extended by integrating two additional sources of syntenic map data, from potato and tomato respectively. First, superscaffolds anchored using the RH UHD genetic and physical maps (van Os *et al.* 2006, de Boer *et al.*

2012) were added. Second, 2,604 sequence-based markers from the tomato EXPEN 2000 derived maps, which are closely collinear with potato (Tanksley *et al.* 1992, Fulton *et al.* 2002, The Tomato Genome Sequencing Consortium 2012) were used to add superscaffolds. In the case of RH, sequence anchoring was derived from the AFLP- and WGP-based hybrid RH physical map (de Boer *et al.* 2012) as well as by direct alignment of RH BAC end sequences and fully sequenced RH seed BACs to the DM sequence. In both cases, the (proxy) marker sequences were aligned to the DM assembly using BLAST, adopting stringent matching criteria. The results were processed into reliable genetic anchor points as described previously for the DM markers.

The RH and tomato based *in silico* anchoring strategies independently anchored 470 (527 Mb, 72.5%) and 402 (417 Mb, 57.4%) superscaffolds, respectively (Table 2). Figure 3 shows the superscaffold anchoring summary for both the linkage (DM map) and the two *in silico* (RH and tomato maps) approaches. The total set of 649 superscaffolds anchored in at least one map were integrated hierarchically, starting with the DMDD-based framework map, placing additional superscaffolds using first the RH and then tomato assignment. The hierarchical 'alignment' of the maps is described below.

Construction of chromosome-scale pseudomolecules

Following anchoring, the superscaffolds were ordered into chromosome-scale pseudomolecules (PMs) in a hierarchical process using genetic, sequence and physical map data. The process is broken into two stages.

Stage-I: In the first stage the STS markers from the DMDD genetic map were aligned to the DM superscaffolds and used to construct the 'backbone' PMs. Additional sequence-linked

and sequence-based markers from the RH and tomato genetic maps were subsequently used to add superscaffolds into the DM backbone PMs (Figure 2). Superscaffolds that were anchored in multiple maps were used as reference points to align the genetic positions in the three different maps. Superscaffolds were added into 'gaps' in the backbone PMs where the positions indicated by the RH and tomato markers were in agreement with the positions initially established by the DMDD map data. The known set of chromosomal inversions on chromosomes 5, 6, 9, 10, 11 and 12 between potato and tomato (Tanksley *et al.* 1992; Iovene *et al.* 2008; Tang *et al.* 2008b) were taken into account when aligning the different genetic maps.

Generally the different anchoring approaches provided direct support for each other with respect to the relative placement of superscaffolds in the PM. With an optimal alignment/agreement for the superscaffold order among the three different maps used for anchoring, 294 of 374 superscaffolds present in at least one map were found to be in the same order as in the other two maps. In some instances, we observed that ordering of superscaffolds derived using RH and tomato maps was inconsistent with that obtained from the DMDD genetic map. The observed differences could be due to many factors, including technical issues such as mapping or assembly errors or biological properties, such as previously unknown structural differences between the compared genomes. However, given the size and complexity of the potato genome, it is encouraging that the placement of 79% of the superscaffolds was corroborated by the different methods employed.

Although superscaffolds were integrated into genomic blocks at this stage, they were unoriented and, due to the difficulty of aligning genetic maps, largely unordered at the chromosome level. To add, orient and refine the order of superscaffolds into an Accessioned

Golden Path (AGP) for constructing chromosome-scale PMs, a separate process was implemented, as described below.

Stage-II: To orient the DM superscaffolds, and to further refine the DMDD linkage based PMs, sequence and physical data from a variety of sources were combined as described in Material and Methods and visualised on a standard GBrowse installation (Figure 4). Custom sequence features were created representing high scoring inter-superscaffold links, allowing the user to ‘click-and-walk’ along the physical evidence from superscaffold to superscaffold in GBrowse. To aid this visualization, the processed RH WGP and tomato alignments, including the aligned sequence markers from the genetic maps used in Stage-I, were added to GBrowse as additional sequence feature tracks.

Using this integrated visualisation tool, three important types of manual improvements to the Stage-I PMs were performed, (i) scaffolding links were used to provide the relative orientation of superscaffolds, (ii) adjacent superscaffolds not previously included in the integrated genetic/physical map were added, and (iii) errors in the assembly were identified. These manual improvements were mainly carried out for the euchromatic (gene-rich) regions and for the euchromatin/heterochromatin borders. In addition to orientating the majority of the anchored superscaffolds, the ‘link-peak’ walk strategy combined with manual curation led to the incorporation of an additional 277 previously un-anchored superscaffolds into the PMs.

During this process 67 chimeric superscaffolds were identified. Of these, 62, 3 and 2 superscaffolds were revealed to have one, two and three misassembly locations, respectively, where false sequence joins had occurred. Many of these errors explained incongruities initially observed in the construction of the backbone PMs from the DMDD

map (Stage-I). Chimeric superscaffolds were manually split and allocated to their respective positions in the PMs. For example, the sequence coordinates 1 to 1117982 bp of PGSC0003DMB000000002 were allocated to Chromosome 4 whereas those from 1117983 to 6562806 bp were allocated to Chromosome 5. These results further illustrate the utility of an integrated genetic and *in silico* anchoring based approach for refining and correcting genome assembly errors.

Included in the refinement process were dot plot alignments of DM chromosome pseudomolecule sequences to pre-release and finished versions of the tomato genome sequence (The Tomato Genome Sequencing Consortium 2012). These alignments focused on the euchromatic regions and the adjacent heterochromatin border regions, where potato and tomato display homology in their sequences. The dot plot alignments to tomato made useful suggestions on how to place as yet unordered potato superscaffolds and superscaffold blocks, after which nearly always BAC end sequence links were identified in potato that confirmed the suggested orientation. Very occasionally, the potato PM description relied on the tomato alignment for placing potato sequence blocks in their presumed orientation, e.g. from PGSC0003DMB000000729 to PGSC0003DMB000000835 at the top of chromosome 1 and from PGSC0003DMB000000692 to PGSC0003DMB000001163 in the south heterochromatin border on chromosome 8.

Inversions with tomato

The potato-tomato dot plot alignments explained the discrepancies that were found between the potato and tomato genetic maps. In the euchromatic regions and the adjacent heterochromatin border regions we collected the sequence positions of the 19 largest

paracentric inversions (with a length of at least 0.3 Mb), which are listed in Table S5 and also indicated in the DM pseudomolecule figures. Newly identified were, among others, a tandem inversion with minor additional rearrangements on potato chromosome arm 1L, a nested inversion on 2L, and an arm inversion on 8S. Furthermore, the known arm inversions on 9S and 11L were found to be tandem inversions, with the second inversion being located in the heterochromatin border. The chromosomal rearrangements on 2L have also been described by Peters *et al.* (2012), who presented a scenario involving four structural conversions between potato and tomato. However, our dot plot sequence alignment for this region is less complex and shows a single, smaller inversion inside a larger inversion. This nested inversion model requires only two structural conversion steps and remains compatible with the cytogenetic results of Peters *et al.* (2012).

No paracentric inversions were identified on chromosome 3. However, on the short arm, the tomato sequence differs from the potato sequence by a 7.0 Mb insertion, which is located at position 2.4 Mb in the DM chromosome 3 pseudomolecule, and which runs from 1.3 to 8.3 Mb in the tomato SL2.40 assembly. In its center, this tomato insert has 4.2 Mb of sequence that is largely devoid of genes (<http://potato.plantbiology.msu.edu/>), while the start and end regions align with gene-containing potato sequence segments from region 42.0 to 50.4 Mb on the south arm of chromosome 3. Although these data suggest a translocation of sequences across the centromere, further investigation is needed to exclude sequence assembly errors.

Validation of link peak-based orientation strategy for chromosome 4

The strategy for PM construction and assembly correction was validated on chromosome 4 by targeted sequencing of 82 DM BAC clones that were selected to overlap candidate links as well as 10 of the 15 putative chimeric superscaffolds mapped to this chromosome. Thirty-one BAC clones could be assembled with contigs which spanned multiple superscaffolds and provided full coverage between the BAC end sequence matches to the superscaffolds, both validating the assembly and providing direct evidence for all 10 chimeric breakpoints. Seven of these sequenced BACs allow the inclusion of further superscaffolds that had not previously been assigned to a pseudomolecule, and one provides evidence for a superscaffold that had been erroneously included.

In addition to the complete assemblies described above, most other clones could be assembled to a series of contigs which did not span multiple superscaffolds and which have not been included in the BAC pool summary (Table S6). Details of the BAC analysis are given in the Methods and a representative example validating a potential break-point in Chromosome 4 is illustrated in Figure 5. A list of putative erroneous superscaffold assembly locations (breakpoints), and the BACs which provide validation for them is given in supplementary table S7.

Demarcating centromeres and pericentromeric boundaries in the pseudomolecules

The putative centromere locations for 7 of the 12 potato chromosomes were identified in the pseudomolecule sequences based on data published by Gong *et al.* 2012 (Table S8). Six centromere locations were identified from chromatin immunoprecipitated sequences. Of the seven published centromeric satellite repeat sequences (Gong *et al.* 2012), only the St24 repeat specific for the chromosome 1 centromere identified DM sequences with a high

repeat copy number characteristic of centromeric regions. With the other six centromeric repeat sequences we could not find reliable centromeric targets in the DM assembly, since these sequences only identified locations with very few repeat copies, which sometimes occurred on a chromosome other than that expected from their designated centromeres.

Pericentromeric boundaries were deduced by comparing the SNP based D84 and DRH genetic maps of Felcher *et al.* (2012) to the current version of PMs. For all chromosomes the typical pattern of distinctly reduced recombination in pericentromeric regions, and increased, varying recombination rates in euchromatic regions was observed (Figure 6). These patterns were used as the primary information source to demarcate putative pericentromeric regions in the PMs, and the boundaries of these regions were well supported, and where needed refined, by the RH genetic maps (van Os *et al.*, 2006). Figure 7 and Figure S2 depict the centromere and pericentromeric locations in the PMs. The pachytene chromosome idiograms in these figures are adapted from The Potato Genome Sequencing Consortium (2011).

Current status of the reference pseudomolecules

The genome anchoring, ordering and orienting process, as described above, led to the joining of 951 genome superscaffolds, or non-chimeric segments thereof, into 144 larger, contiguous sequence blocks, and enabled construction of an AGP assembly for the reference DM potato genome. These chromosome-scale pseudomolecules, version 4.03, contain 93% (as compared to 86%; Potato Genome Sequencing Consortium 2011) of the assembled genome comprising 674 Mb in 951 superscaffolds and include 37,482 (~96%) of the 39,031 predicted genes. A total of 938 superscaffolds (655 Mb or 90 % of the assembled genome

sequence) are assigned absolute or relative orientations within the PMs, whereas the remaining 13 superscaffolds (19 Mb) are assigned with a random orientation. For 279 Mb of superscaffold sequence blocks from the heterochromatin, the exact chromosome position and absolute orientation could not be determined. These partially unordered regions are marked yellow in the pseudomolecule figures (Figures 7, S2). No attempts were made to estimate gaps sizes between the superscaffolds, and in the PM sequences all superscaffolds are separated from each other by a fixed gap sequence of 50,000 N's. The N₉₀ of the DM potato genome assembly is 0.25 Mb and contains 622 superscaffolds, of which 28 (equalling 17 Mb, ~2% of the assembled genome sequence) remain unanchored. The longest anchored superscaffold is 7.1 Mb (PGSC0003DMB000000001; chromosome 1) and the longest unanchored superscaffold (PGSC0003DMB000000064) is 2.2 Mb. The increase in average N₅₀ from 1.5 Mb to 4.1 Mb in DM version 4.03 (Table 3) further supports the enhanced quality of the constructed PMs. The current version of the PMs/AGP is provided in Table S9 and includes the list of unanchored superscaffolds (Chromosome 0) and chimeric superscaffolds.

For visualizing the differences and improvements in the constructed PMs, we compared dotplots of the current PMs (ver 4.03) to the earlier version 2.1.11 (Figure 8). Superscaffold misplacements were apparent as horizontal or vertical shifts in parts of synteny blocks in all pairwise comparisons. The overall structural integrity of the constructed PMs is visible from the expected gradual transition from gene rich to gene poor regions which in turn are well complemented by the normal high repeat region density patterns in the pericentromeric locations gradually declining towards the gene rich euchromatic regions (Figure 6). The PMs along with integrated DMDD and RH genetic maps were visualised using DMAP as described in the Methods section. Figure 7 shows a

representative illustration for chromosome 1 (Chromosomes 2-12 are shown in Figure S2). Good correspondence between DMDD and RH genetic maps and the pseudomolecules was observed.

Although the DMDD map-based strategy was critical in providing the basic anchoring to the DM genome, it had its limitations. Certain superscaffolds lacked sufficient polymorphic STS markers for genomic anchoring and were possibly affected by homozygosity, segregation distortion or other issues (Figure S1). This mainly occurred in pericentromeric/heterochromatin regions (marked by dashed lines, Figures 7, S2) which generally displayed a sparse coverage with DMDD markers, possibly due to the customised marker design strategy that precluded the design of markers in highly repetitive, relatively gene poor regions. For example SNPs were designed against coding regions using RNA-Seq data (Hamilton *et al.* 2011) and, thus, were mainly localised to gene rich regions, which occupy a different 'genomic space' to the gene poor high repeat content regions (Figure 6). The DM based 'PM series' SSRs were designed from repeat-masked genome sequence to avoid placement in repetitive DNA. The DArT methodology also employs genome complexity reduction and has been shown to target the low copy fraction of a plant genome through judicious selection of certain restriction enzymes (Jaccoud *et al.* 2001). Thus, the unavoidable bias towards non-repetitive sequences in the STS markers employed in the DMDD map resulted in many unanchored superscaffolds. This issue was resolved by using additional resources that we refer to as the '*in silico* anchoring' approach. For example, the large block of 'orphaned' superscaffolds, not directly connected to the DMDD map, stretching from DMB 394 to DMB 705 (with the exception of DMBs 193, 15, 59, 100 and 200) on chromosome 1 (see Figure 7) was anchored by the evidence derived from the WGP/AFLP-

based RH map and the tomato EXPEN map and further extended by the 'link-peak walk' strategy, illustrating the importance of the multi-layered anchoring approach adopted here.

Potato genomic resources are provided as tracks/features in the GBrowse for the DM genome (hosted at [Spud DB site "http://potato.plantbiology.msu.edu/"](http://potato.plantbiology.msu.edu/)). One such resource, widely adopted by the potato community, is the 'Illumina Potato 8303 SNP Infinium array' (Felcher *et al.* 2012) released after our map was constructed. This SNP array was employed by Felcher *et al.* (2012) to construct two genetic maps, both involving DM as the female parent. Although the homozygosity of DM precluded segregation of DM loci in these populations, they showed good congruence for most linkage groups to the pre-release version (a 'modified ver 2.1.10' latterly referred to as 'ver 2.1.11') of the DM pseudomolecules. Version 4.03 of the PMs provides an improved correspondence with the genetic maps of Felcher *et al.* (2012) (Figure 6). An updated annotation of the Illumina Potato 8303 SNP Infinium array is provided in Table S3. The DMDD genetic map and associated data files are available at <http://solgenomics.net/>, and includes hyperlinks to the MSU Genome Browser. All of the supplementary data, wherever applicable, is available to download as GFF format files from Spud DB site "<http://potato.plantbiology.msu.edu/>". The potato GBrowse including all of the hosted genomic resources/tracks/features have also been updated to the latest version (PM 4.03) of the DM pseudomolecules.

Conclusions

The integrated genetic and physical reference map presented here comprising nearly 2,500 markers, which are mostly sequence-tagged sites, provides a platform for exploiting the potato reference genome. The most obvious and immediate application is the ability to

position any sequence-based marker locus to a precise location in the DM genome. This will revolutionize trait analysis, although progress will be dependent on the complexity of the trait concerned, population size, replication and accuracy of phenotypic data and other factors that impinge on map resolution. Once mapped, the genome sequence around the locus can be used to design additional genetic markers for fine-scale mapping, and to identify putative candidate genes using the genome annotation. Such genes can be re-sequenced from informative plants showing phenotypic variation for the target trait. This ability to move directly from 'map to genome to gene' will hasten the identification of genes responsible for traits. However, the automated annotation still includes many genes of 'unknown function' and there are likely to be as yet unannotated genes in the genome sequence. Moreover, the DM genome represents only one haplotype in a species known to exhibit abundant sequence diversity.

The conversion of ~93% of the assembled genome sequence to well-structured, oriented and annotated pseudomolecules has made potato more amenable to modern genomic/genotyping approaches, such as genotyping-by-sequencing (GBS) (Uitdewilligen *et al.* 2013). The clear and irreversible shift towards sequence based polymorphism in place of 'fragment based' markers will have the effect of augmenting centimorgan (cM) positions with genome sequence co-ordinates, providing a means for verifying the accuracy of mapping studies. The integrated DMDD map complements the published potato genome sequence and adds to a growing number of resources for genetic and genomic analyses.

The integrated map presented here and associated resources will help to alleviate many of the complicating aspects of potato as a genetic system. Potato is the most economically important crop where cultivars are highly heterozygous polyploids that suffer severe inbreeding depression on self-pollination. Such breeding systems make breeding and

genetical studies difficult and cultivar development generally requires simultaneous recurrent selection for several traits over many years of evaluation. Introduction of traits that would make such crops more sustainable, e.g., drought and salinity tolerance as well as nutrient use efficiency, will be targeted as we confront global climate change and dwindling natural resources (Levy *et al.* 2013). Moreover, attempts to convert the cross-pollinated tetraploid breeding system into an F₁ hybrid diploid based scheme are also in progress (Lindhout *et al.* 2011). The isolation of genes coding for key traits, and characterisation of their functional allelic diversity will be greatly facilitated by the resources provided in this study. A recent example is the identification of a gene largely responsible for the adaptation of Andean-derived potato germplasm to the longer day-lengths of temperate latitudes (Kloosterman *et al.* 2013).

The work presented here has generated a greatly improved ordering of the potato reference genome superscaffolds into chromosomal ‘pseudomolecules’. The reconfigured pseudomolecules and their links with genetic maps provide a major new resource for the research community. They form the basis by which geneticists can identify genes underlying important traits, and through which comparative genomics can be further exploited in diversity assessment, phylogenetic inference and plant breeding.

ACKNOWLEDGEMENTS

We acknowledge the funding made available by the Potato Genome Sequencing grant - UK [Scottish Government Rural and Environmental Science and Analytical Services Division (RESAS), Department for Environment, Food and Rural Affairs (DEFRA), Agriculture and Horticulture Development Board (AHDB) - Potato Council, Biotechnology and Biological

Sciences Research Council (BBSRC, Grant BB/F012640)]; New Zealand Institute for Crop & Food Research Ltd Strategic Science Initiative and the New Zealand Institute for Plant & Food Research Ltd Capability Fund - New Zealand; NMEA (Netherlands Ministry of Economic Affairs, CBSG (Centre for BioSystems Genomics), STW (Netherlands Technology Foundation grant 07796) - Netherlands; Teagasc Core Funding, DAFF-Research Stimulus Fund - Ireland; International Potato Center (CIP-CGIAR) / CRP RTB, Peru; CONICYT (Fondap 1509007, Basal CMM, PBCT-PSD-03), CIRIC INRIA, INIA-Ministry of Agriculture of Chile - Chile; FEMCIDI OEA, PE/09/02 MINCyT-CONCyTEC, 2010-2011, Instituto Nacional de Tecnología Agropecuaria (INTA-Core Funds) and Ministerio de Ciencia y Tecnología (MINCyT) – Argentina; Proyecto FEMCIDI-OEA SEDI/AE- 305 /09 (2008-2012), Proyecto Bilateral Argentina - Perú; FINCyT (099-FINCyT-EQUIP-2009) / (076-FINCyT-PIN-2008), Prestamo BID no. 1663/OC-PE, Instituto Nacional de Innovación Agraria - Ministry of Agriculture of Peru, Peruvian Ministry of Agriculture - Technical Secretariat of coordination with the CGIAR, Consejo Nacional de Ciencia, Tecnología e Innovación Tecnológica - Peru (CONCYTEC), Special Multilateral Fund of the Inter-American Council for Integral Development (FEMCIDI-Peru). We thank Andrzej Kilian (Diversity Arrays Technology, Australia) for DArT genotyping of the DMDD mapping population. We acknowledge Peter E Hedley and Clare Booth (The James Hutton Institute, UK) for help with SNP genotyping. We thank S. B. Divito (Instituto Nacional de Tecnología Agropecuaria, Balcarce, Argentina) for technical assistance. We are also grateful to Luke Ramsay and Peter E. Hedley (The James Hutton Institute, UK) for comments on the manuscript. AFLP and WGP are (registered) trademarks owned by KeyGene N.V.

LITERATURE CITED

- Buntjer, J. B. 1999. Cross Checker. Vol. 291. Department of Plant Breeding, Wageningen University and Research Centre; Wageningen.
- Creste, S., A. T. Neto, and A. Figueira. 2001. Detection of single sequence repeat polymorphisms in denaturing polyacrylamide sequencing gels by silver staining. *Plant Mol. Biol. Rep.* 19: 299-306.
- de Boer, J. M., T. J. A. Borm, T. Jesse, B. Brugmans, L. Wiggers-Perebolte, *et al.* 2012. A hybrid BAC physical map of potato: a framework for sequencing a heterozygous genome (vol 12, 594, 2011). *BMC Genomics* 13: 423.
- Fan, J. B., A. Oliphant, R. Shen, B. G. Kermani, F. Garcia, K. L. Gunderson, M. Hansen, F. Steemers, S. L. Butler, P. Deloukas, *et al.* 2003. Highly parallel SNP genotyping. *Cold Spring Harbor Symposia on Quantitative Biology.* 68: 69-78. doi: 10.1101/sqb.2003.68.69.
- Feingold, S., J. Lloyd, N. Norero, M. Bonierbale, and J. Lorenzen. 2005. Mapping and characterization of new EST-derived microsatellites for potato (*Solanum tuberosum* L.). *Theor. Appl. Genet.* 111: 456-466.
- Felcher, K. J., J. J. Coombs, A. N. Massa, C. N. Hansey, J. P. Hamilton, *et al.* 2012. Integration of two diploid potato linkage maps with the potato genome sequence. *PLoS ONE* 7: e36347.
- Fulton, T. M., R. Van der Hoeven, N. T. Eannetta, and S. D. Tanksley. 2002. Identification, analysis, and utilization of conserved ortholog set markers for comparative genomics in higher plants. *Plant Cell* 14: 1457-1467.

- Gebhardt, C., E. Ritter, A. Barone, T. Debener, B. Walkemeier, *et al.* 1991. RFLP maps of potato and their alignment with the homoeologous tomato genome. *Theor. Appl. Genet.* 83: 49-57.
- Ghislain, M., J. Núñez, M. R. Herrera, J. Pignataro, F. Guzman, *et al.* 2009. Robust and highly informative microsatellite-based genetic identity kit for potato. *Mol. Breed.* 23: 377-388.
- Gong, Z., Y. Wu, A. Koblízková, G. A. Torres, K. Wang, M. Iovene, P. Neumann, W. Zhang, P. Novák, C. R. Buell, J. Macas, and J. Jiang. 2012. Repeatless and repeat-based centromeres in potato: Implications for centromere evolution. *Plant Cell* 24: 3559-3574.
- Green, E. D. and P. Green. 1991. Sequence-tagged site (STS) content mapping of human chromosomes: theoretical considerations and early experiences. *PCR Methods Appl* 1: 77-90.
- Hamilton, J. P. and C. R. Buell. 2012. Advances in plant genome sequencing. *Plant J.* 70: 177-190.
- Hamilton, J. P., C. N. Hansey, B. R. Whitty, K. Stoffel, A. N. Massa, *et al.* 2011. Single nucleotide polymorphism discovery in elite North American potato germplasm. *BMC Genomics* 12: 302.
- Herrera, M. R. and M. Ghislain. 2000. *Molecular biology laboratory protocols: plant genotyping*, 3rd edn. Crop Improvement and Genetic Resources Department, International Potato Center (CIP), Lima.
- International Rice Genome Sequencing Project. 2005. The map-based sequence of the rice genome. *Nature* 436: 793-800.

- Iovene, M., S. M. Wielgus, P. W. Simon, C. R. Buell, and J. M. Jiang. 2008. Chromatin structure and physical mapping of chromosome 6 of potato and comparative analyses with tomato. *Genetics* 180:1307-1317.
- Istrail, S., G. G. Sutton, L. Florea, A. L. Halpern, C. M. Mobarry, *et al.* 2004. Whole-genome shotgun assembly and comparison of human genome assemblies. *Proc. Nat. Acad. Sci. USA* 101: 1916-1921.
- Jaccoud, D., K. Peng, D. Feinstein, and A. Kilian. 2001. Diversity arrays: a solid state technology for sequence information independent genotyping. *Nucleic. Acid. Res.* 29: E25.
- Kloosterman, B., J. A. Abelenda, M. d. M. C. Gomez, M. Oortwijn, J. M. de Boer, *et al.* 2013. Naturally occurring allele diversity allows potato cultivation in northern latitudes. *Nature* 495: 246-250.
- Kurtz, S., A. Phillippy, A. L. Delcher, M. Smoot, M. Shumway, *et al.* 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5:R12.
- Levy, D., W. K. Coleman, and R. E. Veilleux. 2013. Adaptation of potato to water shortage: irrigation management and enhancement of tolerance to drought and salinity. *American Journal of Potato Research* 90: 186-206.
- Lindhout, P., D. Meijer, T. Schotte, R. C. B. Hutten, R. G. F. Visser, and H. J. van Eck. 2011. Towards F 1 Hybrid Seed Potato Breeding. *Potato Research* 54 (4): 301-312.
- Margulies, M., M. Egholm, W. E. Altman, S. Attiya, J. S. Bader, *et al.* 2006. Genome sequencing in microfabricated high-density picolitre reactors (vol 437, pg 376, 2005). *Nature* 441: 120-120.

- Milbourne, D., R. C. Meyer, A. J. Collins, L. D. Ramsay, C. Gebhardt, *et al.* 1998. Isolation, characterisation and mapping of simple sequence repeat loci in potato. *Mol. Gen. Genet.* 259: 233-245.
- Miller, J. R., S. Koren, and G. Sutton. 2010. Assembly algorithms for next-generation sequencing data. *Genomics* 95: 315-327.
- Ning, Z. M., A. J. Cox, and J. C. Mullikin. 2001. SSAHA: A fast search method for large DNA databases. *Genome Res.* 11: 1725-1729.
- Ovchinnikova, A., E. Krylova, T. Gavrilenko, T. Smekalova, M. Zhuk, *et al.* 2011. Taxonomy of cultivated potatoes (*Solanum* section *Petota*: Solanaceae). *Bot. J. Linn. Soc.* 165: 107-155.
- Park, T. H., J. B. Kim, R. C. Hutten, H. J. van Eck, E. Jacobsen, and R. G. Visser. 2007. Genetic positioning of centromeres using half-tetrad analysis in a 4x-2x cross population of potato. *Genetics* 176: 85-94.
- Paz, M. M. and R. E. Veilleux. 1999. Influence of culture medium and in vitro conditions on shoot regeneration in *Solanum phureja* monoploids and fertility of regenerated doubled monoploids. *Plant Breed.* 118: 53-57.
- Peters, S. A., J. W. Bargsten, D. Szinay, J. van de Belt, R. G. Visser, Y. Bai, and H. de Jong. 2012. Structural homology in the Solanaceae: analysis of genomic regions in support of synteny studies in tomato, potato and pepper. *Plant J* 71(4): 602-614.
- Potato Genome Sequencing Consortium. 2011. Genome sequence and analysis of the tuber crop potato. *Nature* 475: 189-195.

- Rezvoy, C. M., D. Charif, L. Gue'guen, and G. A. B. Marais. 2007. MareyMap: an R-based tool with graphical interface for estimating recombination rates. *Bioinformatics* 23: 2188-2189.
- Sliwka, J., H. Jakuczun, M. Chmielarz, A. Hara-Skrzypiec, I. Tomczyn'ska, A. Kilian, and E. Zimnoch-Guzowska. 2012. A resistance gene against potato late blight originating from *Solanum x michoacanum* maps to potato chromosome VII. *Theor. Appl. Genet.* 124 (2): 397–406.
- Spooner, D. M., J. Núñez, G. Trujillo, M. D. Herrera, F. Guzmán, *et al.* 2007. Extensive simple sequence repeat genotyping of potato landraces supports a major reevaluation of their gene pool structure and classification. *Proc. Nat. Acad. Sci. USA* 104: 19398-19403.
- Tang, J. F., S. J. Baldwin, J. M. E. Jacobs, C. G. van der Linden, R. E. Voorrips, *et al.* 2008a. Large-scale identification of polymorphic microsatellites using an *in silico* approach. *BMC Bioinformatics* 9: 374.
- Tang, X., D. Szinay, C. Lang, M. S. Ramanna, E. A. van der Vossen, E. Datema, R. K. Lankhorst, J. De Boer, S. A. Peters, C. Bachem, W. Stiekema, R. G. Visser, H. De Jong, and Y. Bai. 2008b. Cross-Species Bacterial Artificial Chromosome-Fluorescence *in Situ* hybridization painting of the tomato and potato chromosome 6 reveals undescribed chromosomal rearrangements. *Genetics* 180: 1319-1328.
- Tang, X., J. M. de Boer, H. J. van Eck, C. Bachem, R. G. Visser, and H. de Jong. 2009. Assignment of genetic linkage maps to diploid *Solanum tuberosum* pachytene chromosomes by BAC-FISH technology. *Chromosome Res.* 17: 899-915.

- Tanksley, S. D., M. W. Ganai, J. P. Prince, M. C. Devicente, M. W. Bonierbale, *et al.* 1992. High-density molecular linkage maps of the tomato and potato genomes. *Genetics* 132: 1141-1160.
- The Arabidopsis Genome Initiative. 2000. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408: 796-815.
- The French-Italian Public Consortium for Grapevine Genome Characterization. 2007. The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature* 449: 463-U465.
- The Tomato Genome Sequencing Consortium. 2012. The tomato genome sequence provides insights into fleshy fruit evolution. *Nature* 485: 635-641.
- Uitdewilligen, J. G. A. M. L., A.-M. A. Wolters, B. B. D'hoop, T. J. A. Borm, R. G. F. Visser, *et al.* 2013. A next-generation sequencing method for genotyping-by-sequencing of highly heterozygous autotetraploid potato. *PLoS ONE* 8: e62355.
- Van Ooijen, J. W. 2006. Joinmap 4: Software for the Calculation of Genetic Linkage Maps. Kyazma B. V., Wageningen, Netherlands.
- van Os, H., S. Andrzejewski, E. Bakker, I. Barrena, G. J. Bryan, *et al.* 2006. Construction of a 10,000-marker ultradense genetic recombination map of potato: Providing a framework for accelerated gene isolation and a genomewide physical map. *Genetics* 173: 1075-1087.
- Veilleux, R. E., L. Y. Shen, and M. M. Paz. 1995. Analysis of the genetic composition of anther-derived potato by randomly amplified polymorphic DNA and simple sequence repeats. *Genome* 38:1153-1162.

Vos, P., R. Hogers, M. Bleeker, M. Reijans, T. Vandelee, *et al.* 1995. AFLP: a new technique for DNA-fingerprinting. *Nucleic Acids Res.* 23: 4407-4414.

Wenzl, P., J. Carling, D. Kudrna, D. Jaccoud, E. Huttner, *et al.* 2004. Diversity Arrays Technology (DArT) for whole-genome profiling of barley. *Proc. Nat. Acad. Sci. USA* 101: 9915-9920.

Figure legends

Figure 1 Pipeline for anchoring of markers to the potato genome assembly.

Figure 2 Step-wise linkage group assignment and ordering of DM superscaffolds, using genetic anchoring information successively from the DM, RH and tomato genetic maps.

Figure 3 Summary of DM genome assembly anchoring using three different map resources. The number of uniquely and jointly anchored superscaffolds for each resource is given in the appropriate intersection. Cumulative size (Mb) in each category of superscaffolds anchored is shown in parenthesis. The total number of 649 anchored superscaffolds represents 623 Mb of the assembled DM potato genome. Figure updated from PGSC (2011).

Figure 4 Depiction of 'Link-peak' walk strategy taking superscaffold 'PGSC0003DMB000000159' as an example. (a) Custom GBrowse 'Link Peak' intensity track features (shown as red and blue arrows) provided ordered navigation through superscaffolds using the aggregated paired-end/mate-pair data (PEMP). Link peaks to the right (red arrow) indicate 'suggested path' downstream of the AGP while those to the left (blue arrow) indicate converse. Reversal of this trend indicates a negative strand for the superscaffold in question. Traversing from one superscaffold to another by taking leads from these 'link-peak' intensity tracks assisted in manually curating all 12 pseudomolecules. (b) Visualization of the underlying PEMP data.

Figure 5 Assembled BAC sequence for LuSP197F07. Each scaffold assembly is derived from paired end sequences of a combined pool of 82 DM BACs (spanning scaffolding gaps on chromosome 4) and single end sequence at higher read depth from one of the six sub-pools derived from the same BACs. The assemblies show a direct sequence running from PGSC0003DMB000000278 (- orientation, full length, cyan) through into PGSC0003DMB000000051 (+ orientation, blue) in accordance with the accessioned golden path (AGP) and fully validating the decision to split PGSC0003DMB000000278 at position 824768 and to split PGSC0003DMB000000051 at position 1859342 as indicated in the AGP file. Regions of good alignment (>98% identity, >1000 bases) are indicated as thick lines. Thin lines indicate no good alignment between the superscaffold and BAC sequences. The BAC end sequences are labelled with their Genbank IDs and are indicated at each end of the plot by black arrows. Breakpoints in the BAC sequences are indicated by orange diagonal lines and annotated with the assigned breakpoints coordinate from the AGP.

Figure 6 Enhanced accuracy of the current DM pseudomolecules (PMs). Panels A and E show anchoring of superscaffolds to the PM versions 4.03 and 2.1.11, respectively. Superscaffolds with known and unknown orientations are depicted in alternating shades of blue and red, respectively. Gaps in between the superscaffolds are marked in grey. Black areas in panel E represent unanchored superscaffolds (version 2.1.11) that were eventually anchored and ordered in PM version 4.03. Panels B and C show gene and repeat region densities, respectively, in 1 MB bins of PM version 4.03. Gene and repeat region densities ranges from 0 – >150 genes/MB and 0 – >900 repeats/MB, respectively. Panels D and F show the correspondence of the genetic maps (D84, green; DRH, black), adapted from Felcher *et al.* (2012), to PM versions 4.03 and 2.1.11, respectively. Graphs show the genetic (cM) positions plotted against the physical coordinates (Mb) for the SolCAP SNP markers; panels G (D84) and H (DRH) show elaborated examples of good correspondence from chromosome 9.

Figure 7 Illustration of the chromosome 1 pseudomolecule (PM) integrated with the DM and RH genetic maps. STS and AFLP markers anchor sequence locations in the chromosome 1 PM to the DMDD and RH genetic maps, respectively. The AFLP marker positions in the PM were identified through sequence tag alignment of BAC clones from the RH WGP physical map. Superscaffolds comprising the PM are shown as alternating grey and white rectangular blocks. The layout of the PM for each of the genetic maps is shown separately but is identical with superscaffold IDs depicted in the middle. The pachytene idiogram is adapted from the potato reference genome publication (Potato Genome Sequencing Consortium 2011). The putative centromere region and pericentromeric/heterochromatic boundaries are demarcated by asterisk and dashed lines, respectively. Each DMDD marker type is colour coded: blue = DArTs, yellow = SNPs, green = SSRs. Blue and magenta lines emerging from the RH genetic map represent AFLP anchors and the intensity of green color corresponds to the AFLP marker density per bin as reported by Van Os *et al.* (2006). Magenta lines represent AFLP markers with a

relatively inaccurate mapping position on the RH genetic map, covering an interval of 5 or more bins. Regions in the central heterochromatin where superscaffold order and orientation are not completely resolved are indicated in yellow. Inversions with the tomato sequence are indicated with red interval bars.

Figure 8 NUCmer sequence alignment dotplots for the twelve potato chromosomes using current (ver4.03, plotted on x-axis) and previous (ver2.1.11, plotted on y-axis) versions of DM pseudomolecules. Sequences aligned in forward and reverse orientations are represented by red and blue lines, respectively. Scaffold misplacements are shown as horizontal or vertical shifts in parts of the aligned blocks.

Supplementary Figure legends

Figure S1 Genome-wide patterns of marker segregation distortion in DMDD population for 1830 STS markers from different segregation categories plotted as a function of Chi-square value (y-axis) against marker physical position (x-axis) on each of the 12 potato chromosomes. Dotted, dashed and dotted-dashed lines represent Chi-square significance values at $p = 0.01$ for marker segregation categories with two, three and four genotypic classes, respectively.

Figure S2 Illustration of the chromosome 2 - 12 pseudomolecules (PMs) integrated with the DM and RH genetic maps. STS and AFLP markers anchor sequence locations in the chromosome PMs to the DMDD and RH genetic maps, respectively. The AFLP marker positions in the PM were identified through sequence tag alignment of BAC clones from the RH WGP physical map. Superscaffolds comprising the PM are shown as alternating grey and white rectangular blocks. The layout of the PM for each of the genetic maps is shown separately but is identical with superscaffold IDs depicted in the middle. The pachytene idiogram is adapted from the potato reference genome publication (Potato Genome Sequencing Consortium 2011). The putative centromere region and pericentromeric/heterochromatic boundaries are demarcated by asterisks and dashed lines, respectively. Each DMDD marker type is colour coded: blue = DArTs, yellow = SNPs, green = SSRs. Blue and magenta lines emerging from the RH genetic map represent AFLP anchors and the intensity of green color corresponds to the AFLP marker density per bin as reported by Van Os *et al.* (2006). Magenta lines represent AFLP markers with a relatively inaccurate mapping position on the RH genetic map, covering an interval of 5 or more bins. Regions in the central heterochromatin where superscaffold order and orientation are not completely resolved are indicated in yellow. Inversions with the tomato sequence are indicated with red interval bars.

Table Legends

Table 1 Distribution of 1,751 markers comprising four different classes, across the 12 chromosomes in the DMDD population, with the corresponding map and interval lengths (cM) for each chromosome.

Table 2 Anchoring statistics by chromosome for the three different physical maps, *de novo* (DM map) and *in silico* (RH-WGP map and Tomato-SGN map).

Table 3 Improvements in DM pseudomolecules before and after execution of the link peak-based orientation strategy.

Supplementary Table Legends

Table S1 Details of (A) Simple sequence repeat (SSR), (B) Single nucleotide polymorphism (SNP) and (C) Amplified fragment length polymorphism (AFLP) markers employed in DMDD genotyping.

Table S2 Location of sequence-tagged site (STS) markers employed in DMDD genotyping on the DM version 3 superscaffolds and DM version 4.03 pseudomolecules. STS markers include DArTs, SSRs and SNPs.

Table S3 Revised annotation details for the Infinium 8.3k Potato Array SNPs (Felcher *et al.* 2012) on DM version 4.03 pseudomolecules.

Table S4 Genetic and physical locations of STS markers (DArTs, SSRs and SNPs) mapped in DMDD and anchored in DM version 4.03 pseudomolecules.

Table S5 Paracentric inversions between potato and tomato chromosomes detected by dot plot alignments between the chromosome pseudomolecules V4.03 of potato line DM and V2.40 of tomato cv. 'Heinz 1706'.

Table S6 Summary of six BAC pools sequence assembly data comprising 82 DM BAC clones used for validating link peak-based orientation strategy for chromosome 4.

Table S7 BAC pool assembly and validation details for chromosome 4 pseudomolecule version 4.03.

Table S8 Centromere localisation in DM V3 sequence assembly.

Table S9 Accessioned Golden Path (AGP) for the reference DM chromosome-scale pseudomolecules version 4.03. File also includes revised annotation details for potato genes and repeat regions (Potato Genome Sequencing Consortium 2011) and a list of chimeric superscaffolds.

Table 1 Distribution of 1,751 markers comprising four different classes, across the 12 chromosomes in the DMDD population, with the concomitant map and interval lengths (cM) for each chromosome.

Chr ^a	Mapped markers ^b	Map length (cM)	Interval spacing (cM/interval) ^c
01	201	93.0	0.46
02	221	77.4	0.35
03	134	101.8	0.77
04	143	99.7	0.70
05	107	64.1	0.61
06	134	70.5	0.53
07	108	67.1	0.63
08	176	67.8	0.39
09	152	87.9	0.58
10	144	68.9	0.48
11	108	62.9	0.59
12	123	75.2	0.62
All	1751	936.2	0.54

^aBased on the SSRs mapped in previous studies and further confirmed by using *in silico* approaches.

^bExcluding 718 co-segregating markers; when the segregation pattern of two or more markers was identical, only a single marker per set of identical markers was retained to generate the maps. As well as excluding 128 ungrouped markers (including 15 unassigned co-segregating markers) that did not fit any linkage group.

^cCalculated as the map length divided by the number of intervals (mapped markers minus 1, for 'total' it is mapped markers minus 12).

Table 2 Anchoring statistics by chromosome for the three different physical maps, *de novo* (DM) and *in silico* (RH and tomato).

Chromosome	DM Map			RH-WGP Map			Tomato-SGN Map		
	DMB anchored	Cumulative length (Mb)	No. of markers*	DMB anchored	Cumulative length (Mb)	No. of markers	DMB anchored	Cumulative length (Mb)	No. of markers
01	39	45	162	69	80	208	43	41	271
02	35	43	175	35	43	120	33	40	233
03	19	24	108	28	27	73	41	45	194
04	34	47	138	51	57	168	40	39	174
05	20	27	74	33	45	137	25	30	112
06	29	34	108	44	46	119	34	34	133
07	26	24	89	35	39	122	32	31	136
08	32	32	152	24	23	57	40	32	129
09	27	28	109	34	33	91	40	39	136
10	31	38	106	34	44	102	26	32	110
11	20	26	113	36	38	110	22	26	116
12	22	26	72	47	52	164	26	28	109
Total	334	394	1406	470	527	1471	402	417	1853

* Only markers mapped in DMDD and uniquely and reliably anchored to DM assembly are included.

Table 3 Improvements in DM pseudomolecules before and after execution of the link peak-based orientation strategy.

Chr	Stage-I*		Stage-II**		DMB oriented ^a	
	<i>DMB anchored</i>		<i>DMB anchored</i>		<i>DMB oriented^a</i>	
	No (size in Mb)	N ₅₀	No (size in Mb)	N ₅₀	No (size in Mb)	%age
01	83 (79.7)	1.7	123 (82.6)	2.6	121 (79.8)	96.6
02	51 (45.0)	1.3	68 (45.3)	2.2	68 (45.3)	100.0
03	53 (45.3)	1.6	103 (57.2)	4.3	103 (57.2)	100.0
04	73 (60.9)	1.2	120 (66.3)	2.9	119 (62.1)	93.7
05	41 (44.8)	1.7	52 (49.5)	2.9	47 (40.4)	81.6
06	63 (54.0)	1.4	90 (55.1)	2.7	90 (55.1)	100.0
07	52 (50.6)	1.8	78 (52.9)	7.2	78 (52.9)	100.0
08	51 (41.6)	1.2	91 (52.4)	4.9	91 (52.4)	100.0
09	61 (50.6)	1.2	86 (57.3)	8.3	85 (55.9)	97.7
10	50 (51.4)	1.5	77 (56.0)	4.1	74 (55.4)	99.0
11	35 (34.4)	1.4	60 (42.5)	5.7	60 (42.5)	100.0
12	61 (58.5)	1.5	77 (57.4)	1.9	76 (56.0)	97.7
Total	674 (616.8)^b	1.5^c	1025^{b,d} (674.4)^b	4.1^c	1012^{b,d} (655.1)^b	97.2^c

*Refers to the status of PMs before execution of the 'Link-peak' walk strategy

**Refers to the status of PMs after execution of the 'Link-peak' walk strategy

^aOnly attempted at Stage-II

^bTotal

^cAverage

^dChimeric superscaffolds have been included more than once (net number of DMBs anchored=951)

Potato
Assembly

Marker
Sequences

Align


Remove alignments
caused by
repetitive sequences


Tile
high-scoring segment pairs (HSPs)
into hits

Pair and filter
forward and reverse sequences
(as applicable)

Remove ambiguous markers

Marker-assembly
anchor points





A

Potato Superscaffolds: 1.388 Mbp from PGSC0003DMB000000159:1..1,388,432

Browser [Select Tracks](#) [Upload and Share Tracks](#) [Preferences](#)

Search

Landmark or Region:

PGSC0003DMB000000159:1.. Search


Examples: PGSC0003DMB000000031, PGSC0003DMB000000115, PGSC0003DMB000000507, PGSC0003DMB000000687, PGSC0003DMB000000725, PGSC0003DMB000000757, PGSC0003DMB000001135, PGSC0003DMB000001301.

Data Source

Potato Superscaffolds

Scroll/Zoom: Show 1.388 Mbp Flip

Overview



B



Breakpoint Validation - LuSp197F07

