



## Artículo

SUPERNUMERARY TEETH IN *Necromys lasiurus*  
(RODENTIA, CRICETIDAE): THE FIRST RECORD  
IN SIGMODONTINAE

Gustavo Simões Libardi\* and Alexandre Reis Percequillo

Departamento de Ciências Biológicas, Escola Superior de Agricultura "Luiz de Queiroz". Universidade de São Paulo, Av. Pádua Dias, 11, Caixa Postal 9, 13418-900, Piracicaba, São Paulo, Brazil [correspondence: Alexandre Percequillo <[percequillo@usp.br](mailto:percequillo@usp.br)>].

\* Present address: Unidad de Investigación Diversidad, Sistemática y Evolución, Centro Nacional Patagónico, Boulevard Almirante Brown 2915, Casilla de Correo 128, 9120 Puerto Madryn, Chubut, Argentina.

**ABSTRACT.** There are several kinds of anomalies related to the teeth reported in mammals, and one of them is known as supernumerary teeth and is commonly noticed in the literature, being reported in nearly all orders of mammals. Here, we report the occurrence of supernumerary molars in the sigmodontine rodent *Necromys lasiurus*, a common and widespread small rodent of tribe Akodontini. In order to assess the patterns of morphometric variation of *N. lasiurus* in eastern South America, we examined the skulls of 1763 specimens. The supernumerary molars were found in 2 individuals, which represent a frequency of 0.11%. The origin of supernumerary teeth started a long discussion on tooth homology, whether this anomaly is a result of heredity, a mutation or an atavism. Based on the morphologies presented by the extra teeth encountered in *N. lasiurus* and on the evidences and hypothesis available in the literature, we consider that the presence of these supernumerary molars in one of the studied animals could be considered an atavism, and we also consider the extra molar presented by the other specimen is a malformation, due to some kind of random process occurred in later stages of the development of distal dental germ. To the best of our knowledge, this represents the first record of supernumerary molars in the subfamily Sigmodontinae and one of the few known cases in rodents.

**RESUMO.** Dentes supranumerários em *Necromys lasiurus* (Rodentia, Cricetidae): primeiro registro em Sigmodontinae. Diversos tipos de anomalias dentárias já foram descritas para mamíferos e uma delas é conhecida como dente supranumerário comumente reportada na literatura para quase todas as ordens de mamíferos. No presente trabalho apresentamos a ocorrência de molares extras em *Necromys lasiurus*, uma espécie de pequeno roedor sigmodontíneo comum e amplamente distribuído, membro da tribo Akodontini. Durante o levantamento de dados para avaliação morfométrica da variação da espécie no leste da América do Sul, foram examinados os crânios de 1763 espécimes. Os dentes supranumerários foram encontrados em dois indivíduos, representando 0.11% da amostra investigada. A ocorrência dos dentes supranumerários há muito tempo levanta questões acerca da homologia dentária, das mutações e dos atavismos que podem estar relacionados à origem desta anomalia. Baseados na morfologia apresentada pelos molares extra numerários de *N. lasiurus*, nós consideramos que a presença dos mesmos em um dos espécimes é um atavismo, e no outro, trata-se de uma malformação devido a um processo de origem tardia no desenvolvimento das gemas dentárias. Com base em nosso conhecimento de outras espécies e em toda a revisão bibliográfica, a ocorrência dos dentes supranumerários em *N. lasiurus* trata-se dos primeiros registros da anomalia para a subfamília Sigmodontinae e um dos poucos já registrados para a ordem Rodentia.

**Key words:** Akodontini. Brazil. Development. Extra teeth. Molar. Premolar.

**Palavras-chave:** Akodontini. Brasil. Dentes extras. Desenvolvimento. Molar. Pré-molar.

Recibido 18 abril 2014. Aceptado 3 julio 2014. Editor asociado: UFJ Pardiñas

## INTRODUCTION

There are several kinds of anomalies related to the teeth reported in mammals captured in nature, such as congenital agenesis (Smith et al., 1977; Beaver et al., 1981; Stewart and Stewart, 1987); irregular placement (Heran 1970; Beaver et al., 1981; Smith et al., 1977); divided tooth and other crown anomalies (Van Gelder and McLaughlin, 1961; Wolsan, 1983; Stewart and Stewart, 1987; Smith et al., 1977; Angelici and Luiselli, 1999); post trauma repair (Graipel et al., 1997), and many others (see Smith et al., 1977, and Winer et al., 2012, for more examples).

One of them is known as supernumerary teeth and is commonly noticed in the literature, being reported in nearly all orders of mammals, mostly in Didelphimorphia (Takahashi, 1974; Moraes et al., 2001), Pilosa (McAfee and Naples, 2012), Primates (Berkovitz and Musgrave, 1971; Wolsan, 1984; see also Smith et al., 1977), Soricomorpha (Hooper, 1946; Feldhamer and Stober, 1993), Chiroptera (Ramírez-Pulido and Müdspacher, 1987; Rui and Drehmer, 2004), Carnivora (Paradiso, 1966; Beaver et al., 1981; Stewart and Stewart, 1987; Drehmer and Ferigolo, 1996; Graipel et al., 1997; Anderson and Ozolins, 2000; Peters et al., 2013; Winer et al., 2012), Artiodactyla (Knowlton and Glazener, 1965; Chaplin and Atkinson, 1968; Pekelharing, 1968; Zurowski, 1970; Steele and Parama, 1979; Natsume et al., 2005), and Rodentia, both extant (Johnson, 1952; Krutzsch, 1953; Dearden, 1954; Harris and Fleharty, 1962; Sheppe, 1964; Sofaer and Shaw, 1971; Goodwin, 1998; Angelici and Luiselli, 1999) and extinct (Goodwin, 1998; Arnal and Vucetich, 2011). The occurrence of supernumerary teeth in rodents is generally related to extra molars placed posteriorly the toothrow (Johnson, 1952; Sofaer and Shaw, 1971) and is comparatively rarer in natural populations other than Carnivora (Wolsan, 1984), which is the group with higher number of documented cases in literature.

In the present contribution we report the occurrence of supernumerary molars in the sigmodontine rodent *Necromys lasiurus* (Lund, 1840). This species is a common and wide-

spread small rodent of tribe Akodontini that inhabits preferentially open vegetation areas, on the Caatinga, Cerrado, and Chaco biomes, as well as grasslands and shrublands that occur in the central portion of South America (Musser and Carleton, 2005). However, being an extremely versatile rodent, this species is also found in transitional areas, margins of forests and secondary forests, as well as anthropogenic habitats, like plantations and pastures (Mares et al., 1986; Henriques et al., 1997; Pardiñas et al., in press).

## MATERIALS AND METHODS

In order to assess the patterns of morphometric variation of *N. lasiurus* in Brazil, we examined the skulls of 1763 specimens deposited in the following Brazilian collections: Universidade Estadual do Mato Grosso, Cuiabá (UFMT); Universidade Federal de Lavras, Lavras (CMUFLA); Universidade Federal de Minas Gerais, Belo Horizonte (UFMG); Museu de Ciências Naturais, Pontifícia Universidade Católica, Belo Horizonte (MCN-M); Museu de História Natural Capão da Imbuia, Curitiba (MHNCI); Museu Nacional da Universidade Federal do Rio de Janeiro, Rio de Janeiro (MN); Museu de Zoologia da Universidade de São Paulo, São Paulo (MZUSP); Museu de Zoologia da Universidade Estadual de Campinas (ZUEC); and Museu de Ciências Naturais, Universidade Luterana do Brasil, Canoas (MCNU).

The nomenclature employed for dental morphology follows Reig (1977) and dental measurements (length of upper molar series) are accordingly to Percequillo et al. (2008). For age determination we followed Libardi (2013) and Voss (1991), and recognized 5 classes defined by tooth eruption and wear of major cusps, lophs and flexi.

In order to compare the length of the supernumerary toothrows to regular series presented by the same individuals and by other specimens, we calculated the descriptive statistics (mean, standard deviation, minimum and maximum values) and designed a boxplot graph, using the descriptive information of length of toothrow from individuals with supernumerary molars and from some selected samples. We gathered individuals of the same age class and same or nearby localities of the specimens with extra teeth to avoid the effects of other variation factors that could mislead the interpretation of the comparisons of the toothrow length, once the species seems to present age variation and geographic structure in toothrow and other skull dimensions, as highlighted

by Macêdo and Mares (1987) and Libardi (2013). A list of the specimens employed in comparisons from the same and nearby localities where the individuals with supernumerary teeth were found is provided in the **Appendix**.

We calculated the frequency of occurrence of extra teeth in *N. lasiurus* by simply dividing the number of specimens captured in nature presenting the oral feature by the total number of analyzed individuals, and therefore multiplying by 100 in order to obtain a value expressed in percentage. We also provide a list of frequencies and extra teeth positions on several mammalian orders, including Rodentia, in order to compare our findings to some described throughout in past decades for mammals. Some orders were not assessed due to the few information available or due to the unique nature of their dentition, as Cingulata and Cetacea.


We expressed the value of the frequencies of the revised literature, when available, in percentage as it appears in the article; when it was not directly informed, we applied here the same method as to *N. lasiurus*; when no information on total number of analyzed individuals were available, we did not calculate any frequency, providing then a “?” symbol.

The positions of extra teeth in each mammalian species revised are based on the accuracy of the information of the descriptions and figures provided by the authors. It is important to emphasize that we used the same nomenclature employed for the mammalian species in the articles.

## RESULTS

The supernumerary molars were found in 2 individuals out from a sampling of 1763 specimens of *N. lasiurus*, which represent a frequency of 0.11%. The specimen MN 44035 is an adult male age class 2, collected at Gleba José Nani (also Bairro Santa Luzia), Caçapava municipality, São Paulo state, Brazil (23°06'S, 47°36'W; see Weksler and Bonvicino, 2005), collected on May 7, 1973, during the field survey performed by the Campaign for Combating Schistosomiasis (CACEsq). The MN 44035 present 2 extra teeth, one at each upper tooththrow, at the distal end of dental series (**Figs. 1A** and **1B**). The extra teeth are molar-like, covered by enamel, present reduced wear and no dentine exposition, peg-like, with no loph or flexi discernible on molar topography and completely aligned with the other molars of both tooththrows; these supernumer-

ary teeth exhibit the crown positioned slightly higher than the remaining teeth (**Fig. 2A**). All remaining teeth show reduced wear and present a typical configuration of other specimens of *N. lasiurus* from other localities observed with similar molar wear. The total length of the upper right tooththrow is 5.29 mm (ca. 4.60 mm without the extra molar) and of the upper left tooththrow is 5.25 mm (ca. 4.48 mm without the extra molar). The mandible does not show any supernumerary teeth and all molars present the typical morphology. Both upper and lower series occluded perfectly, with the supernumerary teeth being positioned posteriorly to the lower third molars. It was also noted that the presence of the extra teeth did not cause any kind of injury on the other teeth, as well as no evidence of excessive wear on the posterior



**Fig. 1.** *Necromys lasiurus*: Occlusal view of the left upper tooththrow of specimens MN 44035 (A, right; B, left), from Caçapava, São Paulo; and UFMT 839 (C, right; D, left), from Rio Juruena, Mato Grosso, presenting the supernumerary molars, distally to other tooth; length of molar series provided in **Table 1**. Scale = 1 mm.

surface of the lower third molar, or even on the hard palate.

The individual UFMT 839 is an adult, age class 3, presents no information on sex, and was collected at Rio Juruena, Sapezal municipality, Mato Grosso state, Brazil (13°27'24"S, 59°00'07"W) on August 22, 2006. In this specimen, there is only 1 extra tooth in the left upper tooththrow, also positioned at the distal end of this dental series (Figs. 1C and 1D). The extra tooth is molar-like, enameled, with some wear on occlusal surface and small dentine exposition, with no lophs or flexi visible and its crown is positioned slightly higher than the remaining of molar series (Fig. 2B). The extra tooth is erupted transversally towards the labial direction, and the entire tooththrow is unaligned, producing no precise occlusion of the upper and lower molar series. Moreover, the

alveolus of the third tooth and the supernumerary tooth are confluent and very wide, probably with some maxillary bone reabsorption. The lack of alignment of the tooththrow, especially the supernumerary tooth, suggests that there could have been some injury to the gum and the roots during the chewing movements. The total length of the upper right tooththrow is 4.56 mm and of the upper left tooththrow is 5.17 mm (ca. 4.56 mm without the extra molar).

We compared the length of the molar series of the specimens MN 44035 and UFMT 839 to the tooththrow length of syntopic individuals, from Caçapava and Rio Juruena with the same molar wear (Table 1 and Fig. 3). Due to the existence of only 1 specimen with normal dentition from Caçapava, we decided to generate a boxplot considering also the individuals from Estação Biológica de Boracéia, Salesópolis, São Paulo state, the best available sample in the region, and which is nearly 60 km to the south of Caçapava. Table 1 shows the results of the descriptive statistics: in all comparisons, the molar series with extra molars are consistently longer than regular molar series with three teeth. It is also interesting to notice that the measurements of the right normal series of specimen UFMT 839 and of the molar series of MN 44035 excluding the extra teeth are within the limits of variation of samples employed in comparisons.




Fig. 2. *Necomys lasiurus*: Lingual lateral view of the upper left tooththrow of specimens MN 44035, from Caçapava, São Paulo, and UFMT 839, from Rio Juruena, Mato Grosso, both presenting the supernumerary distal molars. It is possible to observe the occlusal surface of the extra teeth positioned above the surface of the other teeth; length of molar series provided in Table 1. Scale = 1 mm.


## DISCUSSION

The frequency of occurrence of the supernumerary teeth in *N. lasiurus* is low (0.11%), and we are confident that this represents information for the species (and not for a restricted sample), since the sample we studied is quite robust and presents specimens throughout the distribution of the species. It is noteworthy that this frequency is smaller than those frequencies exhibited by many other species: apparently in felids, which experienced drastic reductions on the number of post canine teeth, the expression of supernumerary teeth (especially molars) is more common than in other groups; in marsupials, which present complete dentitions with all possible teeth, the frequency is similar to that observed in *N. lasiurus*. The position of the

**Table 1**

Descriptive statistics of the tooththrow lengths of the samples of *Necomys lasiurus* which presented normal molars and of the molar rows of the individuals with supernumerary molars. The “\*” indicates absolute values; N = sampling number; SD = standard deviation. All measurements are provided in millimeters.

Sample or specimen	N	Mean ± SD	Range
MN 44035 (right tooththrow)	-	5.29*	-
MN 44035 (left tooththrow)	-	5.25*	-
UFMT 839 (right tooththrow)	-	4.56*	-
UFMT 839 (left tooththrow)	-	5.17*	-
MN 44033 (from Caçapava, with normal tooththrows)	-	4.91*	-
Boracéia sample	17	4.55 ± 0.168	4.22-4.87
Rio Juruena sample	4	4.79 ± 0.092	4.69-4.91



**Fig. 3.** Individual values and box-plot of the length of the upper tooththrows of MN 44035 and UFMT 839 and of individuals from the same (Caçapava and Rio Juruena) and near localities (Boracéia) without supranumerary teeth for *Necomys lasiurus*. The black dashes represent the individual values for individual specimens or the mean value for samples; the gray boxes represent the quartiles and the whiskers represent the maximum and minimum values of each sample.

extra teeth found in *N. lasiurus* is also congruent to the other cases in Rodentia (Table 2).

The origin of supernumerary teeth started a long discussion on tooth homology (Wolsan, 1984; Peterkova et al., 2006; see also Hinton, 1923 for former hypothesis), whether this anomaly is a result of heredity, a mutation or an atavism. The present dentition of murid rodents is currently understood as a result of the loss of the canines, incisors and premolars, and only the molars and a pair of incisors remained (see Peterkova et al., 2006). Nevertheless, the occurrence of extra molars

posterior to the current third molar, raised the hypothesis that the first molar could be a modified premolar and the dental formula of rodents would be 1 incisor, 1 premolar and 2 molars (premolar theory): therefore, the extra tooth would represent the expression of a suppressed third molar during development (Johnson, 1952; Sheppe, 1964). Based on the timing of developmental patterns of the cheekteeth of embryos of *Mus musculus* in comparison to the timing of human embryos, Peterkova et al. (2006) suggest that the rodent M1, M2 and M3 teeth could correspond to the human dP3, dP4 and M1, respectively. This configuration would explain the presence of supernumerary teeth in the posteriormost portion of the maxillary.

The supernumerary teeth found in MN 44035 are nearly conventional, being aligned and symmetrically bilateral. This configuration could result of heredity (see Sofaer and Shaw, 1971),



**Table 2**

Frequencies and kinds of extra teeth found in other mammalian taxa disponible in literature. The superscripted and subscripted numbers indicate the upper and lower teeth, respectively, when it was posible to determine their position precisely; n.c. = not comparable.

Species	Frequency (%)	Tooth	Literature
<b>DIDELPHIMORPHIA</b>			
<i>Caluromys philander</i>	1.2	distal lower right molar	Moraes et al., 2001
<i>Chironectes minimus</i>	0.7	distal lower right molar	Moraes et al., 2001
<i>Didelphis</i> sp.	3.2	incisors	Takahashi, 1974
<i>Didelphis albiventris</i>	0.5	distal molars	Moraes et al., 2001
<i>D. aurita</i>	0.3	distal upper molars	Moraes et al., 2001
<i>D. marsupialis</i>	1.0	molars	Moraes et al., 2001
<i>Philander andersoni</i>	2.8	upper left molar	Moraes et al., 2001
<i>P. frenata</i>	0.8	distal upper molars	Moraes et al., 2001
<i>P. opossum</i>	0.3	distal upper molars	Moraes et al., 2001
<b>DIPROTODONTIA</b>			
<i>Petrogale concinna</i>	n.c.	distal lower molars	Rodrigues et al., 2011
<b>SIRENIA</b>			
<i>Trichecus</i> sp.	n.c.	distal lower molars	Domning and Hayek, 1984; Rodrigues et al., 2011
<b>PILOSA</b>			
<i>Choloepus didactylus</i>	?	right upper molar	McAfee and Naples, 2012
<i>Choloepus hoffmanni</i>	?	duplicated M1 and right M2	McAfee and Naples, 2012
<b>PRIMATES</b>			
<i>Callimico goeldii</i>	?	distal upper left molar	Schwartz, 1984
<i>Cebus</i> sp.	?	lower premolars	Schwartz, 1984
<i>Leontopithecus rosalia</i>	?	distal lower molars	Schwartz, 1984
<i>Gorilla</i> sp.	?	distal lower molars	Schwartz, 1984
<i>Hylobates</i> sp.	?	distal molars	Schwartz, 1984
<i>Pongo</i> sp.	?	distal lower and upper molars	Schwartz, 1984
<i>Pongo pygmaeus</i>	?	upper premolars	Berkovitz and Musgrave, 1971
<b>RODENTIA</b>			
† <i>Democricetodon</i>	?	distal lower molars	Mein, 1986
† <i>Sciamys principalis</i>	1.51	distal lower right molar	Arnal and Vucetich, 2011
† <i>Spermophilus richardsonii</i>	0.5	distal upper molars	Goodwin, 1998
<i>Heliophobius argenteocinereus</i>	n.c.	distal lower molars	Rodrigues et al., 2011
<i>Hystrix cristata</i>	0.55	distal upper molars	Angelici and Luiselli, 1999
<i>H. leucura</i>	?	distal molar	Johnson, 1952
<i>Mesembriomys gouldi gouldi</i>	?	distal upper right molar	Johnson, 1952
<i>Microtus agrestis</i>	?	distal molar	Johnson, 1952; Hinton, 1923
<i>Necromys lasiurus</i>	0.113	distal upper molars	this study

Species	Frequency (%)	Tooth	Literature
<i>Peromyscus</i> sp.	0.071	distal upper molars/distal upper left molar	Sheppe, 1964
<i>Proechimys</i> sp.	?	distal molar	Johnson, 1952
<i>Saccostomus hildae</i>	?	distal molar	Johnson, 1952; Hinton, 1923
<i>Spermophilus richardsonii</i>	0.105	distal upper molars	Goodwin, 1998
hybrid <i>Spermophilus</i>	5.88	distal upper molars	Goodwin, 1998
<i>Zapus princeps</i>	?	distal upper molars	Krutzsch, 1953
SORICOMORPHA			
<i>Blarina brevicauda</i>	0.5	distal upper molars	Feldhamer and Stober, 1993
<i>Cryptotis parva</i>	0.8	upper right premolar	Feldhamer and Stober, 1993
<i>Sorex saussurei</i>	?	upper right unicuspid	Hooper, 1946
CHIROPTERA			
<i>Artibeus fimbriatus</i>	2.27	upper right post-canine	Rui and Drehmer, 2004
<i>A. lituratus</i>	1.92	upper incisor/right M3	Rui and Drehmer, 2004
<i>A. l. palmarum</i>	?	upper incisors	Ramirez-Pulido and Müdespacher, 1987
<i>Balantiopteryx plicata</i>	4.12	upper incisors	Ramirez-Pulido and Müdespacher, 1987
<i>Carollia subrufa</i>	1.56	lower right incisor	Ramirez-Pulido and Müdespacher, 1987
<i>Choeronycteris mexicana</i>	26.5	upper incisors/upper canines	Ramirez-Pulido and Müdespacher, 1987
<i>Glossophaga soricina</i>	2.03	incisors/upper premolars	Ramirez-Pulido and Müdespacher, 1987
<i>Myotis velifer</i>	0.64	upper right incisor/upper left premolars	Ramirez-Pulido and Müdespacher, 1987
CARNIVORA			
<i>Arctocepalus australis</i>	?	upper right post-canine	Drehmer and Ferigolo, 1996
<i>A. tropicalis</i>	?	lower post-canines	Drehmer and Ferigolo, 1996
<i>Canis latrans</i>	1.4	premolars and molars	Paradiso, 1966
<i>C. lupus</i>	5.3-6.83	several places	Andersone and Ozolins, 2000; Buchalkzyc et al., 1981
<i>Conepatus</i> sp.	1.0	upper left incisor	Bateman, 1970
<i>Felis bengalensis</i>	78.1	P2	Wolsan, 1984
<i>Lutra canadensis</i>	4.3	P1/near P1/distal upper molar	Dearden, 1954; Beaver et al., 1981
<i>Lynx lynx</i>	3.0-30.0	M2	Kvam, 1985; Wolsan, 1984
<i>Meles meles</i>	?	premolars/M2	Ruprecht, 1965
<i>Mustela</i> sp.	13.7	upper incisors	Bateman, 1970
<i>Mustela putorius</i>	?	P3	Ruprecht, 1965
<i>Otaria byronia</i>	0.77	lower post-canines	Drehmer et al., 2004
<i>Otocyon megalotis</i>	3.17	lower and upper molars	Van Valen, 1964
<i>Puma concolor</i>	5.26	microdont tooth between upper canine and P2	Graipel et al., 1997
ARTIODACTYLA			
<i>Alces alces</i>	3.7	lower left incisors/lower right molar	Steele and Parama, 1979

Species	Frequency (%)	Tooth	Literature
<i>Capreolus capreolus</i>	0-16.7	upper canines	Chaplin and Atkinson, 1968
<i>Cervus canadensis</i>	0.77	duplicated left M3	Pekelharing, 1968
<i>C. elaphus</i>	0.17	duplicated M3	Pekelharing, 1968
<i>Naemorhedus crispus</i>	1.3	several places	Natsume et al., 2005
<i>Odocoileus virginianus</i>	0.013-17.9	upper canines	Knowlton and Glazener, 1965
<i>O. v. borealis</i>	?	lower left incisor	Fowle and Passmore, 1948

once the symmetry in position and morphology between the extra teeth suggests that their formation occurred before the differentiation of both sides of maxillae, so resulting from a genetically inducted formation process of the tooth germ instead of developmental interferences, mutation or random split of one single germ (Wolsan, 1984; Goodwin, 1998). Although the possibility of these extra teeth to have hereditary origins, we did not find any other *N. lasiurus* specimen from Caçapava and nearby localities, which counted 75 individuals (**Appendix**). This information, together with all the records we gathered from available bibliography (**Table 2**), is in agreement with the statement that supernumerary teeth are rare in natural populations of sigmodontine rodents.

In the other hand, the asymmetric eruption of the extra molar and the lack of alignment of tooththrow found in UFMT 839 seem to be related to other random developmental or mutation processes (Wolsan, 1984:131; Goodwin, 1998). The occurrence of abnormal morphologies and unusual patterns of eruption is far more common in literature than the configuration observed in specimen MN 44035 (e. g., Harris and Fleharty, 1962; Smith et al., 1977; Graipel et al., 1997; Anderson and Ozolins, 2000; Winer et al., 2012).

Based on the morphologies presented by the extra teeth encountered in *N. lasiurus* and on the evidences and hypothesis available in the literature, we consider more likely that the supernumerary molars of specimen MN 44035 is in fact a result of genetic heredity (based on Sofaer and Shaw, 1971), once its alignment with the molar series, which is also well developed, suggests they originated at the earlier stages of

embryonic development. Thus, if the hypothesis advocated by Johnson (1952) and Peterkova et al. (2006), in which the present dentition of muroid rodents is composed by the dP3, dP4 and M1 of the basic eutherian dentition, are true, the presence of these supernumerary molars in this specimen of *N. lasiurus* could be considered an atavism, i.e., the reappearance of a lost ancestral character (Peterkova et al., 2006). We also consider that the extra molar presented by the specimen UFMT 839 is a malformation, due to some kind of stochastic process during later stages of development of the distal tooth germ (see Glasstone, 1952; Wolsan, 1984:131), rather than earlier developmental process like reported by Sofaer and Shaw (1971). We cannot hypothesize if in this case the origin of supernumerary tooth is related or not to any kind of atavism, and if it is related to the hypothesis proposed by Johnson (1952) or Peterkova et al. (2006).

As observed in other species of rodents (**Table 2**), the positions of supernumerary teeth in both *N. lasiurus* specimens are posterior to the “normal” molar series, and therefore if we accept the conventional hypothesis that rodent molars are M1, M2 and M3, these teeth would represent a “fourth molar” due to the origin of a new dental germ, an unlikely event. However, if we accept the hypothesis regarding the dental formula of rodents, composed by the dP3, dP4 and M1, as valid (Johnson, 1952; Peterkova et al., 2006), these extra teeth would be the M2. As such, their presence could be more likely explained and would adjust adequately to the recurrent expression of supernumerary teeth in the same position for the order Rodentia, in natural populations; experimental breeding



revealed that this seems to be the tendency in *Oryzomys palustris* (Sofaer and Shaw, 1971). The fewer cases of occurrence of extra teeth in other positions reported for Rodentia (Goodwin, 1998:1166) are more difficult to relate to this pattern and also to atavistic origins, compared to the occurrence of the distal extra molars. Nevertheless, Rodrigues et al. (2011) have reported a special case of continuous dental replacement in the African mole-rat *Heliophobius argenteocinereus* (Rodentia: Bathyergidae), which is related to the supernumerary molars. As the authors claim, this kind of unique teeth replacement system, which occurs also only in the species of the manatee genus *Trichechus* and in the pygmy rock-wallaby *Petrogale concinna*, is a convergent mechanism which combines specific dental traits, as dental mesial drift, delayed eruption and the occurrence of supernumerary molars. In the *H. argenteocinereus* case, the supernumerary molars emerge in the distal portion of tooththrows continuously and drift anteriorly as the frontal teeth are worn and atrophied and, subsequently, reabsorbed (see Rodrigues et al., 2011). Although it is difficult to determine the number of teeth, the occurrence of extra teeth agrees with the general pattern observed in Rodentia, with extra teeth emerging distally.

The occurrence of supernumerary teeth in *N. lasiurus* seems to be an odd event in the subfamily Sigmodontinae. Although one of the authors (ARP) personally analyzed nearly 7000 individuals of several species of the tribe Oryzomyini, no extra teeth were observed. U. F. J. Pardiñas (pers. comm.) reported to us the occurrence of extra molars in the same position and presenting similar morphology as presented the individual MN 44035, in a partial skull of *Abrothrix olivacea* (Sigmodontinae, Abrotrichini) found by D. Procopio in an owl pellet from Southern Patagonia. The only other records on literature are from experimental colonies of rice rats (Sofaer and Shaw, 1971).

## CONCLUSIONS

To the best of our knowledge, this represents the first record of supernumerary molars in wild-caught specimens in the subfamily Sig-

modontinae and one of the few known cases in rodents, and it is very likely that these teeth represent a vestigial structure. Considering the complex and poorly understood history of the evolutionary pathways on the origin and evolution of mammalian dentition, we hope that this contribution provide useful information on the knowledge of this important subject.

## ACKNOWLEDGEMENTS

We are thankful to all curators and responsible staff of the collections visited, and especially to J. A. Oliveira and R. V. Rossi, curators of MN-UFRJ and UFMT, respectively, for providing us the material with extra teeth for detailed analysis. We are also grateful to Ulyses F. J. Pardiñas for sharing his information on extra teeth in genus *Abrothrix*. At last, we would like to acknowledge L. Soilbenzon and 2 anonymous reviewers for their suggestion for the improvement of this contribution. This work received financial support from CAPES and FAPESP (Process 09/16009-1).

## LITERATURE CITED

- ANDERSON Z and J OZOLINS. 2000. Craniometrical characteristics and dental anomalies in wolves *Canis lupus* from Latvia. *Acta Theriologica* 45:549-558.
- ANGELICI FM and L LUISELLI. 1999. Extra teeth and dental anomalies in the crested porcupine *Hystrix cristata*, from Sicily. *Acta Theriologica* 44:219-223.
- ARNAL M and MG VUCETICH. 2011. First record of supernumerary teeth in South American fossil rodents. *Journal of Vertebrate Paleontology* 31:925-927.
- BATEMAN JA. 1970. Supernumerary incisors in Mustelids. *Mammal Review* 1(3):81-86.
- BEAVER TD, GA FELDHAMER, and JA CHAPMAN. 1981. Dental and cranial anomalies in the river otter (Carnivora: Mustelidae). *Brimleyana* 7:101-109.
- BERKOVITZ BKB and JH MUSGRAVE. 1971. A rare dental abnormality in an adult male Orangutan (*Pongo pygmaeus*): Bilateral supernumerary maxillary premolars. *Journal of Zoology*, London 164:266-268.
- BUCHALCZYK T, J DYNOWSKI, and S SZTEYN. 1981. Variations in number of teeth and asymmetry of the skull in the wolf. *Acta Theriologica* 26:23-30.
- CHAPLIN RE and J ATKINSON. 1968. The occurrence of upper canine teeth in Roe deer (*Capreolus capreolus*) from England and Scotland. *Journal of Zoology*, London 155:141-144.
- DEARDEN LC. 1954. Extra premolars in the river otter. *Journal of Mammalogy* 35:125-126.
- DOMNING DP and L-AC HAYEK. 1984. Horizontal tooth replacement in the Amazonian manatee (*Trichechus inunguis*). *Mammalia* 48:105-127.
- DREHMER CJ and J FERIGOLO. 1996. Anomalias e patologias dentárias em *Arctocephalus* G. Saint-Hilaire & Cuvier (Pinnipedia, Otariidae) da costa do Rio Grande do Sul, Brasil. *Revista Brasileira de Zoologia* 13:857-865.

- DREHMER CJ, ME FABIÁN, and JO MENEGHETI. 2004. Dental anomalies in the Atlantic population of South American sea lion, *Otaria bryonia* (Pinnipedia, Otariidae): Evolutionary implications and ecological approach. *Latin American Journal of Aquatic Mammals* 3:7-18.
- FELDHAMER GA and TL STOBBER. 1993. Dental anomalies in five species of North America shrews. *Mammalia* 57:115-121.
- FOWLE CD and RC PASSMORE. 1948. A supernumerary incisor in the white-tailed deer. *Journal of Mammalogy* 29:301.
- GLASSTONE S. 1952. The development of halved tooth germs. A study in experimental embryology. *Journal of Anatomy* 86:12-15.
- GOODWIN HT. 1998. Supernumerary teeth in Pleistocene, recent, and hybrid individuals of the *Spermophilus richardsonii* complex (Sciuridae). *Journal of Mammalogy* 79:1161-1169.
- GRAIPEL ME, JJ CHEREM, and DR TAMES. 1997. Anomalia dentária em *Puma concolor*. *Estudos de Biologia (Curitiba)* 42:21-24.
- HARRIS AH and ED FLEHARTY. 1962. Extra tooth in the long-tailed vole. *Journal of Mammalogy* 43:267-268.
- HENRIQUES RPB, MXA BEZERRIL, and T KOHLSDORF. 1997. Abundância, riqueza e seleção de habitat de pequenos mamíferos dos Cerrados do Brasil Central. Pp. 127-130, in: *Contribuição ao conhecimento ecológico do Cerrado* (LL Leite and CH Saito, eds.). Universidade de Brasília, Brasília.
- HERÁN I. 1970. Anomalies in the position of lower teeth in the fisher, *Martes pennanti* Erxl. (Mammalia; Mustelidae). *Canadian Journal of Zoology* 48:1465.
- HINTON MAC. 1923. The dental formula of the Muridae, with special reference to the "mp. 4 Theory." *Annals and Magazine of Natural History, Series 9*, 11:162-170.
- HOOPEL ET. 1946. Extra teeth in a shrew. *Journal of Mammalogy* 27:394.
- IBGE - Instituto Brasileiro de Geografia e Estatística. Cidades. [http:// www.cidades.ibge.gov.br/](http://www.cidades.ibge.gov.br/). Accessed in November 2013.
- JOHNSON DH. 1952. The occurrence and significance of extra molar teeth in rodents. *Journal of Mammalogy* 33:70-72.
- KNOWLTON FF and WC GLAZENER. 1965. Incidence of maxillary teeth in white-tailed deer from San Patricio County, Texas. *Journal of Mammalogy* 46:352.
- KRUTZSCH PH. 1953. Supernumerary molars in the jumping mouse (*Zapus princeps*). *Journal of Mammalogy* 34:265-266.
- KVAM T. 1985. Supernumerary teeth in the European lynx, *Lynx lynx lynx*, and their evolutionary significance. *Journal of Zoology, London A* 206:17-22.
- LIBARDI GS. 2013. Variação não-geográfica em *Necomys lasiurus* (Lund, 1840) (Cricetidae: Sigmodontinae) no Brasil. MSc Dissertation, Escola Superior de Agricultura "Luiz de Queiroz" and Centro de Energia Nuclear na Agricultura, Universidade de São Paulo, Piracicaba, São Paulo, Brazil.
- MACÊDO RH and MA MARES. 1987. Geographic variation in the South American cricetine rodent *Bolomys lasiurus*. *Journal of Mammalogy* 68:578-594.
- MARES MA, KA ERNEST, and DD GETTINGER. 1986. Small mammal community structure and composition in the Cerrado Province of central Brazil. *Journal of Tropical Ecology* 2:289-300.
- MCAFEE RK and VL NAPLES. 2012. Notice on the occurrence of supernumerary teeth in the two-toed sloths *Choloepus didactylus* and *C. hoffmanni*. *Mastozoología Neotropical* 19:163-178.
- MEIN P. 1986. Quelques dents fossiles de morphologie aberrante. *Memmoires du Museum National d'Histoire Naturelle, Serie C* 53:277-284.
- MORAES DA, B LEMOS, and R CERQUEIRA. 2001. Supernumerary molars in Neotropical opossums (Didelphimorphia, Didelphidae). *Mammalian Biology* 66:193-203.
- MUSSER GG and MD CARLETON. 2005. Family Muridae. Pp. 501-755, in: *Mammal species of the world: A taxonomic and geographic reference* (DE Wilson and DA Reeder, eds.); 3<sup>rd</sup> Edition, Smithsonian Institution Press, Washington.
- NATSUME A, K KOYASU, H HANAMURA, H NAKAGAKI, and S ODA. 2005. Variations in the number of teeth in wild Japanese serow (*Naemorhedus crispus*). *Archives of Oral Biology* 50:849-860.
- PARADISO JL. 1966. Notes on supernumerary and missing teeth in the coyote. *Mammalia* 30:120-128.
- PARDIÑAS UFJ, P TETA, PE ORTIZ, P JAYAT and J SALAZAR-BRAVO. In press. Genus *Necomys*. Pp.739-785, in: *Mammals of South America, Volume 2: Rodents* (JL Patton, UFJ Pardiñas and G D'Elia); The University of Chicago Press, Chicago.
- PERCEQUILLO AR. 1998. Sistemática de *Oryzomys* Baird, 1858 do leste do Brasil (Muroidea, Sigmodontinae). MSc Dissertation, Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil.
- PERCEQUILLO AR, E HINGST-ZAHER, and CR BONVICINO. 2008. Systematic review of genus *Cerradomys* Weksler, Percequillo and Voss, 2006 (Rodentia: Cricetidae: Sigmodontinae: Oryzomyini), with description of two new species from Eastern Brazil. *American Museum Novitates* 3622:1-46.
- PEKELHARING CJ. 1968. Molar duplication in red deer and wapiti. *Journal of Mammalogy* 49:524-526.
- PETERKOVA R, H LESOT, and M PETERKA. 2006. Phylogenetic memory of developing mammalian dentition. *Journal of Experimental Zoology* 306B:234-250.
- PETERS FB, EL COELHO, BCE VASCONCELOS, PRO ROTH, and AU CHRISTOFF. 2013. Dental anomalies in *Lontra longicaudis* (Carnivora: Mustelidae) collected in Southern Brazil. *IUCN Otter Specialist Group Bulletin* 30:31-36.
- RAMÍREZ-PULIDO R and C MÜDESPACHER. 1987. Fórmulas dentarias anormales en algunos murciélagos mexicanos. *Acta Zoológica Mexicana, nueva serie* 23:1-54.
- REIG OA. 1977. A proposed unified nomenclature for the enamelled components of the molar teeth of the Cricetidae (Rodentia). *Journal of Zoology (London)* 181:227-241.
- RODRIGUES HG, P MARANGONI, R SUMBERA, P TAFFOREAU, W WENDELEN, and L VIRIOT. 2011.

- Continuous dental replacement in a hyper-chisel tooth digging rodent. Proceedings of the National Academy of Sciences 108:17355-17359.
- RUI AM and CJ DREHMER. 2004. Anomalias e variações na fórmula dentária em morcegos do gênero *Artibeus* Leach (Chiroptera, Phyllostomidae). Revista Brasileira de Zoologia 21:639-648.
- RUPRECHT A. 1965. Supernumerary premolar in *Mustela putorius* Linnaeus, 1758. Acta Theriologica 10:242.
- SHEPPE W. 1964. Supernumerary teeth in deer mouse, *Peromyscus*. Zeitschrift für Säugetierkunde 29:33-36.
- SCHWARTZ JH. 1984. Supernumerary teeth in anthropoid Primates and models of tooth development. Archives of Oral Biology 29:833-842.
- SMITH JD, HH GENOWAYS, and JK JONES JR. 1977. Cranial and dental anomalies in three species of platyrrhine monkeys from Nicaragua. Folia Primatologica 28:1-42.
- SOFAER JA and JH SHAW. 1971. The genetics and development of fused and supernumerary molars in rice rat. Journal of Embryology and Experimental Morphology 26:99-109.
- STEELE DG and WD PARAMA. 1979. Supernumerary teeth in moose and variations in tooth number in North American Cervidae. Journal of Mammalogy 60:852-854.
- STEWART REA and BE STEWART. 1987. Mandibular dental anomalies of Northwest Atlantic harp seals, *Phoca groenlandica*. Canadian Journal of Zoology 65:769-771.
- TAKAHASHI F. 1974. Ocorrência de polidontia em *Didelphis* L. 1758 e sua possível correlação a uma forma ancestral. Anais da Academia Brasileira de Ciências 46:283-285.
- VAN GELDER RG and CA MCLAUGHLIN. 1961. An unusual incisor in *Mephitis mephitis*. Journal of Mammalogy 42:422-423.
- VAN VALEN L. 1964. Nature of the supernumerary molars of *Otocyon*. Journal of Mammalogy 45:284-286.
- VOSS RS. 1991. An introduction to the Neotropical muroid rodent genus *Zygodontomys*. Bulletin of the American Museum of Natural History 210:1-113.
- WEKSLER M and CR BONVICINO. 2005. Taxonomy of pigmy rice rats genus *Oligoryzomys* Bangs, 1900 (Rodentia, Sigmodontinae) of the Brazilian Cerrado, with the description of two new species. Arquivos do Museu Nacional 63:113-130.
- WINER JN, SM LIONG, and FJM VERSTRAETE. 2012. The dental pathology of Southern sea otters (*Enhydra lutris nereis*). Journal of Comparative Pathology 149:346-355.
- WOLSAN M. 1983. Ancestral characters in the dentition of the weasel *Mustela nivalis* L. (Carnivora, Mustelidae). Annales Zoologici Fennici 20:47-51.
- WOLSAN M. 1984. The origin of extra teeth in mammals. Acta Theriologica 29:128-133.
- ZUROWSKI W. 1970. Additional molar in European red deer. Acta Theriologica 15:523-525.

## APPENDIX 1

Specimens examined of genus *Necomys* employed in quantitative and qualitative comparisons in this study; the symbol \* indicates that the specimen was used in the boxplot comparisons.

Caçapava (23°06'S; 45°43'W; Weksler and Bonvicino, 2005): MN 24193, 24194; also includes Gleba José Nani and José Nani, Bairro Sta. Luzia MN 44033\*, 44035\*.

Fazenda Mombaça, Pindamonhangaba (22°55'S; 45°27'W in Pindamonhangaba - IBGE): MN 44036.

Faz. Sto. Ângelo, Represa de Americana, Sumaré (22°49'S; 47°15'W in Sumaré - IBGE): MN 24187.

Guararema (23°25'S; 46°02'W; Weksler and Bonvicino, 2005): ZUEC 1402, 1410, 1414.

Ribeirão da Lagoa, Boracéia, Salesópolis (23°38'S; 45°52'W; Percequillo, 1998): MZUSP 10746, 10757, 10765\*, 10766, 10770\*, 10771\*, 10815\*, 10824, 10828\*, 10831, 10832, 10834\*, 10835\*, 10836, 10837, 10840, 10842, 10844\*, 10846, 10848, 10851, 10855, 10857\*, 10861, 10862\*, 10863\*, 10865, 10867\*, 10868\*, 10877, 10879\*, 10886, 10892\*, 10893, 10894, 10897, 10899\*, 10905, 10918\*, 10919, 10920, 10927, 10945, 21196.

Rio Juruena, Sapezal (13°27'24" S, 59°00'07" W): UFMT 832\*, 839\*, 849\*, 856\*, 4012\*.

São Luís do Paraitinga (23°13'S; 45°18'W in São Luís do Paraitinga - IBGE): ZUEC 2421, 2422, 2423, 2424, 2425.

Taubaté (23°01'S; 45°32'W in Taubaté - IBGE): ZUEC 1405, 1409, 1413; also includes Bairro do Barranco: MZUSP 11211, 11227; Faz. Antonio Taino (Foco 30): MN 24185; Gleba Paulo Japonês, Bairro Barranco: MN 44029, 44030, 44031, 44034, 44037, 44039, 44040, 44043, 44044, 44050, 60426.