

## Vibrational spectra and normal coordinate calculations of para chloro benzoic acid

S Mohan\* and D Arul Dhas

Raman School of Physics, Pondicherry University, Pondicherry-605 014, India

Received 19 April 1993 accepted 12 July 1993

**Abstract** : The FTIR and laser Raman spectra of para chloro benzoic acid have been recorded in the regions 200-4000  $\text{cm}^{-1}$  and 30-4000  $\text{cm}^{-1}$  using Shimadzu FTIR 8101 spectrophotometer and carry model 82 grating spectrophotometer respectively. The vibrational analysis has been carried out by assuming  $C_{2v}$  symmetry. The observed frequencies were assigned to various modes of vibrations on the basis of intensity, frequencies from allied molecules and the normal coordinate calculations. The potential energy distribution associated with normal modes are also reported here. The assignment of fundamental vibrational frequencies for para chloro benzoic acid agree well with the calculated frequencies.

**Keywords** : Vibrational spectra, normal coordinate calculation, para chloro benzoic acid

**PACS Nos.** : 33 10 Gx, 33 20 Ba, 33 20 Fb, 78 30 Cp

### 1. Introduction

The present study is a part of our on going discussions on the vibrational spectra of the derivatives of benzoic acid. The derivatives of benzoic acid are widely used in several applications such as miticides, contrast media in urology, cholecystographic examinations and in the manufacture of pharmaceuticals. It also finds application in dyes, in curing tobacco, in preserving fruits juice, in many esters, as a mordant in cloth printing and as a reference standard in volumetric analysis. Vibrational spectra of benzoic acid and substituted benzoic acids have been studied by various workers [1-3]. Rastogi *et al* [1] have reported about the vibrational study of 3,5-dinitrobenzoic acid. However, there is no report on the vibrational spectra of para chloro benzoic acid in the literature. Hence, an attempt has been made in the present work to record the FTIR and laser Raman spectra of para chloro benzoic acid and to study the complete vibrational analysis on the basis of  $C_{2v}$  point group.

### 2. Experimental details

Pure chemical, para chloro benzoic acid, white solid powder was obtained in the solid form, from SISCO-CHEM Industries, Bombay. The FTIR spectrum was recorded in the region

\* To whom all correspondence should be made

200–4000  $\text{cm}^{-1}$  on Shimadzu FTIR 8101 spectrophotometer in KBr, the IR spectrum was recorded in the region 600–4000  $\text{cm}^{-1}$  on Perkin-Elmer IR 983 and the laser Raman spectrum was recorded in the region 30–4000  $\text{cm}^{-1}$  on Carry Model 82 grating spectrophotometer the 488 nm radiation from an argon ion laser operating at 4 W was used for excitation. The spectral width was 2  $\text{cm}^{-1}$  and the scanning speed was 30  $\text{cm}^{-1} \text{ min}^{-1}$ . The frequencies for all sharp bands are accurate to  $\pm 1 \text{ cm}^{-1}$ .

### 3. Normal coordinate treatment

The calculation of normal vibration was made by the method of Wilson [4]. The potential energy was expressed by a simplified general valence force field (SGVFF) for the following reasons: (i) SGVFF has been shown to be very effective in normal co-ordinate analysis of benzene derivatives. (ii) Valence force constants can be transferred between the related molecules which is very useful in normal co-ordinate analysis [5].

The structural parameters employed in this work were taken from Mohan and Ravikumar [6]. The normal co-ordinate calculations were performed using the modified schachtschneider programme [7]. The initial set of the valence force constants and the corresponding off-diagonal constants were transferred from related benzene derivatives reported recently [5,8]. A zero order calculation with the transferred force constants was performed and the result showed a reasonable agreement between the calculated and observed frequencies for most of the normal modes, except for the low frequency skeletal modes. Hence, the initial set of force constants were refined by least square technique in accordance with the position of bands in the infrared and Raman spectra. The final set of the refined force constants are shown in Table 1.

**Table 1.** Final set of force constants of para chloro benzoic acid (in units of  $\text{m dyne A}^{-1} \text{ Rad}^{-1}$  and  $\text{m dyne A}^{-1} \text{ Rad}^2$ ).

Types of constants	Parameter	Coordinates involved	Final	
Diagonal constants	Stretching	$f_d$	HC	4.5
		$f_D$	CC	3.8
		$f_l$	CC	2.8
		$f_q$	CC	3.9
		$f_n$	CC	4.6
		$f_s$	CC	3.6
		$f_r$	CC	4.1
	Bending	$f_p$	CC	3.6
		$f_m$	CCI	3.7
		$f_\alpha$	H $\hat{C}C$	0.28
		$f_\gamma$	C $\hat{C}C$	0.8
		$f_\phi$	C $\hat{C}C$	1.0
		$f_\theta$	C $\hat{C}C$	1.0

Table 1. (Contd.)

Types of constants		Parameter	Coordinates involved	Final
Interaction constants	Stretch-stretch	$f_{\delta}$	C $\hat{C}$ Cl	0.51
		$f_{\beta}$	C $\hat{C}$ C	1.0
		$f_{dD}$	HC CC	0.19
	Stretch-bend	$f_{Dq}$	CC CC	0.06
		$f_{qn}$	CC CC	0.08
		$f_{sm}$	CC CCl	0.28
		$f_{d\alpha}$	HC H $\hat{C}$ C	0.25
		$f_{D\gamma}$	CC C $\hat{C}$ C	0.36
		$f_{q\gamma}$	CC C $\hat{C}$ C	0.34
		$f_{m\delta}$	CCl C $\hat{C}$ Cl	0.11
		$f_{m\phi}$	CC C $\hat{C}$ C	0.41
		$f_{p\theta}$	CC C $\hat{C}$ C	0.41
	Bend-bend	$f_{d\theta}$	CH C $\hat{C}$ C	0.33
		$f_{\gamma\phi}$	C $\hat{C}$ C C $\hat{C}$ C	0.14
		$f_{\alpha\gamma}$	H $\hat{C}$ C C $\hat{C}$ C	0.12
		$f_{\beta\delta}$	C $\hat{C}$ C C $\hat{C}$ Cl	0.28
		$f_{\gamma\theta}$	C $\hat{C}$ C C $\hat{C}$ C	0.13

#### 4. Results and discussion

The observed frequencies with their relative intensities and assignments of para chloro benzoic acid are given in the Table 2. This compound belongs to  $C_s$  point group. In this point

Table 2. Observed and calculated frequencies ( $\text{cm}^{-1}$ ) and potential energy distribution for para chloro benzoic acid

Species	Observed frequencies and intensity		Calculated frequency	Assignments/ <i>P.E.D.</i>
	IR	FTIR LR		
	3940 W			3090+850
	3879 W			3060+820
	3822 W	3835 W		2 $\times$ 1597+640
	3620 M			2938+680
	3490 M			2 $\times$ 1270+955
	3475 M			O-H stretching*
	3105 M		3101	C-H stretching 92
	3090 M	3090 W	3086	C-H stretching 94
	3060 M	3050 W	3058	C-H stretching 95
	2938 M		2928	C-H stretching-88
		2700 W		2 $\times$ 1186+320
	1748 W			2 $\times$ 1200-642
	1698 M			C=O stretching*
	1684 M	1680 S	1675	$\nu_{\text{C}}$ -C stretching 91

Table 2. (Contd.)

Species	Observed frequencies and intensity			Calculated frequency	Assignments/P.E.D.
	IR	FTIR	LR		
	1652 M				3105 - 2x 725
<i>a'</i>			1628 M	1619	C=C stretching 87
<i>a'</i>		1590 S	1597 S	1586	C=C stretching 94
<i>a'</i>	1575 M	1573 W			1095 + 472
		1492 W			1175 + 325
	1435 W	1429 M			O-H in-plane bending*
<i>a'</i>			1358 W	1351	C-C stretching 82
		1295 W	1299 W		C-O stretching*
<i>a</i>	1270 M	1282 W		1272	C-C stretching 79
<i>a</i>	1200 M			1186	C-C stretching 81
<i>a'</i>		1175 M	1186 M	1169	C-COOH stretching 56
<i>a'</i>		1130 M	1143 M	1128	C-H in-plane bending 48
<i>a'</i>		1095 VS	1105 M	1087	C-H in-plane bending 55
<i>a'</i>	1041 W			1030	C-H in-plane bending 61
<i>a'</i>	1013 W	1013 S		1005	C-H in-plane bending 65
<i>a'</i>	989 M			974	C-H out-of-plane bending 58
<i>a'</i>	955 M			945	C-H out-of-plane bending 51
<i>a'</i>	850 M	855 S		842	C-H out-of-plane bending 42
<i>a'</i>	820 M		815 S	819	C-H out-of-plane bending 49
	810 M	810 W	800 VW		O-H out-of-plane bending*
		762 VS	750 M		O C O in-plane bending*
	732 M				C=O out-of-plane bending*
<i>a'</i>	725 M	725 W		714	C-Cl stretching 72
<i>a''</i>	680 VS	685 S		681	C-C-C out-of-plane bending 65
<i>a</i>	642 W	630 W	640 S	626	C-C-C in-plane bending 70
	628 VS				C-C-O rocking*
<i>a'</i>		549 S		536	C-C-C out-of-plane bending 49
		520 M	520 VW		855 332
<i>a</i>		472 S	480 VW	471	C-C-C in-plane bending 52
		352 M	353 M		C-OH in-plane bending*
<i>a'</i>		342 W		328	C-Cl in-plane bending 39
<i>a'</i>		332 W		317	C-COOH in-plane bending 50
<i>a</i>		325 W	320 W	309	C-C-C in-plane bending 54
<i>a''</i>		307 W		301	C-Cl out-of-plane bending 40
		295 M	290 M		1105-815
<i>a''</i>		272 S		265	C-C-C out-of-plane bending 58
		235 M	228 W		C-OH out-of-plane bending*
<i>a''</i>		215 S		209	C-COOH out-of-plane bending 39
			108 S		Lattice vibration
			67 VS		Lattice vibration

VS - Very strong, VW - Very weak, S - Strong,  
MS - Medium strong, M - Medium, W - Weak, \* - Group vibration

group, there are only two types of vibrations, i.e., planar (*a'*) and non-planar (*a''*). They can be distributed as

$$\Gamma = 21 a' \text{ (planar)} + 9 a'' \text{ (non-planar)}$$




Figure 1. Infrared spectrum of para chloro benzoic acid.




Figure 2. FTIR spectrum of para chloro benzoic acid.

in which  $a'$  gives rise to polarised lines whereas  $a''$  gives depolarised lines in the Raman spectrum. All 30 vibrations are active in the IR and Raman spectra. Structure of para chloro benzoic acid and the spectra recorded are shown in the Figures 1-4.

### Phenyl ring vibrations

#### 4.1. Carbon vibrations :

The C = C vibrations are more interesting if the double bond is in conjugation with the ring. The actual positions are determined not so much by the nature of substituents but by the form

of substituents around the ring Bellamy 1957 [9]. Benzene has two doubly degenerate modes  $e_{2g}$  ( $1596\text{ cm}^{-1}$ ) and  $e_{12g}$  ( $1485\text{ cm}^{-1}$ ) and two non-degenerate modes  $b_{2u}$  ( $1310\text{ cm}^{-1}$ ) and  $a_{12g}$  ( $995\text{ cm}^{-1}$ ) due to skeletal stretching modes of C-C bonds.




Figure 3. Laser Raman spectrum of para chloro benzoic acid




Figure 4. Structure of para chloro benzoic acid

The modes corresponding to C-C stretching in benzene are assigned to the bands at  $1680, 1628, 1590, 1358, 1282$  and  $1200\text{ cm}^{-1}$  in para chloro benzoic acid.

The in-plane carbon bending vibrations, are obtained from the non-degenerate  $b_{1u}$  ( $1010\text{ cm}^{-1}$ ) and degenerate  $e_{2g}$  ( $606\text{ cm}^{-1}$ ) modes of benzene. The  $e_{2g}$  ( $606\text{ cm}^{-1}$ ) degenerate frequency splits into two totally symmetric vibrations under  $C_s$  symmetry and they are observed at  $630$  and  $472\text{ cm}^{-1}$  in para chloro benzoic acid.

The carbon out-of-plane bending vibrations are defined by reference to the non-degenerate  $b_{2g}$  ( $703\text{ cm}^{-1}$ ) and degenerate  $e_{2u}$  ( $404\text{ cm}^{-1}$ ) modes of benzene. The former is found to be constant in substituted benzenes and in para chloro benzoic acid it is observed at  $685\text{ cm}^{-1}$ . The degenerate  $e_{2u}$  ( $404\text{ cm}^{-1}$ ) vibration splits into two non-totally symmetric components and the bands are observed at  $549$  and  $272\text{ cm}^{-1}$ .

#### 4.2. C-H vibrations :

The aromatic structure shows the presence of C-H stretching vibrations in the region  $3000\text{--}3100\text{ cm}^{-1}$ , C-H in-plane bending in the region  $1000\text{--}1100\text{ cm}^{-1}$  and C-H out-of-plane bending in the region  $800\text{--}980\text{ cm}^{-1}$ , which permits ready identification for this structure. In this region the bands are not appreciably affected by the nature of the substituents. The frequencies  $3105$ ,  $3090$ ,  $3060$  and  $2938\text{ cm}^{-1}$  are assigned for C-H stretching modes in para chloro benzoic acid and these assignments are in good agreement with August *et al* 1936 [10], Bailey *et al* 1946 [11]. The frequencies  $1130$ ,  $1095$ ,  $1041$  and  $1013\text{ cm}^{-1}$  are assigned for C-H in-plane bending and are in favourable agreement with values given in the literatures [12,13]. The frequencies  $989$ ,  $955$ ,  $850$  and  $820\text{ cm}^{-1}$  are assigned to C-H out-of-plane bending and these assignments are in good agreement with the literature value [14].

#### 4.3. C-Cl vibration :

The C-Cl stretching frequency is generally observed in the region  $671\text{--}722\text{ cm}^{-1}$ . Hence the strong band at  $725\text{ cm}^{-1}$  is assigned, to C-Cl stretching frequency. The C-Cl in-plane bending vibration and out-of-plane vibrations are generally observed at  $341\text{ cm}^{-1}$  and  $306\text{ cm}^{-1}$ . In para chloro benzoic acid the C-Cl in-plane bending vibrations and out-of-plane bending vibrations occur at  $342\text{ cm}^{-1}$  and  $307\text{ cm}^{-1}$  respectively. The present assignments agree well with the values given in the literatures [15-18].

#### 4.4. C-COOH vibration :

Usually the band around  $1185$ ,  $330$  and  $209\text{ cm}^{-1}$  are assigned to C-COOH stretch, C-COOH in-plane bending and to C-COOH out-of-plane bending respectively. In para chloro benzoic acid the assignments are made at  $1186\text{ cm}^{-1}$  for C-COOH stretching,  $332\text{ cm}^{-1}$  for C-COOH in-plane bending and  $215\text{ cm}^{-1}$  for C-COOH out-of-plane bending respectively.

#### Group vibration

Group vibrations are determined in terms of the motions that the nuclei in a structural group in the molecule undergo during the vibrations and they appear in fairly constant regions in the

#### 4.5. COOH group vibration :

The band observed at  $3475\text{ cm}^{-1}$  has its origin in the O–H stretching vibration. The C=O stretching is a characteristic frequency of carboxylic group. The dipole moment derivatives of the COO in-plane bending mode and C=O stretching mode are very large, splitting with marked frequency differences may be expected for these modes and the out-of-phase C=O stretching mode should have a higher frequency than in-phase mode [19]. The band appearing at  $1698\text{ cm}^{-1}$  is assigned to C=O stretching vibration. No splitting is observed. OH in-plane bending and C=O stretching vibrations are closely coupled. These are appearing at  $1429\text{ cm}^{-1}$  and  $1295\text{ cm}^{-1}$  in the present investigation. The bands at  $762\text{ cm}^{-1}$  and  $628\text{ cm}^{-1}$  are assigned to OCO bending and CCO rocking vibrations. Out-of-plane bending vibration due to O–H give rise to band at  $810\text{ cm}^{-1}$ . The out-of-plane bending vibration due to C=O is given as  $732\text{ cm}^{-1}$ . The OH out-of-plane bending vibration is strongly coupled with C=O out-of-plane bending vibration. The C=O out-of-plane wagging mode of carboxyl group in all likelihood involves motion of both oxygen atoms. The C–OH out-of-plane bending are assigned to frequencies  $352$  and  $235\text{ cm}^{-1}$  respectively. These observation are in close agreement with Ahmad *et al* 1992 [20].

#### 5. Potential energy distribution

To check whether the chosen set of vibrational frequencies contribute maximum to the potential energy distribution, *P.E.D.* has been calculated using the relation,

$$P.E.D. = \frac{F_{ii} L_{ik}^2}{\lambda_k}$$

In the molecule under investigation, the normal modes under symmetry species with comparable magnitudes couple strongly and hence the observed frequencies are the results of mixed modes which is reflected in the potential energy distribution. The highest *P.E.D.* contribution corresponding to each of the observed frequencies are alone listed in the present work. The close agreement between the observed and calculated frequencies confirms the validity of the present assignment.

#### 6. Conclusions

In the present study, a complete vibrational assignment and the analysis of para chloro benzoic acid are available using the frequencies obtained from FTIR and laser Raman spectra on the basis of  $C_s$  symmetry.

#### References

- [1] V K Rastogi, Rajpoot and SN Sharma 1984 *Indian J Phys.* **58B** 311 –
- [2] Włodzimierz, Levendowski and Halina Branska 1986 *J Raman Spectrosc* **17** 17
- [3] M Pagannone, B Fornari and G Mattei 1987 *Spectrochim Acta* **43A** 1
- [4] E B Wilson (Jr.) 1941 *J Chem Phys.* **9** 76
- [5] Danuła Michalska 1993 *Spectrochim. Acta* **49A** 303



- [6] S Mohan and K G Ravikumar 1984 *Acta Phys. Polon.* A **65** 557
- [7] Schachtachneider 1986 *Vibrational Analysis of Polyatomic molecules*, Technical Report nos. 231-64 and 57-65 (Shell Development Co., Emeryville, California)
- [8] S Mohan and R Murugan 1992 *Indian J. Pure Appl. Phys.* **30** 283
- [9] L J Bellamy 1959 *The Infrared Spectra of Complex Molecules* (New York : John Wiley) p 70
- [10] W R Augus, C K Ingold and A H Leckie 1936 *J. Chem. Soc* 925
- [11] C R Bailey, R R Gordan and J B Hale 1946 *J. Chem. Soc.* 299
- [12] H G Silver and J L Wood 1964 *Trans. Farad. Soc.* **60** 5
- [13] E F Mooney 1964 *Spectrochim. Acta* **20** 1343
- [14] S Pinchas, D Samuel and M Weiss-Brodsky 1961 *J. Chem. Soc.* 1688
- [15] S Mohan, T J Bhoopathy and K G Ravikumar 1985 *Acta Phys. Polon.* A**67** 1133
- [16] G Gunasekaran, S Mohan and Kalainathan 1988 *Acta Ciencia Indica* XI V(P) 77
- [17] Vasuki and S Mohan 1990 *Indian J. Pure Appl. Phys.* **28** 200
- [18] Hussein, Abdel-Shaty and K Howard 1977 *J. Mol. Struct.* **42** 37
- [19] Yesook Kim and Katsunosuke Machida 1986 *Spectrochim. Acta* **42A** 881
- [20] Shabbir Ahmad, Samuel Mathew and P K Verma 1982 *Indian J. Pure Appl. Phys.* **30** 764