

Article (refereed) - postprint

Dawson, Alistair. 2018. **Both low temperature and shorter duration of food availability delay testicular regression and affect the daily cycle in body temperature in a songbird.** *Physiological and Biochemical Zoology*, 91 (4). 917-924. <https://doi.org/10.1086/698109>

© 2018 by the University of Chicago

This version available <http://nora.nerc.ac.uk/id/eprint/520376/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. Some differences between this and the publisher's version remain. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <https://www.journals.uchicago.edu/doi/10.1086/698109>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1 Low temperature and a shorter duration of food availability both delay testicular
2 regression and affect the daily cycle in body temperature in a songbird

3

4

5

6

7

8 Running title: Temperature, food and testicular regression

9

10 What is already known:

11 Although photoperiod is the major environmental cue used by birds to time the annual
12 gonadal cycle, food and ambient temperature can modulate photoperiodic through unknown
13 pathways.

14

15 What this study adds:

16 The time of food availability and ambient temperature both modulate photoperiodic
17 responses so that long-photoperiod induced gonadal regression is delayed. Both also affect
18 the daily cycle in body temperature to give it characteristics of a shorter photoperiod,
19 suggesting that the daily cycle in body temperature may play a role in photoperiodic
20 responses.

21 Keywords: starling; testis; annual cycle; body temperature; energy; food; photoperiod;
22 temperature; thyroid

23 Abstract

24 Photoperiodic control of reproduction in birds is based on two processes, a positive effect
25 leading to gonadal maturation and an inhibitory effect subsequently inducing regression.
26 Non-photoperiodic cues can modulate photoperiodic control, particularly the inhibitory
27 process. In previous studies on common starlings *Sturnus vulgaris* (1) restriction of food
28 availability to 8h after dawn had little effect on testicular maturation but dramatically delayed
29 subsequent regression and (2) lower ambient temperature also had little effect during
30 maturation but delayed regression. Could the effects of food restriction and temperature share
31 a common underlying mechanism? Four groups of starlings were kept on a simulated natural
32 cycle in photoperiod in a 2x2 factorial experimental design. Two groups were held under an
33 ambient temperature of 16 °C and the other two at 6 °C. One of each of these groups had
34 food provided *ad lib* and in the other two groups access to food was denied 7h after dawn. In
35 both *ad lib* food groups and food-restricted groups, lower temperature had little effect on
36 testicular maturation but delayed subsequent regression and molt. In both the 16 °C and 6 °C
37 groups, food restriction had no effect on testicular maturation but delayed regression and
38 molt. The daily cycle in body temperature was recorded in all groups when photoperiod had
39 reached 12L:12D, the photoperiod at which regression is initiated. In both 6 °C groups, night-
40 time body temperature was lower than in the 16 °C groups, a characteristic of shorter
41 photoperiods. In the two *ad lib* food groups, high daytime temperature was maintained until
42 dusk whereas in the two food restricted groups, body temperature began to decrease after
43 food withdrawal. Thus both lower temperature and food restriction delayed regression, as if
44 photoperiod was shorter than it actually was, and both resulted in daily cycles in body
45 temperature that reflected cycles under shorter photoperiods. This implies that the daily cycle
46 in body temperature is possibly a common pathway through which non-photoperiodic cues
47 may operate.

48 **1. Introduction**

49 The predominant environmental cue to time the annual cycle in gonadal maturity in
50 birds is photoperiod. However, since gonadal cycles are rarely if ever symmetrical with
51 photoperiod, at least two photoperiodic mechanisms must be implicated. For a considerable
52 time, these have been referred to as photostimulation and photorefractoriness (Burger, 1947),
53 and these were thought to act sequentially to time gonadal maturation and regression
54 respectively. A more recent hypothesis is that there is a stimulatory process and an inhibitory
55 process acting in tandem at all times to control GnRH secretion and the net difference
56 between these two determines the rate of maturation or regression (Dawson, 2015).
57 Differences between species in terms of the timing and duration of gonadal maturity (e.g.
58 absolute and relative photorefractoriness) can be accounted for entirely by differences in
59 sensitivity of the inhibitory process to photoperiod. The neuroendocrine pathway through
60 which long photoperiods induce stimulation of GnRH secretion is fairly well understood e.g.
61 (Ikegami and Yoshimura, 2016), but this is not true of the inhibitory process.

62 Although photoperiod is the primary environmental cue, other factors can have
63 modifying effects. These are more important in species with less predictable breeding
64 schedules (Wingfield et al., 1992) and may act directly on GnRH or through GnIH
65 (Bedecarrats et al., 2016; Ernst et al., 2016; Tsutsui and Ubuka, 2016). Many studies have
66 investigated the effects of ambient temperature on photoperiodically induced testicular
67 maturation in birds. In a few cases, testicular maturation was accelerated by higher
68 temperature (Perfito et al., 2005; Silverin et al., 2008; Wingfield et al., 2003); in the majority
69 it was not (Caro et al., 2013; Caro and Visser, 2009; Dawson, 2005; Perfito et al., 2005;
70 Silverin et al., 2008; Soumalainen, 1938; Visser et al., 2011; Wingfield et al., 1996;
71 Wingfield et al., 1997). However, in all of the studies where subsequent regression was also
72 monitored, high temperature advanced regression (Dawson, 2005; Dawson and Sharp, 2010;

73 Visser et al., 2011; Wingfield et al., 2003; Wingfield et al., 1997). The photoperiodic
74 inhibitory process leading to gonadal regression is therefore more amenable to modulation by
75 temperature than is photoperiodic stimulation.

76 Food restriction also has effects on the reproductive axis (Davies and Deviche 2014).
77 Hahn (1995) photostimulated Red Crossbills *Loxia curvirostra* but restricted their food
78 consumption to short photoperiod levels. This delayed the increase in LH but did not affect
79 testis size after 30 days. Perfito et al. (2008) found that testis size, but not LH, was affected
80 by similar food restriction in Zebra finches *Taeniopygia guttata*. Food-deprived house finches
81 *Haemorhous mexicanus* had under developed testes which was thought to result from
82 decreased GnRH secretion (Valle et al. 2015). The daily timing of food availability does
83 influence the photoperiodic inhibitory process. I did a study a (Dawson, 1986) involving
84 three groups of common starlings (*Sturnus vulgaris*). One was kept under a short photoperiod
85 8L:16D with food provided *ad lib*. Another was transferred to a long photoperiod (16L:8D)
86 with food *ad lib*. The third group was also transferred to a long photoperiod (16L:8L) but
87 food was removed 8 h after dawn, so a feeding day of the same duration as the short
88 photoperiod group. There was little difference in the rate of testicular maturation between the
89 two long photoperiod groups, demonstrating that food had no influence on photoperiodic
90 gonadal stimulation. However, there was a dramatic delay in regression. For most birds,
91 regression was delayed by about 2 weeks (regression started 6 weeks after transfer to long
92 photoperiods as opposed to 4 weeks in birds with *ad lib* food). Remarkably, two birds had
93 shown no regression by the end of the study (12 weeks after transfer). Thus, as with
94 temperature, the photoperiodic inhibitory process is more amenable to modulation by food
95 availability than is photoperiodic stimulation.

96 Lower ambient temperature and decreased food availability had similar effects on the
97 testicular cycle in starlings; neither influenced testicular maturation, but both delayed

98 regression (Dawson, 1986, 2005). It is perhaps counterintuitive that a decrease in food
99 availability and a decrease in ambient temperature do not have negative effects on the
100 reproductive system; they both extend rather than inhibit it.

101 Starlings have a robust and marked daily cycle in body temperature (T_b) which is
102 closely related to photoperiod (Dawson, 2017). T_b starts to increase before dawn, remains
103 high throughout daylight hours, and then decreases rapidly at dusk. The amplitude of the
104 cycle increases as photoperiod decreases. Night-time hypothermia is thought to be an energy
105 conserving mechanism and ambient temperature affects depth of nocturnal hypothermia
106 (Reinertsen and Haftorn, 1986). Several studies have shown that food deprivation, or energy
107 deficiency, can also increase the depth of nocturnal hypothermia (Ben-Hamo et al., 2010;
108 Cooper and Gessaman, 2005; McKechnie and Lovegrove, 2002; Noakes et al., 2013; Nord et
109 al., 2009; Pravosudov and Lucas, 2000; Prinzinger et al., 1991; Reinertsen and Haftorn, 1986;
110 Reinertsen et al., 1988; Waite, 1991; Welton et al., 2002). Importantly, food removal in
111 Japanese quail *Coturnix coturnix japonica* led to diurnal hypothermia (Laurila et al., 2005).

112 Since food restriction and ambient temperature affect both reproduction by delaying
113 gonadal regression, and both can also affect the daily cycle in T_b , could the effects of food
114 and temperature act through a common mechanism, the daily cycle in T_b , and that this
115 influences the timing of gonadal regression? The aim of this study was to confirm that low
116 temperature and food restriction have similar effects on gonadal cycles i.e. little effect on
117 maturation but delaying regression, and to determine whether either or both affect the daily
118 cycle in T_b . The results show that food restriction and ambient temperature both changed the
119 characteristics of the daily cycle in T_b to that of a shorter photoperiod. A shorter photoperiod
120 would also delay gonadal regression. Thus food and ambient temperature may act through a
121 common pathway to influence the plastic photoperiodic inhibitory process, but not the more
122 resilient photoperiodic stimulatory process.

123

124 **2. Materials and methods**

125 *2.1 Birds*

126 Juvenile starlings were caught at Berwick upon Tweed, UK, 55.8 °N during August
127 2014. Only males were kept; females were released. Birds were kept in four environmentally
128 controlled chambers (2.5 m x 2.0 m and 2.2 m high) in which they were allowed to fly freely.
129 They were provided with turkey starter crumbs *ad lib*. All birds were treated with a single
130 dose of anti-helminthic (Flubenvet) in food to eliminate gut parasites and 1% Ivermectin on
131 the skin to remove ectoparasites.

132 Temperature was initially kept at 14 °C with 5 changes of air per hour. The time that
133 lights came on and went off each day was controlled by a Lutron GRAFIK Eye QS system.
134 Light intensity increased from zero to 1000 lux during a 1 min period at dawn and decreased
135 during 1 min at dusk. Photoperiod for all groups was changed daily and simulated the natural
136 changes at 56 °N (minimum at the winter solstice 7L:17D; maximum at summer solstice
137 17L:7D).

138 Food was provided in commercial poultry feeders. Each feeder was placed in a solid
139 enclosure (1m x 0.8m x 0.8m) such that access to the food was through two openings at floor
140 level. In two of the chambers the opening could be closed with a metal plate that slid down at
141 pre-determined times (Chicken Guard, used in free-range chicken coups to prevent nocturnal
142 access by predators).

143

144 *2.2 Experimental design*

145 In early December, birds were laparotomized under isoflurane anaesthesia and the
146 dimensions of the left testis measured to the nearest 0.1 mm using a binocular microscope.

147 Testicular volume was calculated as $\frac{4}{3} \pi a^2 b$ where a is half the width and b is half the
148 length. Birds were laparotomized at approximately three week intervals during the study.

149 At the winter solstice, temperature was decreased to 6 °C in two chambers and in the
150 other two it was increased to 16 °C. In one of the chambers at 6 °C and one at 16 °C, access
151 to food was prevented by the Chicken Guard door closing 7h after dawn (which was at dusk
152 at the winter solstice). In the other two chambers, food was available *ad lib*. Sample numbers
153 in the four groups were: 6 °C *ad lib* food, n = 7; 6 °C restricted food n = 9; 16 °C *ad lib* food,
154 n = 8; 16 °C restricted food, n = 8. Dawn was kept at a fixed time (08:00 GMT), so that the
155 time when access to food was prevented, also occurred at the same time each day (15:00
156 GMT). Photoperiod was increased to simulate the natural changes in photoperiod by
157 extending dusk.

158 The inhibitory photoperiodic process leading to testicular regression (otherwise
159 known as photorefractoriness) starts when photoperiod has increased to 12L: 12D (Dawson,
160 2015). Changes in the daily cycle in T_b which may be involved should therefore be apparent
161 then. The daily cycle in T_b was assessed using DST nano-T temperature data loggers (Star-
162 Oddi Ltd, Iceland) in which recorded data is stored in the logger's internal memory with a
163 real time clock reference for each measurement. The loggers were synchronised with the
164 same computer as the GRAFIK Eye lighting controls. The loggers were 17 mm x 6 mm,
165 weighing 1 g. Each bird was anaesthetised with Isoflurane, a 10 mm incision was made in the
166 skin of the nape of the neck, and a logger inserted sub-dermally. The incision was closed with
167 tissue adhesive (Vetbond). The loggers were implanted 5 days before photoperiod had
168 increased to 12L:12D, and programmed to start recording five days after implantation.
169 Temperature was recorded every 5 min. Birds were again anaesthetised, the loggers removed
170 and data downloaded using the SeaStar program (Star-Oddi Ltd, Iceland).

171 The start of molt is closely associated with the time of testicular regression (Dawson,
172 2006). Since the start of molt can be measured exactly, unlike the gradual decrease in testis
173 size, it provides an accurate quantifiable data point which is related to the timing of
174 regression. Birds began to molt during May. The progress of molt was assessed using a molt
175 scoring system which corrects for primary feather mass (Dawson and Newton, 2004). For
176 each bird, a series of linear regressions was used to calculate molt start date and the date that
177 subsequent increments in molt score occurred.

178 The experimental procedures were licensed by the UK Home Office (Project licence
179 PPL 60/4176).

180

181 *2.3 Statistical analyses*

182 To make the analysis of temperature data more manageable, each block of six 5 min
183 temperature records was averaged to produce a mean for each 30 min period. Data on T_b
184 during 24h and testis volume during the study were then assessed in a repeated measures two-
185 way ANOVA (Graph Pad Prism) followed by Tukey's Multiple Comparison Tests. Molt start
186 dates were analysed using one-factor ANOVA followed by Tukey's Multiple Comparison
187 Test.

188

189 **3. Results**

190 *3.1 Body mass*

191 There were significant differences between the treatment groups: $F_{3, 196} = 3.22$,
192 $P=0.0386$ (Fig. 1). However, there were no consistent differences between any groups. All
193 four groups increased mass significantly between the first two sampling points but there were
194 no subsequent consistent changes. There were two occasions, mid-March and mid-May,

195 when the cold *ad lib* group had body mass significantly higher than the other three groups,
196 but there were no significant differences between any groups at any other times. Therefore,
197 differences in T_b , testis sizes and molt between the 6 °C *ad lib* and restricted food groups, and
198 between the 16 °C *ad lib* and restricted food groups, were not due to under nutrition.

199

200 3.2 Daily cycle in body temperature under 12L:12D

201 There were significant differences between the treatment groups: $F_{3, 1175} = 3.52$,
202 $P=0.0296$. In all groups, T_b increased significantly ($P<0.001$) before dawn and continued to
203 increase after dawn ($P<0.001$). Also, in all four groups, T_b decreased significantly
204 immediately after dusk ($P<0.001$).

205 Comparing the warm and cold *ad lib* groups (Fig. 2 A), T_b was significantly lower
206 ($P<0.01$) in the cold *ad lib* group during the middle of the night but there were no differences
207 at any other time. In the warm restricted group (Fig. 3A) T_b decreased significantly ($P<0.05$)
208 between access to food being denied (13:00) and 15 min before dusk. T_b in this group was
209 significantly lower than in the warm *ad lib* group 15 min before dusk ($P<0.01$) and 15 min
210 after dusk ($P<0.001$). In the cold restricted group (Fig. 4A) T_b also decreased significantly
211 ($P<0.001$) between access to food being denied and 15 min before dusk. T_b in this group was
212 significantly lower than in the cold *ad lib* group 15 min before dusk ($P<0.001$) and 15 min
213 after dusk ($P<0.001$). The decrease in T_b following access to food being denied was greater at
214 the lower ambient temperature. Between 15:00 and dusk, T_b was lower ($P<0.01$) in the cold
215 restricted group than in the warm restricted group (Figs. 3A and 4A).

216 Mean body temperature over 24h did not differ significantly between the groups (Fig.
217 5) although there was a suggestion that mean body temperature was higher in the two groups
218 at 16 °C.

219

220 *3.3 Changes in testis size*

221 There were significant differences between the treatment groups: $F_{3, 196} = 5.94$,
222 $P=0.0029$. Ambient temperature had a slight effect during the period of testicular maturation.
223 Between January and April, testicular volume was slightly, but not significantly greater, in
224 the warm *ad lib* group than in the cold *ad lib* group (Fig. 2B). During early March, testicular
225 volume was greater ($P<0.01$) in the warm restricted group than in the cold restricted group,
226 but not at any other time. In contrast, ambient temperature did have a clear effect on the time
227 of regression in the two *ad lib* groups. Testis volume showed a large decrease ($P<0.001$)
228 between mid and late April in the warm *ad lib* group, but only a slight decrease ($P<0.05$) in
229 the cold *ad lib* group. There was a greater decrease in the cold *ad lib* group between late
230 April and mid-May ($P<0.001$). In both late April and mid-May, testis volume was
231 significantly lower ($P<0.001$) in the warm *ad lib* group than in the cold *ad lib* group. Lower
232 temperature also delayed regression, although less dramatically, in the food-restricted groups.
233 In mid-May, testis size was lower ($P<0.001$) in the cold food-restricted group (Fig. 4B) than
234 in the warm food-restricted group (Fig.3 B).

235 Food restriction had no effect on testicular maturation (Figs. 3B and 4B) but it did
236 have an effect on the timing of regression. Testis volume decreased ($P<0.001$) between mid
237 and late April in the warm *ad lib* group, and between late April and mid-May in the warm
238 restricted group (Fig. 3B). In both late April and mid-May, testis volume was significantly
239 lower ($P<0.001$) in the warm *ad lib* group than in the warm restricted group. Testicular
240 regression began slightly earlier in the cold *ad lib* group than the cold restricted group (Fig.
241 4B). There was a slight decrease in the cold *ad lib* group between mid and late April ($P<0.05$)
242 but no decrease until later in the cold restricted group.

243

244 3.4 Molt

245 There were significant differences between the treatment groups: $F_{3, 25} = 40.28$,
246 $P < 0.0001$. The start of molt closely correlated with the time of testicular regression;
247 treatments affected molt in the same way as regression. Ambient temperature affected the
248 start of molt. Molt started 15 days earlier ($P < 0.001$) in the warm *ad lib* group than in the cold
249 *ad lib* group (Fig. 2C) and molt started 16 days earlier ($P < 0.001$) in the warm restricted group
250 than in the cold restricted group (Figs. 3C and 4C). Food restriction also affected molt. Molt
251 started 7 days earlier ($P < 0.05$) in the warm *ad lib* group than in the warm restricted group
252 (Fig. 3C) and molt started 8 days earlier ($P < 0.01$) in the cold *ad lib* group than in the cold
253 restricted group (Fig. 4C).

254

255 4. Discussion

256 The first aim of this study was to confirm that low temperature and food restriction
257 have similar effects on gonadal cycles i.e. little effect on maturation but delaying regression.
258 There was indeed little effect of temperature, and no effect of food restriction, on testis size
259 during testicular maturation. In contrast, testicular regression and the start of molt were both
260 significantly delayed by both lower ambient temperature and the restriction of food. The
261 lower ambient temperature (10 degrees lower) delayed molt by 15 days in the food *ad lib*
262 groups and by 16 days in the food restricted groups. Food restriction (to 7h after dawn)
263 delayed molt by 7 days in the 16 °C groups and by 8 days in the 6 °C groups. The annual
264 cycle in gonadal maturity is thought to be controlled by two photoperiodic mechanisms
265 acting in tandem throughout the year: a positive drive on GnRH secretion which is
266 predominant during increasing photoperiods and leads to gonadal maturation, and an

267 inhibitory process which becomes predominant during long photoperiods and leads to
268 gonadal regression (Dawson, 2015). The effects of ambient temperature and food restriction
269 appear to modulate photoperiodic control of the latter process rather than the former. There is
270 a potential ecological explanation why lower temperature should lead to prolonged breeding.
271 In cooler years, while food peak food availability may be less, it is likely to be later, and
272 more prolonged.

273 However, it is not possible to be categorical from the data here that the effects were
274 restricted to gonadal regression. Lower temperature caused a slight, although non-significant
275 delay in testicular maturation. Food restriction had no effect on maturation. However, the
276 caveat here is that because of the experimental design, there was little food restriction during
277 the early stages of testicular maturation when photoperiod was still short. Nevertheless,
278 ambient temperature and the time of food availability did both clearly have effects on the
279 photo-induced gonadal cycle.

280 The second aim was to look for a common mechanism through which the two
281 different environmental factors, ambient temperature and food restriction, could operate to
282 impart their similar effects on the gonadal cycle. Obviously both have energetic implications.
283 There is evidence that both, independently, can affect the daily cycle in T_b . In birds with *ad*
284 *lib* food, the daily cycle in T_b is strictly related to photoperiod; high T_b is normally
285 maintained until dusk and this was true of both *ad lib* food groups in this study. In the food
286 restricted birds, T_b began to decrease after access to food was blocked. Therefore the duration
287 of maximal T_b was less than the photoperiod, and the timing of testicular regression was also
288 as if photoperiod was shorter. In the *ad lib* fed birds, lower ambient temperature induced a
289 greater amplitude in the daily cycle of T_b , with lower T_b during darkness. This too is a
290 characteristic of shorter photoperiods (Dawson, 2017). Thus the timing of testicular

291 regression appears, in some way, to be related to the daily cycle in T_b in addition to
292 prevailing photoperiod.

293 It may be controversial to suggest that the daily cycle in T_b modulates photoperiodic
294 responses in birds since there is a wealth of evidence to show that photoperiod is the major
295 cue used to time gonadal maturation and regression e.g. Dawson (2015), and that light acts
296 directly, in the case of birds, through encephalic photoreceptors (Foster and Follett, 1985;
297 Garcia-Fernandez et al., 2015). However, it is well known in plants that temperature can
298 modulate photoperiodic molecular mechanisms to regulate the timing of flowering (Andres
299 and Coupland, 2012; Song, 2016). Furthermore, the present study led to a subsequent study in
300 which starlings were maintained on ultra-short photoperiods, and gonadal responses related to
301 the daily cycle in T_b much more closely than to photoperiod. Nevertheless, this remains a
302 correlation rather than demonstrating a causal relationship.

303 Thyroid hormones may be an important link between energetics, T_b and photoperiodic
304 responses. They regulate metabolic rate (e.g. hypothyroidism is associated with low T_b) and
305 also play a critical role in photoperiodic responses (Dawson, 1993; Yoshimura, 2006).
306 Experimental treatment with exogenous thyroid hormones can mimic long photoperiods
307 (Dawson, 1989; Follett et al., 1988). Wikelski et al (2008) suggested that energy turnover
308 may determine the duration of circannual cycles in house sparrows (*Passer domesticus*); the
309 lower the rate of energy turnover the longer the cycle length. In the present study, higher
310 mean T_b was associated with shorter gonadal cycle. However, the food-restricted birds were
311 not nutritionally stressed – they maintained their body weight. It was apparently the time of
312 food availability rather than total food intake that was important and in house sparrows
313 periodic food availability can act as a *Zeitgeber* for the whole circadian system (Hau and
314 Gwinner, 1996). Future studies could directly address the question of whether the timing
315 rather than the general availability of food is important.

316 In conclusion, reducing the time of food availability and reducing ambient
317 temperature both modulate photoperiodic responses so that long-photoperiod induced
318 gonadal regression is delayed. Both also affect the daily cycle in T_b to give it characteristics
319 of a shorter photoperiod. This suggests the possibility that the daily cycle in T_b may play a
320 role in photoperiodic responses. Although this is a surprising and tentative conclusion,
321 stronger evidence for this was obtained in a subsequent study.

322

323 **Literature cited**

324 Andres, F. and G. Coupland. 2012. The genetic basis of flowering responses to seasonal cues. *Nature*
325 *Rev Gen* 13: 627-639.

326 Bedecarrats, G.Y., M. Baxter, and B. Sparling, B. 2016. An updated model to describe the
327 neuroendocrine control of reproduction in chickens. *Gen Comp Endocrinol* 227: 58-63.

328 Ben-Hamo, M., B. Pinshow, M.D. McCue, S.R. McWilliams and U. Bauchinger. 2010. Fasting
329 triggers hypothermia, and ambient temperature modulates its depth in Japanese quail *Coturnix*
330 *japonica*. *Comp Biochem Physiol A-Mol Integr Physiol* 156: 84-91.

331 Burger, J.W. 1947. On the relation of daylength to the phase of testicular involution and inactivity of
332 the spermatogenic cycle of the starling. *J Exp Zool* 124: 227-239.

333 Caro, S.P., S.V. Schaper, A. Dawson, P.J. Sharp, P. Gienapp and M.E. Visser. 2013. Is
334 microevolution the only emergency exit in a warming world? Temperature influences egg
335 laying but not its underlying mechanisms in great tits. *Gen Comp Endocrinol* 190: 164-169.

336 Caro, S.P. and M.E. Visser. 2009. Temperature-induced elevation of basal metabolic rate does not
337 affect testis growth in great tits. *J Exp Biol* 212: 1994-1998.

338 Cooper, S.J., and J.A. Gessaman. 2005. Nocturnal hypothermia in seasonally acclimatized Mountain
339 Chickadees and Juniper Titmice. *Condor* 107: 151-155.

340 Davies, S., and P. Deviche. 2014. At the crossroads: Food supply and the timing of avian
341 reproduction. *Horm Behav* 66: 41-55.

- 342 Dawson, A. 1986. The effect of restricting the daily period of food availability on testicular growth in
343 starlings, *Sturnus vulgaris*. *Ibis* 128: 572-575.
- 344 Dawson, A. 1989. Pharmacological doses of thyroxine simulate the effects of increased daylength,
345 and thyroidectomy, decreased daylength, on the reproductive system of European starlings. *J*
346 *Exp Zool* 249: 62-67.
- 347 Dawson, A. 1993. Thyroidectomy progressively renders the reproductive system of starlings, *Sturnus*
348 *vulgaris*, unresponsive to changes in daylength. *J Endocrinol* 139: 51-55.
- 349 Dawson, A. 2005. The effect of temperature on photoperiodically regulated gonadal maturation,
350 regression and moult in starlings - potential consequences of climate change. *Funct Ecol* 19:
351 995-1000.
- 352 Dawson, A. 2006. Control of molt in birds: Association with prolactin and gonadal regression in
353 starlings. *Gen Comp Endocrinol* 147: 314-322.
- 354 Dawson, A. 2015. Annual gonadal cycles in birds: Modeling the effects of photoperiod on seasonal
355 changes in GnRH-1 secretion. *Front Neuroendocrinol* 37: 52-64.
- 356 Dawson, A. 2017. Daily cycles in body temperature in a songbird change with photoperiod and are
357 weakly circadian. *J Biol Rhythms* 32: 177-183.
- 358 Dawson, A. and I. Newton. 2004. Use and validation of a molt score index corrected for primary
359 feather mass. *Auk* 121: 372-379.
- 360 Dawson, A. and P.J. Sharp. 2010. Seasonal changes in concentrations of plasma LH and prolactin
361 associated with the advance in the development of photorefractoriness and molt by high
362 temperature in the starling. *Gen Comp Endocrinol* 167: 122–127.
- 363 Ernst, D.K., S.E. Lynn, and G.E. Bentley. 2016. Differential response of GnIH in the brain and
364 gonads following acute stress in a songbird. *Gen Comp Endocrinol* 227: 51-57.
- 365 Follett, B.K., T.J. Nicholls, and C.R. Mayes. 1988. Thyroxine can mimic photoperiodically induced
366 gonadal growth in Japanese quail. *J Comp Physiol B* 157: 837-843.
- 367 Foster, R.G., and B.K. Follett, B.K. 1985. The involvement of a rhodopsin-like photopigment in the
368 photoperiodic response of the Japanese quail. *J Comp Physiol* 157: 519-528.

369 Garcia-Fernandez, J.M., R. Cernuda-Cernuda, W.I.L. Davies, J. Rodgers, M. Turton, S.N. Peirson,
370 B.K. Follett, S. Halford, S. Hughes, M.W. Hankins, and R.G. Foster. 2015. The
371 hypothalamic photoreceptors regulating seasonal reproduction in birds: A prime role for VA
372 opsin. *Front Neuroendocrinol* 37: 13-28.

373 Hahn, T.P. Integration of photoperiodic and food cues to time changes in reproductive physiology by
374 an opportunistic breeder, the Red Crossbill, *Loxia curvirostra* (Aves: Carduelinae). *J. Exp.*
375 *Zool.* 272: 213-226.

376 Hau, M., and E. Gwinner. 1996. Food as a circadian zeitgeber for house sparrows: The effect of
377 different food access durations. *J Biol Rhythms* 11: 196-207.

378 Ikegami, K., and T. Yoshimura. 2016. Comparative analysis reveals the underlying mechanism of
379 vertebrate seasonal reproduction. *Gen Comp Endocrinol* 227: 64-68.

380 Laurila, M., T. Pilto and E. Hohtola. 2005. Testing the flexibility of fasting-induced hypometabolism
381 in birds: effect of photoperiod and repeated food deprivations. *J Thermal Biol* 30: 131-138.

382 McKechnie, A.E., and B.G. Lovegrove. 2002. Avian facultative hypothermic responses: A review.
383 *Condor* 104: 705-724.

384 Noakes, M.J., B. Smit, B.O. Wolf, and A.E. McKechnie. 2013. Thermoregulation in African Green
385 Pigeons (*Treron calvus*) and a re-analysis of insular effects on basal metabolic rate and
386 heterothermy in columbid birds. *J Comp Physiol B-Biochem Syst Environm Physiol* 183:
387 969-982.

388 Nord, A., J.F. Nilsson, M.I. Sandell, and J.A. Nilsson. 2009. Patterns and dynamics of rest-phase
389 hypothermia in wild and captive blue tits during winter. *J Comp Physiol B-Biochem Syst*
390 *Environm Physiol* 179: 737-745.

391 Perfito, N., S., J.M.Y. Kwong, G.E. Bentley, and M. Hau. 2008. Cue hierarchies and testicular
392 development: Is food a more potent stimulus than day length in an opportunistic breeder
393 (*Taeniopygia g. guttata*)? *Horm Behav.* 53: 567-572.

394 Perfito, N., S. Meddle, A.D. Tramontin, P.J. Sharp, and J.C. Wingfield. 2005. Seasonal gonadal
395 recrudescence in song sparrows: Response to temperature cues. *Gen Comp Endocrinol* 143:
396 121-128.

397 Pravosudov, V.V., and J.R. Lucas. 2000. The costs of being cool: a dynamic model of nocturnal
398 hypothermia by small food-caching birds in winter. *JAvian Biol* 31: 463-472.

399 Prinzinger, R., A. Preßmar, and E. Schleucher. 1991. The body temperature in birds. *Comp Biochem*
400 *Physiol* 99A: 499-506.

401 Reinertsen, R.E., and S. Haftorn. 1986. Different metabolic strategies of northern birds for nocturnal
402 survival. *J. Comp. Physiol. B-Biochem Syst Environm Physiol.* 156: 655-663.

403 Reinertsen, R.E., S. Haftorn, and E. Thaler. 1988. Is hypothermia necessary for winter survival of the
404 goldcrest *Regulus regulus*? *J für Ornithol* 129: 433-437.

405 Silverin, B., J. Wingfield, K.-A. Stokkan, R. Massa, A., Järvinen, N.-Å Andersson, M. Lambrechts,
406 A. Sorace, and D. Blomqvist. 2008. Ambient temperature effects on photo induced gonadal
407 cycles and hormonal secretion patterns in Great Tits from three different breeding latitudes.
408 *Horm Behav* 54: 60-68.

409 Song, Y.H. 2016. The effect of fluctuations in photoperiod and ambient temperature on the timing of
410 flowering: Time to move on natural environmental conditions. *Mol Cells* 39: 715-721.

411 Soumalainen, H., 1938. The effect of temperature on sexual activity of non-migratory birds,
412 stimulated by artificial lighting. *Ornis Fennica* 14: 108-112.

413 Tsutsui, K., and T. Ubuka. 2016. GnIH control of feeding and reproductive behaviors. *Front*
414 *Endocrinol* 7. [dio.org/10.3389/fendo.2016.00170](https://doi.org/10.3389/fendo.2016.00170).

415 Valle, S.E., E. Carpentier, B. Vu, K. Tsutsui, and P. Deviche. 2015. Food restriction negatively affects
416 multiple levels of the reproductive axis in male house finches, *Haemorhous mexicanus*. *J.*
417 *Exp. Biol.* 218: 2694-2704.

418 Visser, M.E., S.V. Schaper, L.J. Holleman, A. Dawson, P.J. Sharp, P. Gienapp, and S.P. Caro. 2011.
419 Genetic variation in cue sensitivity involved in avian timing of reproduction. *Functional*
420 *Ecology* 25, 868-877.

421 Waite, T.A. 1991. Nocturnal hypothermia in gray jays *Perisoreus canadensis* wintering in interior
422 Alaska. *Ornis Scand* 22: 107-110.

423 Welton, N.J., A.I. Houston, J. Ekman, and J.M. McNamara. 2002. A dynamic model of hypothermia
424 as an adaptive response by small birds to winter conditions. *Acta Biotheoretica* 50: 39-56.

425 Wikelski, M., L.B. Martin, A. Scheuerlein, M.T. Robinson, N.D. Robinson, B. Heim, M. Hau, M.,
426 and E. Gwinner. 2008. Avian circannual clocks: adaptive significance and possible
427 involvement of energy turnover in their proximate control. *Phil Trans R Soc B-Biol Sci.* 363:
428 411-423.

429 Wingfield, J.C., Hahn, T.P., Levin, R., Honey, P., 1992. Environmental predictability and control of
430 gonadal cycles in birds. *Journal of Experimental Zoology* 261, 214-231.

431 Wingfield, J.C., T.P. Hahn, D.L. Maney, S.J. Schoech, M. Wada, and M.L. Morton. 2003. Effects of
432 temperature on photoperiodically induced reproductive development, circulating plasma
433 luteinizing hormone and thyroid hormones, body mass, fat deposition and molt in mountain
434 white-crowned sparrows, *Zonotrichia leucophrys oriantha*. *Gen Comp Endocrinol* 131: 143-
435 158.

436 Wingfield, J.C., T.P. Hahn, M. Wada, L.B. Astheimer, and S. Schoech. 1996. Interrelationship of day
437 length and temperature on the control of gonadal development, body mass, and fat score in
438 white-crowned sparrows, *Zonotrichia leucophrys gambelii*. *Gen Comp Endocrinol* 101: 242-
439 255.

440 Wingfield, J.C., T.P. Hahn, M. Wada, and S. J. Schoech. 1997. Effects of day length and temperature
441 on gonadal development, body mass, and fat depots in white-crowned sparrows, *Zonotrichia*
442 *leucophrys pugetensis*. *Gen Comp Endocrinol* 107: 44-62.

443 Yoshimura, T. 2006. Molecular mechanism of the photoperiodic response of gonads in birds and
444 mammals. *Comp Biochem Physiol A-Mol Integr Physiol* 144: 345-350.

445

446

447

448 **Figure legends**

449 **Figure 1.**

450 Change in body mass during the study, starting in January and finishing in June. There were
451 four groups of starlings: one group held at 6 °C with food provided ad lib (Cold ad lib, solid
452 blue n = 7); another group held at 6 °C with access to food restricted to 7h after dawn (Cold
453 restricted, open blue n = 9); a group held at 16 °C with food provided ad lib (Warm ad lib,
454 red solid n = 8) and a final group held at 16 °C with access to food restricted to 7h after dawn
455 (Warm restricted, open red n = 8). Each point represents the mean \pm S.E.

456 **Figure 2.**

457 Starlings held under a simulated natural cycle in photoperiod between January and August.
458 Birds were held at 16 °C (solid red, n= 8) or 6 °C (solid blue, n=7) and provided with food *ad*
459 *libitum*. Each point represents the mean \pm S.E. A. Body temperature (T_b) recorded every 30
460 min for 24 h. Photoperiod was 12L:12D and the duration of darkness is shaded. T_b was lower
461 during the night in birds at 6 °C but there were no differences during the day. B. Changes in
462 testicular volume between January and June. There were no significant differences during
463 testicular maturation, but the birds held at 16 °C showed earlier regression. The vertical line
464 represents when photoperiod reached 12:12D when T_b was recorded (A). C. The progress of
465 moult in the two groups. Each point represent the date \pm S.E. that each unit of moult score
466 was reached by each bird. Birds held at 16 °C started to moult sooner.

467 **Figure 3.**

468 Starlings held under a simulated natural cycle in photoperiod between January and August.
469 All birds were held at 16 °C. One group was provided with food *ad libitum* (solid red, n= 8).
470 For the other group, access to food was prevented 7h after dawn (open red, n=8). Each point

471 represents the mean \pm S.E. A. Body temperature (T_b) recorded every 30 min for 24 h.
472 Photoperiod was 12L:12D and the duration of darkness is shaded. The vertical line represents
473 when access to food was prevented, at 13:00 under 12L:12D. T_b began to decrease after
474 access to food was prevented. B. Changes in testicular volume between January and June.
475 There were no significant differences during testicular maturation, but the birds with *ad*
476 *libitum* food showed earlier regression. The vertical line represents when photoperiod reached
477 12:12D when T_b was recorded (A). C. The progress of moult in the two groups. Each point
478 represent the date \pm S.E. that each unit of moult score was reached by each bird. Birds with
479 *ad libitum* food started to moult sooner.

480 **Figure 4.**

481 Starlings held under a simulated natural cycle in photoperiod between January and August.
482 All birds were held at 6 $^{\circ}$ C. One group was provided with food *ad libitum* (solid blue, n= 7).
483 For the other group, access to food was prevented 7h after dawn (open blue, n=9). Each point
484 represents the mean \pm S.E. A. Body temperature (T_b) recorded every 30 min for 24 h.
485 Photoperiod was 12L:12D and the duration of darkness is shaded. The vertical line represents
486 when access to food was prevented, at 13:00 under 12L:12D. T_b began to decrease after
487 access to food was prevented. B. Changes in testicular volume between January and June.
488 There were no significant differences during testicular maturation, but the birds with *ad*
489 *libitum* food showed earlier regression. The vertical line represents when photoperiod reached
490 12:12D when T_b was recorded (A). C. The progress of moult in the two groups. Each point
491 represent the date \pm S.E. that each unit of moult score was reached by each bird. Birds with
492 *ad libitum* food started to moult sooner.

493

494


495 **Figure 5.**

496 Mean body temperature in the four groups of starlings when photoperiod was 12L:12D: A
497 Birds held at 6 °C with food provided ad lib (n = 7); B birds held at 6 °C with access to food
498 restricted to 7h after dawn (n = 9); birds held at 16 °C with food provided ad lib (n = 8) and
499 birds held at 16 °C with access to food restricted to 7h after dawn (n = 8). Each point
500 represents the mean \pm S.E. The differences were not significant.

501

502

503 Fig. 1




504

505

506


507



509

510

511




513

514

515

516 Fig. 4


517



518

519

520 Fig. 5



521