

## Accepted Manuscript

Title: Interactive effects of early life stress and *CACNA1C* genotype on cortisol awakening response

Authors: K. Klaus, K. Butler, H. Gutierrez, S.J. Durrant, K. Pennington



PII: S0301-0511(18)30365-X  
DOI: <https://doi.org/10.1016/j.biopsycho.2018.05.002>  
Reference: BIOPSY 7530

To appear in:

Received date: 19-9-2017  
Revised date: 19-3-2018  
Accepted date: 2-5-2018

Please cite this article as: Klaus, K., Butler, K., Gutierrez, H., Durrant, S.J., Pennington, K., Interactive effects of early life stress and *CACNA1C* genotype on cortisol awakening response. *Biological Psychology* <https://doi.org/10.1016/j.biopsycho.2018.05.002>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Interactive effects of early life stress and *CACNA1C* genotype on cortisol awakening response**

K. Klaus<sup>a</sup>, K. Butler<sup>a</sup>, H. Gutierrez<sup>b</sup>, S.J. Durrant<sup>a</sup>, K. Pennington<sup>a\*</sup>

<sup>a</sup> School of Psychology, University of Lincoln, Brayford Wharf, Lincoln, UK, LN5 7AT

<sup>b</sup> School of Life Sciences, University of Lincoln, Brayford Pool, Lincoln, UK, LN7 6TS

*\*Corresponding author:*

*Dr Kyla Pennington*

*Sarah Swift Building, School of Psychology, University of Lincoln, Brayford Wharf, Lincoln, UK, LN5 7AT*

*Email: kpennington@lincoln.ac.uk*

*Tel. +44 (0)1522 88 6199*

**Highlights**

Investigated cortisol awakening response (CAR) in 109 healthy males

*CACNA1C* rs1006737 genotype mediated the effects of early adversity on CAR

Higher CAR in non-risk genotype carriers may indicate a more resilient HPA axis

More pronounced genotype-trauma interaction when stress experienced prior to adolescence

### Abstract

The rs1006737 (A/G) single nucleotide polymorphism within the gene encoding the Ca<sub>v</sub>1.2 subunit of the L-type voltage-dependent calcium channel (*CACNA1C*) has been strongly implicated in psychiatric disorders. In addition, calcium channels are sensitive to the effects of glucocorticoids and functional variation may contribute to altered stress responsivity. This study aimed to investigate the role of early life stress (ELS) and its interaction with *CACNA1C* rs1006737 in affecting the cortisol awakening response (CAR), an indicator of HPA-axis function. Salivary cortisol was measured in 103 healthy adult males (aged 21-63) on two consecutive days at awakening and 30 minutes later. The ELS measure investigated self-reported adverse life events prior to age 17. The results revealed a marginally significant main effect of *CACNA1C*, a significant main effect of ELS, and a significant genotype-by-ELS interaction on the CAR, whereby non-risk allele carriers (GG) who had experienced early adversity showed higher CAR compared to the other groups. Further exploratory analyses showed that this interaction may have arisen from individuals who had experienced ELS before adolescence (prior to age 13). This study is the first to provide evidence that the effect of ELS on CAR may be partially moderated via *CACNA1C* rs1006737 genotype, whereby the heightened CAR in the GG-ELS group may be an indicator of mental health resilience in response to ELS.

Keywords: *CACNA1C*, calcium channel, early life stress, childhood trauma, cortisol awakening response

## Background

It has been proposed that early life stress (ELS) might contribute to dysregulation in the development of the hypothalamic-pituitary-adrenal (HPA) axis, the body's primary stress response system (McEwen, 1998), therefore rendering genetically at risk individuals vulnerable to trauma in later life and contributing to the development of psychiatric disorders (Aas et al., 2016; Holtzman et al., 2013; van Winkel, Stefanis, & Myin-Germeys, 2008). Indeed, recent meta-analyses indicate that rates of childhood traumatic events are markedly higher in schizophrenia and related disorders (Matheson, Shepherd, Pinchbeck, Laurens, & Carr, 2013; Varese et al., 2012), bipolar disorder (Palmier-Claus, Berry, Bucci, Mansell, & Varese, 2016) and depressive disorders (Mandelli, Petrelli, & Serretti, 2015) compared to healthy controls and numerous studies have reported HPA-axis dysfunction in these conditions (Belvederi Murri et al., 2016; Heim, Newport, Mletzko, Miller, & Hemeroff, 2008; Phillips et al., 2006).

In an attempt to discern biological markers of HPA-axis function, the cortisol awakening response (CAR) has gained considerable interest. The CAR refers to the typical rise in free cortisol levels during the first 30 minutes post-awakening (Clow, Thorn, Evans, & Hucklebridge, 2004). Although the exact function of CAR is uncertain, it might help to prepare the individual for the demands of the upcoming day, or indicate a response to the natural stressor of awakening, therefore reflecting HPA-axis responsivity (Fries, Dettenborn, & Kirschbaum, 2009). Increased CAR has been shown to predict the onset of depression (Adam et al., 2010) and anxiety disorders (Adam et al., 2014), and blunted CAR has been found in patients with severe depression symptomatology (Veen et al., 2011), first-episode psychosis and schizophrenia (Berger et al., 2016), whereas remission from depression after selective serotonin reuptake inhibitor (SSRI) treatment is associated with an increase in CAR, suggesting restoration of HPA-axis activity (Ruhe et al., 2015). However, the results for individuals at-

risk for psychosis have been inconsistent (Cullen et al., 2014; Day et al., 2014; Labad et al., 2015). Furthermore, studies investigating the CAR in the context of early adversity have also produced mixed findings reporting both blunted (Desantis, Kuzawa, & Adam, 2015; Kohrt et al., 2015; Mangold, Wand, Javors, & Mintz, 2010; Meinlschmidt & Heim, 2005) and increased CAR (Butler, Klaus, Edwards, & Pennington, 2017; Engert, Efanov, Dedovic, Dagher, & Pruessner, 2011; Lu et al., 2013; Mondelli et al., 2010). These diverse findings may be the result of factors such as the developmental timing of the ELS and the interaction between stress hormones and gene variants that impact on stress responsivity (Heim & Binder, 2012; Miller, Chen, & Zhou, 2007), with increasing evidence suggesting a role for calcium channel encoding genes (Landgraf, McCarthy, & Welsh, 2014).

Calcium channels are expressed widely across the nervous system (Berger & Bartsch, 2014; Bigos et al., 2010) with intracellular calcium implicated in an array of neuronal processes, including the formation and maintenance of neuronal connections during development, and hormone secretion, neurotransmitter release and gene transcription in adulthood (Heyes et al., 2015; Lidow, 2003). The gene encoding the pore-forming Cav1.2 subunit of the L-type voltage-dependent calcium channel (*CACNA1C*) has gained particular interest in the context of psychopathology (Bhat et al., 2012; Heyes et al., 2015; Ou et al., 2015), with a number of genome-wide association studies (GWAS) supporting its role in major depression, bipolar and schizophrenia spectrum disorders (Ferreira et al., 2008; Liu et al., 2011; Schizophrenia Working Group of the Psychiatric Genomics Consortium, 2014; Sklar et al., 2008). A recent study demonstrated an interaction between two SNPs in *CACNA1C* (rs73248708 and rs116625684) and environmental stressors in increasing depression symptomatology in humans, whereas heterozygous deletion of *CACNA1C* from the forebrain glutamatergic neurons increased stress-induced anxiety-like and depression-like behaviours in mice (Dedic et al., 2018). Within *CACNA1C*, the rs1006737 (A/G) single nucleotide

polymorphism (SNP) positioned in the third intron has been most often investigated. The risk allele (A) has been associated with increased *CACNA1C* mRNA expression in the dorsolateral prefrontal cortex (PFC) in human post-mortem samples (Bigos et al., 2010), reduced hippocampal and striatal activity and diminished hippocampal connectivity during an episodic memory task (Erk et al., 2010; Krug et al., 2014), but increased hippocampal activation during emotional memory tasks at trend level (Bigos et al., 2010). Behaviourally, the risk allele has been associated with schizotypal traits in healthy individuals and the increased risk of schizotypal personality disorder (Roussos, Giakoumaki, Georgakopoulos, Robakis, & Bitsios, 2011; Roussos et al., 2013). Other studies have found increased anxiety, depression, interpersonal sensitivity and neuroticism scores, and consequently it has been proposed that the risk allele carriers may have heightened vulnerability to stress (Erk, Meyer-Lindenberg, Schmierer, et al., 2014; Erk et al., 2010).

Although the exact biological mechanisms leading to increased stress vulnerability in the rs1006737 risk allele carriers remain to be elucidated, animal studies have repeatedly demonstrated the high responsivity of calcium channels to glucocorticoid administration, which has been shown to result in increased expression of calcium channel mRNA and elevated levels of intracellular calcium (Bali, Gupta, Singh, & Jaggi, 2013; Karst et al., 2002; Kerr, Campbell, Thibault, & Landfield, 1992). The  $Ca_v1.2$  subunit encoded by *CACNA1C* is one of the main targets of corticosteroid hormones. Studies on animal models have shown that chronic stress leads to elevated calcium current amplitude and increased  $Ca_v1.2$  mRNA expression in the rat dentate gyrus, amygdala and CA3 areas (Maigaard, Hageman, Jorgensen, Jorgensen, & Wortwein, 2012; Qin, Karst, & Joels, 2004; van Gemert & Joels, 2006). Others have demonstrated increased  $Ca_v1.2$  mRNA expression in the basolateral amygdala neurons after treatment with glucocorticoids *in vitro* (Karst et al., 2002). In addition, increased  $Ca_v1.2$  mRNA expression in the rat hippocampal CA1 area has been reported in an acute stress model (Joels,

Velzing, Nair, Verkuyl, & Karst, 2003) and  $Ca_v1.2$  protein upregulation in the CA1 area in response to corticosterone exposure has also been shown (Van Gemert et al., 2009). As the hippocampus plays a key role in HPA-axis activity and the CAR (Clow, Hucklebridge, Stalder, Evans, & Thorn, 2010; Fries et al., 2009), any deviation in calcium channel activity in this brain region is likely to lead to changes in stress response system functioning.

Increasing evidence, including a study from our research group have shown that genotype-by-ELS interactions impact on cognitive outcomes, potentially via effects on stress-sensitive brain structures (Klaus et al., 2017) and we have further shown that ELS is related to an increased CAR in the same sample of healthy males (Butler et al., 2017). However, no studies to date have investigated the interaction between *CACNA1C* and ELS on the HPA-axis activity. The aim of the current study was to investigate whether the rs1006737 polymorphism of the *CACNA1C* may partially moderate the effects of ELS on HPA-axis function as measured by CAR in a sample of healthy adult males. Recent studies have suggested that a more dynamic CAR in response to external challenges is linked to a more resilient stress response system, whereas severe stressors may lead to HPA-axis exhaustion as evidenced by blunted CAR (Chida & Steptoe, 2009; Jakobsen et al., 2016; Ruhe et al., 2015). We therefore hypothesised that whereas ELS leads to higher CAR overall, the magnitude of CAR would be lower in the rs1006737 risk allele A carriers from the ELS group. Furthermore, as trauma before adolescence is thought to have more profound effects on HPA-axis programming, we expected a larger effect of genotype in those who had experienced ELS before the age of 13 (Butler et al., 2017; Maercker, Michael, Fehm, Becker, & Margraf, 2004) in comparison to those with no ELS.

## Methods

### Participants

The sample overlapped with that described in Butler et al. (2017), but the current analyses were restricted to Caucasian participants only. Briefly, 103 healthy adult males (mean age=34.54 years, SD=10.8, range 21 to 63) were recruited from Lincoln, UK, and the surrounding areas as part of a larger ongoing programme of research. None of the participants had current diagnosis of a psychiatric disorder, drug or alcohol addiction problems, nor used steroid based medication at the time of recruitment as determined through self-report. Symptoms of psychopathology and current perceived stress levels were measured by the Hospital Anxiety and Depression Scale (HADS; Zigmond & Snaith, 1983) and the Perceived Stress Scale (PSS-14; Cohen, Kamarck, & Mermelstein, 1983), respectively. Participants additionally completed custom questionnaires on their psychiatric history, demographic information and sleep history. All participants gave written informed consent to take part and the study was approved by the School of Psychology Research Ethics Committee at the University of Lincoln.

### Salivary cortisol collection and measurement

The methodology for cortisol collection, measurement and characteristics of the cortisol data can be found in Butler et al. (2017). In brief, saliva samples were collected at home using Salivette sampling devices (Sarstedt Ltd., Leicester, UK) over two consecutive days, immediately upon awakening and 30 minutes post-awakening. Inter and intra assay coefficients of variation were 17.14% and 9.67% respectively, which falls within a 20% cut-off used in earlier research (Gassling et al., 2012; Kershaw & Hall, 2016; Sink, Lochmann, & Fecteau, 2008). CAR for day 1 and day 2 was calculated by subtracting the waking cortisol concentration from the 30 minute post-awakening concentration. The CAR from day 1 and day 2 did not significantly differ,  $t(102)=-.218$ ,  $p=.828$ , and the measures were significantly



correlated,  $r=.266$ ,  $p=.007$ , with the re-test reliability estimates similar to those reported previously (Almeida, Piazza, & Stawski, 2009; Hellhammer et al., 2007). Therefore, an overall mean CAR was calculated for each participant.

### **Early life stress**

ELS was assessed using the Childhood Traumatic Events Scale (CTES; Pennebaker & Susman, 1988), which asks participants retrospectively about the occurrence of six categories of trauma: death of a close friend or relative, parental separation or divorce, traumatic sexual experience, physical violence, major illnesses or injuries, or other traumatic experiences prior to the age of 17. Participants recorded the age at which the event occurred, and rated the severity of the traumatic event and the extent that they remember confiding in others about it on a 7-point scale where 7 is extremely traumatic. According to the authors' guidelines, experiences rated as 6 or 7 on the trauma scale were classified as having experienced ELS.

### **CACNA1C genotyping**

DNA was extracted from saliva using Oragene prepIT L2P (DNA Genotek Inc., Ottawa, ON, Canada, <http://www.dnagenotek.com>) according to the manufacturer's protocol. Genotyping for *CACNA1C* rs1006737 was carried out using TaqMan® Pre-Designed SNP Genotyping Assay containing primers and fluorescent allele-specific probes (Applied Biosystems, Warrington, UK). The 25 µl reaction volume contained 30 ng of genomic DNA, 0.66 µl of 40X assay mix, 12.50 µl of TaqMan® Universal PCR Master Mix, and 11.25 µl of DNase-free water (Cat #: 4502, Sigma-Aldrich, Dorset, UK). Amplification was carried out using ABI StepOne™ Plus Real-Time PCR System with 96-well plates. The thermal profile was 60°C for 30 s, 95°C for 10 min, followed by 50 cycles at 92°C for 15 sec and 60°C for 90

sec. PCR software (StepOnePlus™ v2.0) measured SNP-specific fluorescence and genotyped each sample post-PCR.

### Data analysis

A *post hoc* power analysis using G\*Power version 3.1.9.2 (Faul, Erdfelder, Lang, & Buchner, 2007) suggested that the current sample of  $N=103$  provides a power estimate of .71 to detect medium effect size ( $f=.25$ ) in the primary analyses. For the secondary investigations into the developmental timing of ELS, *post hoc* power analysis based on the sample of  $N=103$  suggested a power estimate of .60 to detect medium effect size. Given the limited sample size in the developmental timing analyses, this was treated as an exploratory data analysis. All data were analysed using SPSS version 21 (IBM Corp., Armonk, NY, USA). The main effects of *CACNA1C* genotype and ELS (categorised as ELS/no ELS) and their interaction on CAR were investigated using a two-way analysis of covariance (ANCOVA). Assumptions for a parametric test were checked by inspecting the normality of the residuals visually using histograms and Q-Q plots and by checking skew and kurtosis (Kim, 2013). Although CAR can have high inter-individual variability (Adam et al., 2010), all CAR values in the current fell within the mean  $\pm 4$  standard deviations (as in Carnegie et al., 2014), and no outliers were therefore removed. Awakening cortisol levels, waking time, sleep duration, age and current perceived stress score measured by PSS-14 were included as covariates due to their proposed effect on CAR (Chida & Steptoe, 2009; Stalder et al., 2016). Due to strong correlations between HADS and PSS-14 scores (HADS anxiety:  $r_s=0.718$ ,  $p<0.001$ ; HADS depression:  $r_s=0.543$ ,  $p<0.001$ ), HADS scores were not included as covariates. The same statistical methods were used to investigate the effect of developmental timing of stress and *CACNA1C* variation on CAR by using a cut-off of experiencing preadolescent stress  $<13$  years in accord with previous studies (e.g., Butler et al., 2017; Maercker et al., 2004). The age at which ELS occurred was

not recorded for one participant and so this analysis was performed on 102 participants. *Post hoc* one-way ANCOVAs were used for pairwise comparisons of CAR magnitudes. *Alpha* was set at .05 and this level was used to determine significance in the main ANCOVA analyses. *Post hoc* tests were corrected for multiple comparisons using Bonferroni correction, whereby significance for the primary genotype-by-ELS group comparisons was set at  $p < .008$  and for the secondary developmental timing analyses the significance was set at  $p < .003$ .

## Results

All participants were successfully genotyped for *CACNA1C* rs1006737 and the genotype distribution conformed to the Hardy-Weinberg Equilibrium,  $\chi^2(1, N=103)=.067$ ,  $p=.795$ . As the number of AA homozygotes was low ( $n=10$ ), AA and AG carriers were combined for all analyses, therefore testing a dominant model similarly to a number of previous studies (Erk, Meyer-Lindenberg, Linden, et al., 2014; Erk, Meyer-Lindenberg, Schmierer, et al., 2014; Krug et al., 2010), including the original GWAS that implicated rs1006737 in psychiatric disorders (Ferreira et al., 2008). A two-way ANOVA revealed no difference in age, HADS anxiety and depression scores, PSS-14 scores, waking time, sleep duration or awakening cortisol levels between the two genotype groups,  $p > .05$  in all cases. HADS anxiety and depression scores and PSS-14 scores did differ between ELS groups ( $p < .05$  in all cases) as previously reported in Butler et al. (2017), but this difference did not vary between genotype groups. Chi-squared test showed no significant differences in the genotype distributions between ELS groups,  $\chi^2(1, N=103)=.007$ ,  $p=.934$  (see Table 1 for descriptive statistics). In the combined sample, the cortisol levels at 30 minutes post-awakening ( $M=9.18$  nmol/L,  $SD=5.38$ ) were significantly higher than awakening cortisol levels ( $M=7.40$  nmol/L,  $SD=4.31$ ),  $t(102) = -4.22$ ,  $p < 0.001$ . The overall mean CAR was 1.77 nmol/L ( $SD=4.26$ ; 95% CI: 0.94-2.60).

### Effects of *CACNA1C* genotypic variation and ELS on CAR

Two-way ANCOVA with *CACNA1C* genotype (AA/AG vs GG) and ELS group (ELS vs No ELS) as fixed factors and age, sleep duration, waking time, awakening cortisol levels and PSS-14 score as covariates revealed a marginally significant main effect of genotype,  $F(1, 94)=3.678, p=.058$ , partial  $\eta^2=.038$ , a significant main effect of ELS,  $F(1, 94)=8.977, p=.003$ , partial  $\eta^2=.087$ , and a significant genotype-ELS interaction,  $F(1, 94)=5.060, p=.027$ , partial  $\eta^2=.051$  (see Figure 1). *Post hoc* one-way ANCOVAs showed that GG-ELS group exhibited significantly higher CAR than GG-No ELS group,  $F(1, 40)=7.634, p=.009$ , partial  $\eta^2=.160$ , and AA/AG-No ELS group,  $F(1, 46)=8.658, p=.005$ , partial  $\eta^2=.158$ , and also higher CAR than AA/AG-ELS group at the margin of significance,  $F(1, 32)=3.601, p=.067$ , partial  $\eta^2=.101$ .




Figure 1. Effect of *CACNA1C* genotype (AA/AG vs GG) and early life stress (ELS vs No ELS) before the age of 17 on mean cortisol awakening response (nmol/L). Trauma categories are based on Childhood Traumatic Events Scale (Pennebaker & Susman, 1988), where ELS= a

rating of 6-7 on trauma scale and No ELS= a rating of 0-5. Error bars show +/- 1 standard errors. \*\* $p < .01$

As the developmental timing of ELS has been found to affect HPA-axis function, in particular prior to adolescence (Lupien, McEwen, Gunnar, & Heim, 2009), we further investigated whether the observed genotype-ELS association stemmed from the individuals who had experienced ELS before age 13. Two-way ANCOVA with *CACNA1C* genotype and three ELS groups (No ELS, ELS <13, ELS 13-17) as fixed factors showed a significance of the main effect of genotype at trend level,  $F(1, 91)=2.599$ ,  $p=.110$ , partial  $\eta^2=.028$ , a significant main effect of ELS,  $F(2, 91)=4.716$ ,  $p=.011$ , partial  $\eta^2=.094$ , and a marginally significant genotype-ELS interaction,  $F(2, 91)=2.775$ ,  $p=.068$ , partial  $\eta^2=.057$  (see Figure 2).




Figure 2. Effect of *CACNA1C* genotype (AA/AG vs GG) and early life stress (No ELS, ELS before the age of 13, ELS age 13-17) on mean cortisol awakening response (nmol/L). The number of participants in each group is given above the bars. Trauma categories are based on

Childhood Traumatic Events Scale (Pennebaker & Susman, 1988), where ELS= a rating of 6-7 on trauma scale and No ELS= a rating 0-5. Error bars show +/- 1 standard errors.

### Conclusions

This study aimed to investigate whether *CACNA1C* rs1006737 genotype partially moderates the effect of ELS on the CAR in a sample of healthy males, and further to investigate whether this association is driven by ELS experienced prior to adolescence. In agreement with our hypothesis, we found a genotype-by-ELS interaction whereby CAR profiles in the ELS group differed according to the *CACNA1C* genotype, with the schizophrenia non-risk genotype (GG) carriers showing significantly elevated CAR. The CAR did not differ by *CACNA1C* genotype in the no-ELS group. This builds on our previous findings in this dataset (Butler et al., 2017) and provides further evidence for a role of calcium channel signalling in HPA-axis functioning. Furthermore, secondary exploratory analyses suggested that the significant genotype-ELS interaction may have arisen from the ELS encountered prior to the onset of adolescence. As the analyses on the developmental timing included small groups and the results were only marginally significant, these findings should be treated with caution. However, they are in agreement with previous evidence of the role of calcium signalling in the establishment of neural circuitry and stress sensitivity of neuronal connectivity during key stages of brain development (Heyes et al., 2015; Molet et al., 2016).

This research demonstrates for the first time that increased CAR in response to ELS is not uniform in all individuals, but may be partially moderated by the *CACNA1C* genotype. The analyses also showed a significant main effect of ELS and a marginally significant main effect of *CACNA1C* genotype, but these main effects were largely driven by the non-risk GG homozygotes from the ELS group. It could be argued that the elevated CAR in the GG-ELS group were due to current stress levels (Pruessner, Hellhammer, & Kirschbaum, 1999) or

indicate vulnerability to developing depressive disorders (Adam et al., 2010). Indeed, a recent study demonstrated an association between two *CACNA1C* SNPs and stressful life events on depressive symptomatology in healthy adults (Dedic et al., 2018). However, the results from our ANCOVA model on the association between ELS and *CACNA1C* were significant even when controlling for current perceived stress, and remained unchanged after further controlling for current anxiety and depression levels (data not presented), suggesting that the underlying mechanisms of glucocorticoid response may be determined during development (Lupien et al., 2009). Indeed, the interaction between *CACNA1C* genotype and ELS, whereby the CAR was markedly higher in the GG-ELS group compared to other groups, was predominantly seen when the stressor was encountered prior to adolescence. Although these results are treated as exploratory, these findings may be expected, as voltage-gated calcium channels are essential for early brain development (Heyes et al., 2015), and whereas structures involved in HPA-axis function, such as the hippocampus, amygdala and PFC continue maturing postnatally (Gogtay et al., 2006; Lupien et al., 2009), both the hippocampus (Lupien et al., 2009) and the amygdala may be most sensitive to stress before adolescence (Pechtel, Lyons-Ruth, Anderson, & Teicher, 2014).

Although the exact function of the CAR is still debated, the results presented here may suggest that increased CAR in those GG carriers exposed to early trauma may be thought of as a formerly adaptive response that serves to prepare the individual for the challenges of the upcoming day during childhood and is carried on to adulthood (Lupien et al., 2009). A more dynamic CAR may therefore indicate a flexible HPA axis that responds adaptively to external stressors (Jakobsen et al., 2016). The adaptive nature of a dynamic CAR is further supported by the findings that patients remitting from depression in response to SSRI treatment experience an increase in CAR, suggesting restoration of HPA-axis activity (Ruhe et al., 2015). On the other hand, the risk allele A carriers from the ELS group showed CAR levels

comparable to the No ELS group. As high levels of stress, or chronic stress can lead to CAR levels similar to that of the non-stressed individuals (Wardenaar et al., 2011), this potentially suggests hypocortisolism due to increased vulnerability to stressors in the A carriers (Erk et al., 2010) and lends evidence to the nonlinear nature of cortisol which is further moderated by genetic variation.

The mechanisms by which calcium channels mediate the effects of ELS on HPA-axis dynamics are still to be elucidated and are likely to be complex. Several studies have shown that glucocorticoid exposure increases  $Ca_v1.2$  mRNA and protein expression in the hippocampal areas (Joels et al., 2003; Maigaard et al., 2012; Van Gemert et al., 2009) and elevated L-type calcium channel expression potentiates additional glucocorticoid release, creating a positive feedback loop. Consequently, increased intracellular calcium exposure over an extended period of time may lead to deleterious effects associated with stress, including cell death and associated hippocampal damage (Bali et al., 2013; Erk et al., 2010; van Gemert & Joels, 2006). A more recent study showed that heterozygous *CACNA1C* knockout in the mouse forebrain leads to increased susceptibility to anxiety when subjected to environmental stressors during development, but this study failed to find an effect of calcium channel expression on basal and stress-induced corticosterone levels (Dedic et al., 2018). Studies on humans have shown that *CACNA1C* rs1006737 risk allele is associated with increased *CACNA1C* mRNA expression in the PFC (Bigos et al., 2010), but decreased expression in the cerebellum (Gershon et al., 2014), suggesting different regulatory mechanisms of this SNP depending on the brain region. Further work needs to be conducted on the effects of glucocorticoids and genetic variation in *CACNA1C* on  $Ca_v1.2$  expression in brain regions involved in HPA-axis function.

There were some limitations in the current study. Firstly, the overall sample and the subgroups were limited in size, particularly in the secondary analyses on the developmental timing. Additionally, the power calculation was based on a medium effect size and as such we



were not powered to detect small effect sizes. We therefore acknowledge that a larger sample is needed to replicate these findings with greater power and to detect small effect sizes. Furthermore, owing to the small number of AA homozygotes ( $n=10$ ), we pooled AA homozygotes with AG heterozygotes for the current analyses, which might have obscured any effects attributable to AA homozygosity. In addition, rs1006737 is in complete linkage disequilibrium with several other SNPs positioned in the third intron of *CACNA1C* (Nyegaard et al., 2010) and we therefore cannot disentangle the effects specific to the SNP investigated in this study. We also acknowledge that other genes previously implicated in cortisol function, such as the fk506 binding protein encoding *FKBP5* (Kohrt et al., 2015), brain derived neurotrophic factor encoding *BDNF* (Shalev et al., 2009), and serotonin transporter encoding *5-HTT* (Wust et al., 2009) may, possibly additively, also affect HPA-axis activity. Moreover, both glucocorticoids and calcium channel activity can affect gene transcription (Lupien et al., 2009; West, Griffith, & Greenberg, 2002), meaning that the effects seen here could be due to the regulation of other genes not investigated in this study.

It should also be noted that the mean CAR reported in this paper is rather low. As previously acknowledged, there are many factors which can influence the CAR (Chida & Steptoe, 2009; Stalder et al., 2016). Of these, we did adjust for awakening cortisol levels, waking time, sleep duration, age and current perceived stress levels in our analyses. Yet other factors might have affected CAR, such as socioeconomic status and body mass index (Stalder et al., 2016) which we did not have access to in our dataset. Further discussion on the factors influencing CAR magnitude, such as gender (Pruessner et al., 1997; Vreeburg et al., 2009; Wust et al., 2000), undictated waking times (Kudielka & Kirschbaum, 2003) and the sampling schedule have been discussed in Butler et al. (2017). However, it has also been proposed that negative CAR is a legitimate phenomenon (Stalder et al., 2016). Our previous study demonstrated that CAR magnitude may also be associated with the type of traumatic event

experienced and cumulative trauma exposure (Butler et al., 2017), but due to the small sample size investigation into these aspects of trauma was not feasible and could be addressed in future studies. Additionally, CAR is known to have high intra-individual variability due to both trait and state factors (Hellhammer et al., 2007). Although the CAR re-test reliability in the current study was comparable to previous reports (Almeida et al., 2009; Hellhammer et al., 2007), future studies could collect cortisol data over more days to increase the reliability of the CAR. Finally, as gender differences in HPA-axis function after ELS have been reported (DeSantis et al., 2011; Kudielka & Kirschbaum, 2003; Kudielka & Kirschbaum, 2005), future studies are needed to investigate the ELS-*CACNA1C* interactions on CAR in female samples.

In sum, this is the first study to demonstrate that the effect of ELS on CAR may be partially moderated by *CACNA1C* genotype, whereby the non-risk GG homozygotes who have experienced ELS exhibit higher CAR, potentially supporting the idea of the adaptive nature of heightened CAR (Fries et al., 2009), but the risk allele A carriers subjected to ELS show a pattern suggestive of hypoactive CAR. We further provided some preliminary evidence that this effect may be more profound if the adversity has been experienced before adolescence. In light of the previously controversial findings in this area, the results from the current study emphasise the importance of considering the individual's genetic background when investigating the effect of environmental stressors on HPA-axis function and provide a potential mechanism by which ELS may affect CAR.

## Acknowledgements

This work was supported by the University of Lincoln's College of Social Science Research Fund, the School of Psychology at the University of Lincoln and WiSE Academic Returners' Research Fund (R2F) awarded to KP and a University of Lincoln College of Social Science PhD studentship to KK.

## References

- Aas, M., Henry, C., Andreassen, O. A., Bellivier, F., Melle, I., & Etain, B. (2016). The role of childhood trauma in bipolar disorders. *International Journal of Bipolar Disorders*, *4*, 10. doi: 10.1186/s40345-015-0042-0
- Adam, E. K., Doane, L. D., Zinbarg, R. E., Mineka, S., Craske, M. G., & Griffith, J. W. (2010). Prospective prediction of major depressive disorder from cortisol awakening responses in adolescence. *Psychoneuroendocrinology*, *35*(6), 921-931. doi: 10.1016/j.psyneuen.2009.12.007
- Adam, E. K., Vrshek-Schallhorn, S., Kendall, A. D., Mineka, S., Zinbarg, R. E., & Craske, M. G. (2014). Prospective associations between the cortisol awakening response and first onsets of anxiety disorders over a six-year follow-up--2013 Curt Richter Award Winner. *Psychoneuroendocrinology*, *44*, 47-59. doi: 10.1016/j.psyneuen.2014.02.014
- Almeida, D. M., Piazza, J. R., & Stawski, R. S. (2009). Interindividual differences and intraindividual variability in the cortisol awakening response: An examination of age and gender. *Psychology and Aging*, *24*(4), 819. doi: 10.1037/a0017910
- Bali, A., Gupta, S., Singh, N., & Jaggi, A. S. (2013). Implicating the role of plasma membrane localized calcium channels and exchangers in stress-induced deleterious effects.

- European Journal of Pharmacology*, 714(1-3), 229-238. doi: 10.1016/j.ejphar.2013.06.010
- Belvederi Murri, M., Prestia, D., Mondelli, V., Pariante, C., Patti, S., Olivieri, B., Arzani, C., Masotti, M., Respino, M., Antonioli, M., Vassallo, L., Serafini, G., Perna, G., Pompili, M., & Amore, M. (2016). The HPA axis in bipolar disorder: Systematic review and meta-analysis. *Psychoneuroendocrinology*, 63, 327-342. doi: 10.1016/j.psyneuen.2015.10.014
- Berger, M., Kraeuter, A. K., Romanik, D., Malouf, P., Amminger, G. P., & Sarnyai, Z. (2016). Cortisol awakening response in patients with psychosis: Systematic review and meta-analysis. *Neuroscience and Biobehavioral Reviews*, 68, 157-166. doi: 10.1016/j.neubiorev.2016.05.027
- Berger, S. M., & Bartsch, D. (2014). The role of L-type voltage-gated calcium channels Cav1.2 and Cav1.3 in normal and pathological brain function. *Cell and Tissue Research*, 357(2), 463-476. doi: 10.1007/s00441-014-1936-3
- Bhat, S., Dao, D. T., Terrillion, C. E., Arad, M., Smith, R. J., Soldatov, N. M., & Gould, T. D. (2012). CACNA1C (Cav1.2) in the pathophysiology of psychiatric disease. *Progress in Neurobiology*, 99(1), 1-14. doi: 10.1016/j.pneurobio.2012.06.001
- Bigos, K. L., Mattay, V. S., Callicott, J. H., Straub, R. E., Vakkalanka, R., Kolachana, B., Hyde, T. M., Lipska, B. K., Kleinman, J. E., & Weinberger, D. R. (2010). Genetic variation in CACNA1C affects brain circuitries related to mental illness. *Archives of General Psychiatry*, 67(9), 939-945. doi: 10.1001/archgenpsychiatry.2010.96
- Butler, K., Klaus, K., Edwards, L., & Pennington, K. (2017). Elevated cortisol awakening response associated with early life stress and impaired executive function in healthy adult males. *Hormones and Behavior*, 95, 13-21. doi: 10.1016/j.yhbeh.2017.07.013

- Carnegie, R., Araya, R., Ben-Shlomo, Y., Glover, V., O'Connor, T. G., O'Donnell, K. J., Pearson, R., & Lewis, G. (2014). Cortisol awakening response and subsequent depression: prospective longitudinal study. *British Journal of Psychiatry*, *204*(2), 137-143. doi: 10.1192/bjp.bp.113.126250
- Chida, Y., & Steptoe, A. (2009). Cortisol awakening response and psychosocial factors: a systematic review and meta-analysis. *Biological Psychology*, *80*(3), 265-278. doi: 10.1016/j.biopsycho.2008.10.004
- Clow, A., Hucklebridge, F., Stalder, T., Evans, P., & Thorn, L. (2010). The cortisol awakening response: more than a measure of HPA axis function. *Neuroscience and Biobehavioral Reviews*, *35*(1), 97-103. doi: 10.1016/j.neubiorev.2009.12.011
- Clow, A., Thorn, L., Evans, P., & Hucklebridge, F. (2004). The awakening cortisol response: methodological issues and significance. *Stress*, *7*(1), 29-37. doi: 10.1080/10253890410001667205
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, *24*(4), 385-396.
- Cullen, A. E., Zunszain, P. A., Dickson, H., Roberts, R. E., Fisher, H. L., Pariante, C. M., & Laurens, K. R. (2014). Cortisol awakening response and diurnal cortisol among children at elevated risk for schizophrenia: Relationship to psychosocial stress and cognition. *Psychoneuroendocrinology*, *46*, 1-13. doi: 10.1016/j.psyneuen.2014.03.010
- Day, F. L., Valmaggia, L. R., Mondelli, V., Papadopoulos, A., Papadopoulos, I., Pariante, C. M., & McGuire, P. (2014). Blunted Cortisol Awakening Response in People at Ultra High Risk of Developing Psychosis. *Schizophrenia Research*, *158*(1-3), 25-31. doi: 10.1016/j.schres.2014.06.041
- Dedic, N., Pohlmann, M. L., Richter, J. S., Mehta, D., Czamara, D., Metzger, M. W., Dine, J., Bedenk, B. T., Hartmann, J., Wagner, K. V., Jurik, A., Almlil, L. M., Lori, A.,

- Moosmang, S., Hofmann, F., Wotjak, C. T., Rammes, G., Eder, M., Chen, A., Ressler, K. J., Wurst, W., Schmidt, M. V., Binder, E. B., & Deussing, J. M. (2018). Cross-disorder risk gene CACNA1C differentially modulates susceptibility to psychiatric disorders during development and adulthood. *Molecular Psychiatry*, *23*(3), 533-543. doi: 10.1038/mp.2017.133
- Desantis, A. S., Kuzawa, C. W., & Adam, E. K. (2015). Developmental origins of flatter cortisol rhythms: socioeconomic status and adult cortisol activity. *American Journal of Human Biology*, *27*(4), 458-467. doi: 10.1002/ajhb.22668
- DeSantis, S. M., Baker, N. L., Back, S. E., Spratt, E., Ciolino, J. D., Moran-Santa Maria, M., Dipankar, B., & Brady, K. T. (2011). Gender differences in the effect of early life trauma on hypothalamic-pituitary-adrenal axis functioning. *Depression and Anxiety*, *28*(5), 383-392. doi: 10.1002/da.20795
- Engert, V., Efanov, S. I., Dedovic, K., Dagher, A., & Pruessner, J. C. (2011). Increased cortisol awakening response and afternoon/evening cortisol output in healthy young adults with low early life parental care. *Psychopharmacology*, *214*(1), 261-268. doi: 10.1007/s00213-010-1918-4
- Erk, S., Meyer-Lindenberg, A., Linden, D. E. J., Lancaster, T., Mohnke, S., Grimm, O., Degenhardt, F., Holmans, P., Pocklington, A., Schmierer, P., Haddad, L., Muhleisen, T. W., Mattheisen, M., Witt, S. H., Romanczuk-Seiferth, N., Tost, H., Schott, B. H., Cichon, S., Nothen, M. M., Rietschel, M., Heinz, A., & Walter, H. (2014). Replication of brain function effects of a genome-wide supported psychiatric risk variant in the CACNA1C gene and new multi-locus effects. *Neuroimage*, *94*(Supplement C), 147-154. doi: 10.1016/j.neuroimage.2014.03.007
- Erk, S., Meyer-Lindenberg, A., Schmierer, P., Mohnke, S., Grimm, O., Garbusow, M., Haddad, L., Poehland, L., Muhleisen, T. W., Witt, S. H., Tost, H., Kirsch, P., Romanczuk-

- Seiferth, N., Schott, B. H., Cichon, S., Nothen, M. M., Rietschel, M., Heinz, A., & Walter, H. (2014). Hippocampal and Frontolimbic Function as Intermediate Phenotype for Psychosis: Evidence from Healthy Relatives and a Common Risk Variant in CACNA1C. *Biological Psychiatry*, 76(6), 466-475. doi: 10.1016/j.biopsych.2013.11.025
- Erk, S., Meyer-Lindenberg, A., Schnell, K., Opitz von Boberfeld, C., Esslinger, C., Kirsch, P., Grimm, O., Arnold, C., Haddad, L., Witt, S. H., Cichon, S., Nothen, M. M., Rietschel, M., & Walter, H. (2010). Brain function in carriers of a genome-wide supported bipolar disorder variant. *Archives of General Psychiatry*, 67(8), 803-811. doi: 10.1001/archgenpsychiatry.2010.94
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G\*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175-191.
- Ferreira, M. A., O'Donovan, M. C., Meng, Y. A., Jones, I. R., Ruderfer, D. M., Jones, L., Fan, J., Kirov, G., Perlis, R. H., Green, E. K., Smoller, J. W., Grozeva, D., Stone, J., Nikolov, I., Chambert, K., Hamshere, M. L., Nimgaonkar, V. L., Moskvina, V., Thase, M. E., Caesar, S., Sachs, G. S., Franklin, J., Gordon-Smith, K., Ardlie, K. G., Gabriel, S. B., Fraser, C., Blumenstiel, B., Defelice, M., Breen, G., Gill, M., Morris, D. W., Elkin, A., Muir, W. J., McGhee, K. A., Williamson, R., MacIntyre, D. J., MacLean, A. W., St. C. D., Robinson, M., Van Beck, M., Pereira, A. C., Kandaswamy, R., McQuillin, A., Collier, D. A., Bass, N. J., Young, A. H., Lawrence, J., Ferrier, I. N., Anjorin, A., Farmer, A., Curtis, D., Scolnick, E. M., McGuffin, P., Daly, M. J., Corvin, A. P., Holmans, P. A., Blackwood, D. H., Gurling, H. M., Owen, M. J., Purcell, S. M., Sklar, P., Craddock, N., & Wellcome Trust Case Control, C. (2008). Collaborative genome-

wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder.

*Nature Genetics*, 40(9), 1056-1058. doi: 10.1038/ng.209

Fries, E., Dettenborn, L., & Kirschbaum, C. (2009). The cortisol awakening response (CAR): facts and future directions. *International Journal of Psychophysiology*, 72(1), 67-73. doi: 10.1016/j.ijpsycho.2008.03.014

Gassling, V., Holterhus, P. M., Herbers, D., Kulle, A., Niederberger, U., Hedderich, J., Wiltfang, J., & Gerber, W. D. (2012). Stress-coping and cortisol analysis in patients with non-syndromic cleft lip and palate: an explorative study. *PloS One*, 7(7), e41015. doi: 10.1371/journal.pone.0041015

Gershon, E. S., Grennan, K., Busnello, J., Badner, J. A., Ovsiew, F., Memon, S., Alliey-Rodriguez, N., Cooper, J., Romanos, B., & Liu, C. (2014). A rare mutation of CACNA1C in a patient with bipolar disorder, and decreased gene expression associated with a bipolar-associated common SNP of CACNA1C in brain. *Molecular Psychiatry*, 19(8), 890-894. doi: 10.1038/mp.2013.107

Gogtay, N., Nugent, T. F., 3rd, Herman, D. H., Ordonez, A., Greenstein, D., Hayashi, K. M., Clasen, L., Toga, A. W., Giedd, J. N., Rapoport, J. L., & Thompson, P. M. (2006). Dynamic mapping of normal human hippocampal development. *Hippocampus*, 16(8), 664-672. doi: 10.1002/hipo.20193

Heim, C., & Binder, E. B. (2012). Current research trends in early life stress and depression: review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Experimental Neurology*, 233(1), 102-111. doi: 10.1016/j.expneurol.2011.10.032

Heim, C., Newport, D. J., Mletzko, T., Miller, A. H., & Hemeroff, C. B. (2008). The link between childhood trauma and depression: Insights from HPA axis studies in humans. *Psychoneuroendocrinology*, 33(6), 693-710. doi: 10.1016/j.psyneuen.2008.03.008



- Hellhammer, J., Fries, E., Schweisthal, O. W., Schlotz, W., Stone, A. A., & Hagemann, D. (2007). Several daily measurements are necessary to reliably assess the cortisol rise after awakening: state- and trait components. *Psychoneuroendocrinology*, *32*(1), 80-86. doi: 10.1016/j.psyneuen.2006.10.005
- Heyes, S., Pratt, W. S., Rees, E., Dahimene, S., Ferron, L., Owen, M. J., & Dolphin, A. C. (2015). Genetic disruption of voltage-gated calcium channels in psychiatric and neurological disorders. *Progress in Neurobiology*, *134*, 36-54. doi: 10.1016/j.pneurobio.2015.09.002
- Holtzman, C. W., Trotman, H. D., Goulding, S. M., Ryan, A. T., Macdonald, A. N., Shapiro, D. I., Brasfield, J. L., & Walker, E. F. (2013). Stress and neurodevelopmental processes in the emergence of psychosis. *Neuroscience*, *249*, 172-191. doi: 10.1016/j.neuroscience.2012.12.017
- Jakobsen, G. R., Fisher, P. M., Dyssegaard, A., McMahon, B., Holst, K. K., Lehel, S., Svarer, C., Jensen, P. S., Knudsen, G. M., & Frokjaer, V. G. (2016). Brain serotonin 4 receptor binding is associated with the cortisol awakening response. *Psychoneuroendocrinology*, *67*(Supplement C), 124-132. doi: 10.1016/j.psyneuen.2016.01.032
- Joels, M., Velzing, E., Nair, S., Verkuyl, J. M., & Karst, H. (2003). Acute stress increases calcium current amplitude in rat hippocampus: temporal changes in physiology and gene expression. *European Journal of Neuroscience*, *18*(5), 1315-1324.
- Karst, H., Nair, S., Velzing, E., Rumpff-van Essen, L., Slagter, E., Shinnick-Gallagher, P., & Joels, M. (2002). Glucocorticoids alter calcium conductances and calcium channel subunit expression in basolateral amygdala neurons. *European Journal of Neuroscience*, *16*(6), 1083-1089.

- Kerr, D. S., Campbell, L. W., Thibault, O., & Landfield, P. W. (1992). Hippocampal glucocorticoid receptor activation enhances voltage-dependent Ca<sup>2+</sup> conductances: relevance to brain aging. *Proceedings of the National Academy of Sciences of the United States of America*, 89(18), 8527-8531.
- Kershaw, J. L., & Hall, A. J. (2016). Seasonal variation in harbour seal (*Phoca vitulina*) blubber cortisol - A novel indicator of physiological state? *Scientific Reports*, 6, 21889. doi: 10.1038/srep21889
- Kim, H.-Y. (2013). Statistical notes for clinical researchers: assessing normal distribution (2) using skewness and kurtosis. *Restorative dentistry & endodontics*, 38(1), 52-54.
- Klaus, K., Butler, K., Durrant, S. J., Ali, M., Inglehearn, C. F., Hodgson, T. L., Gutierrez, H., & Pennington, K. (2017). The effect of COMT Val158Met and DRD2 C957T polymorphisms on executive function and the impact of early life stress. *Brain and behavior*, 7(5), e00695.
- Kohrt, B. A., Worthman, C. M., Ressler, K. J., Mercer, K. B., Upadhaya, N., Koirala, S., Nepal, M. K., Sharma, V. D., & Binder, E. B. (2015). Cross-cultural gene– environment interactions in depression, post-traumatic stress disorder, and the cortisol awakening response: FKBP5 polymorphisms and childhood trauma in South Asia: GxE interactions in South Asia. *International Review of Psychiatry*, 27(3), 180-196.
- Krug, A., Nieratschker, V., Markov, V., Krach, S., Jansen, A., Zerres, K., Eggermann, T., Stocker, T., Shah, N. J., Treutlein, J., Muhleisen, T. W., & Kircher, T. (2010). Effect of CACNA1C rs1006737 on neural correlates of verbal fluency in healthy individuals. *Neuroimage*, 49(2), 1831-1836. doi: 10.1016/j.neuroimage.2009.09.028
- Krug, A., Witt, S. H., Backes, H., Dietsche, B., Nieratschker, V., Shah, N. J., Nothen, M. M., Rietschel, M., & Kircher, T. (2014). A genome-wide supported variant in CACNA1C influences hippocampal activation during episodic memory encoding and retrieval.

*European Archives of Psychiatry and Clinical Neuroscience*, 264(2), 103-110. doi: 10.1007/s00406-013-0428-x

Kudielka, B., & Kirschbaum, C. (2003). Awakening cortisol responses are influenced by health status and awakening time but not by menstrual cycle phase. *Psychoneuroendocrinology*, 28(1), 35-47.

Kudielka, B. M., & Kirschbaum, C. (2005). Sex differences in HPA axis responses to stress: a review. *Biological Psychology*, 69(1), 113-132. doi: 10.1016/j.biopsycho.2004.11.009

Labad, J., Stojanovic-Pérez, A., Montalvo, I., Solé, M., Cabezas, Á., Ortega, L., Moreno, I., Vilella, E., Martorell, L., & Reynolds, R. M. (2015). Stress biomarkers as predictors of transition to psychosis in at-risk mental states: roles for cortisol, prolactin and albumin. *Journal of Psychiatric Research*, 60, 163-169.

Landgraf, D., McCarthy, M. J., & Welsh, D. K. (2014). Circadian clock and stress interactions in the molecular biology of psychiatric disorders. *Curr Psychiatry Rep*, 16(10), 483. doi: 10.1007/s11920-014-0483-7

Lidow, M. S. (2003). Calcium signaling dysfunction in schizophrenia: a unifying approach. *Brain Research: Brain Research Reviews*, 43(1), 70-84.

Liu, Y., Blackwood, D. H., Caesar, S., de Geus, E. J., Farmer, A., Ferreira, M. A., Ferrier, I. N., Fraser, C., Gordon-Smith, K., Green, E. K., Grozeva, D., Gurling, H. M., Hamshere, M. L., Heutink, P., Holmans, P. A., Hoogendijk, W. J., Hottenga, J. J., Jones, L., Jones, I. R., Kirov, G., Lin, D., McGuffin, P., Moskvina, V., Nolen, W. A., Perlis, R. H., Posthuma, D., Scolnick, E. M., Smit, A. B., Smit, J. H., Smoller, J. W., St Clair, D., van Dyck, R., Verhage, M., Willemsen, G., Young, A. H., Zandbelt, T., Boomsma, D. I., Craddock, N., O'Donovan, M. C., Owen, M. J., Penninx, B. W., Purcell, S., Sklar, P., Sullivan, P. F., & Wellcome Trust Case-Control, C. (2011). Meta-analysis of genome-

- wide association data of bipolar disorder and major depressive disorder. *Molecular Psychiatry*, *16*(1), 2-4. doi: 10.1038/mp.2009.107
- Lu, S., Gao, W., Wei, Z., Wu, W., Liao, M., Ding, Y., Zhang, Z., & Li, L. (2013). Reduced cingulate gyrus volume associated with enhanced cortisol awakening response in young healthy adults reporting childhood trauma. *PloS One*, *8*(7), e69350. doi: 10.1371/journal.pone.0069350
- Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews: Neuroscience*, *10*(6), 434-445. doi: 10.1038/nrn2639
- Maercker, A., Michael, T., Fehm, L., Becker, E. S., & Margraf, J. (2004). Age of traumatisation as a predictor of post-traumatic stress disorder or major depression in young women. *British Journal of Psychiatry*, *184*(6), 482-487. doi: DOI 10.1192/bjp.184.6.482
- Maigaard, K., Hageman, I., Jorgensen, A., Jorgensen, M. B., & Wortwein, G. (2012). Electroconvulsive stimulations prevent chronic stress-induced increases in L-type calcium channel mRNAs in the hippocampus and basolateral amygdala. *Neuroscience Letters*, *516*(1), 24-28. doi: 10.1016/j.neulet.2012.03.043
- Mandelli, L., Petrelli, C., & Serretti, A. (2015). The role of specific early trauma in adult depression: A meta-analysis of published literature. Childhood trauma and adult depression. *European Psychiatry*, *30*(6), 665-680. doi: 10.1016/j.eurpsy.2015.04.007
- Mangold, D., Wand, G., Javors, M., & Mintz, J. (2010). Acculturation, childhood trauma and the cortisol awakening response in Mexican-American adults. *Hormones and Behavior*, *58*(4), 637-646. doi: 10.1016/j.yhbeh.2010.06.010
- Matheson, S. L., Shepherd, A. M., Pinchbeck, R. M., Laurens, K. R., & Carr, V. J. (2013). Childhood adversity in schizophrenia: a systematic meta-analysis. *Psychological Medicine*, *43*(2), 225-238. doi: 10.1017/S0033291712000785

- McEwen, B. S. (1998). Protective and damaging effects of stress mediators. *New England Journal of Medicine*, 338(3), 171-179. doi: 10.1056/NEJM199801153380307
- Meinlschmidt, G., & Heim, C. (2005). Decreased cortisol awakening response after early loss experience. *Psychoneuroendocrinology*, 30(6), 568-576. doi: 10.1016/j.psyneuen.2005.01.006
- Miller, G. E., Chen, E., & Zhou, E. S. (2007). If it goes up, must it come down? Chronic stress and the hypothalamic-pituitary-adrenocortical axis in humans. *Psychological Bulletin*, 133(1), 25.
- Molet, J., Maras, P. M., Kinney-Lang, E., Harris, N. G., Rashid, F., Ivy, A. S., Solodkin, A., Obenaus, A., & Baram, T. Z. (2016). MRI uncovers disrupted hippocampal microstructure that underlies memory impairments after early-life adversity. *Hippocampus*, 26(12), 1618-1632. doi: 10.1002/hipo.22661
- Mondelli, V., Dazzan, P., Hepgul, N., Di Forti, M., Aas, M., D'Albenzio, A., Di Nicola, M., Fisher, H., Handley, R., Marques, T. R., Morgan, C., Navari, S., Taylor, H., Papadopoulos, A., Aitchison, K. J., Murray, R. M., & Pariante, C. M. (2010). Abnormal cortisol levels during the day and cortisol awakening response in first-episode psychosis: the role of stress and of antipsychotic treatment. *Schizophrenia Research*, 116(2-3), 234-242. doi: 10.1016/j.schres.2009.08.013
- Nyegaard, M., Demontis, D., Foldager, L., Hedemand, A., Flint, T. J., Sorensen, K. M., Andersen, P. S., Nordentoft, M., Werge, T., Pedersen, C. B., Hougaard, D. M., Mortensen, P. B., Mors, O., & Borglum, A. D. (2010). CACNA1C (rs1006737) is associated with schizophrenia. *Molecular Psychiatry*, 15(2), 119-121. doi: 10.1038/mp.2009.69
- Ou, X., Crane, D. E., MacIntosh, B. J., Young, L. T., Arnold, P., Ameis, S., & Goldstein, B. I. (2015). CACNA1C rs1006737 genotype and bipolar disorder: Focus on intermediate

- phenotypes and cardiovascular comorbidity. *Neuroscience and Biobehavioral Reviews*, 55, 198-210. doi: 10.1016/j.neubiorev.2015.04.022
- Palmier-Claus, J. E., Berry, K., Bucci, S., Mansell, W., & Varese, F. (2016). Relationship between childhood adversity and bipolar affective disorder: systematic review and meta-analysis. *British Journal of Psychiatry*, 209(6), 454-459. doi: 10.1192/bjp.bp.115.179655
- Pechtel, P., Lyons-Ruth, K., Anderson, C. M., & Teicher, M. H. (2014). Sensitive periods of amygdala development: the role of maltreatment in preadolescence. *Neuroimage*, 97, 236-244. doi: 10.1016/j.neuroimage.2014.04.025
- Pennebaker, J. W., & Susman, J. R. (1988). Disclosure of traumas and psychosomatic processes. *Social Science and Medicine*, 26(3), 327-332.
- Phillips, L. J., McGorry, P. D., Garner, B., Thompson, K. N., Pantelis, C., Wood, S. J., & Berger, G. (2006). Stress, the hippocampus and the hypothalamic-pituitary-adrenal axis: implications for the development of psychotic disorders. *Australian and New Zealand Journal of Psychiatry*, 40(9), 725-741. doi: 10.1080/j.1440-1614.2006.01877.x
- Pruessner, J. C., Hellhammer, D. H., & Kirschbaum, C. (1999). Burnout, perceived stress, and cortisol responses to awakening. *Psychosomatic Medicine*, 61(2), 197-204.
- Pruessner, J. C., Wolf, O. T., Hellhammer, D. H., Buske-Kirschbaum, A., von Auer, K., Jobst, S., Kaspers, F., & Kirschbaum, C. (1997). Free cortisol levels after awakening: a reliable biological marker for the assessment of adrenocortical activity. *Life Sciences*, 61(26), 2539-2549.
- Qin, Y., Karst, H., & Joels, M. (2004). Chronic unpredictable stress alters gene expression in rat single dentate granule cells. *Journal of Neurochemistry*, 89(2), 364-374. doi: 10.1111/j.1471-4159.2003.02332.x

- Roussos, P., Giakoumaki, S. G., Georgakopoulos, A., Robakis, N. K., & Bitsios, P. (2011). The CACNA1C and ANK3 risk alleles impact on affective personality traits and startle reactivity but not on cognition or gating in healthy males. *Bipolar Disorders*, *13*(3), 250-259.
- Roussos, P., McClure, M. M., Hazlett, E. A., New, A. S., Siever, L. J., Bitsios, P., & Giakoumaki, S. G. (2013). CACNA1C as a risk factor for schizotypal personality disorder and schizotypy in healthy individuals. *Psychiatry Research*, *206*(1), 122-123. doi: 10.1016/j.psychres.2012.08.039
- Ruhe, H. G., Khoenkhoen, S. J., Ottenhof, K. W., Koeter, M. W., Mocking, R. J., & Schene, A. H. (2015). Longitudinal effects of the SSRI paroxetine on salivary cortisol in Major Depressive Disorder. *Psychoneuroendocrinology*, *52*(Supplement C), 261-271. doi: 10.1016/j.psyneuen.2014.10.024
- Schizophrenia Working Group of the Psychiatric Genomics Consortium. (2014). Biological insights from 108 schizophrenia-associated genetic loci. *Nature*, *511*(7510), 421-427.
- Shalev, I., Lerer, E., Israel, S., Uzefovsky, F., Gritsenko, I., Mankuta, D., Ebstein, R. P., & Kaitz, M. (2009). BDNF Val66Met polymorphism is associated with HPA axis reactivity to psychological stress characterized by genotype and gender interactions. *Psychoneuroendocrinology*, *34*(3), 382-388. doi: 10.1016/j.psyneuen.2008.09.017
- Sink, T. D., Lochmann, R. T., & Fecteau, K. A. (2008). Validation, use, and disadvantages of enzyme-linked immunosorbent assay kits for detection of cortisol in channel catfish, largemouth bass, red pacu, and golden shiners. *Fish Physiology and Biochemistry*, *34*(1), 95-101. doi: 10.1007/s10695-007-9150-9
- Sklar, P., Smoller, J. W., Fan, J., Ferreira, M. A., Perlis, R. H., Chambert, K., Nimgaonkar, V. L., McQueen, M. B., Faraone, S. V., Kirby, A., de Bakker, P. I., Ogdie, M. N., Thase, M. E., Sachs, G. S., Todd-Brown, K., Gabriel, S. B., Sougnez, C., Gates, C.,

- Blumenstiel, B., Defelice, M., Ardlie, K. G., Franklin, J., Muir, W. J., McGhee, K. A., MacIntyre, D. J., McLean, A., VanBeck, M., McQuillin, A., Bass, N. J., Robinson, M., Lawrence, J., Anjorin, A., Curtis, D., Scolnick, E. M., Daly, M. J., Blackwood, D. H., Gurling, H. M., & Purcell, S. M. (2008). Whole-genome association study of bipolar disorder. *Molecular Psychiatry*, *13*(6), 558-569. doi: 10.1038/sj.mp.4002151
- Stalder, T., Kirschbaum, C., Kudielka, B. M., Adam, E. K., Pruessner, J. C., Wust, S., Dockray, S., Smyth, N., Evans, P., Hellhammer, D. H., Miller, R., Wetherell, M. A., Lupien, S. J., & Clow, A. (2016). Assessment of the cortisol awakening response: Expert consensus guidelines. *Psychoneuroendocrinology*, *63*, 414-432. doi: 10.1016/j.psyneuen.2015.10.010
- Van Gemert, N. G., Carvalho, D. M., Karst, H., Van Der Laan, S., Zhang, M., Meijer, O. C., Hell, J. W., & Joels, M. (2009). Dissociation between rat hippocampal CA1 and dentate gyrus cells in their response to corticosterone: effects on calcium channel protein and current. *Endocrinology*, *150*(10), 4615-4624.
- van Gemert, N. G., & Joels, M. (2006). Effect of chronic stress and mifepristone treatment on voltage-dependent Ca<sup>2+</sup> currents in rat hippocampal dentate gyrus. *Journal of Neuroendocrinology*, *18*(10), 732-741. doi: 10.1111/j.1365-2826.2006.01472.x
- van Winkel, R., Stefanis, N. C., & Myin-Germeys, I. (2008). Psychosocial stress and psychosis. A review of the neurobiological mechanisms and the evidence for gene-stress interaction. *Schizophrenia Bulletin*, *34*(6), 1095-1105. doi: 10.1093/schbul/sbn101
- Varese, F., Smeets, F., Drukker, M., Lieveise, R., Lataster, T., Viechtbauer, W., Read, J., van Os, J., & Bentall, R. P. (2012). Childhood Adversities Increase the Risk of Psychosis: A Meta-analysis of Patient-Control, Prospective- and Cross-sectional Cohort Studies. *Schizophrenia Bulletin*, *38*(4), 661-671. doi: 10.1093/schbul/sbs050



- Veen, G., van Vliet, I. M., DeRijk, R. H., Giltay, E. J., van Pelt, J., & Zitman, F. G. (2011). Basal cortisol levels in relation to dimensions and DSM-IV categories of depression and anxiety. *Psychiatry Research*, *185*(1-2), 121-128. doi: 10.1016/j.psychres.2009.07.013
- Vreeburg, S. A., Kruijtzter, B. P., van Pelt, J., van Dyck, R., DeRijk, R. H., Hoogendijk, W. J. G., Smit, J. H., Zitman, F. G., & Penninx, B. W. J. H. (2009). Associations between sociodemographic, sampling and health factors and various salivary cortisol indicators in a large sample without psychopathology. *Psychoneuroendocrinology*, *34*(8), 1109-1120. doi: 10.1016/j.psyneuen.2009.04.024
- Wardenaar, K. J., Vreeburg, S. A., van Veen, T., Giltay, E. J., Veen, G., Penninx, B. W., & Zitman, F. G. (2011). Dimensions of depression and anxiety and the hypothalamo-pituitary-adrenal axis. *Biological Psychiatry*, *69*(4), 366-373. doi: 10.1016/j.biopsych.2010.09.005
- West, A. E., Griffith, E. C., & Greenberg, M. E. (2002). Regulation of transcription factors by neuronal activity. *Nature Reviews: Neuroscience*, *3*(12), 921-931. doi: 10.1038/nrn987
- Wust, S., Kumsta, R., Treutlein, J., Frank, J., Entringer, S., Schulze, T. G., & Rietschel, M. (2009). Sex-specific association between the 5-HTT gene-linked polymorphic region and basal cortisol secretion. *Psychoneuroendocrinology*, *34*(7), 972-982. doi: 10.1016/j.psyneuen.2009.01.011
- Wust, S., Wolf, J., Hellhammer, D. H., Federenko, I., Schommer, N., & Kirschbaum, C. (2000). The cortisol awakening response-normal values and confounds. *Noise and Health*, *2*(7), 79.
- Zigmond, A. S., & Snaith, R. P. (1983). The hospital anxiety and depression scale. *Acta Psychiatrica Scandinavica*, *67*(6), 361-370.

ACCEPTED MANUSCRIPT

Table 1. Age, HADS and PSS-14 scores, waking time, sleep duration, awakening and +30 cortisol levels divided by ELS group (before age 17) and *CACNA1C* genotype.

Characteristic	No ELS		ELS	
	AA/AG (n=35)	GG (n=29)	AA/AG (n=21)	GG (n=18)
Age (Years)	35.23(10.84)	32.90(10.88)	35.24(10.44)	35.06(11.62)
HADS Anxiety	5.71(3.30)	5.90(3.31)	8.00(4.01)	7.28(4.30)
HADS Depression	2.71(2.14)	2.59(2.61)	3.67(2.82)	4.5(2.68)
PSS-14	18.83(8.35)	20.06(7.38)	23.10(9.16)	22.78(7.57)
Waking Time	7:29(1:11)	7:14(0:54)	7:16(0:53)	7:46(1:34)
Sleep Duration (Hours)	6.77(1.17)	6.66(1.09)	6.63(1.67)	6.63(1.43)
Awakening Cortisol (nmol/L)	7.30(4.70)	7.86(4.16)	6.15(2.10)	8.34(5.46)
+30 Cortisol (nmol/L)	8.20(5.34)	8.67(3.45)	8.24(3.52)	12.97(8.00)

*Notes.* Data shown are means and standard deviations for early life stress experienced before the age of 17. ELS = Early life stress, HADS = Hospital Anxiety and Depression Scale, PSS-14 = Perceived Stress Scale.