



**Lu, Szu-Ching and Xiu, Kaihua and Li, Ke and Marquardt, Tamara L and Evans, Peter J and Li, Zong-Ming (2017) Effects of carpal tunnel syndrome on force coordination and muscle coherence during precision pinch. Journal of Medical and Biological Engineering, 37 (3). pp. 328-335. ISSN 1609-0985 , <http://dx.doi.org/10.1007/s40846-017-0232-6>**

This version is available at <https://strathprints.strath.ac.uk/64552/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)

The Strathprints institutional repository (<https://strathprints.strath.ac.uk>) is a digital archive of University of Strathclyde research outputs. It has been developed to disseminate open access research outputs, expose data about those outputs, and enable the management and persistent access to Strathclyde's intellectual output.

# Effects of Carpal Tunnel Syndrome on Force Coordination and Muscle Coherence during Precision Pinch

Szu-Ching Lu<sup>1</sup>, Kaihua Xiu<sup>1</sup>, Ke Li<sup>4</sup>, Tamara L. Marquardt<sup>1</sup>, Peter J. Evans<sup>2</sup>, Zong-Ming Li<sup>1,2,3\*</sup>

<sup>1</sup>Department of Biomedical Engineering, Cleveland Clinic, Cleveland, Ohio, USA

<sup>2</sup>Department of Orthopaedic Surgery, Cleveland Clinic, Cleveland, Ohio, USA

<sup>3</sup>Department of Physical Medicine and Rehabilitation, Cleveland Clinic, Cleveland, Ohio, USA

<sup>4</sup>Department of Biomedical Engineering, School of Control Science and Engineering, Shandong University, Jinan, Shandong, China

## Emails:

SCL: [shinlu1984@gmail.com](mailto:shinlu1984@gmail.com)

KHX: [kaihua.xiu@gmail.com](mailto:kaihua.xiu@gmail.com)

KL: [kli@sdu.edu.cn](mailto:kli@sdu.edu.cn)

TLM: [marquat@ccf.org](mailto:marquat@ccf.org)

PJE: [evansp2@ccf.org](mailto:evansp2@ccf.org)

ZML: [liz4@ccf.org](mailto:liz4@ccf.org)

## \*Corresponding author

Zong-Ming Li

E-mail: [liz4@ccf.org](mailto:liz4@ccf.org)

Fax: 216-444-9198

## Word Count:

Abstract: 249

Main Text: 3298

## Funding Sources:

National Institutes of Health (NIAMS R01AR056964)

*For Double-Blind Peer Review*

# Effects of Carpal Tunnel Syndrome on Force Coordination and Muscle Coherence during Precision Pinch

Word Count:

Abstract: 249

Main Text: 3298

## Abstract

Carpal tunnel syndrome (CTS), caused by entrapment of the median nerve in the carpal tunnel, impairs hand function including dexterous manipulation. The purpose of this study was to investigate the effects of CTS on force coordination and muscle coherence during low-intensity sustained precision pinch while the wrist assumed different postures. Twenty subjects (10 CTS patients and 10 asymptomatic controls) participated in this study. An instrumented pinch device was used to measure the thumb and index finger forces while simultaneously collecting surface electromyographic activities of the abductor pollicis brevis (APB) and first dorsal interosseous (FDI) muscles. Subjects performed a sustained precision pinch at 10% maximum pinch force for 15 sec with the wrist stabilized at 30° extension, neutral, or 30° flexion using customized splints. The force discrepancy and the force coordination angle between the thumb and index finger forces were calculated, as well as the  $\beta$ -band (15-30 Hz) coherence between APB and FDI. The index finger applied greater force than the thumb ( $p < 0.05$ ); this force discrepancy was increased with wrist flexion ( $p < 0.05$ ), but was not affected by CTS ( $p > 0.05$ ). The directional force coordination was not significantly affected by wrist posture or CTS ( $p > 0.05$ ). In general, digit force coordination during precision pinch seems to be sensitive to wrist flexion, but is not affected by CTS. The  $\beta$ -band muscular coherence was increased by wrist flexion for CTS patients ( $p < 0.05$ ), which could be a compensatory mechanism for the flexion-induced exacerbation of CTS symptoms. In summary, this study observed the effect of wrist posture on the force discrepancy and the wrist posture  $\times$  CTS interaction effect on the  $\beta$ -band coherence during precision pinch.

**Keywords:** carpal tunnel syndrome; force coordination; muscle coherence; precision pinch

# 1 Introduction

Carpal tunnel syndrome (CTS) is a common compression neuropathy of the upper extremity, with high prevalence in the general population. Due to entrapment of the median nerve in the carpal tunnel, CTS patients experience symptoms of hand tingling, numbness, and pain. These symptoms are exacerbated in the extreme wrist flexion position (i.e. Phalen's maneuver), which is commonly used as a provocative test for diagnostic purposes [1,2]. The median nerve supplies sensory input to the palmar side of the thumb, index finger, middle finger, and the radial half of the ring finger. The motor branch of the median nerve innervates the opponens pollicis, abductor pollicis brevis (APB) and superficial head of flexor pollicis brevis, as well as first and second lumbricals. CTS is known to impair sensory functions of the hand as commonly evaluated by two-point discrimination, Semmes Weinstein monofilament testing, and sensory latency [3,4]. Motor function of the hand is also shown to be affected by CTS as demonstrated by weakness of grasp and pinch strength [5-7], although motor capability of the thumb has been found to be relatively preserved [8,9].

Precision pinch with the thumb and index finger is a dexterous manual task involving sensorimotor coordination of the two digits. CTS patients commonly demonstrate lack of dexterity in activities of daily living, such as inexplicably dropping objects. As a potential compensatory strategy to overcome sensorimotor deficits and prevent objects from unintentionally slipping, patients with CTS apply excessive pinch force while lifting objects [10-12]. CTS also impairs digit force accuracy and stability during precision pinch, especially when the force application lacks accompanying visual feedback [13].

During precision pinch, extrinsic and intrinsic hand muscles work together to ensure successful manipulation, and therefore, muscle coordination is necessary for successful pinch

performance. Coherence analysis of electromyographic (EMG) signals has been used to quantify muscle coordination, which could shed a light on the modulation of the neural inputs to the coordinated muscles [14] Specifically for pinching tasks, EMG-EMG coherence among hand and forearm muscles in the 15-30 Hz range ( $\beta$ -band) was shown to be associated with maintaining a steady force [15,16]. Impaired sensory input by digital nerve anaesthesia and deafferentation was also shown to modulate hand muscle coherence in the  $\beta$ -band [17,18]. Due to the sensorimotor deficits of the thumb and index finger associated with CTS, muscular coherence may be impaired in CTS patients completing precision pinch tasks.

Although grip and pinch strength have been extensively investigated in CTS patients, there is limited understanding of precision pinch forces and associated muscle activities. Therefore, the purpose of this study was to investigate the effects of CTS on force coordination and muscle coherence during precision pinch. For muscle coordination and coherence, the APB and the first dorsal interosseous (FDI) were chosen as they are key intrinsic hand muscles involved in pinching tasks. In addition, force and muscle coordination were examined with the wrist in different flexion/extension positions to understand the postural effects on hand function. We hypothesized that the CTS patients would present less effective force coordination and stronger coupling between the APB and FDI muscles than the controls.

## **2 Materials and Methods**

### **2.1 Participants**

A total of 20 right-hand dominant participants were recruited for this study, including 10 CTS patients ( $50.8 \pm 9.6$  years old; 8 females and 2 males) and 10 asymptomatic controls ( $47.9 \pm$

13.3 years old; 8 females and 2 males). All participants provided informed consent prior to study participation in accordance with the Institutional Review Board at Cleveland Clinic.

The inclusion criteria for the CTS group included satisfying at least three of the following criteria: (1) history of pain and/or numbness in the median-innervated territory of the right hand for at least 3 months; (2) positive provocative maneuvers with Tinel's sign, Phalen's maneuver, and/or median nerve compression test; (3) abnormal electrodiagnostic test results demonstrating median nerve neuropathy in the right hand; (4) an overall Boston Carpal Tunnel Syndrome Questionnaire score greater than 1.5 [19]; and (5) confirmation of CTS according to clinical discretion [20]. For the control group, the inclusion criteria included absence of CTS-like symptoms. The exclusion criteria for the CTS and control groups were: (1) left-hand dominance; (2) existence of any central nervous system disease; (3) diabetes; (4) pregnancy; (5) arthritis in the right hand or wrist; (6) steroid injection to the right hand within three months of study participation; and (7) history of musculoskeletal injury or surgery to the right hand or wrist.

## **2.2 Experimental Set-up**

A pinch apparatus consisting of two six-component force/torque transducers (Mini40, ATI Industrial Automation, Inc., Apex, NC, USA) was used to measure the thumb and index finger forces. Each transducer was attached to an aluminum mounting support which was rigidly fixed to a stainless steel plate. The pinch contact surfaces were covered with 100-grit sandpaper and the pinch span was 1.8 cm. The force/torque signals were amplified and multiplexed using a custom interface box (ATI Industrial Automation, Inc., Apex, NC, USA) and converged to an 18-bit analog digital converter (PXI-6289, National Instruments, Austin, TX, USA). A surface EMG system (MyoSystem 1400, Noraxon USA, Inc., Scottsdale, AZ, USA) was used to record

the activity of the APB and FDI muscles. The system had a 12-bit resolution and a hardware band pass filter of 10-500 Hz. In addition, a 22-inch computer monitor was positioned 50 cm in front of the participant to graphically provide real-time force information.

## 2.3 Experimental Protocol

Participants washed their hands with soap and water prior to the experiment. In addition, the skin on the right hand was prepared using sandpaper and an alcohol swab before the application of EMG electrodes. A dual Ag/AgCl electrode with a center-to-center distance of 2.75 cm (Noraxon U.S.A., Inc., Scottsdale, AZ) was attached to the skin surface above both the APB and FDI muscles according to literature recommendations [21]. A ground electrode was also attached to the styloid process of radius. Then, each participant was seated comfortably on a height-adjustable chair by the testing table with their right arm abducted 30° in the frontal plane and flexed 30° in the sagittal plane. The forearm was rested on the table with the elbow flexed 90°. Customized splints were worn by the subject to stabilize the wrist in postures of 30° extension, anatomical neutral (0°), and 30° flexion. The pinch apparatus was fixed to the testing table at predetermined orientations so that the angle between the palm and pinch contact surfaces were 60°, allowing participants to perform pinching comfortably at each wrist posture (Fig. 1). The participants were instructed not to place the long, ring and little fingers against the pinch device nor the index finger.

[Figure 1]

**Fig. 1 Experimental setup for measuring digit forces and muscle activities during precision pinch at various wrist postures**



The study consisted of two tasks. First, the participants were instructed to pinch the apparatus using the thumb and index finger with their maximum effort. Verbal encouragement was given, guiding participants to reach their maximum pinch force within 5 sec. Three maximum pinch trials were performed for each wrist posture. For each trial, the maximum pinch force was defined as the maximum value of the averaged thumb and index finger normal forces. Then, the three maximum pinch force values for each posture were averaged and 10% of the average value was set as the target force for the subsequent submaximal pinch trials. During the submaximal pinch task, graphical information of the real-time pinch force, including a target line, was provided on the computer monitor. Participants were asked to match their pinch force to the target line as accurately as possible. Each submaximal pinch trial was 15 sec in duration and the participants were encouraged to match the target as soon as possible. A total of 10 trials were performed for each wrist posture. The three wrist postures were randomized, and a 1-minute rest was given between consecutive trials. A customized *LabVIEW* (National Instruments, Austin, TX, USA) program collected the force and EMG data at a sampling rate of 1000 Hz.

## **2.4 Data Processing**

For the submaximal task, the force and EMG data in the first 5 sec of each trial were excluded from analyses to avoid the effects of non-stationarity. The remaining 10 sec of data for each trial were analyzed using customized *MATLAB* (The MathWorks, Natick, MA, USA) programs.

To quantify the force matching accuracy, the root mean square error (RMSE) between the measured force and the target force [13] was calculated as:

$$RMSE = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - x_t)^2} \quad (1)$$

184 where  $n$  is the number of force samples,  $x_i$  is the instant mean of the thumb and index finger  
 185 normal forces, and  $x_t$  is the target force (i.e. 10% of the averaged maximum pinch force). The  
 186 force discrepancy between the two digits was calculated as the percentage difference between the  
 187 resultant force magnitudes of the index finger and the thumb normalized by the force magnitude  
 188 of the thumb. The 3D force vectors of the thumb and index finger were transformed to a common  
 189 coordinate system [22,23], and then the angle between the two force vectors was calculated using  
 190 the following equation:

$$\theta = \cos^{-1} \frac{\vec{F}_1 \cdot \vec{F}_2}{|\vec{F}_1| \cdot |\vec{F}_2|} \quad (2)$$

191 where  $\vec{F}_1$  and  $\vec{F}_2$  are the force vectors of the thumb and index finger, respectively. The angle  
 192 between the thumb and index finger was defined as the force coordination angle, which ranges  
 193 from 0 to 180°, where 0° means the two force vectors are in phase and 180° means that they are  
 194 in opposite directions. The coordination angle between the digits was averaged over the 10-sec  
 195 period.

196 To calculate the coherence within the  $\beta$ -band between APB and FDI, first the EMG data  
 197 were filtered and rectified using a 4<sup>th</sup> order band-pass (5-100 Hz) Butterworth filter. Then, the  
 198 signals were further processed using a bivariate autoregressive model and a boxcar window to  
 199 generate coefficients for coherence estimation [24]. The coherence between two EMG signals  
 200 was calculated as:

$$C_{xy}(f) = \frac{S_{xy}(f)}{\sqrt{S_{xx}(f)S_{yy}(f)}} \quad (3)$$

where  $f$  is a given frequency,  $S_{xy}$  is the cross spectrum of associated signals, and  $S_{xx}$  and  $S_{yy}$  are the auto spectra of the associated signals. The estimated coherence was then transformed using a Fisher Z-transformation. The mean coherence between APB and FDI in the  $\beta$ -band was calculated. The coherence value ranges from 0 to 1, where higher value means greater coupling between the muscles.

## 2.5 Statistical Analysis

Two-way repeated measures ANOVAs, with one factor repeated, were performed to test the main effects of group (CTS and control) and posture (extension, neutral, and flexion), as well as the interaction effect of group and posture. The effects on the maximum pinch force, force matching accuracy, force discrepancy, force coordination angle, and mean  $\beta$ -band coherence were examined. Post hoc Tukey's tests were completed for pairwise comparisons. Statistical analyses were performed using *SigmaStat 3.5* (Systat Software, San Jose, CA) and the significance level of  $\alpha = 0.05$ .

## 3 Results

The demographic data of the 10 participants in the CTS group are shown in Table 1. Their CTS symptom duration ranged from 10 months to 21 years. The Boston CTS Questionnaire scores of the patients ranged from 1.58 to 3.58. Eight of them received the provocative maneuver tests, and they presented positive results with Tinel's sign or Phalen's Test.

**Table 1 The age, gender, symptom duration, Boston CTS Questionnaire score, and the result of the provocative maneuver tests of 10 participants in the CTS group**

Subject #	Age (y/o)	Gender	CTS Symptom Duration	Boston CTS Questionnaire Score	Tinel's Sign / Phalen's Test
CTS 01	64	Female	5 years	2.21	Positive
CTS 02	56	Female	16 years	3.11	Positive
CTS 03	55	Female	1.5 years	2.89	Positive
CTS 04	46	Female	21 years	2.79	N/A
CTS 05	28	Male	10 months	2.16	N/A
CTS 06	58	Male	1 year	2.47	Positive
CTS 07	50	Female	1 year	2.00	Positive
CTS 08	50	Female	6 years	3.58	Positive
CTS 09	53	Female	1.5 years	1.58	Positive
CTS 10	48	Female	1.5 years	2.63	Positive

The maximum pinch force and force matching accuracy are presented in Table 2. The maximum pinch force was significantly affected by the factor of wrist posture ( $p < 0.001$ ), but not by the group factor ( $p = 0.514$ ) or the posture  $\times$  group interaction ( $p = 0.077$ ). The maximum pinch force in the flexed wrist posture was significantly less than that at the neutral ( $p = 0.002$ ) and extended ( $p = 0.003$ ) postures. The force matching accuracy for submaximal pinching was not significantly affected by group ( $p = 0.436$ ), posture ( $p = 0.956$ ), or the posture  $\times$  group interaction ( $p = 0.199$ ).

**Table 2 Maximum pinch force and submaximal force accuracy for the control and CTS groups at different wrist postures (mean  $\pm$  standard deviation)**

	Control (n = 10)		CTS (n = 10)	
	Maximum (N)	Accuracy (N)	Maximum (N)	Accuracy (N)
Extension	37.9 ± 14.5	0.12 ± 0.06	39.4 ± 11.5	0.13 ± 0.09
Neutral	37.0 ± 14.1	0.10 ± 0.04	40.7 ± 9.1	0.15 ± 0.11
Flexion	32.9 ± 13.0	0.11 ± 0.05	38.3 ± 8.1	0.13 ± 0.08

For the submaximal pinch task, there existed a force discrepancy between the digits with the index finger resultant force being greater than that of the thumb for both the control and CTS groups ( $p < 0.05$ ). This force discrepancy was significantly affected by posture ( $p < 0.05$ ), but not by group ( $p = 0.916$ ). Wrist flexion led to an increased force discrepancy between the digits. In the wrist extension posture, the force discrepancies were  $19.0 \pm 23.5\%$  for the control group and  $20.0 \pm 29.5\%$  for the CTS group. At the neutral posture, the discrepancies were  $24.4 \pm 29.0\%$  and  $23.4 \pm 34.2\%$  for the control and CTS groups, respectively. As the wrist deviated to a more flexed posture of  $30^\circ$ , the force discrepancy for the control group was  $28.2 \pm 32.7\%$  and for the CTS group was  $23.7 \pm 36.4\%$ .

The force coordination angle between the digits ranged from  $157^\circ$  to  $164^\circ$  (Fig. 2). The angle was not significantly affected by group ( $p = 0.096$ ), posture ( $p = 0.191$ ), or the group  $\times$  posture interaction ( $p = 0.828$ ). The average coordination angle across the three wrist postures was  $158.2 \pm 5.9^\circ$  for the control group and  $163.0 \pm 7.3^\circ$  for the CTS group.

### [Figure 2]

**Fig. 2 Force coordination angles (mean  $\pm$  standard deviation) at difference wrist postures for the CTS and control groups**

The coherence in the  $\beta$ -band was not significantly affected by group ( $p = 0.684$ ) or posture ( $p = 0.269$ ); however, there was a significant group  $\times$  posture interaction ( $p < 0.05$ ). The coherence values for the control group were  $0.153 \pm 0.041$ ,  $0.158 \pm 0.045$ , and  $0.152 \pm 0.035$  at wrist extension, neutral, and flexion, respectively. For the CTS group, the coherence values were  $0.144 \pm 0.018$  for extension,  $0.144 \pm 0.019$  for neutral, and  $0.158 \pm 0.03$  for flexion. The interaction effect was associated with a significant difference of pairwise comparisons within the CTS group (Fig. 3).

[Figure 3 here]

**Fig. 3  $\beta$ -band coherence at different wrist postures for CTS and control groups**

## **4 Discussion**

Our finding of decreased maximal pinch force at a flexed wrist posture is consistent with previous reports that pinch/grip strength is sensitive to wrist posture, particularly in the direction of flexion/extension with weak strength in flexion [25-29]. The weakness can be explained by the length-tension relationship of the extrinsic flexors of the digits. The muscular compartments are at an optimal length for maximum active force production at the slightly extended, functional wrist position. As the wrist joint flexes, the associated muscular compartment becomes less optimal, leading to an impairment of pinch force production. The result that CTS patients had similar maximal pinch force to healthy controls corroborates with previous findings that motor capability is relatively preserved in the CTS patient population [8,9].

For the sustained submaximal pinch, the CTS-associated sensory and motor deficits were expected to increase the force matching error. However, no statistical difference in the force matching accuracy was observed between the two groups or among wrist postures. In the current study, the precision pinch was performed on a stabilized object and visual feedback of force application was provided. It is possible that the sensorimotor deficits associated with CTS were compensated by the visual information to generate accurate pinch force [13]. In addition, 10% of the maximum pinch force was set as the target force in this study, which could be too low to see the effects of CTS or wrist posture. However, higher pinch force may induce muscle fatigue and affect the results of force matching.

The result that the index finger generated more force than the thumb during submaximal precision pinch is consistent with a previous study [13]. Force discrepancy has been postulated as a consequence of anatomical structure and neural control. In the current study, no group effect was found on the inter-digit force difference, but there was a wrist posture effect. Wrist flexion caused increased force discrepancy between the thumb and index finger. The increase in the index finger contribution may be due to the more diminished thumb flexor function in a flexed wrist position [30].

The directional coordination of the precision pinch forces was not influenced by CTS or wrist posture. Biomechanically, the thenar muscles tend to abduct the thumb and generate shear force, making the thumb force vector deviated from the opposing direction perpendicular to the pinching surface. Alteration of thenar muscle function implicated by median nerve dysfunction could change directional force coordination. The fact that coordination angle was not changed by CTS might be explained by (a) the task required relatively low exertion effort and (b) the preservation of motor capability in the patient group. It is possible that pinch force

dyscoordination will be more salient in patients with more severe CTS, especially during maximal pinching force production. The coordination angle, regardless of wrist posture and subject group, deviated about 20° from perfect opposition of 180°, which is consistent with previous findings [23,22]. The individual digits did not apply forces perpendicular to the pinch surface, nor did they orient their forces parallel in space. For pinching on a stabilized object without the requirement of force equilibrium, each digit may apply a force that favors its own anatomical structure and independent neural control.

We found that CTS patients demonstrated different  $\beta$ -band coherence of the APB and FDI muscles in flexion from that in extension. Coherence of hand muscles was shown in the  $\beta$ -band while maintaining steady pinch force [16], and increased coherence was observed when manipulating more compliant objects because of the increased sensorimotor integration required to adjust digit force and position for compliant object control [15]. It has also been shown that muscle coherence is affected by impaired sensory inputs due to deafferentation [18] and digital nerve anaesthesia [17]. In addition to sensory loss, the changes in somatosensory feedback due to varied wrist posture impacts muscle coherence across intrinsic and extrinsic muscles [31]. Furthermore, extrinsic muscles were found to have a stronger coherence than the intrinsic muscles [32,33]. The lower coherence has been interpreted as a benefit for intrinsic muscle function [34]. Intrinsic muscles are specifically important for fine modulation of digit forces, for this reason, a less degree of coupling might be better for individual muscle control. In the current study, participants performed an isometric pinch on a stabilized object, thus the independence of muscle activation may not be required for this task. Although varied wrist posture does not change the length of the APB and FDI muscles, wrist flexion may exacerbate the symptoms of CTS. In contrast to the reduced coherence after digital nerve block [17], we observed that the



CTS patients had increased coherence in wrist flexion. Compared to the acute sensory loss after digital nerve anaesthesia, patients with CTS experience altered sensation in a longer process. Different modulations in the muscle coherence may reflect the different natures of sensory modifications. It could be postulated that the increase in  $\beta$ -band coherence in CTS patients with flexed wrists is a compensatory mechanism for the flexion-induced exacerbation of symptoms.

Some limitations in this study should be considered. First, moderate changes in wrist posture were applied. Slight wrist extension ( $10\text{-}30^\circ$ ) has been widely used for functional positioning [35,36] and no significant change was found in the carpal tunnel pressure between the neutral wrist position and  $30^\circ$  extension [37]. These may explain the findings in the current study that the hand has similar functionality at the neutral position and  $30^\circ$  extension. Although hand function is more sensitive to wrist flexion, the  $30^\circ$  wrist flexion posture employed in the current study was relatively moderate in comparison to the clinically used wrist flexion for Phalen's maneuver. Perhaps a more flexed wrist position would reveal greater differences between CTS patients and controls. Second, this study had a relatively small sample size and the CTS patients were not classified by severity of symptoms. Though most CTS patients have sufficient thenar muscle function, the wasting or weakness of thenar muscles could be demonstrated in patients with more severe symptoms [38]. The force coordination between digits and the muscle coherence between APB and FDI could be more affected in CTS cases with thenar muscle atrophy. At last, a stabilized, non-deformable pinch device was used in the current study for the investigation of force coordination and muscle coherence. However, translating and manipulating movable objects are the skills often required in daily life. As greater muscle coherence was found in handling a more compliant object [15], neuromuscular coordination could be affected by the compliance of objects. Also, the disturbance of gravity when holding

objects in the air could be another factor to influence force and muscle coordination. Therefore, different task designs and object materials may lead to varied force coordination and muscle coherence. In addition, muscle fatigue may also affect force coordination and muscle coherence. However, the target was set as 10% of the maximum pinch force and a 1-minute rest was given between trials, the effect of muscle fatigue was assumed to be limited in this study.

In conclusion, this study observed the wrist posture effect on the force discrepancy that the index finger generated significantly higher force than the thumb when the wrist was flexed. Also, the wrist posture  $\times$  group interaction effect was observed that, within the CTS group, the  $\beta$ -band coherence was significantly higher in the wrist flexion condition than in the wrist extension condition.

## 5 Acknowledgments

This publication was made possible by Grant R01AR056964 from NIAMS/NIH. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIAMS or NIH.

## 6 References

1. MacDermid, J. C., & Wessel, J. (2004). Clinical diagnosis of carpal tunnel syndrome: a systematic review. *J Hand Ther*, 17(2), 309-319, doi:10.1197/j.jht.2004.02.015.
2. Wiesman, I. M., Novak, C. B., Mackinnon, S. E., & Winograd, J. M. (2003). Sensitivity and specificity of clinical testing for carpal tunnel syndrome. *Can J Plast Surg*, 11(2), 70-72.
3. Havton, L. A., Hotson, J. R., & Kellerth, J. O. (2007). Correlation of median forearm conduction velocity with carpal tunnel syndrome severity. *Clin Neurophysiol*, 118(4), 781-785, doi:10.1016/j.clinph.2006.12.011.
4. Szabo, R. M., Gelberman, R. H., & Dimick, M. P. (1984). Sensibility testing in patients with carpal tunnel syndrome. *J Bone Joint Surg Am*, 66(1), 60-64.
5. Baker, N. A., Moehling, K. K., Desai, A. R., & Gustafson, N. P. (2013). Effect of carpal tunnel syndrome on grip and pinch strength compared with sex- and age-matched normative data. *Arthritis Care Res (Hoboken)*, 65(12), 2041-2045, doi:10.1002/acr.22089.

6. Gehrmann, S., Tang, J., Kaufmann, R. A., Goitz, R. J., Windolf, J., & Li, Z. M. (2008). Variability of precision pinch movements caused by carpal tunnel syndrome. *J Hand Surg Am*, 33(7), 1069-1075, doi:10.1016/j.jhsa.2008.02.030.
7. Tamburin, S., Cacciatori, C., Marani, S., & Zanette, G. (2008). Pain and motor function in carpal tunnel syndrome: a clinical, neurophysiological and psychophysical study. *J Neurol*, 255(11), 1636-1643, doi:10.1007/s00415-008-0895-6.
8. Agabegi, S. S., Freiberg, R. A., Plunkett, J. M., & Stern, P. J. (2007). Thumb abduction strength measurement in carpal tunnel syndrome. *J Hand Surg Am*, 32(6), 859-866, doi:10.1016/j.jhsa.2007.04.007.
9. Li, Z. M., Harkness, D. A., & Goitz, R. J. (2005). Thumb strength affected by carpal tunnel syndrome. *Clin Orthop Relat Res*, 441, 320-326.
10. Hsu, H. Y., Kuo, L. C., Kuo, Y. L., Chiu, H. Y., Jou, I. M., Wu, P. T., et al. (2013). Feasibility of a novel functional sensibility test as an assisted examination for determining precision pinch performance in patients with carpal tunnel syndrome. *PLoS One*, 8(8), e72064, doi:10.1371/journal.pone.0072064.
11. Lowe, B. D., & Freivalds, A. (1999). Effect of carpal tunnel syndrome on grip force coordination on hand tools. *Ergonomics*, 42(4), 550-564, doi:10.1080/001401399185469.
12. Yen, W. J., Kuo, Y. L., Kuo, L. C., Chen, S. M., Kuan, T. S., & Hsu, H. Y. (2014). Precision pinch performance in patients with sensory deficits of the median nerve at the carpal tunnel. *Motor Control*, 18(1), 29-43, doi:10.1123/mc.2013-0004.
13. Li, K., Evans, P. J., Seitz, W. H., Jr., & Li, Z. M. (2015). Carpal tunnel syndrome impairs sustained precision pinch performance. *Clin Neurophysiol*, 126(1), 194-201, doi:10.1016/j.clinph.2014.05.004.
14. Baker, S. N., Olivier, E., & Lemon, R. N. (1997). Coherent oscillations in monkey motor cortex and hand muscle EMG show task-dependent modulation. *J Physiol*, 501 ( Pt 1), 225-241.
15. Kilner, J. M., Baker, S. N., Salenius, S., Hari, R., & Lemon, R. N. (2000). Human cortical muscle coherence is directly related to specific motor parameters. *J Neurosci*, 20(23), 8838-8845.
16. Kilner, J. M., Baker, S. N., Salenius, S., Jousmaki, V., Hari, R., & Lemon, R. N. (1999). Task-dependent modulation of 15-30 Hz coherence between rectified EMGs from human hand and forearm muscles. *J Physiol*, 516 ( Pt 2), 559-570.
17. Fisher, R. J., Galea, M. P., Brown, P., & Lemon, R. N. (2002). Digital nerve anaesthesia decreases EMG-EMG coherence in a human precision grip task. *Exp Brain Res*, 145(2), 207-214, doi:10.1007/s00221-002-1113-x.
18. Kilner, J. M., Fisher, R. J., & Lemon, R. N. (2004). Coupling of oscillatory activity between muscles is strikingly reduced in a deafferented subject compared with normal controls. *J Neurophysiol*, 92(2), 790-796, doi:10.1152/jn.01247.2003.
19. Levine, D. W., Simmons, B. P., Koris, M. J., Daltroy, L. H., Hohl, G. G., Fossel, A. H., et al. (1993). A self-administered questionnaire for the assessment of severity of symptoms and functional status in carpal tunnel syndrome. *J Bone Joint Surg Am*, 75(11), 1585-1592.
20. Keith, M. W., Masear, V., Chung, K. C., Maupin, K., Andary, M., Amadio, P. C., et al. (2009). American Academy of Orthopaedic Surgeons Clinical Practice Guideline on diagnosis of carpal tunnel syndrome. *J Bone Joint Surg Am*, 91(10), 2478-2479, doi:10.2106/jbjs.i.00643.

21. Basmajian, J. V., & Blumenstein, R. (1980). *Electrode placement in EMG biofeedback*: Williams & Wilkins.
22. Marquardt, T. L., & Li, Z. M. (2013). Quantifying Digit Force Vector Coordination during Precision Pinch. *J Mech Med Biol*, 13(2), 1350047, doi:10.1142/s0219519413500474.
23. Li, K., Nataraj, R., Marquardt, T. L., & Li, Z. M. (2013). Directional coordination of thumb and finger forces during precision pinch. *PLoS One*, 8(11), e79400, doi:10.1371/journal.pone.0079400.
24. Pasluosta, C. F., Domalain, M. M., Fang, Y., Yue, G. H., & Li, Z. M. (2013). Influence of nerve supply on hand electromyography coherence during a three-digit task. *J Electromyogr Kinesiol*, 23(3), 594-599, doi:10.1016/j.jelekin.2013.01.006.
25. Imrhan, S. N. (1991). The influence of wrist position on different types of pinch strength. *Appl Ergon*, 22(6), 379-384.
26. Lamoreaux, L., & Hoffer, M. M. (1995). The effect of wrist deviation on grip and pinch strength. *Clin Orthop Relat Res*(314), 152-155.
27. Mathur, K., Pynsent, P. B., Vohra, S. B., Thomas, B., & Deshmukh, S. C. (2004). Effect of wrist position on power grip and key pinch strength following carpal tunnel decompression. *Journal of hand surgery*, 29(4), 390-392, doi:10.1016/j.jhsb.2004.02.012.
28. O'Driscoll, S. W., Horii, E., Ness, R., Cahalan, T. D., Richards, R. R., & An, K. N. (1992). The relationship between wrist position, grasp size, and grip strength. *J Hand Surg Am*, 17(1), 169-177.
29. Li, Z. M. (2002). The influence of wrist position on individual finger forces during forceful grip. *J Hand Surg Am*, 27(5), 886-896.
30. Harvey, L., Herbert, R. D., & Stadler, M. (2010). Effect of wrist position on thumb flexor and adductor torques in paralysed hands of people with tetraplegia. *Clin Biomech (Bristol, Avon)*, 25(3), 194-198, doi:10.1016/j.clinbiomech.2009.11.010.
31. Jesunathadas, M., Laitano, J., Hamm, T. M., & Santello, M. (2013). Across-muscle coherence is modulated as a function of wrist posture during two-digit grasping. *Neurosci Lett*, 553, 68-71, doi:10.1016/j.neulet.2013.08.014.
32. Johnston, J. A., Winges, S. A., & Santello, M. (2009). Neural control of hand muscles during prehension. *Adv Exp Med Biol*, 629, 577-596, doi:10.1007/978-0-387-77064-2\_31.
33. Poston, B., Danna-Dos Santos, A., Jesunathadas, M., Hamm, T. M., & Santello, M. (2010). Force-independent distribution of correlated neural inputs to hand muscles during three-digit grasping. *J Neurophysiol*, 104(2), 1141-1154, doi:10.1152/jn.00185.2010.
34. Winges, S. A., Kornatz, K. W., & Santello, M. (2008). Common input to motor units of intrinsic and extrinsic hand muscles during two-digit object hold. *J Neurophysiol*, 99(3), 1119-1126, doi:10.1152/jn.01059.2007.
35. Lannin, N. A., Horsley, S. A., Herbert, R., McCluskey, A., & Cusick, A. (2003). Splinting the hand in the functional position after brain impairment: a randomized, controlled trial. *Arch Phys Med Rehabil*, 84(2), 297-302, doi:10.1053/apmr.2003.50031.
36. Pizzi, A., Carlucci, G., Falsini, C., Verdesca, S., & Grippo, A. (2005). Application of a volar static splint in poststroke spasticity of the upper limb. *Arch Phys Med Rehabil*, 86(9), 1855-1859, doi:10.1016/j.apmr.2005.03.032.
37. Coppieters, M. W., Schmid, A. B., Kubler, P. A., & Hodges, P. W. (2012). Description, reliability and validity of a novel method to measure carpal tunnel pressure in patients with carpal tunnel syndrome. *Man Ther*, 17(6), 589-592, doi:10.1016/j.math.2012.03.005.

461 38. Ebata, T., Imai, K., Tokunaga, S., Takahasi, Y., & Abe, Y. (2014). Thumb opposition in  
462 severe carpal tunnel syndrome with undetectable APB-CMAP. *Hand Surg*, 19(2), 199-  
463 204, doi:10.1142/s0218810414500208.

464

465