

## RESEARCH ARTICLE

## Open Access



# Cancer-related health behaviours of young people not in education, employment or training ('NEET'): a cross-sectional study

Catherine H. Stewart<sup>1\*</sup>, Philip Berry<sup>2</sup>, Dunja Przulj<sup>3</sup> and Charlene Treanor<sup>4</sup>

## Abstract

**Background:** Links between participating in unhealthy behaviours, e.g. smoking, and an increased risk of developing some cancers are well established. Unemployed adults are more likely to participate in cancer-related health behaviours than their employed counterparts. However, evidence of whether this is true in young adults not in education, employment or training (NEET) compared to their 'non-NEET' peers is either limited or inconclusive. Using cross-sectional health data from across the UK, this study aims to investigate whether participation in cancer-related health behaviours varies by NEET status.

**Methods:** Data for 16–24 year olds were extracted from the 2010–12 Health Surveys for England (HSE) and Scottish Health Surveys (SHeS). Information on economic activity in the last week was used to determine NEET status. Data on whether respondents had been seeking employment within the last four weeks and availability to start within the next two weeks allowed NEETs to be further identified as unemployed (UE) or economically inactive (EI). Logistic regression modelled the effect of being NEET on odds of being a current smoker; heavy drinker; not participating in sport; having eaten less than five portions of fruit or vegetables the day before survey interview and having an unhealthy body mass index (BMI). Analyses were performed before and after exclusion of EI NEETs.

**Results:** Data were extracted for 4272 individuals, of which 715 (17%) were defined as NEET with 371 (52%) and 342 (48%) further classified as UE and EI respectively. Two NEETs could not be further defined as UE or EI due to missing information. Relative to non-NEETs, NEETs were significantly more likely to be current smokers, not participate in sport and have an 'unhealthy' BMI. These results held after adjustment for socio-demographic characteristics both before and after exclusion of EI NEETs. Before exclusion of EI NEETs, NEETs were significantly less likely to be heavy drinkers than non-NEETs. There was no significant difference in likelihood of heavy drinking between NEETs and non-NEETs when excluding EI NEETs.

**Conclusions:** NEETs were generally at an increased risk of participating in cancer-related health behaviours than non-NEETs. As the likelihood of becoming NEET is greater in socioeconomically-disadvantaged groups, interventions to discourage unhealthy behaviours in NEETs may contribute to a reduction in health inequalities.

**Keywords:** NEET, Cancer, Health behaviours, Young adults, Unemployed, Smoking, Alcohol, Exercise, BMI, Lifestyle

\* Correspondence: [catherine.stewart@glasgow.ac.uk](mailto:catherine.stewart@glasgow.ac.uk)

<sup>1</sup>MRC/CSO Social & Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB, UK

Full list of author information is available at the end of the article



© The Author(s). 2017 **Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

## Background

Young people are defined as 'NEET' if they are aged 16–24 years old and Not In Education, Employment or Training (NEET) [1]. In the second quarter of 2016 there were an estimated 865,000 'NEETs' in the United Kingdom (UK) [2]. Research has demonstrated both medium and long-term economic effects of becoming NEET at time of school-leaving with such individuals being more likely to still be unemployed up to five years later as well as being at an increased risk of being unemployed or in a low-paid job up to ten years later [3]. NEETs who do eventually find employment are more likely to face a lifetime of poorer income [4], lower social class [5] and lower levels of job satisfaction [6].

However, consequences of being NEET are not restricted to poorer economic outcomes. Unemployment at younger ages has been demonstrated to have immediate adverse effects on health including increased rates of poorer mental wellbeing [7], depression [8] and suicidal behaviours [9] amongst those who are NEET. Moreover, limited research has also shown some negative effects of unemployment at younger ages on long-term health. Functional somatic symptoms [10], chronic limiting illness [5] and psychological symptoms [11] in adulthood are all reported to be consequences of youth unemployment.

The association between unemployment and poor health can be explained somewhat by increased participation in unhealthy behaviours, such as smoking and drinking alcohol amongst unemployed individuals [12, 13]. However, the evidence on whether participation in unhealthy behaviours among NEETs is greater than among their 'non-NEET' peers is either limited or inconclusive [14]. Whilst some studies have reported significant associations between NEET status and smoking [7], Baggio et al. [15] found that although smoking was likely to increase the risk of becoming NEET, the pathway from NEET status to tobacco use was not significant. Similarly, significant associations between being NEET and increased drinking or alcohol abuse/dependence have been found in some studies [7, 9], but not in others [14]. Additionally, the correlation between unemployment and increased alcohol consumption found by Janlert and Hammarström [16] only applied to longer periods of unemployment. There have also been reports of lower levels of involvement in sport or exercise amongst NEETs [17].

Participation in unhealthy behaviours has been linked to an increased risk of developing a range of cancers [18, 19]. In 2012, the most common cancers in Europe, representing half of the overall burden of cancer, were breast, colorectal, prostate and lung cancer [20]. Previous research has attributed some of the risk of developing each of these four cancers to participation in unhealthy behaviours including smoking (lung) [21], alcohol consumption (breast) [22], low fruit and vegetable intake (lung, colorectal) [23],

physical inactivity (breast, colorectal, prostate) [24] and excess body weight (colorectal) [25]. Given the association between unemployment and increased participation in unhealthy behaviours and well-established links between participating in such behaviours and cancer, NEETs may be at an increased risk of cancer. Although links between unemployment and cancer have been shown to exist [26, 27], studies have either tended to focus on unemployment in middle age or use cohorts spanning a wide range of ages. Studies focusing primarily on unemployment in early adulthood as a risk factor for cancer and which also cover the other dimensions included in the NEET definition, i.e. not in education or training, are lacking. This study aims to develop such an evidence base by investigating whether NEETs have higher rates of participation in cancer-related health behaviours compared to non-NEETs.

## Methods

### Aims of the study

Using cross-sectional health survey data for samples of 16–24 year olds, the aims of this study were: (i) to compare socio-demographic and mental and physical health-related characteristics of NEETs and non-NEETs; (ii) to investigate whether participation in cancer-related health behaviours were greater amongst NEETs and; (iii) whether any association between NEET status and such health behaviours persisted even after adjustment for socio-demographic and mental and physical health-related factors.

### Design & setting of the study

Data for all 16–24 year olds who participated in the Scottish Health Survey (SHeS) and Health Survey for England (HSE) over the years 2010–2012 were downloaded from the UK Data Service [28–33]. The SHeS and HSE were designed to provide nationally-representative samples of adults (aged 16 years and over) and children (aged 0–15 years) in the general population living in private households in Scotland and England. Both were based on a two-stage stratified random sample design. Postcode sectors in each constituent country were ordered by region (Health Board in Scotland and Local Authority in England) and deprivation. The first stage of the design involved creating a sample of randomly-selected postcode sectors. At the second stage, a sample of addresses was randomly drawn from each selected postcode sector based on the Postcode Address File (PAF). All adults and up to two children at each address were eligible for inclusion in the survey. If there were more than two children within a household, then two were randomly selected for inclusion [34–37]. The health surveys were chosen as they contained data on a wide range of socio-demographic variables, including economic destination of respondents, as well as information on cancer-related health behaviours. Using data from Scotland and England provided a more representative

view of NEETs across the UK and allowed for testing of independent effects of each constituent country on health outcomes.

### Health behaviour outcomes

Binary indicator variables (yes/no) were created to reflect the following cancer-related health behaviours: current smoker; heavy drinker (defined as >14 units of alcohol per week for females and >21 units for males); participation in sport; <5 portions of fruit/vegetables the day before survey interview and unhealthy BMI. Unhealthy BMI referred to 'underweight' (BMI <18.5 kg/m<sup>2</sup>), overweight (BMI 25–29.99 kg/m<sup>2</sup>) or obese (BMI ≥30 kg/m<sup>2</sup>).

### NEET status

Survey data on economic activity in the last week were used to create a NEET indicator variable. Respondents were defined as 'NEET' if activity included the following: unpaid work for their own or a relative's business; waiting to take up paid work; looking for paid work or a government training scheme; intending to seek work, but temporarily sick or injured; permanently unable to work or looking after home or family. The UK Government further classifies NEETs into unemployed or economically inactive [1]. NEETs are defined as unemployed (UE) if they have been actively seeking work within the last 4 weeks and would be available to start work within the next two weeks [1]. Otherwise, NEETs are defined as economically inactive (EI) if they have not been seeking work within the last 4 weeks and/or would not be able to start work within the next two weeks [1]. The EI definition captures long-term sick/disabled individuals or individuals looking after family/children. Survey data contained information on whether respondents had been seeking employment within the last 4 weeks and whether they would be available to start within the next 2 weeks, thus allowing a variable to be created to further identify NEETs as UE or EI.

### Socio-demographic & health-related characteristics

Information obtained from survey data included sex, age, ethnicity, marital status, car/van access, top academic qualification, housing tenure, receipt of means-tested benefits, total annual household income and a measure of socio-economic position using the National Statistics Socio-economic classification (NS-SEC) [38]. Measures of physical and mental health status were also available including limiting long-term illness, self-assessed general health and non-psychiatric morbidity assessed using the 12-item General Health Questionnaire (GHQ-12) [39]. Higher scores on the GHQ-12 indicate a greater likelihood of probable psychiatric morbidity. Finally, variables were created to indicate year of survey interview and country of survey to investigate

whether there was any change in likelihood of participation in cancer-related health behaviours in young people over time and if there were differences between Scotland and England.

### Statistical analysis

Socio-demographic characteristics of NEETs and non-NEETs were compared by regressing NEET status on each of the socio-demographic characteristics in a univariate logistic regression model. Logistic regression was used to model the effect of being NEET on the odds of being a current smoker; heavy drinker; not participating in sport; having eaten less than five portions of fruit or vegetables the day before survey interview and having an unhealthy BMI, before and after adjustment for the other independent variables. NEET status was included in the model even if the effect was not significant. Since health behaviours of UE NEETs and EI NEETs are likely to be different, logistic regressions for health behaviours were performed before and after exclusion of EI NEETs. Further, as individuals aged under 18 years are not legally permitted to purchase alcohol or tobacco in Scotland or England, analyses for smoking and alcohol-related outcomes were restricted to survey respondents aged 18 years and over. Missing data were imputed using regression imputation. All analyses were conducted in IBM SPSS Statistics 21 [40].

## Results

### Characteristics of survey respondents

Data were available for 1717 SHeS respondents and 2555 HSE respondents, giving a total sample size of 4,272 respondents. Characteristics of respondents by NEET status are presented in Table 1. In this sample of young people aged 16–24 years, 715 (17%) were classified as being NEET. Of the 715 respondents classified as NEET, 371 (52%) were further defined as unemployed NEETs and 342 (48%) as economically inactive. Two NEETs could not be defined as unemployed or economically inactive due to missing information.

### Socio-demographic characteristics of NEETs

Results from the univariate logistic regression (Table 2) showed that, before exclusion of economically inactive NEETs, NEETs were significantly more likely ( $p < 0.001$ ) to be female; older in age; be married or cohabiting with a partner; have no car/van access; be educated only to standard grade level or below or, have foreign or no qualifications; not own their home either outright or with a mortgage; receive means-tested benefits; have lower total annual household income; have NS-SEC category other than managerial/professional; have a limiting long-term illness; have fair-bad self-assessed general health and a GHQ-12 score of three or more. There was

**Table 1** Characteristics and chi-square analysis of health survey respondents by NEET status

Variable	Categories	All NEETs Included	Economically Inactive NEETs Excluded	Total <sup>a</sup> (%)
		% NEET	% NEET	
Sex	1 Male	**14	*11	1881 (44)
	2 Female	20	8	2391 (56)
Age <sup>b</sup>		**Mean age NEET = 21 Mean age non-NEET = 20	*Mean age NEET = 20 Mean age non-NEET = 20	
Ethnicity	1 White UK & Irish	17	10	3571 (84)
	2 Other (incl. gypsy/traveller)	16	10	684 (16)
Marital status	1 Married/cohabiting	**27	*12	821 (19)
	2 Other (incl. single/separated/divorced)	15	9	3450 (81)
Access to car/van	1 Yes	**11	**6	3024 (71)
	2 No	33	19	1248 (29)
Top academic qualification	1 Degree or higher	**12	**8	536 (13)
	2 HNC/D or equiv (higher education below degree)	12	6	386 (9)
	3 Higher/A-level or equiv (upper school qualification)	9	5	1496 (35)
	4 Standard grade/O-level or equiv (lower school qualification)	21	13	1314 (31)
	5 Foreign or other qualification	34	18	170 (4)
	6 No educational qualification	41	23	347 (8)
Housing tenure	1 Own outright/with mortgage	**8	**5	1955 (46)
	2 Other (incl. part rent/part mortgage, renting, rent-free, squatting)	25	14	2313 (54)
Receipt of means-tested benefits	0 No	**7	**5	1706 (40)
	1 Yes	24	13	2566 (60)
Total annual household income	1 < £15,600	**36	**22	1057 (25)
	2 £15,600-£25,999	17	10	736 (17)
	3 £26,000-£36,399	10	6	510 (12)
	4 £36,400-£51,999	5	3	517 (12)
	5 £52,000-£69,999	5	2	344 (8)
	6 £70,000-£150,000+	7	4	386 (9)
	7 Refused	15	9	426 (10)
	8 Don't know	14	8	296 (7)
NSSEC	1 Managerial & professional	**6	**4	439 (10)
	2 Intermediate	16	8	652 (15)
	3 Routine & manual	19	11	2262 (53)
	4 Other or never worked/long-term unemployed	18	11	919 (22)
Limiting long-term illness	1 Limiting long-term illness	**32	13	446 (10)
	2 Non-limiting long-term illness	14	8	409 (10)
	3 No limiting long-term illness	15	9	3411 (80)
Self-assessed general health	1 Very good/good	**15	**9	3757 (88)
	2 Fair/bad/very bad	34	19	512 (12)

**Table 1** Characteristics and chi-square analysis of health survey respondents by NEET status (Continued)

GHQ-12 score	0 Score 0	**15	**8	2250 (53)
	1 Score 1–2	16	9	1144 (27)
	2 Score 3–4	19	11	476 (11)
	3 Score 5–12	29	19	402 (9)
Survey	1 Health Survey for England (HSE)	17	9	2555 (60)
	2 Scottish Health Survey (SHeS)	17	10	1717 (40)
Survey year	1 2010	18	9	1502 (35)
	2 2011	17	9	1527 (36)
	3 2012	16	10	1243 (29)

\* $p < 0.05$ \*\* $p < 0.001$ 

<sup>a</sup>The value for Total may not exactly equal the sum of NEET and non-NEET counts due to missing data within categories. Frequencies and percentages in this column are based on all NEETs

<sup>b</sup>Mean age at survey interview between NEETs and non-NEETs was compared using a 2-sample *t*-test. 95% confidence interval for  $\mu(\text{Age}_{\text{non-neet}}) - \mu(\text{Age}_{\text{neet}})$  was  $(-1.2, -0.8)$  when all NEETs were included and  $(-0.9, -0.3)$  when excluding economically inactive NEETs

no difference in the odds of being NEET by ethnicity or between Scottish and English health survey respondents and the odds of being NEET was not significantly different across the three survey years ( $p = 0.843$  and  $p = 0.505$  respectively). When EI NEETs were excluded, limiting long-term illness was no longer significantly associated with the odds of being NEET ( $p = 0.085$ ) and the direction of the effect of sex on NEET status was reversed with males significantly more likely to be NEET. The direction of the effects of survey (SHeS/HSE) and survey year was also reversed when economically inactive NEETs were excluded; however, the effects of these variables remained non-significant ( $p = 0.166$  and  $p = 0.634$  respectively). Parameter estimates for remaining variables were either similar or attenuated compared to those before exclusion of EI NEETs.

#### Health outcomes by NEET status

Frequencies and column percentages of participation in cancer-related unhealthy behaviours by NEET status before and after exclusion of economically inactive NEETs are presented in Table 3 with results from logistic regressions of investigating the effect of NEET status on health outcomes in Tables 4, 5, 6, 7 and 8.

Results from Table 4 demonstrated that NEETs were significantly more likely to be current smokers than non-NEETs ( $p < 0.001$ ) both before (odds ratio (OR) = 2.38, 95% confidence interval (CI) = 1.99–2.84) and after (OR = 2.34, 95% CI = 1.85–2.96) EI NEETs were excluded (i.e. when considering UE NEETs only). This result persisted even after adjustment for significant socio-demographic and health-related confounders. Odds ratios for heavy drinking (Table 5) demonstrated that NEETs were less likely to be heavy drinkers than non-NEETs; however, this decreased risk was only significant before exclusion of EI NEETs (OR = 0.73, 95% CI = 0.59–0.90). Adjusting for significant socio-demographic characteristics did not alter results. NEETs were significantly more likely to report not taking part in

sporting activities (Table 6) (OR = 2.12, 95% CI = 1.80–2.50 when all NEETs were included). The effect was attenuated, but still significant, after excluding EI NEETs (OR = 1.54, 95% CI = 1.23–1.92). The increased risk of reporting not taking part in any sporting activities amongst NEETs remained significant even in the fully-adjusted model. The likelihood of reporting non-participation in sporting activities significantly decreased over time ( $p < 0.05$ ) as demonstrated by the odds ratios for survey year.

NEETs were significantly more likely to report not having eaten the UK Government-recommended five portions of fruit or vegetables the day before survey interview ( $p < 0.05$ ) than non-NEETs both before (OR = 1.34, 95% CI = 1.09–1.64) and after (OR = 1.46, 95% CI = 1.11–1.93) exclusion of EI NEETs (Table 7). Results were attenuated in the fully-adjusted model and the increased risk was no longer significant both before or exclusion of EI NEETs. The odds of reporting not having eaten at least five portions significantly increased over time ( $p < 0.001$ ) as demonstrated by the odds ratios for survey year.

Finally, NEETs were also highly significantly more likely to have an 'unhealthy' BMI ( $p < 0.001$ ) than non-NEETs both before (OR = 1.57, 95% CI = 1.33–1.84) and after (OR = 1.62, 95% CI = 1.23–1.88) excluding EI NEETs (Table 8). The increased risk amongst NEETs remained significant even in the fully-adjusted model, but the effect was slightly stronger before exclusion of EI NEETs.

#### Health outcomes by survey region

No participation in sport and fruit and vegetable consumption were the only outcomes for which there was a significant effect of country of survey. However, survey region was only significant ( $p < 0.05$ ) after excluding EI NEETs for the 'no participation in sport' outcome (Table 6). In the fully-adjusted model, respondents to the SHeS were significantly less likely to report no participation in sport than respondents to the HSE (OR = 0.84, 95% CI = 0.72–0.97)

**Table 2** Results<sup>a</sup> from univariate logistic regressions for the effect of socio-demographic characteristics on NEET status

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
Sex		
Male	1.00**	1.00*
Female	1.51 (1.28–1.78)	0.72 (0.58–0.90)
Age	1.16** (1.12–1.20)	1.09** (1.05–1.14)
Ethnicity		
White UK & Irish	1.00	1.00
Other (incl. gypsy/traveller)	0.94 (0.75–1.17)	1.00 (0.75–1.34)
Marital status		
Married/cohabiting	2.13 (1.78–2.56)	1.37 (1.06–1.78)
Other (incl. single/separated/divorced)	1.00**	1.00*
Access to car/van		
Yes	1.00**	1.00**
No	4.13 (3.50–4.88)	3.54 (2.85–4.41)
Top academic qualification		
Degree or higher	1.00**	1.00**
HNC/D or equiv (higher education below degree)	1.00 (0.66–1.50)	0.69 (0.40–1.18)
Higher/A-level or equiv (upper school qualification)	0.77 (0.56–1.06)	0.64 (0.43–0.94)
Standard grade/O-level or equiv (lower school qualification)	1.99 (1.48–2.68)	1.59 (1.11–2.28)
Foreign or other qualification	3.77 (2.48–5.71)	2.48 (1.45–4.25)
No educational qualification	5.18 (3.68–7.28)	3.22 (2.10–4.94)
Housing tenure		
Own outright/with mortgage	0.26 (0.22–0.32)	0.33 (0.26–0.42)
Other (incl. part rent/part mortgage, renting, rent-free, squatting)	1.00**	1.00**
Receipt of means-tested benefits		
No	0.26 (0.21–0.32)	0.34 (0.27–0.44)
Yes	1.00**	1.00**
Total annual household income		
< £15,600	1.00**	1.00**
£15,600–£25,999	0.37 (0.29–0.46)	0.39 (0.29–0.53)
£26,000–£36,399	0.21 (0.15–0.29)	0.21 (0.14–0.33)
£36,400–£51,999	0.09 (0.06–0.14)	0.13 (0.08–0.21)
£52,000–£69,999	0.10 (0.06–0.16)	0.08 (0.04–0.17)
£70,000–£150,000+	0.14 (0.09–0.21)	0.16 (0.10–0.27)
Refused	0.32 (0.23–0.42)	0.35 (0.23–0.51)
Don't know	0.29 (0.20–0.41)	0.33 (0.21–0.52)
NSSEC		
Managerial & professional	1.00**	1.00**
Intermediate	3.04 (1.94–4.75)	2.23 (1.22–4.06)
Routine & manual	3.69 (2.45–5.56)	3.38 (1.98–5.77)
Other or never worked/long-term unemployed	3.36 (2.18–5.17)	2.24 (1.85–5.68)
Limiting long-term illness		
Limiting long-term illness	2.53 (2.02–3.16)	1.42 (1.01–1.99)



**Table 2** Results<sup>a</sup> from univariate logistic regressions for the effect of socio-demographic characteristics on NEET status (Continued)

Non-limiting long-term illness	0.91 (0.68–1.22)	0.88 (0.60–1.29)
No limiting long-term illness	1.00**	1.00
GHQ-12 score		
Score 0	1.00**	1.00**
Score 1–2	1.08 (0.89–1.32)	1.07 (0.82–1.39)
Score 3–4	1.37 (1.06–1.77)	1.44 (1.03–2.01)
Score 5–12	2.31 (1.80–2.95)	2.53 (1.85–3.46)
Self-assessed general health		
Very good/good	1.00**	1.00**
Fair/bad/very bad	2.99 (2.43–3.67)	2.43 (1.84–3.20)
Survey		
Health Survey for England (HSE)	1.00	1.00
Scottish Health Survey (SHeS)	0.98 (0.84–1.16)	1.16 (0.94–1.44)
Survey year		
2010	1.00	1.00
2011	0.93 (0.77–1.12)	1.02 (0.79–1.32)
2012	0.89 (0.73–1.09)	1.13 (0.87–1.47)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>Odds ratios (OR) and 95% confidence intervals (95% CI) are presented

after exclusion of EI NEETs. For fruit and vegetable consumption (Table 7), respondents to the SHeS were significantly more likely to report not having eaten at least five portions ( $p < 0.001$  before and after exclusion of EI NEETs) compared to respondents to HSE.

## Discussion

This study aimed to contribute to the limited evidence base on whether not being in education, employment or training was associated with a greater likelihood of participating in cancer-related behaviours.

## Socio-demographic characteristics of NEETs

Increasing age was significantly associated with increased odds of being NEET; however, this effect was stronger before exclusion of EI NEETs. This result, along with females being at an increased risk of being NEET before excluding EI NEETs, possibly reflects females taking time out of education or employment to start a family as they get older. The fact that effects of gender and age are reversed or attenuated when EI NEETs are excluded would appear to support this belief. Findings also support previous reports of NEETs being from socioeconomically-disadvantaged backgrounds [41].

**Table 3** Frequencies (column %) of participation in unhealthy behaviours by NEET status

Outcome	All NEETs Included		Economically Inactive NEETs Excluded		Total (%)	
	NEET (%)	non-NEET (%)	NEET (%)	non-NEET (%)	All NEETs Included	Economically Inactive NEETs Excluded
Current smoker	297 (46)	684 (27)	146 (46)	684 (27)	981 (30)	830 (29)
Heavy drinker <sup>a</sup>	128 (20)	656 (25)	75 (24)	656 (25)	784 (24)	731 (25)
No participation in sport	330 (46)	1003 (29)	143 (38)	1003 (29)	1333 (32)	1146 (30)
Less than 5 portions fruit/vegetables <sup>b</sup>	579 (81)	2652 (76)	307 (82)	2652 (76)	3231 (77)	2959 (77)
Unhealthy BMI <sup>c</sup>	387 (54)	1498 (43)	199 (53)	1498 (43)	1885 (45)	1697 (44)

<sup>a</sup>Heavy drinking refers to consuming >14 units of alcohol per week for females and >21 units for males <sup>b</sup>Refers to whether the respondent reported eating less than 5 portions of fruit or vegetables the day before survey interview (yes/no)

<sup>c</sup>Refers to whether the respondent was defined as having an unhealthy BMI (yes/no). Unhealthy BMI refers to being 'underweight' (BMI <18.5 kg/m<sup>2</sup>); 'overweight' (BMI 25–29.99 kg/m<sup>2</sup>) or 'obese' (BMI >=30 kg/m<sup>2</sup>)

**Table 4** Odds ratios and 95% CIs for effect of NEET status on Current Smoking<sup>a,b</sup>

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
UNIVARIATE <sup>c</sup>		
NEET		
No	1.00**	1.00**
Yes	2.38 (1.99–2.84)	2.34 (1.85–2.96)
FULLY ADJUSTED <sup>c</sup>		
NEET		
No	1.00*	1.00*
Yes	1.36 (1.10–1.68)	1.49 (1.14–1.94)
Sex		
Male	1.26 (1.07–1.49)	1.23 (1.03–1.47)
Female	1.00*	1.00*
Age		
	1.06* (1.01–1.11)	1.06* (1.01–1.11)
Ethnicity		
White UK & Irish	1.00**	1.00*
Other (incl. gypsy/traveller)	0.57 (0.45–0.72)	0.65 (0.50–0.84)
Access to car/van		
Yes	1.00**	1.00**
No	1.56 (1.29–1.89)	1.41 (1.15–1.73)
Top academic qualification		
Degree or higher	1.00**	1.00**
HNC/D or equiv (higher education below degree)	1.75 (1.24–2.46)	1.71 (1.21–2.43)
Higher/A-level or equiv (upper school qualification)	1.39 (1.04–1.86)	1.29 (0.96–1.73)
Standard grade/O-level or equiv (lower school qualification)	2.68 (1.99–3.61)	2.52 (1.85–3.42)
Foreign or other qualification	2.98 (1.91–4.63)	3.47 (2.15–5.60)
No educational qualification	3.04 (2.05–4.51)	3.42 (2.19–5.32)
Housing tenure		
Own outright/with mortgage	1.00**	1.00**
Other (incl. part rent/part mortgage, renting, rent-free, squatting)	1.97 (1.62–2.39)	1.95 (1.59–2.39)
NSSEC		
Managerial & professional	1.00**	1.00**
Intermediate	1.24 (0.90–1.71)	1.28 (0.92–1.77)
Routine & manual	1.38 (1.04–1.84)	1.34 (1.01–1.79)
Other or never worked/long-term unemployed	0.58 (0.39–0.87)	0.52 (0.33–0.80)
Self-assessed general health		
Very good/good	1.00**	1.00**
Fair/bad/very bad	1.97 (1.57–2.47)	2.09 (1.63–2.68)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>The outcome is whether the respondent reported being a current smoker (yes/no)<sup>b</sup>As the legal minimum age for buying tobacco in Scotland and England is 18 years of age, 16 and 17 year-olds have been excluded from analysis<sup>c</sup>Univariate refers to the model containing NEET status only and fully adjusted is the model containing all significant socio-demographic and health-related characteristics



**Table 5** Odds ratios and 95% CIs for effect of NEET status on Heavy Drinking<sup>a,b</sup>

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
<b>UNIVARIATE<sup>c</sup></b>		
<b>NEET</b>		
No	1.00*	1.00
Yes	0.73 (0.59–0.90)	0.90 (0.69–1.19)
<b>FULLY ADJUSTED<sup>c</sup></b>		
<b>NEET</b>		
No	1.00*	1.00
Yes	0.71 (0.56–0.91)	0.91 (0.68–1.22)
<b>Sex</b>		
Male	0.71 (0.60–0.85)	0.69 (0.58–0.82)
Female	1.00**	1.00**
<b>Ethnicity</b>		
White UK & Irish	1.00**	1.00**
Other (incl. gypsy/traveller)	0.43 (0.33–0.56)	0.41 (0.31–0.54)
<b>Access to car/van</b>		
Yes	1.00*	1.00*
No	1.24 (1.03–1.49)	1.29 (1.06–1.56)
<b>Marital status</b>		
Married/cohabiting	0.65 (0.53–0.80)	0.67 (0.54–0.83)
Other (incl. single/separated/divorced)	1.00**	1.00**
<b>Receipt of means-tested benefits</b>		
No	1.00*	1.00*
Yes	0.80 (0.67–0.95)	0.80 (0.67–0.96)
<b>NSSEC</b>		
Managerial & professional	1.00*	1.00*
Intermediate	1.06 (0.78–1.45)	1.07 (0.78–1.46)
Routine & manual	1.40 (1.08–1.83)	1.40 (1.07–1.84)
Other or never worked/long-term unemployed	1.13 (0.80–1.61)	1.18 (0.82–1.70)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>The outcome is whether the respondent was defined as being a heavy drinker (yes/no). Heavy drinking refers to consuming >14 units of alcohol per week for females and >21 units for males<sup>b</sup>As the legal minimum age for buying alcohol in Scotland and England is 18 years of age, 16 and 17 year-olds have been excluded from analysis<sup>c</sup>Univariate refers to the model containing NEET status only and fully adjusted is the model containing all significant socio-demographic and health-related characteristics

In terms of health characteristics, fair-bad self-assessed general health and short-term non-psychotic psychiatric morbidity, (GHQ-12 score), were significantly associated with an increased risk of being NEET. These are known indicators of poorer mental health, which has been previously associated with becoming NEET [15].

Other differences noted between EI and UE NEETs related to having a limiting long-term illness, which was no longer significant after excluding EI NEETs. This result is expected since excluding EI NEETs would remove individuals with long-term illness/disability. Further, there was an increased likelihood of being NEET amongst SHes versus HSE respondents after excluding EI NEETs. Although the difference was not statistically significant, this would suggest greater rates of unemployment amongst young people in Scotland compared to England. Increasing the sample size by adding data from more recent health surveys as they become available may confirm significant differences in the likelihood of being NEET across different regions of the UK.

#### Cancer-related health behaviours of NEETs

This study found a greater tendency for NEETs to participate in cancer-related unhealthy behaviours compared to non-NEETs. However, there were some differences in the effect of NEET status before and after exclusion of EI NEETs.

There are several possible explanations as to why participation in unhealthy behaviours may be greater in NEETs compared to non-NEETs. As confirmed by this study and in previous studies, NEETs are more likely to be poorly educated [42]. Poor education may diminish knowledge of how to live a healthy life [43] and reduce decision-making abilities for making healthy choices [44]. However, there remained an independent effect of NEET status on participation on some unhealthy behaviours even after adjustment for top academic qualification. Similarly, as demonstrated in this study and in previous research, NEETs were more likely to have reduced income [41]. Reduced income may restrict healthy dietary options or the ability to participate in healthy recreational activities [45]. This could explain associations between being NEET and reduced fruit and vegetable consumption, participation in sport and an unhealthy BMI. Indeed, in addition to other socio-demographic and health-related confounders, total annual household income explained the effect of NEET status on fruit and vegetable consumption before and after EI NEETs were excluded. However, being NEET remained independently associated with reduced participation in sport and an unhealthy BMI even after adjustment for total annual household income. Being NEET also remained significantly associated with being a current smoker after adjustment for total annual household income. It could be expected that reduced income may lead to decreased participation in unhealthy behaviours such as smoking and drinking due to the financial cost associated with these behaviours, but there is a well-known link between unemployment and smoking in young people [46]. As well as the addiction to nicotine, smoking may be a coping mechanism as a way of

**Table 6** Odds ratios and 95% CIs for effect of NEET status No Participation in Sport<sup>a</sup>

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
UNIVARIATE <sup>b</sup>		
NEET		
No	1.00**	1.00**
Yes	2.12 (1.80–2.50)	1.54 (1.23–1.92)
FULLY ADJUSTED <sup>b</sup>		
NEET		
No	1.00**	1.00*
Yes	1.52 (1.26–1.82)	1.30 (1.03–1.65)
Sex		
Male	0.65 (0.57–0.75)	0.67 (0.58–0.77)
Female	1.00**	1.00**
Age		
	1.04* (1.01–1.07)	1.04* (1.01–1.07)
Ethnicity		
White UK & Irish	1.00*	
Other (incl. gypsy/traveller)	1.22 (1.02–1.47)	
Access to car/van		
Yes	1.00**	1.00*
No	1.31 (1.13–1.53)	1.29 (1.10–1.52)
Top academic qualification		
Degree or higher	1.00**	1.00**
HNC/D or equiv (higher education below degree)	1.62 (1.18–2.23)	1.54 (1.11–2.13)
Higher/A-level or equiv (upper school qualification)	1.84 (1.42–2.37)	1.76 (1.36–2.28)
Standard grade/O-level or equiv (lower school qualification)	2.29 (1.76–2.99)	2.22 (1.69–2.91)
Foreign or other qualification	2.77 (1.87–4.10)	2.79 (1.82–4.26)
No educational qualification	2.49 (1.79–3.46)	2.41 (1.68–3.46)
Limiting long-term illness		
Limiting long-term illness	1.22 (0.96–1.54)	
Non-limiting long-term illness	0.81 (0.64–1.03)	
No limiting long-term illness	1.00*	
Self-assessed general health		
Very good/good	1.00*	1.00*
Fair/bad/very bad	1.44 (1.15–1.79)	1.39 (1.11–1.73)
Survey year		
2010	1.00*	1.00*
2011	0.94 (0.81–1.10)	0.91 (0.77–1.08)
2012	0.79 (0.67–0.94)	0.75 (0.63–0.90)
Survey		
Health Survey for England		1.00*
Scottish Health Survey		0.84 (0.72–0.97)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>The outcome is whether the respondent reported no participation in sporting activities (yes/no)<sup>b</sup>Univariate refers to the model containing NEET status only and fully adjusted is the model containing all significant socio-demographic and health-related characteristics

**Table 7** Odds ratios and 95% CIs for effect of NEET status on Fruit & Vegetable Consumption<sup>a</sup>

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
UNIVARIATE <sup>b</sup>		
NEET		
No	1.00*	1.00*
Yes	1.34 (1.09–1.64)	1.46 (1.11–1.93)
FULLY ADJUSTED <sup>b</sup>		
NEET		
No	1.00	1.00
Yes	1.10 (0.87–1.39)	1.23 (0.91–1.66)
Fruit consumption the same as usual		
Less than usual	1.00**	1.00**
More than usual	0.24 (0.18–0.32)	0.24 (0.18–0.32)
About the same as usual	0.41 (0.34–0.50)	0.39 (0.32–0.48)
Vegetable consumption the same as usual		
Less than usual	1.00**	1.00**
More than usual	0.71 (0.55–0.92)	0.74 (0.56–0.96)
About the same as usual	0.59 (0.50–0.70)	0.59 (0.49–0.70)
Sex		
Male	1.19 (1.02–1.39)	1.19 (1.02–1.40)
Female	1.00*	1.00*
Ethnicity		
White UK & Irish	1.00**	1.00**
Other (incl. gypsy/traveller)	0.61 (0.51–0.75)	0.64 (0.52–0.79)
Top academic qualification		
Degree or higher	1.00**	1.00**
HNC/D or equiv (higher education below degree)	1.24 (0.90–1.70)	1.24 (0.89–1.71)
Higher/A-level or equiv (upper school qualification)	1.22 (0.96–1.54)	1.18 (0.92–1.50)
Standard grade/O-level or equiv (lower school qualification)	1.97 (1.51–2.56)	1.90 (1.45–2.49)
Foreign or other qualification	1.89 (1.20–2.96)	1.95 (1.20–3.19)
No educational qualification	1.51 (1.07–2.14)	1.45 (0.99–2.12)
Receipt of means-tested benefits		
No	1.00*	1.00*
Yes	1.19 (1.00–1.40)	1.20 (1.01–1.43)
Total annual household income		
< £15,600	1.00*	1.00*
£15,600–£25,999	0.87 (0.68–1.12)	0.92 (0.70–1.20)
£26,000–£36,399	1.06 (0.79–1.41)	1.14 (0.85–1.55)
£36,400–£51,999	0.81 (0.61–1.07)	0.88 (0.66–1.17)
£52,000–£69,999	0.68 (0.50–0.93)	0.74 (0.54–1.02)
£70,000–£150,000+	0.57 (0.43–0.76)	0.62 (0.46–0.84)
Refused	0.88 (0.66–1.17)	0.98 (0.72–1.32)
Don't know	0.85 (0.62–1.18)	0.99 (0.70–1.40)

**Table 7** Odds ratios and 95% CIs for effect of NEET status on Fruit & Vegetable Consumption<sup>a</sup> (Continued)

Self-assessed general health		
Very good/good	1.00*	1.00*
Fair/bad/very bad	1.34 (1.03–1.73)	1.34 (1.01–1.77)
Survey		
Health Survey for England (HSE)	1.00**	1.00**
Scottish Health Survey (SHeS)	1.82 (1.54–2.16)	1.83 (1.53–2.18)
Survey year		
2010	1.00**	1.00**
2011	1.08 (0.90–1.29)	1.09 (0.90–1.31)
2012	1.59 (1.28–1.97)	1.56 (1.25–1.94)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>The outcome is whether the respondent reported eating less than 5 portions of fruit or vegetables the day before survey interview (yes/no)<sup>b</sup>Univariate refers to the model containing NEET status only and fully adjusted is the model containing all significant socio-demographic and health-related characteristics

dealing with the stresses associated with unemployment [46–48]. Indeed, the association between NEET status and current smoking strengthened when considering unemployed NEETs only by excluding EI NEETs.

An inverse association between NEET status and heavy drinking was observed when considering all NEETs; however, this association became non-significant on exclusion of EI NEETs. Available socio-demographic and health-related significant risk factors could not explain the negative association between being NEET and heavy drinking when considering all NEETs. For example, although some previous research has reported lower levels of alcohol consumption amongst young mothers [16, 49] and young people who are disabled [50], adjusting for gender did not alter the effect of NEET status on heavy drinking and limiting long-term illness was not significantly associated with heavy drinking in this study. However, this study did not control for whether respondents had children. Early parenthood has been shown to moderate alcohol consumption [49]. As the likelihood of having children may be greater amongst young people who are economically inactive, particularly as analysis of the heavy drinking outcome is restricted to those aged 18 and over, the NEET status variable, before exclusion of EI NEETs, may be accounting for residual confounding resulting from not controlling for whether the respondent had children.

The effect of NEET status on reporting of no participation in sport was also attenuated on exclusion of EI NEETs, but the effect remained significant. Again, this result is not unexpected given previous reports of lower levels of sport amongst young mothers [51] and young people who are permanently or temporarily sick, disabled or injured [52]. Although physical inactivity has been linked with certain cancers [24], the 'no participation in sport' measure used in this study only reflects

one dimension of physical inactivity. Other aspects of physical activity (PA), such as occupational-, transport- and domestic-related domains, which, to some extent, have also been shown to be protective for health [24], including some cancers [53, 54], have not been considered here. Using a more comprehensive measure of PA may have altered the effect of NEET status on this outcome. In particular, the observed attenuation in the effect of NEET status after excluding EI NEETs may have been smaller if a measure of occupational-related activity had also been included. Alternatively, if a domestic-related measure of PA had been included then there could plausibly have been a greater attenuation of the effect of NEET status after excluding EI NEETs as a result of excluding young, economically inactive females. Such individuals may be exposed to higher levels of PA through the physical demands of looking after the home and young family.

The association between being NEET and reduced fruit and vegetable consumption was stronger when considering UE NEETs only. Although both unemployment and disability have been associated with an increased risk of food poverty, including insufficient consumption of fruit and vegetables [55], this finding would suggest that the risk is greater in unemployed young people than those who are economically inactive when compared to their 'non-NEET' peers. However, the effect of NEET status on this outcome could be explained by socio-demographic and health-related confounders both before and after exclusion of EI NEETs.

The increase in reporting of not eating at least five portions of fruit and vegetables over time was interesting given there had been a decrease in reporting of non-participation in sport over time. It would therefore appear that this study does not support previously-reported associations between sedentary behaviours and

**Table 8** Odds ratios and 95% CIs for effect of NEET status on BMI<sup>a</sup>

Variable	All NEETs Included OR (95% CI)	Economically Inactive NEETs Excluded OR (95% CI)
UNIVARIATE <sup>b</sup>		
NEET		
No	1.00**	1.00**
Yes	1.57 (1.33–1.84)	1.62 (1.23–1.88)
FULLY ADJUSTED <sup>b</sup>		
NEET		
No	1.00**	1.00*
Yes	1.36 (1.15–1.61)	1.34 (1.07–1.68)
Age	1.05** (1.03–1.09)	1.07** (1.03–1.10)
Ethnicity		
White UK & Irish		1.00*
Other (incl. gypsy/traveller)		0.81 (0.68–0.98)
Marital status		
Married/cohabiting	1.29 (1.09–1.53)	1.22 (1.01–1.48)
Other (incl. single/separated/divorced)	1.00*	1.00*
Top academic qualification		
Degree or higher		1.00*
HNC/D or equiv (higher education below degree)		1.31 (0.99–1.74)
Higher/A-level or equiv (upper school qualification)		1.17 (0.93–1.46)
Standard grade/O-level or equiv (lower school qualification)		1.42 (1.11–1.81)
Foreign or other qualification		1.56 (1.04–2.31)
No educational qualification		1.54 (1.11–2.14)
NSSEC		
Managerial & professional		1.00*
Intermediate		1.03 (0.97–1.33)
Routine & manual		0.80 (0.64–1.00)
Other or never worked/long-term unemployed		0.94 (0.71–1.24)
Self-assessed general health		
Very good/good	1.00**	1.00**
Fair/bad/very bad	1.51 (1.24–1.82)	1.56 (1.26–1.93)

\* $p < 0.05$ \*\* $p < 0.001$ <sup>a</sup>The outcome is whether the respondent was defined as having an unhealthy BMI (yes/no). Unhealthy BMI refers to being 'underweight' (BMI <18.5 kg/m<sup>2</sup>); 'overweight' (BMI 25–29.99 kg/m<sup>2</sup>) or 'obese' (BMI >=30 kg/m<sup>2</sup>)<sup>b</sup>Univariate refers to the model containing NEET status only and fully adjusted is the model containing all significant socio-demographic and health-related characteristics

less healthy eating patterns in young adults [56]. However, the increase in participation in sport in 2012 found in this study coincides with London hosting the 2012 Olympic Games. The recognised peak in participation in sport in 2012 has since declined [57], therefore findings may not have been similar if more recent data had been used. Since the decline has been shown to be greater amongst more socioeconomically-disadvantaged groups [58], NEETs may be more vulnerable to this decline.

### Strengths and limitations

Merging of health surveys allowed for cross-national comparisons of health behaviours associated with cancer. A particular strength was that the comprehensive data on economic profiles of respondents collected by these surveys allowed NEETs to be classed further as EI NEETs or UE NEETs. This was important since previous studies reported differences in socio-demographic characteristics and health behaviours of EI and UE NEETs [59, 60]; a finding that was also confirmed by this study.

A strength of using the Scottish Health Survey is that information on respondents can be further linked to Scottish Morbidity Records, including the Scottish Cancer Registry (SMR06). The SMR06 scheme collects information on all residents in Scotland that have had a diagnosis of cancer. Future research could investigate the role of NEET status on developing cancer after adjustment for cancer-related behaviours and other socio-demographic characteristics, for a subset of the participants used in this study.

Cancer-related health behaviours, such as smoking, have also been associated with a greater risk of becoming NEET [15]. Due to the cross-sectional nature of the data, this study could not determine the temporal sequence of becoming NEET and participating in cancer-related behaviours and it was not possible to determine a causal effect of being NEET on cancer-related health behaviours, therefore reverse causation is possible.

## Conclusions

This study has shown that NEETs were at an increased risk of exhibiting cancer-related behaviours compared to non-NEETs, including smoking, not participating in sport and having an unhealthy BMI. Attempts to reduce participation in such behaviours amongst NEETs may contribute to a reduction in cancers associated with these behaviours. However, policymakers should be aware of differences between unemployed and economically and inactive NEETs. This was particularly relevant for heavy drinking. As the likelihood of becoming NEET is greater in socioeconomically-disadvantaged groups, interventions to tackle unhealthy behaviours among NEETs may contribute to a reduction in health inequalities.

## Abbreviations

CI: Confidence interval; EI: Economically inactive; GHQ-12: 12-item general health questionnaire; HSE: Health survey for England; NEET: Not in education employment or training; NS-SEC: National statistics socio-economic classification; OR: Odds ratio; SHS: Scottish health survey; UE: Unemployed

## Acknowledgements

We would like to acknowledge the Bupa Foundation Fund Innovation Grant sandpit workshop for providing the opportunity to form the research team and create our research proposal. We also thank Dr Lucy Davies (Research Funding Manager at CRUK) for her continued help and guidance during the duration of the project.

## Funding

This study was funded by Cancer Research UK (C53258/A19682) and the Medical Research Council/ Chief Scientist Office Social & Public Health Sciences Unit, University of Glasgow under the Measuring and Analysing Socioeconomic Inequalities in Health programme (MC\_UU\_12017/13 & SPHSU13). Cancer Research UK approved the final study design/proposal. There was no involvement of the funding bodies in data collection, analysis and interpretation and in writing the manuscript.

## Availability of data and materials

The datasets generated and analysed during the current study are available in the Health and health behaviour repository on the UK Data Service website [<https://www.ukdataservice.ac.uk/get-data/themes/health/>] [28–33].

## Authors' contributions

CS, PB, DP and CT all devised the study. CS cleaned, merged and analysed the data. CS wrote the first draft of the report. PB, DP and CT critically reviewed the paper and suggested revisions. All authors read and approved the final manuscript.

## Competing interests

The authors declare that they have no competing interests.

## Consent for publication

Not applicable.

## Ethics approval and consent to participate

Original ethical approval for the 2010–11 and 2012 Scottish Health Surveys was obtained from the Multicentre Research Ethics Committee for Wales and the Research Committee for Wales respectively [37, 61]. For the 2010 and 2011–12 Health Surveys for England, original ethical approval was obtained from the Oxford B Research Ethics Committee and Oxford A Research Ethics Committee respectively [62–64]. Respondents sign consent forms at time of survey. No further ethical approval or consent was required when downloading the anonymised datasets for statistical and research purposes from the UK Data Service; however, registration with the site is required.

## Author details

<sup>1</sup>MRC/CSO Social & Public Health Sciences Unit, University of Glasgow, Top Floor, 200 Renfield Street, Glasgow G2 3QB, UK. <sup>2</sup>Northern Institute for Cancer Research, Newcastle University, Framlington Place, Newcastle NE2 4HH, UK. <sup>3</sup>Health & Lifestyles Research Unit, Wolfson Institute of Preventive Medicine, Queen Mary University of London, 2 Stayner's Road, London E1 4AH, UK. <sup>4</sup>Centre for Public Health, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Institute of Clinical Sciences B, Royal Victoria Hospital Site, Grosvenor Road, Belfast BT12 6BJ, UK.

Received: 23 August 2016 Accepted: 24 February 2017

Published online: 02 March 2017

## References

- Office for National Statistics. Young People Not in Education, Employment or Training (NEET): February 2017. 2017. <https://www.ons.gov.uk/employmentandlabourmarket/peoplenotinwork/unemployment/bulletins/youngpeoplenotineducationemploymentortrainingneet/feb2017>. Accessed 28 Feb 2017.
- Delebarre J. NEET: Young People Not in Education, Employment or Training. 2016. <http://researchbriefings.parliament.uk/ResearchBriefing/Summary/SN06705>. Accessed 9 Aug 2016.
- Crawford C, Duckworth K, Vignoles A, et al. Young people's education and labour market choices aged 16/17 to 18/19: UK Government. 2011.
- Mroz TA, Savage TH. The long-term effects of youth unemployment. *J Hum Resour.* 2006;41(2):259–93.
- Bartley M, Plewis I. Accumulated labour market disadvantage and limiting long-term illness: data from the 1971–1991 office for national statistics' longitudinal study. *Int J Epidemiol.* 2002;31(2):336–41.
- Bell DNF, Blanchflower DG. Recession and Unemployment in the OECD. CESifo Forum. Munich: Ifo Institute for Economic Research; 2010.
- Hagquist C, Starrin B. Youth unemployment and mental health – gender differences and economic stress. *Scand J Soc Welfare.* 1996;5(4):215–28.
- Siegrist J, Benach J, McKnight A, et al. Employment arrangements, work conditions and health inequalities. In: Report on new evidence on health inequality reduction, produced by Task 2 for the Strategic review of health inequalities post 2010. 2010.
- Fergusson DM, John Horwood L, Woodward LJ. Unemployment and psychosocial adjustment in young adults: causation or selection? *Soc Sci Med.* 2001;53(3):305–20.
- Brydsten A, Hammarström A, Strandh M, et al. Youth unemployment and functional somatic symptoms in adulthood: results from the Northern Swedish cohort. *Eur J Public Health.* 2015;25(5):796–800.
- Hammarström A, Janlert U. Early unemployment can contribute to adult health problems: results from a longitudinal study of school leavers. *J Epidemiol Community Health.* 2002;56(8):624–30.



12. Montgomery SM, Cook DG, Bartley MJ, et al. Unemployment, cigarette smoking, alcohol consumption and body weight in young British men. *Eur J Public Health*. 1998;8(1):21–7.
13. Voss M, Nylén L, Floderus B, et al. Unemployment and early cause-specific mortality: a study based on the Swedish twin registry. *Am J Public Health*. 2004;94(12):2155–61.
14. Hammer T. Unemployment and use of drug and alcohol among young people: a longitudinal study in the general population. *Br J Addict*. 1992; 87(11):1571–81.
15. Baggio S, Iglesias K, Deline S, et al. Not in education, employment, or training status among young Swiss Men. Longitudinal associations with mental health and substance use. *J Adolesc Health*. 2015;56(2):238–43.
16. Janlert U, Hammarström A. Alcohol consumption among unemployed youths: results from a prospective study. *Br J Addict*. 1992;87(5):703–14.
17. Nardi B, Lucarelli C, Talamonti M, et al. NEETs versus EETs: an observational study in Italy on the framework of the HEALTH25 European project. *Res Post-Compulsory Educ*. 2015;20(4):377–99.
18. Parkin DM, Boyd L, Walker LC. 16. The fraction of cancer attributable to lifestyle and environmental factors in the UK in 2010. *Br J Cancer*. 2011; 105(52):S77–81.
19. Danaei G, Vander Hoon S, Lopez AD, et al. Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors. *Lancet*. 2005;366(9499):1784–93.
20. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, et al. Cancer incidence and mortality patterns in Europe: Estimates for 40 countries in 2012. *Eur J Cancer*. 2013;49(6):1374–403.
21. Ezzati M, Lopez AD. Smoking and oral tobacco use. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*: world health organization. 2004. p. 883–958.
22. Rehm J, Room R, Monteiro M, et al. Alcohol use. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*: world health organization. 2004. p. 959–1108.
23. Lock K, Pomerleau J, Causer L, et al. Other nutrition-related risk factors and physical inactivity: Low fruit and vegetable consumption. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*: world health organization. 2004. p. 597–728.
24. Bull FC, Armstrong TP, Dixon T, et al. Physical inactivity. In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*: world health organization. 2004. p. 729–882.
25. James WPT, Jackson-Leach R, Mhurchu CN, et al. Overweight and obesity (high body mass index). In: Ezzati M, Lopez AD, Rodgers A, Murray CJL, editors. *Comparative quantification of health risks: global and regional burden of disease attributable to selected major risk factors*: world health organization. 2004. p. 497–596.
26. Morris JK, Cook DG, Shaper AG. Loss of employment and mortality. *Br Med J*. 1994;308(6937):1135–39.
27. Lyng E. Unemployment and cancer: a literature review. In: Kogevinas M, Pearce N, Susser M, Boffetta P, editors. *Social inequalities and cancer*: IARC publication. 1997. p. 343–51.
28. ScotCen Social Research, Department of Epidemiology and Public Health University College London. *Scottish Health Survey, 2010*. [data collection]. 2nd Edition. 2016. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-6987-2>. Accessed 9 Aug 2016.
29. ScotCen Social Research, Department of Epidemiology and Public Health University College London, MRC/CSO Social and Public Health Sciences Unit University of Glasgow. *Scottish Health Survey, 2011*. [data collection]. 4th Edition. 2016. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-7247-4>. Accessed 9 Aug 2016.
30. ScotCen Social Research, Department of Epidemiology and Public Health University College London, MRC/CSO Social and Public Health Sciences Unit University of Glasgow. *Scottish Health Survey 2012*. [data collection]. 3rd Edition. 2016. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-7417-3>. Accessed 9 Aug 2016.
31. NatCen Social Research, Department of Epidemiology and Public Health Royal Free and University College Medical School. *Health Survey for England, 2010*. [data collection]. 3rd Edition. 2015. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-6986-3>. Accessed 9 Aug 2016.
32. NatCen Social Research, Department of Epidemiology and Public Health University College London. *Health Survey for England, 2011*. [data collection]. 2013. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-7260-1>. Accessed 9 Aug 2016.
33. NatCen Social Research, Department of Epidemiology and Public Health University College London. *Health Survey for England, 2012*. [data collection]. 2014. UK Data Service. <http://dx.doi.org/10.5255/UKDA-SN-7480-1>. Accessed 9 Aug 2016.
34. Gray L, Batty GD, Craig P, et al. Cohort profile: the Scottish health surveys cohort: linkage of study participants to routinely collected records for mortality, hospital discharge, cancer and offspring birth characteristics in three nationwide studies. *Int J Epidemiol*. 2010;39(2):345–50.
35. Mindell J, Biddulph JP, Hirani V, et al. Cohort profile: the health survey for England. *Int J Epidemiol*. 2012;41(6):1585–93.
36. Stewart C. *Multilevel modelling of event history data: comparing methods appropriate for large datasets*. Glasgow: University of Glasgow; 2010.
37. Scottish Government. *The Scottish health survey 2010 volume 2: technical report*. 2011.
38. Office for National Statistics. *The National Statistics Socio-economic classification (NS-SEC)*. <https://www.ons.gov.uk/methodology/classificationsandstandards/otherclassifications/thenationalstatistics socioeconomicclassificationnssecbasedonsoc2010>. Accessed 9 Aug 2016.
39. Goldberg DP, Williams P. *A User's guide to the general health questionnaire: NFER-nelson*. 1991.
40. IBM Corp. Released 2012. *IBM SPSS Statistics for Windows*. Version 21.0 [program]. Armonk: IBM Corp.; 2012.
41. UCL Institute of Health Equity. *Local action on health inequalities: Reducing the number of young people not in employment, education or training (NEET)*. Health equity briefing 3. London: Public Health England; 2014.
42. Social Exclusion Unit. *Bridging the Gap: New opportunities for 16–18 year olds not in education, employment or training*. 1999.
43. Arendt JN. Does education cause better health? A panel data analysis using school reforms for identification. *Econ Educ Rev*. 2005;24(2):149–60.
44. Cutler DM, Lleras-Muney A. *Education and health: evaluating theories and evidence*. In: National bureau of economic research. 2006.
45. Adler NE, Newman K. *Socioeconomic disparities in health: pathways and policies*. *Health Aff (Millwood)*. 2002;21(2):60–76.
46. Hammarström A, Janlert U. Unemployment and change of tobacco habits: a study of young people from 16 to 21 years of age. *Addiction*. 1994;89(12): 1691–96.
47. De Vogli R, Santinello M. Unemployment and smoking: does psychosocial stress matter? *Tob Control*. 2005;14(6):389–95.
48. Marmot M, Wilkinson R. *Social patterning of individual health behaviours: the case of cigarette smoking*. *Social Determinants of Health*. New York: Oxford University Press Inc; 2005.
49. Seaman P, Ikegwonu T. *Drinking to belong: Understanding young adults' alcohol use within social networks*. 2010.
50. Centre for Longitudinal Studies. *Research summary three: Growing up with a disability*. London: Trajectories and transitions of disabled children and young people; 2014.
51. Brown WJ, Trost SG. Life transitions and changing physical activity patterns in young women. *Am J Prev Med*. 2003;25(2):140–43.
52. Jones H, Millward P, Buraimo B. *Adult participation in sport: analysis of the taking part survey*: department for culture, media and sport. 2011.
53. Shi Y, Li T, Wang Y, et al. Household physical activity and cancer risk: a systematic review and dose–response meta-analysis of epidemiological studies. *Sci Rep*. 2015;5:14901.
54. Dosemeci M, Hayes RB, Vetter R, et al. Occupational physical activity, socioeconomic status, and risks of 15 cancer sites in Turkey. *Cancer Causes Control*. 1993;4(4):313–21.
55. Mwatsama M, Stewart L. *Food poverty and health: briefing statement*: faculty of public health of the royal colleges of physicians of the united kingdom. 2005.
56. Lake AA, Townshend T, Alvanides S, et al. Diet, physical activity, sedentary behaviour and perceptions of the environment in young adults. *J Hum Nutr Diet*. 2009;22(5):444–54.
57. Department for Culture MS. *A New strategy for sport: consultation paper*. 2015.
58. Sport England. *Adult participation in 30 min, moderate intensity sport*: National (England), by key demographic subgroups: Excel table. 2005.



59. Sissons P, Jones K. Lost in transition? The changing labour market and young people not in employment, education or training: The Work Foundation. London: Lancaster University; 2012.
60. Yates S, Payne M. Not so NEET? a critique of the use of 'NEET' in setting targets for interventions with young people. *J Youth Stud.* 2006;9(3):329–44.
61. Scottish Government. The Scottish health survey 2012 edition volume 1: main report. 2013.
62. NHS Information Centre for Health and Social Care. Health survey for England - 2010, respiratory health: volume 2 methods and documentation. 2011.
63. NHS Information Centre for Health and Social Care. Health survey for England - 2011, health, social care and lifestyles: volume 2 methods and documentation. 2012.
64. NHS Information Centre for Health and Social Care. Health survey for England - 2012: volume 2 methods and documentation. 2013.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

