

RESEARCH NOTE

INTERNATIONAL MICROBIOLOGY (2007) 10:147-150
DOI: 10.2436/20.1501.01.21 ISSN: 1139-6709 www.im.microbios.org



Novel 10-bp deletion in the translational attenuator of a constitutively expressed *erm(A)* gene from *Staphylococcus epidermidis*

Leticia Millán,¹ Pilar Goñi,¹ Paula Cerdá,¹ M. Carmen Rubio,^{1,2}
Rafael Gómez-Lus^{1*}

¹Department of Microbiology, Zaragoza University School of Medicine, Spain. ²Microbiology Service, Lozano Blesa University Clinic Hospital, Zaragoza, Spain

Received 14 February 2007 · Accepted 2 May 2007

Summary. Alterations in the *erm(A)* regulatory region of six clinical isolates of *Staphylococcus epidermidis* and one of *Staphylococcus haemolyticus* displaying a constitutive resistance phenotype were investigated. A novel deletion of 10 bp with respect to the corresponding sequence of Tn554 was identified in the attenuator of a constitutively expressed *erm(A)* gene of one of the *S. epidermidis* isolates. Thus far, this is the smallest deletion conferring constitutive resistance in the translational attenuator of *erm(A)* in a naturally occurring *S. epidermidis* strain of human origin. [*Int Microbiol* 2007; 10(2):147-150]

Key words: *Staphylococcus epidermidis* · *erm(A)* regulatory region · deletions · macrolide, lincosamide, streptogramin B (MLS_B) resistance

Introduction

Resistance to macrolide, lincosamide, and streptogramin B antibiotics (MLS_B resistance) in staphylococci is mainly due to the dimethylation of an adenine residue (A2058) located in a highly conserved region of the 23S rRNA [6,24]. In staphylococci, four major methylase gene classes, *erm(A)*, *erm(B)*, *erm(C)*, and *erm(F)*, all of which play a role in MLS_B resistance [6,12,23], have been identified. Expression of the predominant *erm* genes in staphylococci, *erm(A)* and *erm(C)*, is either constitutive or inducible by 14- and 15-membered macrolides. Resistance is not related to the class of the *erm* determinant, but depends on the sequence of the regulatory region upstream of the methylase gene and is explained by a translation attenuation mechanism [6,13,24]. Constitutively

expressed *erm(A)* and *erm(C)* genes are of particular clinical relevance since they also confer resistance to 16-membered macrolides, lincosamides, streptogramin B antibiotics, and the new ketolide drugs. However, in inducible MLS_B resistance, the strains are resistant to 14- and 15-membered macrolides only, whereas 16-membered macrolides, lincosamides, streptogramin B antibiotics, and ketolides remain active [6,19]. The *erm(A)* gene, which is part of transposon Tn554, is usually integrated at a specific site (*att554*) in the *Staphylococcus aureus* chromosome. However, second insertions have also been described in chromosomal and plasmid sites, both in *S. aureus* and coagulase-negative staphylococci (CoNS) [6,9,10,21,22,25]. Structural alterations in the *erm(A)* regulatory region have been reported for constitutive mutants derived from in-vitro selection experiments and for constitutively expressed naturally occurring staphylococcal plasmids [9,14,25].

In this work, we studied alterations in the regulatory region of the constitutively expressed naturally occurring *erm(A)* gene from six clinical isolates of *Staphylococcus epidermidis* and one clinical isolate of *Staphylococcus haemolyticus*.

*Corresponding author: R. Gómez-Lus
Department of Microbiology
Zaragoza University School of Medicine
Domingo Miral, s/n
50009 Zaragoza, Spain
Tel. +34-976761692. Fax +34-976761693
E-mail: gomezlus@unizar.es

Materials and methods

Bacterial strains and culture conditions. Six clinical strains of *S. epidermidis* (0467, 4747, 6906, 6926, 7847 and 8377) and one of *S. haemolyticus* (2505) were collected. All strains exhibited a constitutive resistance phenotype and carried the *erm(A)* gene. The strains were isolated from the respiratory tracts of patients in the Microbiology Service of the University Clinical Hospital Lozano Blesa (Zaragoza, Spain) between October 2000 and November 2002. Initial identification was based on colony morphology, microscopy appearance, and agglutination tests with the Pastorex Staph Plus kit (Bio-Rad, La Coquette, France). Species were identified using the WIDER Computerized System (Dade Behring, West Sacramento, CA, USA). The isolates were cultured on Mueller-Hinton agar (MHA) (Bio-Rad, La Coquette, France) at 37°C and stored frozen (–80°C) in sterile skimmed milk.

Macrolide-lincosamide resistance testing. All strains had the constitutive resistance phenotype, as confirmed by a triple-disk diffusion technique carried out using a modification of the assay described by Seppälä et al. [18]. The disks contained erythromycin (15 µg), azithromycin (15 µg), clindamycin (2 µg) (Bio-Rad, La Coquette, France), and miocamycin (50 µg, Neo-Sensitabs, Taastrup, Denmark).

PCR amplification. The presence of the *erm(A)*, *erm(B)*, and *erm(C)* genes was tested by PCR, as reported by Sutcliffe et al. [20]. To detect structural alterations in the *erm(A)* and/or *erm(C)* regulatory regions, the PCR assays described by Werckenthin and Schwarz [25] and by Lodder et al. [8] were used. The expected PCR products were 592 bp for the *erm(A)* regulatory region and 295 bp for that of *erm(C)*. All PCR amplifications were carried out in a Perkin-Elmer Cetus DNA Thermal Cycler (Perkin-Elmer, Norwalk, CT, USA), and the products were analyzed by electrophoresis through 1.5% agarose gels. PCR reagents were purchased from Promega (Madison, WI, USA). Three of the PCR amplicons of the *erm(A)* regulatory region were purified (GFX PCR DNA and Gel Band Purification Kit, Amersham Biosciences, Freiburg, Germany) and then sequenced (DYEnamic ET Dye Terminator Cycle Sequencing Kit for MegaBACE DNA Analysis Systems, Amersham). The software BLAST was used to compare the sequences of the PCR products to that of the inducible *erm(A)* gene of Tn554, available at the website of the National Center for Biotechnology Information [<http://www.ncbi.nlm.nih.gov>]. A clinical control strain with an inducible phenotype, *S. aureus* 4850, which harbored *erm(A)* and *erm(C)* with a complete attenuator, was used. The *erm(A)* regulatory region of the control strain was identical to that of Tn554 (database accession no. X03216), whereas its *erm(C)* regulatory region was identical to that of plasmid pE5 carrying *erm(C)* (database accession no. M17990).

Nucleotide sequence accession number. The nucleotide sequence of the *erm(A)* regulatory region and partial *erm(A)* gene of strain *S. epidermidis* 6926 was submitted to the GenBank database and assigned accession no. DQ112022.

Results and Discussion

Analysis of the presence of the macrolide resistance genes *erm(A)*, *erm(B)*, and *erm(C)* showed that four *S. epidermidis* (strains 0467, 4747, 6926, and 7847) and one *S. haemolyticus* (strain 2505) harbored both *erm(A)* and *erm(C)*, whereas the two remaining *S. epidermidis* isolates (6906 and 8377) carried *erm(A)*, *erm(B)*, and *erm(C)*. Sequence analysis of the

PCR fragment obtained from *S. epidermidis* 6926 identified a novel 10-bp deletion in the *erm(A)* regulatory region. In *S. epidermidis* 8377, a 121-bp deletion was observed in that region. The remaining strains (*S. epidermidis* 0467, 4747, 6906, 7847, and *S. haemolyticus* 2505) contained neither deletions nor tandem duplications [14] in the regulatory region. *erm(A)* PCR amplicons of strains *S. epidermidis* 6926, *S. epidermidis* 8377, and *S. epidermidis* 6906 were sequenced.

The 10-bp deletion of *S. epidermidis* 6926 comprised nucleotides 5358–5367 (AGTTCGTTAT), which corresponded to part of the IR3 sequence located in the open reading frame (ORF) of the 19-amino acid (aa) peptide (Fig. 1, database accession no. DQ112022). Since truncated IR3 is not able to form a stable mRNA secondary structure with IR4, the latter will pair with IR5 and thereby render IR6 permanently accessible to ribosomes, allowing constitutive *erm(A)* gene expression to occur. In addition, this strain also harbored an *erm(C)* gene, but sequence analysis of the PCR amplicon corresponding to the gene's regulatory region showed that it shared 100% identity with the regulatory region of the *S. aureus* plasmid pE5, which carries *erm(C)* and confers inducible MLS_B resistance [11] (data not shown). To our knowledge, this is the first report of a 10-bp deletion in the regulatory region of an *erm(A)* gene in a *S. epidermidis* clinical strain with a constitutive resistance phenotype. Schmitz et al. [17] detected a 14-bp deletion that comprised part of the ORF of the 19-aa peptide and part of IR3 (nucleotides 5344–5357). The deletion confers constitutive resistance in *S. aureus* mutants—previously carrying an inducible *erm(A)* gene—selected in vitro in the presence of non-inducer antibiotics. Our work continues and extends this study as it describes a smaller deletion in this region from a naturally occurring *S. epidermidis* strain of human origin, which suggests selective pressure in nature due to the widespread use of MLS_B antibiotics.

The 121-bp deletion of strain *S. epidermidis* 8377 comprised the ORF of the 19-aa peptide as well as the region immediately downstream, which included the IR4 and IR5 sequences. This deletion, which has also been reported by others [14], renders IR6 always accessible to ribosomes, independently of the presence or absence of inducer.

The *erm(A)* regulatory region of *S. epidermidis* 6906 showed complete identity with that of Tn554. Contrary to previously described [9,17,25], point mutations that could have accounted for constitutive expression of *erm(A)* were not detected. This strain also harbored *erm(B)* and *erm(C)* determinants. PCR amplification and sequence of the *erm(C)* regulatory region showed complete identity with that of the sequence of plasmid pE5 (data not shown). Thus, constitutive expression of the methylase could be due to *erm(B)*, as the latter is located on Tn551 in staphylococci and expressed constitutively [2]. By contrast, in streptococci and certain

ring *in vivo* and *in vitro* after selection in the presence of non-inducer antibiotics have been reported. Non-inducing MLS_B antibiotics are still widely used in human and in veterinary medicine. The development of constitutively resistant strains will strongly reduce the efficacy of these drugs in the control of pathogenic staphylococci. Thus, the spectrum of resistance changes from limited resistance to 14- and 15-membered macrolides to extended resistance to 14-, 15-, and 16-membered macrolides as well as lincosamides and type B streptogramins. Consequently, the use of MLS_B antibiotics is not recommended for the control of staphylococci with an inducible phenotype [4,7–9,14–17,25,26]. Note also that alterations in the regulatory region of *erm(A)* are not only confined to staphylococci; rather, a 44-bp duplication, a 68-bp deletion, point mutations and a single base insertion have also been described in this region in several *Streptococcus pyogenes* clinical isolates displaying constitutive MLS_B resistance [1]. Thus, our results confirm that deletions in the *erm(A)* regulatory region not only are obtained in the laboratory but they also occur in nature, which suggests selective pressure resulting from the use of MLS_B antibiotics.

Acknowledgements. This study was supported by project FIS PI05/2310 (Ministerio de Sanidad y Consumo, Instituto Aragonés de Ciencias de la Salud, Spain), and project B24 (Departamento de Ciencia, Tecnología y Universidad, Diputación General de Aragón, Spain). L. Millán and P. Cerdá were the recipients of fellowships from the Diputación General de Aragón, Spain.

References

- Doktor SZ, Shortridge V (2005) Differences in the DNA sequences in the upstream attenuator region of *erm(A)* in clinical isolates of *Streptococcus pyogenes* and their correlation with macrolide/lincosamide resistance. *Antimicrob Agents Chemother* 49:3070-3072
- Eady EA, Ross JL, Tipper JL, Walters CE, Cove JH, Noble WC (1993) Distribution of genes encoding erythromycin ribosomal methylases and an erythromycin efflux pump in epidemiologically distinct groups of staphylococci. *J Antimicrob Chemother* 31:211-217
- Horinouchi S, Byeon WH, Weisblum B (1983) A complex attenuator regulates inducible resistance to macrolides, lincosamides, and streptogramin type B antibiotics in *Streptococcus sanguis*. *J Bacteriol* 154:1252-1262
- Lampson BC, Parisi JT (1986) Naturally occurring *Staphylococcus epidermidis* plasmid expressing constitutive macrolide-lincosamide-streptogramin B resistance contains a deleted attenuator. *J Bacteriol* 166:479-483
- Lampson BC, Parisi JT (1986) Nucleotide sequence of the constitutive macrolide-lincosamide-streptogramin B resistance plasmid pNE131 from *Staphylococcus epidermidis* and homologies with *Staphylococcus aureus* plasmids pE194 and pSN2. *J Bacteriol* 167:888-892
- Leclercq R, Courvalin P (1991) Bacterial resistance to macrolide, lincosamide, and streptogramin antibiotics by target modification. *Antimicrob Agents Chemother* 35:1267-1272
- Lodder G, Schwarz S, Gregory P, Dyke K (1996) Tandem duplication in *ermC* translational attenuator of the macrolide-lincosamide-streptogramin B resistance plasmid pSES6 from *Staphylococcus equorum*. *Antimicrob Agents Chemother* 40:215-217
- Lodder G, Werckenthin C, Schwarz S, Dyke K (1997) Molecular analysis of naturally occurring *ermC*-encoding plasmids in staphylococci isolated from animals with and without previous contact with macrolide/lincosamide antibiotics. *FEMS Immunol Med Microbiol* 18:7-15
- Murphy E (1985) Nucleotide sequence of *ermA*, a macrolide-lincosamide-streptogramin B determinant in *Staphylococcus aureus*. *J Bacteriol* 162:633-640
- Murphy E, Löfdahl S (1984) Transposition of Tn554 does not generate a target duplication. *Nature* 307:292-294
- Projan SJ, Monod M, Narayanan CS, Dubnau D (1987) Replication properties of pIM13, a naturally occurring plasmid found in *Bacillus subtilis*, and of its close relative pE5, a plasmid native to *Staphylococcus aureus*. *J Bacteriol* 169:5131-5139
- Roberts MC, Sutcliffe J, Courvalin P, Jensen LB, Rood J, Seppälä H (1999) Nomenclature for macrolide-lincosamide-streptogramin B resistance determinants. *Antimicrob Agents Chemother* 43:2823-2830
- Schmitz FJ, Verhoef J, Fluit AC, The Sentry Participants Group (1999) Prevalence of resistance to MLS antibiotics in 20 European university hospitals participating in the European SENTRY surveillance programme. *J Antimicrob Chemother* 43:783-792
- Schmitz FJ, Petridou J, Astfalk N, Scheuring S, Köhrer K, Verhoef J, Fluit AC, Schwarz S (2001) Structural alterations in the translational attenuator of constitutively expressed *erm(A)* genes in *Staphylococcus aureus*. *Antimicrob Agents Chemother* 45:1603-1604
- Schmitz FJ, Witte W, Werner G, Petridou J, Fluit AC, Schwarz S (2001) Characterization of the translational attenuator of 20 methicillin-resistant, quinupristin/dalfopristin-resistant *Staphylococcus aureus* isolates with reduced susceptibility to glycopeptides. *J Antimicrob Chemother* 48:931-942
- Schmitz FJ, Petridou J, Astfalk N, Köhrer K, Scheuring S, Schwarz S (2002) Molecular analysis of constitutively expressed *erm(C)* genes selected *in vitro* by incubation in the presence of the noninducers quinupristin, telithromycin, or ABT-773. *Microb Drug Resist* 8:171-177
- Schmitz FJ, Petridou J, Jagusch H, Astfalk N, Scheuring S, Schwarz S (2002) Molecular characterization of ketolide-resistant *erm(A)*-carrying *Staphylococcus aureus* isolates selected *in vitro* by telithromycin, ABT-773, quinupristin and clindamycin. *J Antimicrob Chemother* 49:611-617
- Seppälä H, Nissinen A, Yu Q, Huovinen P (1993) Three different phenotypes of erythromycin-resistant *Streptococcus pyogenes* in Finland. *J Antimicrob Chemother* 32:885-891
- Shortridge VD, Zhong P, Cao Z, Beyer JM, Almer LS, Ramer NC, Doktor SZ, Flamm RK (2002) Comparison of *in vitro* activities of ABT-773 and telithromycin against macrolide-susceptible and -resistant streptococci and staphylococci. *Antimicrob Agents Chemother* 46:783-786
- Sutcliffe J, Grebe T, Tait-Kamradt A, Wondrack L (1996) Detection of erythromycin-resistant determinants by PCR. *Antimicrob Agents Chemother* 40:2562-2566
- Thakker-Varia S, Jenssen WD, Moon-McDermott L, Weinstein MP, Dubin DT (1987) Molecular epidemiology of macrolides-lincosamides-streptogramin B resistance in *Staphylococcus aureus* and coagulase-negative staphylococci. *Antimicrob Agents Chemother* 31:735-743
- Tillotson LE, Jenssen WD, Moon-McDermott L, Dubin DT (1989) Characterization of a novel insertion of the macrolides-lincosamides-streptogramin B resistance transposon Tn554 in methicillin-resistant *Staphylococcus aureus* and *Staphylococcus epidermidis*. *Antimicrob Agents Chemother* 33:541-550
- Weisblum B (1995) Erythromycin resistance by ribosome modification. *Antimicrob Agents Chemother* 39:577-585
- Weisblum B (1995) Insights into erythromycin action from studies of its activity as inducer of resistance. *Antimicrob Agents Chemother* 39:797-805
- Werckenthin C, Schwarz S (2000) Molecular analysis of the translational attenuator of a constitutively expressed *erm(A)* gene from *Staphylococcus intermedius*. *J Antimicrob Chemother* 46:785-788
- Werckenthin C, Schwarz S, Westh H (1999) Structural alterations in the translational attenuator of constitutively expressed *ermC* genes. *Antimicrob Agents Chemother* 43:1681-1685