

RESEARCH ARTICLE

INTERNATIONAL MICROBIOLOGY (2009) 12:115-121
DOI: 10.2436/20.1501.01.88 ISSN: 1139-6709 www.im.microbios.org

Differential toxicity of antifungal protein AFP against mutants of *Fusarium oxysporum*

Magdalena Martín-Urdiroz,^{1,3} Ana L. Martínez-Rocha,^{1,3} Antonio Di Pietro,¹
Álvaro Martínez-del-Pozo,² M. Isabel G. Roncero^{1*}

¹Department of Genetics, University of Córdoba, Córdoba, Spain. ²Department of Biochemistry and Molecular Biology I, Faculty of Chemical Sciences, Complutense University of Madrid, Madrid, Spain. ³School of Biosciences, University of Exeter, Exeter, United Kingdom

Received 23 February 2009 · Accepted 15 May 2009

Summary. Antifungal protein (AFP) from *Aspergillus giganteus* was assayed for toxicity against the *Fusarium oxysporum* wild-type strain and mutants in genes involved in cell signaling ($\Delta pacC$, $pacC^c$ $\Delta fmk1$) or cell-wall biogenesis ($\Delta chsV$, $\Delta chs7$, $\Delta gas1$). The mutants were classified into two groups according to their sensitivity to AFP: $\Delta pacC$, $\Delta gas1$ and $\Delta chs7$, which were significantly more resistant to AFP than the wild-type, and $pacC^c$, $\Delta fmk1$ and $\Delta chsV$, which were more sensitive. Western blot analysis revealed increased binding of AFP to the three resistant mutants, $\Delta pacC$, $\Delta gas1$ and $\Delta chs7$, but also to $\Delta chsV$, indicating that differential binding may not be a key determinant for sensitivity. Addition of Ca^{2+} or K^+ dramatically reduced antifungal activity and binding of AFP, suggesting that these cations compete for the same targets as AFP at the surface of the fungal cell. [*Int Microbiol* 2009; 12(2):115-121]

Keywords: *Fusarium oxysporum* · antifungal protein · chitin synthase · cell wall

Introduction

The control of fungal infections is largely based on the use of chemicals with a more or less narrow spectrum of action. The emergence of resistant pathogenic strains upon prolonged application has triggered considerable interest in the isolation of new antifungal compounds that are more specific and with fewer side effects on human health and the environment. By exploiting the differential activity of a given antifungal compound towards a fungal wild-type strain and defined mutants, insight into its mechanism of action can be gained [8], allowing the development of even more targeted and efficient products.

Antimicrobial peptides are host defense molecules that are ubiquitous in multicellular plants and animals, as well as in many single cell organisms [32]. Most antimicrobial peptides are cationic molecules, with a net positive charge, that bind to microbial membranes, which are generally negatively charged. The *Aspergillus giganteus* antifungal protein (AFP), a small polypeptide of 51 amino acids, is secreted as an inactive precursor containing six extra amino acid residues at the NH_2 -terminal end, which are processed in the extracellular medium [16]. AFP has been assayed against a wide variety of microorganisms, including prokaryotes and eukaryotes, and its inhibitory effect on the growth of many filamentous fungi has been well documented [12,29,31]. Interestingly, AFP is not active against *Penicillium chrysogenum* and *Aspergillus niger*, two organisms producing antifungal proteins similar to AFP (PAF and Anapf, respectively) [29]. AFP binds to phospholipid membranes [12] and to nucleic acids [17]. The in vitro interaction with nucleic acids, resulting from the oligonucleotide/oligosaccharide (OB) fold structure of AFP, suggests that this property might be related to its biological activity [17]. Nonetheless, at present, the

*Corresponding author: M.I.G. Roncero
Departamento de Genética
Campus de Rabanales. Universidad de Córdoba
Edificio Gregor Mendel, Campus de Rabanales
14071 Córdoba, Spain
Tel. +34-957218981. Fax +34-957212072
E-mail: ge1gorom@uco.es

antifungal mode of action of AFP remains largely unknown (for a recent review see [19]).

Fusarium oxysporum is a soil-borne plant pathogen that causes vascular wilt disease on many different plant species worldwide [6]. With the exception of grasses and most tree crops, few of the widely cultivated crops are not hosts to a pathogenic form of this species. Isolates from *F. oxysporum* have been classified into more than 120 different formae speciales according to their host range. More recently, the species has also been reported as an emerging human pathogen in immunocompromised individuals [23,25].

Successful infection of the host plant by *F. oxysporum* requires a series of highly regulated processes. Several groups, including our own, are studying this vascular wilt fungus in attempts to unravel key mechanisms of the fungal infection process by creating mutants in specific genes through targeted gene knockout. This approach has begun to shed light on the mechanisms underlying infection and the development of vascular wilt disease [6]. The targeted inactivation of *fmk1*, which encodes a mitogen-activated protein kinase (MAPK) orthologous to yeast Fus3/Kss1, produced mutants that were unable to penetrate the roots of tomato plants and failed to cause disease symptoms [5]. By contrast, mutants carrying loss-of-function ($\Delta pacC$) or gain-of-function ($pacC^C$) alleles of the pH signal transcription factor PacC showed increased or reduced virulence, respectively, compared to the wild-type strain [3].

A different mechanism of tolerance to plant defense compounds involves the fungal cell wall. The *F. oxysporum* wall contains an inner layer of chitin and glucan that acts as a support for an external layer of glycoproteins [27]. Chitin synthases catalyze the polymerization of β -1,4-linked *N*-acetyl-D-glucosamine to chitin, a major structural component of the fungal cell wall [26]. Through random insertional mutagenesis screening, a mutant showing complete loss of virulence on tomato was isolated, and the affected gene *chsV* was found to encode a class V chitin synthase [14]. A likely mechanism of ChsV in virulence is to prevent diffusion of plant antifungal compounds to their cellular targets. The *chs7* gene encodes a chaperone-like protein of *F. oxysporum* orthologous to *Saccharomyces cerevisiae* Chs7p [30] and is required for full virulence on tomato plants [15]. The *gas1* gene encodes a β -1,3-glucanotransferase of *F. oxysporum*, which is involved in the processing of β -1,3-glucan, another major component of the fungal wall. Mutants lacking Gas1 showed dramatically reduced virulence on tomato plants, both in a root infection assay and in a fruit invasion model [2].

In this study, we used a well-characterized collection of *Fusarium* mutants to test their sensitivity towards AFP. The observation that AFP is not inhibitory to yeasts and has dif-

ferential effects on filamentous fungal species [29] strongly suggests that the compound has specific target(s) in the fungal cell. *Fusarium* belongs to the group of fungi that are susceptible to AFP [11,12,31]. The goal of this study was to gain further insight into the mode of action of AFP by comparing its inhibitory effect against the *F. oxysporum* wild-type strain and a collection of defined gene knockout mutants. The antifungal activity of AFP was previously reported to be sensitive to cations [29]. Therefore, its inhibitory effect against resistant and sensitive *F. oxysporum* mutants was tested in the presence of 0.1 M CaCl₂ or 0.2 M KCl.

Materials and methods

Protein purification. AFP was purified to homogeneity from *Aspergillus giganteus* cultures as described previously [12,16,17,24]. Protein concentration was determined from absorbance measurements based on an E^{0.1%} (1-cm optical path) at 278 nm of 1.76 [12], using a Nanodrop ND-1000 spectrophotometer.

Fungal strains and culture conditions. Fungal strains used in this work are listed in Table 1. *Fusarium oxysporum* f. sp. *lycopersici* wild-type strain 4287 and mutants derived thereof were stored as microconidial suspensions with 30% glycerol at -80°C . Fungal strains were grown in potato dextrose broth (PDB) on a rotary shaker at 170 rpm and 28°C as described [7].

AFP susceptibility assays. Microconidia were collected from PDB cultures by filtration and centrifugation, washed in sterile water, counted, and transferred to fresh PDB or YPG medium (pH 4.5) [11]. The susceptibility of the six *Fusarium oxysporum* f. sp. *lycopersici* mutants to AFP was compared to that of the wild-type. Aliquots containing 5×10^5 or 10^3 microconidia were added to 200 μl PDB or YPG, respectively. When indicated, the media contained AFP at concentrations ranging from 0 to 4 $\mu\text{g/ml}$ in PDB or from 0 to 200 $\mu\text{g/ml}$ in YPG. The inhibitory effect of AFP on *Fusarium* growth in YPG was evaluated after 42 h at 28°C by determining the optical density at 600 nm (OD₆₀₀) in a Tecan SpectraFluor Plus spectrophotometer. Experiments were carried out in triplicate in PDB or in duplicate in YPG and repeated at least twice, with similar results. Growth was expressed as percentage of the control without AFP (100%) for each strain.

Table 1. *Fusarium oxysporum* strains used in this study

Strain	Genotype	Source/reference
4287 (FGSC 9935)	<i>F. o. f. sp. lycopersici</i> , wild type	Fungal Genetics Stock Center
$\Delta fmk1$	<i>fmk1::Phleo^R</i>	[5]
$\Delta chsV$	<i>chsV::Hyg^R</i>	[14]
$\Delta pacC$	<i>pacC::Hyg^R</i>	[3]
<i>pacC^C</i>	<i>pacC^C, Hyg^R</i>	[3]
$\Delta chs7$	<i>chs7::Hyg^R</i>	[2]
$\Delta gas1$	<i>gas1::Hyg^R</i>	[15]

Assays for the effect of cations on AFP. The effect of cations on the inhibitory activity of AFP towards *F. oxysporum* was tested in PDB supplemented with 0.1 M CaCl₂ or 0.2 M KCl and containing 0–4 µg AFP/ml. Fungal growth was determined after 36 h as described above and expressed as percentage of the control (PDB with 0.1 M CaCl₂ or 0.2 M KCl and without AFP). Experiments were performed in duplicate and repeated at least twice with similar results.

Western blot analyses. To determine the binding capacity of AFP to different *F. oxysporum* strains, 2.5 × 10⁶ microconidia/ml of the wild-type strain were pre-germinated in PDB at 28°C, 170 rpm during 4 h. After this time, different concentrations of AFP (0.02–0.5 µg/ml) were added to 100 µl aliquots of the germling suspensions, and samples were incubated at 170 rpm, 28°C, for 2 h. After centrifugation of the samples, the supernatants were removed carefully and concentrated to 20 µl in a Speed-Vac SVC100 (Savant). To quantify unbound AFP, the samples were separated by SDS-PAGE. As a control, 20 µl of the different AFP concentrations without microconidia were loaded.

AFP was detected by Western blot hybridization using a primary rabbit polyclonal anti-AFP antibody [16] at a dilution of 1/3000. After a 1-h hybridization at room temperature, the blots were washed and IgG-conjugated anti-rabbit secondary antibody (Cell Signaling Technology, Beverly, MA, USA) was added at a 1/2000 dilution, according to the instructions of the manufacturer.

The effect of Ca²⁺ on AFP binding to microconidia of the different strains was determined by pre-germinating 2.5 × 10⁶ microconidia/ml in PDB at 2°C and 170 rpm for 4 h, followed by the addition of 0.1 M CaCl₂ (final concentration) and a 5-min incubation. The pre-germinated microconidia were then washed with water and incubated with 0.25 µg AFP/ml at 170 rpm, 28°C, for 2 h. After centrifugation of the samples, supernatants were removed, concentrated to 20 µl in a Speed-Vac, and subjected to SDS-PAGE to determine the unbound AFP fraction by Western blot as described above.

Results

Different sensitivities to AFP of *Fusarium oxysporum* mutants. The mutants could be divided into two groups according to their AFP sensitivity (Fig. 1). The first group consisted of mutants $\Delta pacC$, $\Delta gas1$, and $\Delta chs7$, which were significantly more resistant to AFP than the wild-type strain. For example, growth of the wild-type strain in the presence of 0.75 µg AFP/ml was inhibited by 90% compared to the untreated control, whereas in the resistant mutants inhibition was only between 30 and 50% (Fig. 1A). Relative differences within the group of resistant mutants were also observed. The growth of $\Delta pacC$ was almost totally inhibited by 100 µg AFP/ml, whereas $\Delta gas1$ and $\Delta chs7$ were less affected, even at concentrations of 200 µg/ml (Fig. 2). The second group of mutants consisted of $pacC^c$, $\Delta fmk1$, and $\Delta chsV$, which showed higher sensitivity to AFP than the wild-type strain (Fig. 1B). Within this group, $\Delta chsV$ was the most sensitive strain, followed by $\Delta fmk1$ and $pacC^c$ (Fig. 2). Results similar to those in YPG were also obtained in PDB medium. Resistant mutants $\Delta pacC$, $\Delta gas1$, and $\Delta chs7$ showed no decrease in growth at 0.5 µg AFP/ml,




Fig. 1. Antifungal activity of AFP against *Fusarium oxysporum* strains. Results for each strain are expressed as percentage of A_{600} in the control without AFP. (A) Wild-type strain (×) and the resistant mutants $\Delta pacC$ (closed circles), $\Delta gas1$ (closed squares) and $\Delta chs7$ (closed triangles). (B) Wild-type strain (×) and the sensitive mutants $pacC^c$ (open circles), $\Delta fmk1$ (open triangles) and $\Delta chsV$ (open squares). Mean values and standard deviations were calculated from two independent experiments.

whereas sensitive strains $pacC^c$, $\Delta fmk1$ and $\Delta chsV$ hardly grew at all at this concentration (results not shown).

All mutants except $\Delta fmk1$ had higher membrane permeability than the wild-type strain when incubated with 0.2 µM Sytox Green. Fluorescence microscopy analysis did not show detectable differences in chitin distribution between the different mutants and the wild-type strain. Similarly, the distribution of cell-wall mannoproteins, as revealed by Concanavalin A staining, did not differ significantly between mutant and wild-type strains (data not shown).




Fig. 2. Growth of *Fusarium oxysporum* strains in the presence of increasing concentrations of AFP ($\mu\text{g/ml}$). Lack of turbidity in the medium indicates the absence of hyphal growth.

Ca^{2+} and K^{+} cations abolish the antifungal activity of AFP. The addition of either cation strongly reduced or almost abolished antifungal activity of AFP against the wild-type strain and the sensitive mutants $\Delta fmk1$ and $\Delta chsV$ (Fig. 3). CaCl_2 was more effective than KCl in

reducing AFP antifungal activity. No effect of the two cations was detected in any of the resistant mutants because even the highest concentration of AFP used in the experiment ($4 \mu\text{g/ml}$) had no inhibitory activity (the $\Delta chs7$ mutant is shown in Fig. 3 as a representative example).




Fig. 3. Suppression of the antifungal activity of AFP on *Fusarium oxysporum* by cations. Aliquots of 5×10^5 microconidia of the indicated strains were added to 96-well microtiter plates containing PDB medium supplemented with the indicated concentrations of AFP in the absence (closed diamonds) or presence of 0.1 M CaCl_2 (open triangles) or 0.2 M KCl (open circles), and incubated for 36 h at 28°C . Hyphal growth was measured by determining optical density at 600 nm. Results for each strain are expressed as percentage of A_{600} in the control without AFP. Mean values and standard deviations were calculated from two independent experiments.




Fig. 4. Binding of AFP does not correlate strictly with sensitivity. (A) Solutions containing the indicated concentrations of AFP were incubated for 2 h in the absence (–) or presence (+) of 2.5×10^6 microconidia of the *F. oxysporum* wild type strain/ml. Conidia were removed by centrifugation and supernatants fractionated by SDS PAGE, blotted to membranes, and hybridized with anti-AFP antibody. (B) Solutions containing 0.25 mg AFP/ml AFP were incubated for 2 h with 2.5×10^6 microconidia of the indicated strains/ml and analyzed by Western blot as indicated in (A), but using a longer exposure time. The absence of a hybridizing band in the supernatant indicates increased binding of AFP to the fungal surface.

Ability of AFP to bind to *F. oxysporum*, and inhibition of binding by Ca^{2+} . When decreasing concentrations of AFP were added to pre-germinated microconidia of the wild-type strain, depletion of AFP from the supernatant, due to binding of the protein to fungal cells, was clearly detected at concentrations of 0.25 $\mu\text{g/ml}$ and 0.1 $\mu\text{g/ml}$ (Fig. 4A). Accordingly, 0.25 $\mu\text{g/ml}$ was then used to determine AFP binding to the different mutant strains. As seen in Fig. 4B, all resistant mutants showed significantly increased binding to AFP compared to binding by the wild-type strain whereas binding by the two sensitive mutants ΔchsV and Δfmk1 differed. Specifically, the ΔchsV strain bound more avidly to AFP, while binding of Δfmk1 was comparable to that observed for the wild-type. These results suggest that binding of AFP to the fungal spore is not a key determinant of sensitivity or resistance to this antifungal peptide.

As shown above, the antifungal activity of AFP was abolished in the presence of cations such as Ca^{2+} and K^{+} . When 0.25 μg AFP/ml was added after incubation of pre-germinated microconidia with 0.1 M CaCl_2 , significantly more unbound AFP was detected in both classes of mutants as well as in the wild-type strain (Fig. 5), suggesting that cations interfere with AFP binding to *F. oxysporum*.

Discussion

AFP is a naturally derived molecule that efficiently inhibits the growth of a wide range of filamentous fungi, including




Fig. 5. Ca^{2+} inhibits AFP binding to *Fusarium oxysporum* microconidia. Solutions containing 0.25 mg AFP/ml were incubated for 2 h with 2.5×10^6 microconidia the indicated strains /ml in the absence (–) or presence (+) of 0.1 M CaCl_2 and analyzed by Western blot as indicated in Fig. 4.

agronomically important plant pathogens such as *Magnaporthe grisea* [21], *Botrytis cinerea* [20], *F. oxysporum* [11, this work] and others [31], as well as the human pathogen *A. fumigatus* [29]. Its fungitoxic properties and apparent lack of cytotoxicity towards mammalian cells [28] make AFP a promising candidate for antifungal intervention. AFP expression in transgenic rice plants, either from a constitutive or a pathogen-inducible promoter, has been shown to confer enhanced resistance to rice blast disease [4,22]. Besides AFP, other small fungal proteins with similar characteristics have been identified, including PAF from *Penicillium chrysogenum* [18] and Anafp from *A. niger* [10]. This family of AFP-like proteins represents a promising and largely untapped resource for agronomical and medical applications [19].

An essential prerequisite for exploiting the biotechnological potential of AFP is the precise elucidation of its cellular targets and mode of action. A number of mechanisms have been proposed for AFP's antifungal activity, including binding and destabilization of the plasma membrane [21,29], inhibition of chitin synthesis [11], nuclear localization and binding to nucleic acids [17,21], or a combination thereof. So far, none of these mechanisms has been conclusively linked to antifungal activity [19], and other modes of action therefore cannot be ruled out.

In the present study, significant differences in strain susceptibility were noted. For instance, *F. oxysporum* mutants lacking the β -1,3-glucanosyltransferase Gas1 or the chaperone-like protein Chs7 were resistant to AFP concentrations as high as 200 $\mu\text{g/ml}$, while growth of the wild-type was almost completely inhibited by 10 $\mu\text{g/ml}$ (see Fig. 2). By contrast, susceptible mutants lacking the MAPK Fmk1 or the chitin synthase ChsV were inhibited by 2 and 1 μg AFP/ml, respectively. The increased susceptibility of the ΔchsV mutant was surprising, since a previous study had reported enhanced resistance of this strain to AFP [11]. The reason for this discrepancy is unclear, but we consider it unlikely that it is due to differences in methodology, because the experimental con-

ditions (microtiter plates, OD₆₀₀) and the medium (YPG) used in the present work were comparable to those of the previous study. To confirm the results obtained in YPG, the experiments were repeated in a different medium (PDB), yielding essentially the same results. Thus the higher sensitivity of the *Fusarium* Δ *chsV* mutant to AFP appears to be significant. Note that Δ *chsV* was previously shown to have increased sensitivity to the structurally unrelated plant saponin α -tomatine [14].

Our results demonstrate that mutations in a single gene can dramatically affect the interaction between AFP and its unknown cellular target(s). However, since the mutants tested in this study have mostly pleiotropic phenotypes, at present we can only speculate on the mechanisms underlying the differences in AFP susceptibility/resistance. A feature common to these strains is the altered structure of the cell wall. Some of the mutations (Δ *chsV*, Δ *gas1*, Δ *chs7*) directly affect genes involved in cell wall biogenesis [2,14,15], while others (Δ *fmk1*, Δ *pacC*) target signaling components that regulate a wide array of cellular functions, including cell-wall structure [1,5]. The results from Western analysis, showing that AFP binds differentially to the surface of these strains, support the idea that differences in sensitivity between the mutants are related to cell-wall alterations. The fact that no clear correlation between sensitivity and binding affinity was detected—both resistant (Δ *gas1*, Δ *chs7*) and sensitive (Δ *chsV*) mutants showed increased binding to AFP—suggests that binding itself is not a major determinant for AFP activity, although it may be a prerequisite for inhibition.

Why mutations in certain cell-wall genes (*gas1*, *chs7*) lead to enhanced AFP resistance while others (*chsV*) increase susceptibility is currently an open question. One possible explanation is that some of these genes carry out divergent functions in cell-wall architecture and maintenance. In the Δ *gas1* mutant, which is highly resistant to AFP, transcript levels of the *chsV* gene are much higher than in wild-type cells [2]. If increased expression of *chsV* were related to AFP resistance, then a mutant lacking *chsV* should be more sensitive to the antifungal peptide. In this context, it is worth noting that the Δ *gas1* mutant is also highly resistant to proteolytic enzymes [2], whereas Δ *fmk1* and Δ *chsV* are more sensitive than the wild-type strain (Martínez-Rocha et al., unpublished data). The structural bases for these differences are currently unknown, but sensitivity to cell-wall-degrading enzymes and to AFP might be functionally related. In fact, AFP can bind chitin in vitro and, similar to certain chitinases, harbors a putative chitin-binding domain at its N-terminus [11,13].

To determine whether cations inhibited the binding of AFP to the different fungal strains, the amount of non-bound AFP was assayed by Western blot analysis in the absence and presence of 0.1 M CaCl₂ in the wild-type, two resistant

strains (Δ *gas1* and Δ *chs7*), and one sensitive (Δ *fmk1*) strain. The results showed that the antifungal activity of AFP against the wild-type strain and the susceptible mutants was essentially abolished in the presence of cations (Ca²⁺, K⁺). This finding confirms those of a previous report showing that the growth inhibitory activity of AFP, as well as its membrane permeabilization effect, was sensitive to cations [29]. Moreover, our Western blot data demonstrate that cations interfere with AFP binding to the fungal surface. Since AFP itself also has a positive net charge, cations might exert their inhibitory effect on AFP activity by competing for putative binding sites at the fungal cell surface.

In summary, the present study shows that mutations in single genes have significant effects on the sensitivity and binding of fungal strains to the antifungal protein AFP. Pinpointing the molecular bases underlying these differences should provide further insight into the antifungal mode of action of AFP and extend our understanding of the biological role of this conserved family of proteins in filamentous fungi.

Acknowledgements. The authors gratefully acknowledge Esther Martínez for her skillful technical assistance (University of Córdoba). This research was supported by grants BIO2007-62661 and BFU2006-04404 from the Spanish Ministry of Science and Innovation, CVI-138 from the Autonomous Government of Andalusia, and Programa Propio de Investigación from the University of Córdoba. M.M.U. and A.M.R. had PhD fellowships from the Spanish Ministry of Education and Science.

References

- Caracuel Z, Casanova C, Roncero MIG, Di Pietro A, Ramos J (2003) pH response transcription factor PacC controls salt stress tolerance and expression of the P-Type Na⁺-ATPase Ena1 in *Fusarium oxysporum*. *Eukaryot Cell* 2:1246-1252
- Caracuel Z, Martínez-Rocha AL, Di Pietro A, Madrid MP, Roncero MIG (2005) *Fusarium oxysporum gas1* encodes a putative β -1,3-glucanoyltransferase required for virulence on tomato plants. *Mol Plant-Microbe Interact* 18:1140-1147
- Caracuel Z, Roncero MIG, Espeso EA, González-Verdejo CI, García-Maceira FI, Di Pietro A (2003) The pH signalling transcription factor PacC controls virulence in the plant pathogen *Fusarium oxysporum*. *Mol Microbiol* 48:765-779
- Coca M, Bortolotti C, Rufat M, et al. (2004) Transgenic rice plants expressing the antifungal AFP protein from *Aspergillus giganteus* show enhanced resistance to the rice blast fungus *Magnaporthe grisea*. *Plant Mol Biol* 54:245-259
- Di Pietro A, García-Maceira FI, Mègelecz E, Roncero MIG (2001) A MAP kinase of the vascular wilt fungus *Fusarium oxysporum* is essential for root penetration and pathogenesis. *Mol Microbiol* 39:1140-1152
- Di Pietro A, Madrid MP, Caracuel Z, Delgado-Jarana J, Roncero MIG (2003) *Fusarium oxysporum*: exploring the molecular arsenal of a vascular wilt fungus. *Mol Plant Pathol* 4:315-326
- Di Pietro A, Roncero MIG (1998) Cloning, expression, and role in pathogenicity of *pgl* encoding the major extracellular endopolysaccharuronase of the vascular wilt pathogen *Fusarium oxysporum*. *Mol Plant-Microbe Interact* 11:91-98

8. Fernández-Ortuño D, Torés JA, de Vicente A, Pérez-García A (2008) Mechanisms of resistance to Qol fungicides in phytopathogenic fungi. *Int Microbiol* 11:1-9
9. García-Pedrajas MD, Roncero MIG (1996) A homologous and self-replicating system for efficient transformation of *Fusarium oxysporum*. *Curr Genet* 29:191-198
10. Gun Lee D, Shin SY, Maeng CY, Jin ZZ, Kim KL, Hahn KS (1999) Isolation and characterization of a novel antifungal peptide from *Aspergillus niger*. *Biochem Biophys Res Commun* 263:646-651
11. Hagen S, Marx F, Ram AF, Meyer V (2007) The antifungal protein AFP from *Aspergillus giganteus* inhibits chitin synthesis in sensitive fungi. *Appl Environ Microbiol* 73:2128-2134
12. Lacadena J, Martínez del Pozo A, Gasset M, et al. (1995) Characterization of the antifungal protein secreted by the mould *Aspergillus giganteus*. *Arch Biochem Biophys* 324:273-281
13. Liu RS, Huang H, Yang Q, Liu WY (2002) Purification of alpha-sarcin and an antifungal protein from mold (*Aspergillus giganteus*) by chitin affinity chromatography. *Protein Expr Purif* 25:50-58
14. Madrid MP, Di Pietro A, Roncero MIG (2003) Class V chitin synthase determines pathogenesis in the vascular wilt fungus *Fusarium oxysporum* and mediates resistance to plant defence compounds. *Mol Microbiol* 47:257-266
15. Martín-Urdíroz M, Madrid MP, Roncero MIG (2004) Role of chitin synthase genes in *Fusarium oxysporum*. *Microbiology* 150:3175-3187
16. Martínez-Ruiz A, Martínez del Pozo A, Lacadena J, Mancheño JM, Onaderra J, Gavilanes JG (1997) Characterization of a natural larger form of the antifungal protein (AFP) from *Aspergillus giganteus*. *Biochim Biophys Acta* 1340:81-87
17. Martínez del Pozo A, Lacadena V, Mancheño JM, Olmo N, Onaderra M, Gavilanes JG (2002) The antifungal protein AFP of *Aspergillus giganteus* is an oligonucleotide/oligosaccharide binding (OB) fold-containing protein that produces condensation of DNA. *J Biol Chem* 277:46179-46183
18. Marx F, Haas H, Reindl M, Stoffler G, Lottspeich F, Redl B (1995) Cloning, structural organization and regulation of expression of the *Penicillium chrysogenum paf* gene encoding an abundantly secreted protein with antifungal activity. *Gene* 167:167-171
19. Meyer V (2008) A small protein that fights fungi: AFP as a new promising antifungal agent of biotechnological value. *Appl Microbiol Biotechnol* 78:17-28
20. Moreno AB, Martínez del Pozo A, Borja M, San Segundo B (2003) Activity of the antifungal protein from *Aspergillus giganteus* against *Botrytis cinerea*. *Phytopathology* 93:1344-1353
21. Moreno AB, Martínez del Pozo A, San Segundo B (2006) Biotechnologically relevant enzymes and proteins: antifungal mechanism of the *Aspergillus giganteus* AFP against the rice blast fungus *Magnaporthe grisea*. *Appl Microbiol Biotechnol* 72:883-895
22. Moreno AB, Peñas G, Rufat M, Bravo JM, Estopà M, Messeguer J, San Segundo B (2005) Pathogen-induced production of the antifungal AFP protein from *Aspergillus giganteus* confers resistance to the blast fungus *Magnaporthe grisea*. *Mol Plant-Microbe Interact* 18:960-972
23. Nucci M, Anaissie E (2007) *Fusarium* infections in immunocompromised patients. *Clin Microbiol Rev* 20:695-704
24. Olson BH, Goerner GL (1965) Alpha sarcin, a new antitumor agent. I. Isolation, purification, chemical composition, and the identity of a new amino acid. *Appl Microbiol* 13:314-321
25. Ortoneda M, Guarro J, Madrid MP, Caracuel Z, Roncero MIG, Mayayo E, Di Pietro (2004) *Fusarium oxysporum* as a multihost model for the genetic dissection of fungal virulence in plants and mammals. *Infect Immun* 72:1760-1766
26. Roncero C (2002) The genetic complexity of chitin synthesis in fungi. *Curr Genet* 41:367-378
27. Schoffemeer EA, Klis FM, Sietsma JH, Cornelissen BJ (1999) The cell wall of *Fusarium oxysporum*. *Fungal Genet Biol* 27:275-282
28. Szappanos H, Szigeti GP, Pal B, et al. (2006) The antifungal protein AFP secreted by *Aspergillus giganteus* does not cause detrimental effects on certain mammalian cells. *Peptides* 27:1717-1725
29. Theis T, Wedde M, Meyer V, Stahl U (2003) The antifungal protein from *Aspergillus giganteus* causes membrane permeabilization. *Antimicrob Agents Chemother* 47:588-593
30. Trilla JA, Cos T, Duran A, Roncero C (1997) Characterization of *CHS4* (*CAL2*), a gene of *Saccharomyces cerevisiae* involved in chitin biosynthesis and allelic to *SKT5* and *CSD4*. *Yeast* 13:795-807
31. Vila L, Lacadena J, Fontanet P, Martínez del Pozo A, San Segundo B (2001) A protein from the mold *Aspergillus giganteus* is a potent inhibitor of fungal plant pathogens. *Mol Plant-Microbe Interact* 14:1327-1331
32. Zasloff M (2002) Antimicrobial peptides of multicellular organisms. *Nature* 415:389-395