

## RESEARCH REVIEW

INTERNATIONAL MICROBIOLOGY (2007) 10:233-243  
DOI: 10.2436/20.1501.01.32 ISSN: 1139-6709 [www.im.microbios.org](http://www.im.microbios.org)



# Hydration of vinyl ether groups by unsaturated glycoside hydrolases and their role in bacterial pathogenesis

Wataru Hashimoto,<sup>1\*</sup> Takafumi Itoh,<sup>1</sup> Yukie Maruyama,<sup>1</sup> Bunzo Mikami,<sup>2</sup>  
Kousaku Murata<sup>1</sup>

<sup>1</sup>Division of Food Science and Biotechnology, Graduate School of Agriculture, Kyoto University, Uji, Kyoto, Japan.

<sup>2</sup>Division of Applied Life Sciences, Graduate School of Agriculture, Kyoto University, Uji, Kyoto, Japan

Received 15 September 2007 · Accepted 30 November 2007

**Summary.** Many pathogenic microorganisms invade mammalian and/or plant cells by producing polysaccharide-degrading enzymes (lyases and hydrolases). Mammalian glycosaminoglycans and plant pectins that form part of the cell surface matrix are typical targets for these microbial enzymes. Unsaturated glycoside hydrolase catalyzes the hydrolytic release of an unsaturated uronic acid from oligosaccharides, which are produced through the reaction of matrix-degrading polysaccharide lyase. This enzymatic ability suggests that unsaturated glycoside hydrolases function as virulence factors in microbial infection. This review focuses on the molecular identification, bacterial distribution, and structure/function relationships of these enzymes. In contrast to general glycoside hydrolases, in which the catalytic mechanism involves the retention or inversion of an anomeric configuration, unsaturated glycoside hydrolases uniquely trigger the hydrolysis of vinyl ether groups in unsaturated saccharides but not of their glycosidic bonds. [*Int Microbiol* 2007; 10(4):233-243]

**Key words:** bacterial lyases · bacterial hydrolases · glycoside hydrolase family · glycosaminoglycan · pathogenic/saprophytic bacteria

## Introduction

The adhesion of bacteria to eukaryotic cells may be a primary mechanism of bacterial infection, in addition to playing a role in secondary effects of the infectious process. Polysaccharides, including glycosaminoglycans and pectins, that form part of the cell surface matrix are typical targets of microbial pathogens that invade host cells, and many specific interactions between pathogens and these polysaccharides have been described [16]. In this context, bacterial polysaccharide lyases function as virulence factors.

Several different enzymes were found to act on the unsaturated saccharides produced through the reactions of polysaccharide lyases. These enzymes included other types of lyases as well as hydrolases. For example, *Erwinia chrysanthemi* oligogalacturonate lyase (Ogl) [48] and *Sphingomonas* sp. A1 oligoalginate lyase (A1-IV) [12] degrade pectin- and alginate-unsaturated oligosaccharides produced through reactions of pectin and alginate lyases. Ogl and A1-IV release an unsaturated monosaccharide and unsaturated saccharide from substrate oligosaccharides through a  $\beta$ -elimination reaction. A novel glycoside hydrolase acting on unsaturated saccharides has been found in *Bacillus* sp. strain GL1 [11] and *Flavobacterium heparinum*, also called *Pedobacter heparinus* [27]. The enzyme specifically hydrolyzes the glycosidic bond in unsaturated saccharides between the unsaturated glucuronyl residue ( $\Delta$ GlcA) at the nonreducing termi-

\*Corresponding author: W. Hashimoto  
Division of Food Science and Biotechnology  
Graduate School of Agriculture, Kyoto University  
Uji, Kyoto 611-0011, Japan  
Tel. +81-774383756. Fax +81-774383767  
Email: [whasimot@kais.kyoto-u.ac.jp](mailto:whasimot@kais.kyoto-u.ac.jp)

nus and the saccharide linked to the residue, with subsequent release of  $\Delta$ GlcA and saturated saccharide from substrate oligosaccharides. The enzyme was thus designated an unsaturated glucuronyl hydrolase (UGL) [11]. Although the enzyme showed no homology with other proteins in DNA and protein databases, as determined following the cloning and sequencing of the enzyme gene in *Bacillus* [11], recent complete genome sequence analyses indicate that homologous genes are encoded in pathogenic bacteria, such as streptococci and vibrio [37]. In addition to UGL, an unsaturated galacturonyl hydrolase (UGH) has recently been identified in *Bacillus subtilis* [20]. In unsaturated saccharides, the enzyme specifically hydrolyzes the glycosidic bond between the unsaturated galacturonyl residue ( $\Delta$ GalA) at the nonreducing terminus and the saccharide linked to the residue.

UGL and UGH belong to two huge novel families GH-88 and GH-105, respectively, which (as of Sept. 2007) comprise 76 (GH-88) and 61 (GH-105) bacterial and fungal members. Recent structural analyses showed that UGL and UGH share a unique catalytic mechanism, one that differs from the retention or inversion mechanism of general glycoside hydrolases [17,21]. Here, we detail the enzyme characteristics, biological roles, and structure/function relationships of the GH-88 and GH-105 families of unsaturated glycoside hydrolases, including, respectively, UGL and UGH.

## Classification of carbohydrate-related enzymes

Carbohydrate-related enzymes, such as glycoside hydrolases, polysaccharide lyases, glycosyl transferases, and carbohydrate esterases, are categorized into more than 200 families based on amino acid sequences in the Carbohydrate-Active enZymes (CAZy) database [<http://afmb.cnrs-mrs.fr>] [3]. Among these enzymes, glycoside hydrolases account for the greatest number of families, numbering over 100 (families GH-1 to GH-110, as of Sept. 2007). Tertiary structures of hydrolases in 69 families have been determined and their structure/function relationships well-analyzed. Almost all glycoside hydrolases have two catalytic residues as general acid and base/nucleophile catalysts. Most of the catalytic residues consist of acidic amino acids, Asp and/or Glu. Catalytic mechanisms of glycoside hydrolases are generally divided into two types based on the anomeric configuration of the substrate in the reaction products, i.e., retention or inversion. These glycoside hydrolase mechanisms have been extensively studied and reviewed [4,45].

Polysaccharide lyases are classified into 18 families (PL-1 to PL-18, as of Sept. 2007). These enzymes commonly rec-

ognize uronic acid residues in polysaccharides, catalyze a  $\beta$ -elimination reaction, and produce unsaturated saccharides with C=C double bonds at C4 and C5 sites in nonreducing terminal uronic acid residues. The characteristics of lyases indicate that these enzymes share common structural features determining uronate recognition and the  $\beta$ -elimination reaction. Crystal structures of polysaccharide lyases belonging to 11 families have been determined, and structure/function relationships of enzymes, such as lyases for pectate, pectin, alginate, chondroitin, hyaluronan, xanthan, and rhamnogalacturonan (RG), are being analyzed [13,26,32–34,53,54]. In the catalytic reaction carried out by polysaccharide lyases, positively charged residues bind to and/or neutralize the carboxyl group in uronic acid residues. A general base catalyst extracts the proton from C5 of uronic acid residues and a general acid catalyst donates the proton to the glycosidic bond to be cleaved [46].

## Enzyme characteristics

### Unsaturated glucuronyl hydrolase (UGL).

Substrates for UGL are unsaturated saccharides with a  $\Delta$ GlcA residue at the nonreducing terminus, released from polysaccharides through polysaccharide lyase reactions. Typical substrates include mammalian extracellular matrices, glycosaminoglycans such as chondroitin, heparin, and hyaluronan, and bacterial biofilms such as xanthan and gellan. Unsaturated saccharides show an absorbance at 235 nm due to the double bond in the  $\Delta$ GlcA residue at the nonreducing terminus. UGL activity is measured by monitoring the decrease of absorbance at 235 nm, because the resulting  $\Delta$ GlcA is nonenzymatically converted to an  $\alpha$ -keto acid, followed by the loss of the double bond at C4 and C5 in this residue [11,52]. UGL activity was first detected in *F. heparinum*, which degrades chondroitin and heparin [27]. The bacterial enzyme acts on unsaturated chondroitin disaccharide produced from chondroitin through the reaction of chondroitin lyase and releases  $\Delta$ GlcA from the disaccharide [7]. Enzyme activity was also observed in *Bacillus* sp. strain GL1, in which xanthan and gellan are degraded [11]. The bacillus enzyme acts on xanthan trisaccharide, produced from xanthan through reactions of xanthan lyase,  $\beta$ -D-glucanase, and  $\beta$ -D-glucosidase, and gellan tetrasaccharide, produced from gellan through the reaction of gellan lyase [11]. The unsaturated disaccharides derived from chondroitin and hyaluronan become substrates for the bacillus enzyme.

UGLs have been purified from *F. heparinum* and *Bacillus* sp. strain GL1 and enzymatically characterized [7,11,27,37,52]. Except for N-terminal modification and amino acid

composition, the flavobacterial and bacillus enzymes are similar in molecular mass (42 kDa), optimum pH (pH 6.0–6.5) and temperature (37–45°C), the absence of a metal requirement, and substrate specificity (sulfate-free unsaturated disaccharides from chondroitin and hyaluronan, and 6S-monosulfated chondroitin disaccharide). Neither enzyme acts on the substrate *p*-nitrophenyl glucuronide, as do general glucuronidases. The N-terminus of the flavobacterial enzyme was modified uncertainly, while the N-terminal amino acid sequence of the bacillus enzyme was readily determined. Both enzymes specifically release  $\Delta$ GlcA from unsaturated saccharides but not GlcA from polysaccharide. Residual saccharide other than  $\Delta$ GlcA in the resulting reaction products is saturated and has no C=C double bond. These results demonstrate that UGL is a novel exotype glycoside hydrolase rather than a general glucuronidase.

In 1999, the gene encoding the enzyme was cloned in *Bacillus* sp. strain GL1. It consists of 1134 bp coding for a 42,859-kDa protein (377 amino acids) that has no known homology with other proteins in DNA and protein databases [11]. In 2002, the flavobacterial gene (1209 bp) for UGL was also isolated; it encodes a 45,621-kDa protein (402 amino acids) [40]. In the flavobacterial enzyme, the N-terminal 20 amino acid residues probably function as a signal peptide. Although the enzyme properties of bacillus and flavobacterial UGLs are similar, the amino acid sequence identity between the two is not high (30% identity). Recent complete genome sequence analyses indicated that a large number of microorganisms, ranging from bacteria to fungi (over 70 species), have a UGL-homologous gene in their genome. In the CAZy database, UGL and its homologues form a new family, GH-88. Microbial producers of UGL include pathogenic bacteria such as clostridia, streptococci, and vibrios. Genes homologous to that encoding the bacillus enzyme have been found mainly in streptococci: *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, and *Streptococcus suis*. The genetic features of UGL in the streptococcal genome are detailed below.

**Unsaturated galacturonyl hydrolase (UGH).** In 2006, a novel enzyme (YteR) was identified in the saprophytic bacterium *Bacillus subtilis* [20]. YteR is responsible for the metabolism of a plant cell wall polysaccharide, RG, which consists of L-rhamnose (Rha) and GalA. Although YteR was initially regarded as a function-unknown protein, after completion of genomic analysis, determination of the enzyme's tertiary structure showed that it was similar to that of UGL [16]. YteR acts on unsaturated RG disaccharides ( $\Delta$ GalA-Rha) produced from RG through the reaction of RG lyases, and releases  $\Delta$ GalA and Rha through the hydrolytic

reaction. Similar to the UGL reaction, the resulting  $\Delta$ GalA is nonenzymatically converted to  $\alpha$ -keto acid. Based on this reaction, YteR has been designated an unsaturated galacturonyl hydrolase (UGH).

Since GalA is the C4 epimer of GlcA,  $\Delta$ GalA is identical to  $\Delta$ GlcA due to the lack of the hydroxyl group at the C4 site, which in turn is a result of the reaction of polysaccharide lyases. Thus, UGL and UGH release the same molecule derived from unsaturated uronic acid residues through the catalytic reaction, although both strictly recognize the anomeric configuration in those residues.

The YteR gene consists of 1122 bp coding for a 42,969-kDa protein (373 amino acids). Another protein, YesR, which is homologous to YteR (26% sequence identity), is encoded in the *B. subtilis* genome. YesR is a 38,677-kDa UGH consisting of 344 amino acids [20]. Although both YteR and YesR catalyze the hydrolytic reaction of unsaturated RG disaccharides, the optimum pH of the enzymes differs significantly (YteR, pH 4; YesR, pH 6), as does the Michaelis constant for RG (YteR, 100  $\mu$ M; YesR, 719  $\mu$ M), and the  $k_{cat}$  (YteR, 0.28/s; YesR, 13.9/s) [20]. Both enzymes are inert on UGL substrates as well as on unsaturated disaccharides from chondroitin and hyaluronan. They both show an optimum temperature of 50°C and a thermal stability of less than 50°C. *E. chrysanthemi* RhiN is similar to YteR (26% sequence identity) and has been suggested to express UGH activity [14]. UGH-homologous genes are present in the genome of 49 types of microbes, from bacteria to fungi, including plant-related bacteria such as plant-pathogenic *Agrobacterium tumefaciens* and *Erwinia carotovora*, endophytic *Enterobacter* species, and plant symbiont *Rhizobium leguminosarum*. In the CAZy database, UGH and its homologues form a new family, GH-105.

## Biological roles of unsaturated glycoside hydrolases

Unsaturated glycoside hydrolases UGL and UGH are responsible for the metabolism of mammalian extracellular matrices, plant cell walls, and bacterial biofilms. The biological role of each enzyme in its typical bacterial producer, i.e., *Bacillus* sp. strain GL1, *F. heparinum* and *Streptococcus* for UGL, and *B. subtilis* for UGH, are the focus of this section.

***Bacillus* sp. strain GL1.** This bacterium degrades bacterial biofilms made up of xanthan and gellan as sole carbon source [8,9]. Xanthan is an exopolysaccharide produced by the plant-pathogenic bacterium *Xanthomonas campestris*. The polymer has a cellulosic backbone with linear trisaccha-

ride side chains consisting of a pyruvylated mannose (PyrMan)-GlcA-acetylated Man sequence attached at the C3 position of alternate glucosyl (Glc) residues [22]. The xanthan depolymerization system in *Bacillus* sp. strain GL1 was determined through genetic, enzymatic, and glycochemical analyses [41] (Fig. 1A). Extracellular xanthan lyases (97 and 81 kDa) first attack xanthan to release PyrMan from the side chains [9]. A  $\beta$ -D-glucanase (350 kDa) hydrolyzes glycosidic bonds in the main chain and degrades the polymer into the tetrasaccharide ( $\Delta$ GlcA-Man-(Glc)Glc) [41]. The latter product is degraded into its constituent monosaccharides through subsequent reactions of intracellular  $\beta$ -D-glucosidase (51 kDa), UGL (42 kDa), and  $\alpha$ -D-mannosidase (330 kDa) [41].

Gellan is produced by the Gram-negative bacterium *Sphingomonas paucimobilis* and consists of a linear repeating tetrasaccharide (Glc-GlcA-Glc-Rha) with *O*-acetyl and glyceryl moieties on Glc adjacent to GlcA as side chains [23]. Gellan degradation in *Bacillus* sp. strain GL1 is shown in Fig. 1B. Gellan is first depolymerized to a tetrasaccharide-repeating unit ( $\Delta$ GlcA-Glc-Rha-Glc) by extracellular gellan lyases (260 and 130 kDa) [10]. The resulting tetrasaccharide is then degraded into its constituent monosaccharides through subsequent reactions of intracellular UGL (42 kDa),  $\beta$ -D-glucosidase (51 kDa), and  $\alpha$ -L-rhamnosidase (100 kDa) [11].


UGL is inducibly produced in bacillus cells in the presence of xanthan or gellan, while the enzyme is not expressed in bacterial cells grown on glucose or pectin [11]. The UGL substrate specificity and expression profile indicate that the enzyme is essential for complete depolymerization of bacterial biofilms (xanthan and gellan) by *Bacillus* sp. strain GL1.

**Glycosaminoglycans.** *F. heparinum* and some streptococci degrade mammalian glycosaminoglycans from extracellular matrices. Glycosaminoglycans such as chondroitin, heparin, and hyaluronan are linear and negatively charged polysaccharides with a repeating disaccharide unit consisting of a uronic acid residue (GlcA or iduronic acid, IdoA) and an aminosugar residue (glucosamine or galactosamine) [6]. Chondroitin consists of GlcA and *N*-acetyl-D-galactosamine (GalNAc), with a sulfate group at position 4 or 6 or both [15]. Chondroitin sulfates A and C have GalNAc with a sulfate group at positions 4 and 6. Heparin and heparan sulfate are composed of D-glucosamine (GlcN) and GlcA or IdoA [44]. Hyaluronan is composed of GlcA and *N*-acetyl-D-glucosamine (GlcNAc) [5]. Except for hyaluronan, mammalian glycosaminoglycans covalently bind to proteins; the resulting proteoglycans play an important role in cellular architecture and permeability [15]. Hyaluronan is crucial for cell-to-cell association in mammals and for the capsule in streptococcal bacteria [25]. These glycosaminoglycans in extracellular matrices

become a target for pathogens to invade host cells and serve as sites of specific interaction for many pathogens [47].

***Flavobacterium heparinum.*** Gram-negative *F. heparinum*, isolated from a soil sample, degrades chondroitin, heparin, and hyaluronan by the actions of several enzymes [27,30,39,49,51]. In the presence of chondroitin and hyaluronan, the bacterium inducibly produces chondroitin AC lyase (77 kDa), chondroitin B lyase (54 kDa), sulphatase (50 kDa), and UGL (42 kDa) in the periplasm [7]. Chondroitin AC lyase acts on chondroitin, chondroitin sulfate A, chondroitin sulfate C, and hyaluronan, and releases unsaturated disaccharides with  $\Delta$ GlcA at the nonreducing terminus. Dermatan sulfate is depolymerized into unsaturated disaccharide through the reaction of chondroitin B lyase. The resulting unsaturated disaccharides derived from chondroitin, chondroitin sulfate C, and hyaluronan are subsequently degraded into constituent monosaccharides through the reaction of UGL, while the enzyme is inert on unsaturated disaccharide 4-sulfate from chondroitin sulfate A and unsaturated disaccharide from dermatan sulfate [7]. In heparin degradation, *F. heparinum* also produces three heparin lyases: I (43 kDa), -II (85 kDa), and -III (73 kDa) [30]. Unsaturated disaccharides resulting from the reactions of lyases are further degraded by UGL, making it a key enzyme in glycosaminoglycan metabolism in *F. heparinum*.

***Streptococcus* spp.** Infections with certain streptococcal species, such as *S. agalactiae*, *S. pneumoniae*, and *S. pyogenes*, cause severe diseases, e.g., pneumonia, bacteremia, sinusitis, and meningitis [29,36]. Septicemia, meningitis, and pneumonia are induced in pigs through infection by *S. suis* [31]. All of these bacterial species produce polysaccharide lyases, which function virulence factors that degrade glycosaminoglycans [24,28]. As described above, a UGL-homologous gene is present in the genome of these streptococci. Interestingly, this gene is located in a gene cluster for the putative phosphotransferase system (PTS), which imports aminosugar, a component of glycosaminoglycans. The cluster also includes the polysaccharide lyase gene (Fig. 1C), strongly suggesting that UGL functions as another streptococcal virulence factor, one that is responsible for the complete degradation of glycosaminoglycans. Unlike the bacillus and flavobacterial enzymes, streptococcal UGL acts on unsaturated chondroitin disaccharide 4-sulfate [unpublished results], i.e., the streptococcal enzyme is capable of degrading highly sulfated mammalian glycosaminoglycans. This property enables the bacteria to readily invade mammalian cells through the degradation of extracellular sulfated glycosaminoglycans.




**Fig. 1.** Biological functions of unsaturated glycoside hydrolases. (A) Xanthan metabolism in *Bacillus* sp. strain GL1. G, Glc; GA, GlcA; M, Man;  $\Delta$ GA,  $\Delta$ GlcA; Pyr, pyruvate group; Ac, acetyl group. (B) Gellan metabolism in *Bacillus* sp. strain GL1. G, Glc; GA, GlcA; R, Rha;  $\Delta$ GA,  $\Delta$ GlcA. (C) Streptococcal genetic cluster. Genes, *PTS-IIA*, *PTS-IIB*, *PTS-IIC*, and *PTS-IID*, encode the phosphotransferase system for aminosugar import. *ugl* and *hepC* code for unsaturated glucuronyl hydrolase and polysaccharide lyase. (D) RG-I metabolism in *B. subtilis*. GA, GalA; R, Rha;  $\Delta$ GA,  $\Delta$ GalA. Thick, thin, and dotted arrows indicate the metabolic pathway for polysaccharides, cleavage sites of polysaccharide-depolymerizing enzymes, and biosynthetic pathways for the enzymes, respectively.

**Bacillus subtilis.** This bacterium is a typical representative of saprophytic microbes [38]. For such saprophytic and/or plant pathogenic bacteria, plant cell wall degradation is a major event in plant cell invasion. The plant cell wall consists of pectin, cellulose, and hemicellulose [2]. Pectin is further divided into three components, a linear main chain (polygalacturonan) and two branched chains, RG-I and -II [50]. RG-I consists of alternating Rha and GalA as a backbone with additions of arabinans and galactans as side chains [35]. RG-II contains polygalacturonan as a backbone and a complex of approximately 30 monosaccharides as a side chain [43]. In the presence of RG-I, *B. subtilis* induces the expression of three gene clusters, *yes*, *yte*, and *ybc-ybd* [42]. Gene clusters *yes* and *yte* include genes for YesR and YteR. The degradation pathway of RG-I in *B. subtilis* is shown in Fig. 1D. An extracellular RG lyase, YesW (68 kDa), catalyzes the initial cleavage of the glycosidic bond in the RG-I main chain, with the resulting oligosaccharides converted to the unsaturated RG disaccharide through the reaction of extracellular exotype YesX (68 kDa). The unsaturated disaccharide is finally degraded into constituent monosaccharides through the reaction of intracellular YteR (43 kDa) or YesR (39 kDa). Generally, the bacterium cannot assimilate polysaccharides until monosaccharides have been released, indicating that the UGHs YteR and YesR are essential to the metabolism of plant cell wall RG-I in *B. subtilis*.

**Structure/function relationships.** Unsaturated glycoside hydrolases UGL and UGH are unique in that they act on unsaturated saccharides but not on saturated saccharides, and release unsaturated uronic acid. To clarify the structure/function relationships of these enzymes, the three-dimensional structures of *Bacillus* sp. strain GL1 and *S. agalactiae* UGLs and *B. subtilis* UGH were determined by X-ray crystallography [16,20].

**Structures of UGL and UGH.** The overall structure of bacillus UGL is shown in Fig. 2A. The enzyme has approximate dimensions of  $45 \times 45 \times 40$  Å and shows a single globular domain consisting of 12  $\alpha$ -helices (H1-H12), three antiparallel  $\beta$ -sheets composed of two or three strands (SA, SB, and SC), two short  $3_{10}$ -helices, and loops connecting each element. The 12  $\alpha$ -helices constitute an  $\alpha_6/\alpha_6$ -barrel structure with a deep cleft, considered to be the active site. The



**Fig. 2.** Overall structure of unsaturated glycoside hydrolases. **(A)** Ligand-free UGL from *Bacillus* sp. strain GL1. **(B)** Substrate ( $\Delta$ GlcA-GalNAc)-bound UGL (D88N) from *Bacillus* sp. strain GL1. **(C)** Substrate ( $\Delta$ GalA-Rha)-bound YteR (D143N) from *B. subtilis*. Colors denote elements with a secondary structure (pink,  $\alpha$ -helices; cyan,  $\beta$ -strands; and yellow, turns and coils). A green or blue ball-and-stick model indicates the substrate (red, oxygen atom; yellow, carbon atom; and blue, nitrogen atom). **(D)** Molecular surface of the substrate-bound UGL (D88N) from *Bacillus* sp. strain GL1. Colors denote amino acid residues (red, Asp or Glu; blue, Lys or Arg; cyan, His; and magenta, Trp, Tyr, or Phe). A green ball-and-stick model indicates the substrate (red, oxygen atom; yellow, carbon atom; and blue, nitrogen atom).

$\alpha_6/\alpha_6$ -barrel structure is made up of six outer  $\alpha$ -helices (H1, H3, H5, H7, H9, and H11) oriented in roughly the same direction, and six inner  $\alpha$ -helices (H2, H4, H6, H8, H10, and H12) running in the opposite direction. These 12 helices are connected by short and long loops in a nearest-neighbor, up-down pattern. Short loops extend along one side of the barrel. On the other side, long loops constitute the wall of a deep pocket-like cleft, which has a diameter of 15–20 Å at the lip and is 15 Å deep.


In structural comparisons to bacillus UGL, three proteins, i.e., the hypothetical protein YteR from *B. subtilis* [55] (Fig. 2C), glucoamylase from *Thermoanaerobacterium thermosaccharolyticum* [1], and *N*-acyl-D-glucosamine 2-epimerase (AGE) [19] from porcine kidney, showed the highest similarity. These proteins also adopt an  $\alpha_6/\alpha_6$ -barrel structure as a basic scaffold, although they show no sequence similarity at the amino acid level. The crystal structure of YteR was determined as a function-unknown protein by the Midwest Center for Structural Genomics [55]. The high structural similarity between UGL and YteR enabled us to clarify that YteR exhibits UGL-like activity. The detailed intrinsic function of YteR was described in the previous section.

**Catalytic mechanism.** The structure of an enzyme that is complexed with substrates contributes to the identification of its active site and clarification of its mechanisms for catalytic reaction and substrate recognition. Thus, the crystal structure of bacillus UGL complexed with a substrate, unsaturated chondroitin disaccharide ( $\Delta$ GlcA-GalNAc), was determined at 1.9 Å resolution [17] (Fig. 2B). No significant conformational change occurs in the overall structure between substrate-free and bound enzymes. To study the complexed structure, enzyme D88N, which has a mutation at position 88 (Asp→Asn), is used because the wild-type enzyme readily degrades the substrate and thus cannot be isolated in the substrate-bound form. The disaccharide is bound to the deep pocket at the center of the UGL molecule, which indicates that the active site is in the pocket. A molecular surface model of the complex structure shows that the active site is surrounded by aromatic and positively charged residues (Fig. 2D).

$\Delta$ GlcA is bound to residues Trp42, Asn(Asp)88, Phe91, Trp134, Asp149, Gln211, Trp219, Arg221, Trp225, and Tyr338, while GalNAc interacts with residues Trp42, His86, Trp134, His210, Tyr338, and His339 (Fig. 3A). Almost all residues responsible for binding to the disaccharide are well

conserved in family GH-88. The complex structure is unique in that no atom forms hydrogen bonds to the glycosidic oxygen (O3 atom of GalNAc). The arrangement of amino acid residues and water molecules at the active site is detailed as follows (Fig. 3A): a water molecule (Wat) is situated at an “ $\alpha$ -face position” of  $\Delta$ GlcA and binds through hydrogen bonds to Asp149 (3.1 Å), His193 (2.7 Å), Gln211 (2.6 Å), and O5 atom (3.2 Å) of  $\Delta$ GlcA, but not to glycosidic oxygen (4.2 Å); Wat is close to C5 carbon (3.4 Å) of  $\Delta$ GlcA, but not to the C1 carbon atom (3.9 Å); Asp149 is close to the C4 atom (2.9 Å) of  $\Delta$ GlcA, but not to glycosidic oxygen (5.8 Å); Asn88 at the mutation site (wild-type, Asp88) is not close to the C1 carbon (4.3 Å) of  $\Delta$ GlcA and glycosidic oxygen (4.9 Å); no water molecule is present at the “ $\beta$ -face position” of  $\Delta$ GlcA. These characteristics are common to crystal structures of UGL complexed with unsaturated hyaluronan disaccharide ( $\Delta$ GlcA-GlcNAc) and unsaturated gellan tetrasaccharide ( $\Delta$ GlcA-Glc-Rha-Glc) [18] (Fig. 3B, C). This arrangement of amino acid residues and water molecules is unusual in glycoside hydrolases [4,45].

The novel catalytic mechanism of UGL, deduced from its structural characteristics, consists of the following two reaction steps: (i) the water addition reaction (hydration) of the vinyl ether group (C4=C5–O5) in  $\Delta$ GlcA and (ii) subsequent hydrolysis of the glycosidic bond between  $\Delta$ GlcA and GalNAc. In the hydration reaction, Asp149 donates a proton to the double bond (C4 atom) of  $\Delta$ GlcA as a general acid catalyst and then deprotonates Wat as a general base catalyst. The resulting deprotonated water molecule attacks the C5 atom of  $\Delta$ GlcA. The product, a hemiketal, is unstable and readily converted to  $\alpha$ -keto acid (hemiacetal) in an enzyme-independent manner. This hydration is partially caused by the electron-rich double bond in C4=C5, which acts as an “electron sink.” In the hydrolysis reaction, due to aldehyde-hemiacetal equilibrium, the hemiacetal is



**Fig. 3.** Interaction between unsaturated glycoside hydrolases and substrates (stereodiagram). (A) *Bacillus* sp. strain GL1 UGL (D88N) complexed with unsaturated chondroitin disaccharide ( $\Delta$ GlcA-GalNAc). (B) *Bacillus* sp. strain GL1 UGL (D88N) complexed with unsaturated hyaluronan disaccharide ( $\Delta$ GlcA-GlcNAc). (C) *Bacillus* sp. strain GL1 UGL (D88N) complexed with unsaturated gellan tetrasaccharide ( $\Delta$ GlcA-Glc-Rha-Glc). (D) *B. subtilis* YteR (D143N) complexed with unsaturated RG disaccharide ( $\Delta$ GalA-Rha). The bound substrate as well as the surrounding amino acid residues and water molecules (Wat, gray ball) interacting with the substrate or amino acid residues are shown. The substrate is represented by a yellow ball-and-stick model (red, oxygen atom; and cyan, nitrogen atom). Direct hydrogen bonds ( $\leq 3.2$  Å), shown as dotted lines, are formed between the substrate and amino acid residues. (E) Superpositioning of active site structures of the bacillus UGL and YteR (magenta, UGL and blue, YteR).




Fig. 4. Substrate recognition of unsaturated glucuronyl hydrolases. Left: Superpositioning of active site structures of UGLs from *Bacillus* sp. strain GL1 and *S. agalactiae* (brown, bacillus UGL; and green, streptococcal UGL). Right: Sequence alignment (brown, *Bacillus* sp. strain GL1 UGL; green, *S. agalactiae* UGL; purple, *S. pneumoniae* UGL; and blue, *S. pyogenes* UGL).

finally converted to an aldehyde (4-deoxy-L-threo-5-hexosulose-uronate), leaving behind a saccharide through cleavage of the glycosidic bond. Asp88 probably stabilizes the carbenium ion at C5 of  $\Delta$ GlcA during the catalytic reaction. GC/MS analysis of the reaction product in the presence of  $H_2^{18}O$  directly showed hydration of the vinyl ether group in  $\Delta$ GlcA but not of the glycosidic bond [17]. The reaction product has  $^{18}O$  on the C5 atom but not on the C1 atom. Site-directed mutagenesis and kinetic studies support this catalytic mechanism and indicate that the proposed role of each amino acid residue in the active site is plausible [16–18].

The X-ray crystallographic and GC/MS studies provide evidence of a novel catalytic mechanism of UGL, i.e., the enzyme triggers hydration of the vinyl ether group in  $\Delta$ GlcA but not of the glycosidic bond to be cleaved. Thus, UGL is clearly unlike general glycoside hydrolases, in which the catalytic mechanism consists of retention or inversion of an anomeric configuration.

The tertiary structure of the YteR mutant (D143N, Asp143 to Asn) complexed with the substrate ( $\Delta$ GalA-Rha) was also determined by X-ray crystallography [21] (Fig. 2C). Similar to UGL, the enzyme accommodates the substrate in the deep pocket at the center of the molecule.  $\Delta$ GalA interacts with residues His40, Tyr41, Asp88, His132, Trp141, Asn(Asp)143, Met147, Arg213, Trp217, Gly332, Thr333, and His352, while Rha is bound to residues His40, Tyr41, His42, and Gly332 (Fig. 3D). Ionizable residues His132, Asn(Asp)143, His189, and Arg213 interact with the substrate and are conserved in YteR and its homologs. The arrange-

ment of amino acid residues at the active site is similar for YteR and UGL. Asp143 in YteR corresponds to Asp149 in UGL, likewise for His189 to His193 and Arg213 to Arg221. The interaction between YteR and  $\Delta$ GalA-Rha is as follows (Fig. 3D): Wat hydrogen bonds to Asn(Asp)143 (2.9 Å) but not to glycosidic oxygen; Wat is also close to the C5 carbon atom of  $\Delta$ GalA (3.5 Å); Asn(Asp)143 is close to the C4 atom of  $\Delta$ GalA (3.4 Å). These residues of YteR can be superimposed on those of UGL (Fig. 3E), indicating that UGH also catalyzes the first hydration reaction to the vinyl ether group in  $\Delta$ GalA. The catalytic mechanism of YteR is as follows: Asp143 donates a proton to the double bond (C4 atom) of  $\Delta$ GalA as a general acid catalyst. Asp143 then deprotonates Wat as a general base catalyst and the deprotonated Wat molecule attacks the C5 atom of  $\Delta$ GalA. The product (hemiketal) is unstable and readily converted to  $\alpha$ -keto acid (hemiacetal). Due to aldehyde-hemiacetal equilibrium, the resulting hemiacetal is finally converted to aldehyde (4-deoxy-L-threo-5-hexosulose-uronate) and saturated saccharide (Rha) through cleavage of the glycosidic bond.

**Substrate recognition.** In the complex structure of the UGL mutant (D88N) and  $\Delta$ GlcA-GalNAc (Fig. 3A), there are fewer interactions between D88N and GalNAc than between D88N and  $\Delta$ GlcA, indicating that the enzyme interacts strongly with  $\Delta$ GlcA rather than with GalNAc through hydrogen bonds and van der Waals contacts. UGL acts on different unsaturated saccharides derived from chondroitin, hyaluronan, xanthan, and gellan, but these substrates without




exception have  $\Delta$ GlcA residues at their nonreducing termini. The structural feature of UGL binding to its substrates was also examined by determination of the crystal structures of UGL complexed with unsaturated hyaluronan disaccharide and unsaturated gellan tetrasaccharide [18] (Fig. 3B,C). Analysis of the structures of UGL complexed with different substrates showed that  $\Delta$ GlcA is specifically recognized through the formation of several hydrogen bonds and stacking interactions by amino acid residues highly conserved in the GH-88 family of enzymes (Fig. 3A–C). Aromatic residues in particular are responsible for the stacking interactions (Fig. 3). The carboxyl group of  $\Delta$ GlcA interacts with the enzyme through the formation of hydrogen bonds, and interaction is stabilized by the positive end of the inner  $\alpha$ -helix dipole of the  $\alpha_6/\alpha_6$ -barrel (Fig. 2B,C). This structural feature of UGL, crucial to strict recognition of the unsaturated uronic acid residue, is also found in YteR.

Unlike the bacillus and flavobacterial UGLs, streptococcal UGL acts on unsaturated chondroitin disaccharide containing a sulfate group at C4 of GalNAc. The crystal structure of *S. agalactiae* UGL was determined at 1.8 Å resolution by X-ray crystallography to clarify the difference in substrate specificity among UGLs [unpublished results]. The in silico model, based on this crystal structure and the coordinate of unsaturated chondroitin disaccharide 4-sulfate, indicated that Arg236 of streptococcal UGL specifically recognizes the sulfate group (Fig. 4). The bacillus and flavobacterial UGLs have a histidine residue in place of Arg236. The arginine residue is well-conserved in UGLs from *S. agalactiae*, *S. pneumoniae*, and *S. pyogenes* (Fig. 4). The binding ability of Arg236 to the sulfate group probably contributes to the broad substrate specificity of streptococcal UGLs.

## Concluding remarks

Most genes coding for unsaturated glycoside hydrolases UGL and UGH are encoded in the genome of microorganisms producing polysaccharide lyases, since hydrolases act on unsaturated saccharides released from polysaccharides through the reaction of polysaccharide lyases. Many microbial polysaccharide lyases attack extracellular matrices such as glycosaminoglycan or plant cell wall pectin. Mammalian pathogenic and/or symbiont bacteria, including bacteroides, clostridia, enterococci, lactobacilli, streptococci, and vibrios, predominate among UGL producers in the enzyme family GH-88. In enzyme family GH-105, plant pathogenic, endophytic, and/or plant symbiont bacteria, e.g., agrobacteria, *Erwinia*, enterobacteria, and rhizobia, are typical producers



**Fig. 5.** The  $\alpha/\alpha$ -barrel structure shared by carbohydrate-related enzymes. Magenta, UGL; cyan, YteR; yellow, AGE; and green, A1-III.

of UGH. Thus, UGL and UGH, together with polysaccharide lyases, play important roles in allowing bacteria to associate with mammals or plants and function as virulence factors for host cells. Inhibitors of polysaccharide lyases and unsaturated glycoside hydrolases may become potent pharmaceuticals for treating microbial infectious diseases, because most of these enzymes are produced by microorganisms but not by plants or mammals. Clarification of the structure/function relationships of polysaccharide lyases and unsaturated glycoside hydrolases is expected to facilitate the design of novel drugs targeting such diseases. Since the catalytic mechanism of UGL and UGH in particular differs substantially from that of general glycoside hydrolases, inhibitors of these enzymes should be effective without exerting side effects on mammalian and plant host cells.

The  $\alpha/\alpha$ -barrel structure is categorized as a  $\alpha/\alpha$  toroid fold in the SCOP database [<http://scop.mrc-lmb.cam.ac.uk/scop/>]. The  $\alpha/\alpha$  toroid-fold proteins are further divided into five superfamilies, among which the unsaturated glycoside hydrolases UGL and UGH belong to a superfamily of six-hairpin glycoside hydrolases. This superfamily includes carbohydrate-related enzymes such as glucoamylases (Protein Data Bank (PDB) entry, 1GAI), cellulases (PDB entry, 1KWF), maltose phosphorylase (PDB entry, 1H54), AGE (PDB entry, 1FP3), and alginate lyase A1-III (PDB entry, 1QAZ). We found that the basic scaffold, i.e., the  $\alpha/\alpha$ -barrel, of UGL, YteR, AGE, and A1-III, are well-superimposed (Fig. 5).

Structural similarity provides valuable information with which to clarify the intrinsic function of hypothetical proteins. Analogous to the identification of YteR as a UGH, *Salmonella typhimurium* hypothetical protein YihS, containing an  $\alpha/\alpha$ -barrel structure, has recently been identified as a novel carbohydrate-related enzyme that catalyzes isomerization and epimerization [manuscript in preparation]. Regarding the structural similarity observed in  $\alpha/\alpha$ -barrel proteins, we hypothesize that these carbohydrate-related enzymes developed from a common ancestral protein, although they catalyze different reactions, i.e., hydrolase, phosphorylase, epimerase, or lyase, and no similarities among their primary structures have been found. Further study of the molecular evolution of enzymes is expected to reveal common structural features.

**Acknowledgements.** This work was supported in part by Grants-in-Aid (KM, BM, and WH) and the Targeted Proteins Research Program from the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

## References

1. Aleshin AE, Feng PH, Honzatko RB, Reilly PJ (2003) Crystal structure and evolution of a prokaryotic glucoamylase. *J Mol Biol* 327:61-73
2. Carpita NC, Gibeaut DM (1993) Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *Plant J* 3:1-30
3. Davies GJ, Gloster TM, Henrissat B (2005) Recent structural insights into the expanding world of carbohydrate-active enzymes. *Curr Opin Struct Biol* 15:637-645
4. Davies G, Henrissat B (1995) Structures and mechanisms of glycosyl hydrolases. *Structure* 3:853-859
5. DeAngelis PL (1999) Hyaluronan synthases: fascinating glycosyltransferases from vertebrates, bacterial pathogens, and algal viruses. *Cell Mol Life Sci* 56:670-682
6. Ernst S, Langer R, Cooney CL, Sasisekharan R (1995) Enzymatic degradation of glycosaminoglycans. *Crit Rev Biochem Mol Biol* 30:387-444
7. Gu K, Linhardt RJ, Laliberté M, Gu K, Zimmermann J (1995) Purification, characterization and specificity of chondroitin lyases and glycuronidase from *Flavobacterium heparinum*. *Biochem J* 312:569-577
8. Hashimoto W, Maesaka K, Sato N, Kimura S, Yamamoto K, Kumagai H, Murata K (1997) Microbial system for polysaccharide depolymerization: enzymatic route for gellan depolymerization by *Bacillus* sp. GL1. *Arch Biochem Biophys* 339:17-23
9. Hashimoto W, Miki H, Tsuchiya N, Nankai H, Murata K (1998) Xanthan lyase of *Bacillus* sp. strain GL1 liberates pyruvylated mannose from xanthan side chains. *Appl Environ Microbiol* 64:3765-3768
10. Hashimoto W, Sato N, Kimura S, Murata K (1998) Polysaccharide lyase: molecular cloning of gellan lyase gene and formation of the lyase from a huge precursor protein in *Bacillus* sp. GL1. *Arch Biochem Biophys* 354:31-39
11. Hashimoto W, Kobayashi E, Nankai H, Sato N, Miya T, Kawai S, Murata K (1999) Unsaturated glucuronyl hydrolase of *Bacillus* sp. GL1: novel enzyme prerequisite for metabolism of unsaturated oligosaccharides produced by polysaccharide lyases. *Arch Biochem Biophys* 368:367-374
12. Hashimoto W, Miyake O, Momma K, Kawai S, Murata K (2000) Molecular identification of oligoalginate lyase of *Sphingomonas* sp. strain A1 as one of the enzymes required for complete depolymerization of alginate. *J Bacteriol* 182:4572-4577
13. Huang W, Matte A, Li Y, Kim YS, Linhardt RJ, Su H, Cygler M (1999) Crystal structure of chondroitinase B from *Flavobacterium heparinum* and its complex with a disaccharide product at 1.7 Å resolution. *J Mol Biol* 294:1257-1269
14. Hugouvieux-Cotte-Pattat N (2004) The RhaS activator controls the *Erwinia chrysanthemi* 3937 genes *rhiN*, *rhiT* and *rhiE* involved in rhamnogalacturonan catabolism. *Mol Microbiol* 51:1361-1374
15. Iozzo RV (1998) Matrix proteoglycans: from molecular design to cellular function. *Annu Rev Biochem* 67:609-652
16. Itoh T, Akao S, Hashimoto W, Mikami B, Murata K (2004) Crystal structure of unsaturated glucuronyl hydrolase, responsible for the degradation of glycosaminoglycan, from *Bacillus* sp. GL1 at 1.8 Å resolution. *J Biol Chem* 279:31804-31812
17. Itoh T, Hashimoto W, Mikami B, Murata K (2006) Crystal structure of unsaturated glucuronyl hydrolase complexed with substrate: molecular insights into its catalytic reaction mechanism. *J Biol Chem* 281:29807-29816
18. Itoh T, Hashimoto W, Mikami B, Murata K (2006) Substrate recognition by unsaturated glucuronyl hydrolase from *Bacillus* sp. GL1. *Biochem Biophys Res Commun* 344:253-262
19. Itoh T, Mikami B, Maru I, Ohta Y, Hashimoto W, Murata K (2000) Crystal structure of *N*-acyl-D-glucosamine 2-epimerase from porcine kidney at 2.0 Å resolution. *J Mol Biol* 303:733-744
20. Itoh T, Ochiai A, Mikami B, Hashimoto W, Murata K (2006) A novel glycoside hydrolase family 105: the structure of family 105 unsaturated rhamnogalacturonyl hydrolase complexed with a disaccharide in comparison with family 88 enzyme complexed with the disaccharide. *J Mol Biol* 360:573-585
21. Itoh T, Ochiai A, Mikami B, Hashimoto W, Murata K (2006) Structure of unsaturated rhamnogalacturonyl hydrolase complexed with substrate. *Biochem Biophys Res Commun* 347:1021-1029
22. Jansson PE, Kenne L, Lindberg B (1975) Structure of the extracellular polysaccharide from *Xanthomonas campestris*. *Carbohydr Res* 45:275-282
23. Jansson PE, Lindberg B, Sandford PA (1983) Structural studies of gellan gum, an extracellular polysaccharide elaborated by *Pseudomonas elodea*. *Carbohydr Res* 124:135-139
24. Jedrzejewski MJ (2001) Pneumococcal virulence factors: structure and function. *Microbiol Mol Biol Rev* 65:187-207
25. Laurent TC, Fraser JR (1992) Hyaluronan. *FASEB J* 6:2397-2404
26. Li S, Jedrzejewski MJ (2001) Hyaluronan binding and degradation by *Streptococcus agalactiae* hyaluronate lyase. *J Biol Chem* 276:41407-41416
27. Linker A, Hoffman P, Meyer K, Sampson P, Korn ED (1960) The formation of unsaturated disaccharides from mucopolysaccharides and their cleavage to  $\alpha$ -keto acid by bacterial enzymes. *J Biol Chem* 235:3061-3065
28. Liu GY, Nizet V (2004) Extracellular virulence factors of group B Streptococci. *Front Biosci* 9:1794-1802
29. Llull D, Lopez R, Garcia E (2001) Genetic bases and medical relevance of capsular polysaccharide biosynthesis in pathogenic streptococci. *Curr Mol Med* 1:475-491
30. Lohse DL, Linhardt RJ (1992) Purification and characterization of heparin lyases from *Flavobacterium heparinum*. *J Biol Chem* 267:24347-24355
31. Lun ZR, Wang QP, Chen XG, Li AX, Zhu XQ (2007) *Streptococcus suis*: an emerging zoonotic pathogen. *Lancet Infect Dis* 7:201-209
32. Maruyama Y, Hashimoto W, Mikami B, Murata K (2005) Crystal structure of *Bacillus* sp. GL1 xanthan lyase complexed with a substrate: insights into the enzyme reaction mechanism. *J Mol Biol* 350:974-986

33. Mayans O, Scott M, Connerton I, Gravesen T, Benen J, Visser J, Pickersgill R, Jenkins J (1997) Two crystal structures of pectin lyase A from *Aspergillus* reveal a pH driven conformational change and striking divergence in the substrate-binding clefts of pectin and pectate lyases. *Structure* 5:677-689
34. McDonough MA, Kadirvelraj R, Harris P, Poulsen JC, Larsen S (2004) Rhamnogalacturonan lyase reveals a unique three-domain modular structure for polysaccharide lyase family 4. *FEBS Lett* 565:188-194
35. McNeil M, Darvill AG, Albersheim P (1980) Structure of plant cell walls: X. Rhamnogalacturonan I, a structurally complex pectic polysaccharide in the walls of suspension-cultured sycamore cells. *Plant Physiol* 66:1128-1134
36. Mitchell TJ (2003) The pathogenesis of streptococcal infections: from tooth decay to meningitis. *Nat Rev Microbiol* 1:219-230
37. Mori S, Akao S, Nankai H, Hashimoto W, Mikami B, Murata K (2003) A novel member of glycoside hydrolase family 88: overexpression, purification, and characterization of unsaturated  $\beta$ -glucuronidase of *Bacillus* sp. GL1. *Protein Expr Purif* 29:77-84
38. Morikawa M (2006) Beneficial biofilm formation by industrial bacteria *Bacillus subtilis* and related species. *J Biosci Bioeng* 101:1-8
39. Myette JR, Shriver Z, Claycamp C, McLean MW, Venkataraman G, Sasisekharan R (2003) The heparin/heparan sulfate 2-O-sulfatase from *Flavobacterium heparinum*. Molecular cloning, recombinant expression, and biochemical characterization. *J Biol Chem* 278:12157-12166
40. Myette JR, Shriver Z, Kiziltepe T, McLean MW, Venkataraman G, Sasisekharan R (2002) Molecular cloning of the heparin/heparan sulfate  $\Delta$ 4,5 unsaturated glycuronidase from *Flavobacterium heparinum*, its recombinant expression in *Escherichia coli*, and biochemical determination of its unique substrate specificity. *Biochemistry* 41:7424-7434
41. Nankai H, Hashimoto W, Miki H, Kawai S, Murata K (1999) Microbial system for polysaccharide depolymerization: enzymatic route for xanthan depolymerization by *Bacillus* sp. strain GL1. *Appl Environ Microbiol* 65:2520-2526
42. Ochiai A, Itoh T, Kawamata A, Hashimoto W, Murata K (2007) Plant cell wall degradation by saprophytic *Bacillus subtilis* strains: gene clusters responsible for rhamnogalacturonan depolymerization. *Appl Environ Microbiol* 73:3803-3813
43. O'Neill MA, Warrenfeltz D, Kates K, Pellerin P, Doco T, Darvill AG, Albersheim P (1996) Rhamnogalacturonan-II, a pectic polysaccharide in the walls of growing plant cell, forms a dimer that is covalently cross-linked by a borate ester. *In vitro* conditions for the formation and hydrolysis of the dimer. *J Biol Chem* 271:22923-22930
44. Powell AK, Yates EA, Fernig DG, Turnbull JE (2004) Interactions of heparin/heparan sulfate with proteins: appraisal of structural factors and experimental approaches. *Glycobiology* 14:17R-30R
45. Rye CS, Withers SG (2000) Glycosidase mechanisms. *Curr Opin Chem Biol* 4:573-580
46. Rye CS, Withers SG (2002) Elucidation of the mechanism of polysaccharide cleavage by chondroitin AC lyase from *Flavobacterium heparinum*. *J Am Chem Soc* 124:9756-9767
47. Sawitzky D (1996) Protein-glycosaminoglycan interactions: infectious aspects. *Med Microbiol Immunol* 184:155-161
48. Shevchik VE, Condemine G, Robert-Baudouy J, Hugouvieux-Cotte-Pattat N (1999) The exopolysaccharide lyase PelW and the oligogalacturonate lyase Ogl, two cytoplasmic enzymes of pectin catabolism in *Erwinia chrysanthemi* 3937. *J Bacteriol* 181:3912-3919
49. Su H, Blain F, Musil RA, Zimmermann JJ, Gu K, Bennett DC (1996) Isolation and expression in *Escherichia coli* of *hepB* and *hepC*, genes coding for the glycosaminoglycan-degrading enzymes heparinase II and heparinase III, respectively, from *Flavobacterium heparinum*. *Appl Environ Microbiol* 62:2723-2734
50. Thakur BR, Singh RK, Handa AK (1997) Chemistry and uses of pectin—a review. *Crit Rev Food Sci Nutr* 37:47-73
51. Tkalec AL, Fink D, Blain F, Zhang-Sun G, Laliberte M, Bennett DC, Gu K, Zimmermann JJ, Su H (2000) Isolation and expression in *Escherichia coli* of *csIA* and *csIB*, genes coding for the chondroitin sulfate-degrading enzymes chondroitinase AC and chondroitinase B, respectively, from *Flavobacterium heparinum*. *Appl Environ Microbiol* 66:29-35
52. Warnick CT, Linker A (1972) Purification of an unusual  $\alpha$ -glycuronidase from flavobacteria. *Biochemistry* 11:568-572
53. Yoder MD, Lietzke SE, Jurnak F (1993) Unusual structural features in the parallel  $\beta$ -helix in pectate lyases. *Structure* 1:241-251
54. Yoon H-J, Mikami B, Hashimoto W, Murata K (1999) Crystal structure of alginate lyase A1-III from *Sphingomonas* species A1 at 1.78 Å resolution. *J Mol Biol* 290:505-514
55. Zhang R, Minh T, Lezondra L, Korolev S, Moy SF, Collart F, Joachimiak A (2005) 1.6 Å crystal structure of YteR protein from *Bacillus subtilis*, a predicted lyase. *Proteins* 60:561-565